Limits on charge non-conservation from possible seasonal variations of the solar neutrino experiments

Manuel Torres^{*}and Hector Vucetich [†] Instituto de Física, Universidad Nacional Autónoma de México Aptdo. Postal 20-364, México D.F. 01000, México.

November 27, 2017

Abstract

Variable speed of light (VSL) theories generically lead to large violations of charge conservation that can be written in terms of a dimensionless parameter λ . It is shown that the motion of the Earth with respect to the Sun could lead to a seasonal variation for the SAGE and GALLEX-GNO experiments and analyzing the reported counting rates for these experiments, a very stringent bound $\lambda \leq 2 \times 10^{-19}$ is obtained, some 10⁹ times smaller than previous ones. Furthermore, a bound on the lifetime of the $^{71}\text{Ga} \rightarrow ^{71}\text{Ge}$ charge-nonconserving decay in VSL theories is found as: $\tau_{\text{CNC}} \geq 1.4 \times 10^{27}$ years. Similarly a new upper limit for the ratio of the charge-nonconserving to the normal weak decay of the neutron in VSL theories is obtained: $\Gamma(n \rightarrow p + \nu_e + \bar{\nu}_e)/\Gamma(n \rightarrow p + e + \bar{\nu}_e) \leq 2 \times 10^{-27}$.

PACS: 11.30.-j,04.50.+h,26.65.+t

Keywords: Charge conservation; VSL Theories; solar neutrino experiments.

Recently, a growing interest in the time variation of fundamental constants has arisen, driven by the tantalizing observation that $\alpha = e^2/\hbar c$ may have been slightly smaller when the Universe was younger $z > 1, t_0 - t > 8 \times 10^9$ yr [1, 2]. This may be attributed either to a time variation of the electron charge e or of the speed of light c [3, 4, 5].

The second possibility has received much attention, since a varying speed of light may solve the homogeneity, horizon and flatness problems in cosmology [6, 7, 8]. Several variations of these theories have been proposed with varying degree of sophistication [9, 10, 11, 12, 13].

^{*}e-mail: torres@fisica.unam.mx

[†]Corresponding author. e-mail: vucetich@fisica.unam.mx; Fax: (52) 56225015. On leave of absence from: Observatorio Astronómico, Universidad Nacional de La Plata.

It was observed in Ref. [14] that a generic variable speed of light (VSL) theory leads necessarily to a large violation of charge conservation. This is because Maxwell's equations are modified in the form:

$$\frac{1}{c}\partial_{\mu}\left(cF^{\mu\nu}\right) = 4\pi j^{\nu},\tag{1}$$

from which the following relation of charge nonconservation to the variation of c can be derived [15]

$$\frac{\dot{Q}}{Q} = -\frac{\dot{c}}{c},\qquad(2)$$

This equation provides very stringent tests of the variation of c, since there have been many experiments to test the conservation of charge [15].

If it is assumed that the electron charge e is constant, charge non-conservation can only be the result of processes that change charge discontinuously. Models for these kind of processes have been proposed in Refs. [12, 16]. An analysis based on this models yield an upper limit on \dot{c}/c that are much more smaller than those obtained by direct measurements [17]: $|\dot{c}/c| < 10^{-31} yr^{-1}$.

A consistent VSL leads to a dynamically prediction of the value of c via a wave-like equation for $\psi = \ln(c/c_0)$, whose source term is proportional to the trace of the energy-momentum tensor. Hence, in the neighborhood of a quasistatic system, such as a star, a generic expression for the variation of ctakes the form [13, 15]

$$c(t) = c_c(t) \left(1 - \lambda \frac{GM}{c_0^2 r}\right).$$
(3)

Here λ is a constant that depends on the specific VSL theory. In the limit $r \to \infty$ the expression for c reduces to the cosmological one $(c \to c_c(t))$. In this work we are interested in finding a bound to the constant λ that parametrizes the charge nonconservation.

In the static regime, the ∇c contributions are second order and can be neglected. In this case there are two contributions to the charge non-conservation: one accounts for time-variation over cosmological time scales and the other for the motion of the earth with respect to massive bodies

$$\frac{\dot{c}}{c} = \left(\frac{\dot{c}}{c}\right)_{\text{cosmological}} + \left(\frac{\dot{c}}{c}\right)_{\text{local}} = n\frac{\dot{a}}{a} + \frac{\lambda GM}{c_0^2 r^3} \, \boldsymbol{r} \cdot \boldsymbol{v} \,, \tag{4}$$

where v is the velocity of the Earth with respect to the center of mass of nearby bodies. The last term of equation (4) represents a breakdown of charge conservation, induced by the presence of local matter.

Also, charge nonconservation may be associated to the breakdown of Einstein's Equivalence principle, as shown in Ref. [14]. In this case, CNC processes would lead to an expression similar to the second term of (4)

$$\frac{Q}{Q} = \frac{\Gamma_0}{4} \frac{\boldsymbol{g}_0 \cdot \boldsymbol{v}}{c_0^2} \tag{5}$$

where g_0 is the local gravitational acceleration and the parameter Γ_0 characterizes anomalous accelerations and anomalous mass tensors [18]:

$$\delta m_P = 2\Gamma_0 \frac{E_C}{c_0^2},\tag{6}$$

$$\Delta a_C = \frac{\delta m_P}{m} g_0. \tag{7}$$

The SAGE [19, 20, 21] and GALLEX-GNO [22, 23, 24, 25, 26, 27] radiochemical solar neutrino experiments are based on the neutrino capture reaction $\nu_e + {}^{71}\text{Ga} \rightarrow e^- + {}^{71}\text{Ge}$. This reaction has a low energy threshold: $E_{th} = 0.2332 \text{ MeV}$ that makes possible the detection of pp solar neutrinos. However it was pointed out in Ref. [28] that in these experiments the appearance of ${}^{71}\text{Ge}$ produced by the ν_e on the target ${}^{71}\text{Ga}$ can in principle be the result of charge nonconservation reaction. The reason is that as there is no electron emitted in a charge nonconservation reaction such as $n \rightarrow p + \gamma$ or $n \rightarrow p + \nu_e + \bar{\nu}_e$, this leads in principle to the possibility that the ${}^{71}\text{Ga}$ undergoes a charge-nonconserving decay to the ground and to the first excited state of the ${}^{71}\text{Ge}$, that otherwise wold be forbidden by energy conservation. Based on the results form SAGE and GALLEX alone it is not possible to discern whether the ${}^{71}\text{Ge}$ was produce via a charge-nonconserving processes or by a normal weak decay; hence it is possible to test those theories that predict the violation of the charge conservation.

In Ref. [15] these results were applied for the case of the Earth motion relative to Virgo supercluster, in this case $v \approx 1000 \text{ km/s}$ and $GM/c_0^2 r \approx 2 \times 10^{-7}$. The charge nonconservation effect would contribute to the CNC ⁷¹Ga \rightarrow ⁷¹Ge + neutrals processes, hence an upper limit on the parameter λ can be derived:

$$|\lambda| \le 2 \times 10^{-10}.\tag{8}$$

The above limit is based on cosmological considerations. Ref. [13] has stressed the importance of local tests of variable speed of light theories. In this paper, we shall develop such a test, based on changes in the rate of Charge Nonconserving Processes (CNC) induced the motion of the Earth around the Sun, that introduces a periodic variation of c with a one year period [29]:

$$\frac{Q}{Q} \simeq -\lambda \frac{GM_{\odot}}{c_0^2 a_{\oplus}} e_{\oplus} \omega_{\oplus} \sin \omega_{\oplus} (t - t_0)$$
(9)

where M_{\odot} is the mass of the Sun, a_{\oplus} and $e_{\oplus} \approx 0.0167$ are the major axis and eccentricity of the orbit of the Earth, and $\omega_{\oplus} = 2\pi/T_{\oplus}$ the frequency (mean motion) of the Earth motion around the Sun. Since t_0 , the time of perihelion passage, is around January 10, a periodic signal may be discerned (or bounded) comparing spring and autumn results from experiments testing charge conservation. In the case of the Gallium experiments a seasonal effect is expected; the term $\mathbf{r} \cdot \mathbf{v}$ varies along the year in a sinusoidal way. As previously discussed, a charge nonconservation decay ⁷¹Ga to ⁷¹Ge is energetically allowed. According to Eqs. (2) and (4) a contribution to the charge increase required in this situation can only arise during half of the year, because the term $\mathbf{r} \cdot \mathbf{v}$ changes sign during the other half and the reverse process (⁷¹Ge to ⁷¹Ga) is energetically forbidden. Hence the average contribution of the charge nonconserving effect to the ⁷¹Ge counting for the spring period would be expected to be different as compared to that of autumn period.

Combining the relevant terms of Eqs.(2) and (4) the relative charge variation can be written as

$$\frac{Q}{Q} = \frac{\lambda GM}{c_0^2 \, a_{\oplus}} \Lambda \,, \tag{10}$$

Here $GM/c_0^2 a_{\oplus} \approx 1 \times 10^{-8}$, and Λ is the time average of the $r \cdot v/r^2$ factor, that for the spring period can be computed as

$$\Lambda = \frac{2e_{\oplus}h_{\oplus}}{\left(1 + e_{\oplus}\right)^3 a_{\oplus}^2} \left(1 + \frac{e_{\oplus}^2}{3}\right),\tag{11}$$

where h = L/m is the angular momentum per unit mass of the Earth and the rest of the notation is the same as in Eq. (9). Hence $h_{\oplus}/a_{\oplus} \approx 29.79 \,\mathrm{km/s}$ corresponds to the mean orbital velocity.

The SAGE and GALLEX collaborations have reported results of 73 ± 18 and 78 ± 12 solar neutrino units (SNU) respectively, where $1 \text{ SNU} = 10^{-36}$ neutrino captures per target atom per second. Furthermore the SAGE collaboration have reported the neutrino rates for periods of 30 days along the whole year. Unfortunately, the GALLEX-GNO collaboration have not reported monthly or bimonthly averages. The seasonal averages where computed from the data for individual runs in Refs. [22, 23, 24, 25, 26, 27] and from the monthly data in Refs. [19, 20, 21]. Analyzing these results the asymmetry for the spring period as compared with autumn one yields

$$\Delta \Phi_{\text{GALLEX+GNO}} = 11.3 \pm 9.8 \text{ SNU}$$

$$\Delta \Phi_{\text{SAGE}} = -0.5 \pm 10.5 \text{ SNU}$$

$$\Delta \Phi_{\text{Comb}} = 5.8 \pm 7.2 \text{ SNU}$$
(12)

The maximum possible flux asymmetry $\Delta \Phi$ when these two periods are compared can be taken as a 3σ limit:

$$|\Delta \Phi| \le 21.6 \text{ SNU.} \tag{13}$$

From this result we conclude that the charge asymmetry \dot{Q}/Q can be no larger than $21.6 \times 10^{-36} \,\mathrm{s}^{-1}$ per target atom; in combination with the previous results it leads to the following bound for the parameter λ :

$$|\lambda| \le 2 \times 10^{-19} \,. \tag{14}$$

This bound is about 10^9 times smaller than the previous one, based on cosmological considerations.

A similar bound would be found for the breakdown of Einstein's Equivalence Principle:

$$|\Gamma_0| \le 10^{-18}.$$
 (15)

This is somewhat smaller than the expected accuracy of STEP [30, 31].

These results can also be used to derive a bound for the ${}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge} +$ neutrals lifetime, in the case of processes of the type proposed in Refs [12, 16]. From the limits on the $\Delta \Phi = 21.3$ SNU asymmetry we conclude that the chargenonconserving decay rate of ${}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge}$ fulfills the relation

$$\frac{dN\left(^{71}\mathrm{Ga}\right)}{dt} = \lambda_{\mathrm{CNC}}N\left(^{71}\mathrm{Ga}\right) < \Delta\Phi N\left(^{71}\mathrm{Ga}\right)$$
(16)

where $N(^{71}\text{Ga})$ is the number of ^{71}Ga atoms in the neutrino detectors and λ_{CNC} is the decay rate for the $^{71}\text{Ga} \rightarrow ^{71}\text{Ge}$ charge-nonconserving decay. Hence the bound for the lifetime of this processes yields

$$\tau_{\rm CNC} \left({}^{71}{\rm Ga} \to {}^{71}{\rm Ge} \right) \ge 1.4 \times 10^{27} \text{ years} \,. \tag{17}$$

This result should be compared with the previous upper limit $\tau_{\rm CNC} \geq 3.5 \times 10^{26}$ years obtained in Ref. [28]. This bound was obtained by considering the the total counts of 73 ± 18 SNU for SAGE and 78 ± 12 SNU for GALLEX. In the same way, a bound for the branching ratio of CNC processes to beta decay can be obtained as in Ref. [28] from the seasonal asymmetry effect produced by the second term of Eq. (4).

$$\Gamma(n \to p + \nu_e + \bar{\nu}_e) / \Gamma(n \to p + e + \bar{\nu}_e) \le 2 \times 10^{-27}$$
(18)

We have shown that the absence of seasonal spring-autumn variations in Gallium solar neutrino experiments imposes extremely tight bounds on the parameters of varying speed of light theories, some 10^9 times smaller than those obtained from cosmological considerations. This not only confirms the importance of local tests of VSL theories, as stressed in Ref. [13] but also imposes very strict constraints on their structure.

H. V. acknowledges the hospitality of IFUNAM. This work was partially supported by CONACyT grants No. G32736-E and U42046.

References

- J. K. Webb, V. V. Flambaum, C. W. Churchill, M. J. Drinkwater, and J. D. Barrow, Phys. Rev. Lett. 82, 884 (1999), astro-ph/9803165.
- [2] J. K. Webb et al., Phys. Rev. Lett. 87, 091301 (2001), astro-ph/0012539.
- [3] J. Magueijo, J. D. Barrow, and H. B. Sandvik, Phys. Rev. Lett. (2002), astro-ph/0202374.
- [4] J. D. Barrow, J. Magueijo, and H. B. Sandvik, Phys. Lett. B541, 201 (2002), astro-ph/0204357.

- [5] A. Peres (2002), quant-ph/0209114.
- [6] A. Albrecht and J. Magueijo, Phys. Rev. D 59, 043516 (1999).
- [7] J. D. Barrow and J. Magueijo, Phys. Lett. B443, 104 (1998), astro-ph/9811072.
- [8] J. D. Barrow and J. Magueijo, Class. Quant. Grav. 16, 1435 (1999), astro-ph/9901049.
- [9] J. Moffat, Int. J. Mod. Phys. D 2, 351 (1993).
- [10] J. Moffat, Found. Phys. 23, 411 (1993).
- [11] J. D. Barrow, Phys. Rev. D 59, 043515 (1999).
- [12] J. Magueijo, Phys. Rev. D 62, 103521 (2000).
- [13] J. Magueijo, Phys. Rev. D 63, 043502 (2001).
- [14] S. J. Landau, P. D. Sisterna, and H. Vucetich (2001), gr-qc/0105025.
- [15] S. J. Landau, P. D. Sisterna, and H. Vucetich, Phys. Rev. D63, 081303 (2001).
- [16] T. Harko and M. K. Mak, Class. Quantum Grav. 16, 2741 (1999).
- [17] J. D. Prestage, R. L. Tjoelker, and L. Maleki, Physical Review Letters 74, 3511 (1995).
- [18] C. M. Will, Theory and Experiment in Gravitational Physics (C. U. P., Cambridge-London, 1981).
- [19] J. N. Abdurashitov et al. (SAGE), Phys. Rev. Lett. 83, 4686 (1999), astro-ph/9907131.
- [20] J. N. Abdurashitov et al. (SAGE), Phys. Rev. C60, 055801 (1999), astro-ph/9907113.
- [21] J. N. Abdurashitov et al. (SAGE), J. Exp. Theor. Phys. 95, 181 (2002).
- [22] G. Collaboration, Phys. Lett. B **314**, 445 (1993).
- [23] G. Collaboration, Phys. Lett. B 357, 237 (1995).
- [24] G. Collaboration, Phys. Lett. B 388, 384 (1996).
- [25] G. Collaboration, Phys. Lett. B 447, 127 (1999).
- [26] M. Altmann et al. (GNO), Phys. Lett. B490, 16 (2000), hep-ex/0006034.
- [27] C. M. Cattadori (GNO), Nucl. Phys. Proc. Suppl. 110, 311 (2002).

- [28] E. B. Norman, J. N. Bahcall, and M. Goldhaber, Phys. Rev. D53, 4086 (1996).
- [29] H. Vucetich, R. C. Mercader, G. Lozano, G. Mindlin, A. R. L. García, and J. Desimoni, Phys. Rev. D38, 2930 (1988).
- [30] R. Reinhard, Y. Jafry, and R. Laurence, ESA J. 17, 251 (1993).
- [31] J. P. Blaser, A. M. Cruise, T. Damour, J. León, H. J. Paik, and R. Rummel, Tech. Rep., ESA (1994), sCI(94)5.