Second-order Perturbations of the Friedmann World Model

Hyerim Noh^(a) and Jai-chan Hwang^(b)

^(a) Korea Astronomy Observatory, Daejon, Korea

^(b) Department of Astronomy and Atmospheric Sciences, Kyungpook National University, Taegu, Korea

(June 30, 2021)

We consider instability of the Friedmann world model to the second-order in perturbations. We present the perturbed set of equations up to the second-order in the Friedmann background world model with general spatial curvature and the cosmological constant. We consider systems with the completely general imperfect fluids, the minimally coupled scalar fields, the electro-magnetic field, and the generalized gravity theories. We also present the case of null geodesic equations, and the one based on the relativistic Boltzmann equation. In due stage a decomposition is made for the scalar-, vector- and tensor-type perturbations which couple each other to the second-order. Gauge issue is resolved to each order. The basic equations are presented without imposing any gauge condition, thus in a gauge-ready form so that we can use the full advantage of having the gauge freedom in analysing the problems. As an application we show that to the second-order in perturbation the relativistic pressureless ideal fluid of the scalar-type reproduces exactly the known Newtonian result. As another application we rederive the large-scale conserved quantities (of the pure scalar-and tensor-perturbations) to the second order, first shown by Salopek and Bond, now from the exact equations. Several other applications are made as well.

PACS numbers: 04.50.+h, 04.62.+v, 98.80.-k, 98.80.Hw

"the linear perturbations are so surprisingly simple that a perturbation analysis accurate to second order may be feasible"

Sachs and Wolfe (1967)

Contents

L	Introduction	2
II	Basic equationsAADM $(3+1)$ equationsBCovariant $(1+3)$ equationsCMulti-component situation	5 5 6 8
III	Perturbed quantities A Metric and connections B Normal-frame quantities C General (\tilde{u}_a) -frame quantities D Fluid quantities E Frame choice F Spacetime curvatures to the linear-order	9 9 10 11 12 13 14
IV	Perturbed equations A Basic equations with general fluids B Scalar field 1 Covariant equations 2 Perturbations 3 Covariant equations 4 Covariant equations 5 Calar fields 6 Scalar fields 7 Perturbations 8 Covariant equations 9 Perturbations	 17 19 20 21 21 21
	D Generalized gravity theories 1 Covariant equations 2 Perturbed equations	22 22 23

	Ε	Electromagnetic field
		1 Covariant equations
		2 Perturbations
	F	Null-geodesic and temperature anisotropy
	G	Boltzmann equation
		1 Covariant equations
		2 Perturbed equations
\mathbf{V}	\mathbf{De}	composition 32
	А	Three perturbation types
	В	Background equations
	С	Decomposed equations
VI	Ga	uge Issue 36
	А	Gauge transformation
	В	Linear-order
		1 Temporal gauge conditions
		2 Spatial gauge conditions
	С	Second-order
		1 Gauge conditions
		2 Constructing gauge-invariant combinations
		3 Gauge-invariant variables
		4 Spatial gradient variable
VII	Ap	plications 47
	Α	Closed form equations
	В	Solutions to the linear-order
		1 Scalar-type
		2 Vector-type
		3 Tensor-type $\ldots \ldots \ldots$
	С	Pressureless irrotational fluid
		1 Nonlinear equation based on $3 + 1$ formulation $\ldots \ldots 53$
		2 Nonlinear equation based on $1 + 3$ formulation $\ldots \ldots 53$
	_	3 Gravitational wave as a source
	D	Pure scalar-type perturbation
	E	Pure rotation
	F	Pure gravitational wave
	G	Action formulation
VIII	Dis	scussions 57

I. INTRODUCTION

We consider instabilities of the spatially homogeneous and isotropic comological spacetime to the second-order in perturbations. The relativistic cosmological perturbation plays fundamental roles in the modern theory of large-scale cosmic structure formation. The original analysis of linear perturbations based on Einstein gravity with a hydrodynamic fluid was made by Lifshitz in 1946 [1] in an almost complete form. Due to the extremely low level anisotropies of the cosmic microwave background (CMB) radiation, the cosmological dynamics of the structures in the large-scale and in the early universe are generally believed to be small deviations from the homogeneous and isotropic background Friedmann world model [2]. The conventional relativisitic cosmological perturbation analysis considers such deviations small enough so that one could treat them as *linear*. The linear perturbation theory works as the basic framework in handling the cosmological structure formation processes. Recent observations of the CMB anisotropies in all sky by the WMAP satellite and others [3], for example, assure the validity of the basic assumptions used in the cosmological perturbation theory, i.e., the linearity of the relevant cosmic structures [4].

Still, as the observed relatively small-scale structures are apparently nonlinear, the gravitational instability based on the pure linear theory is not enough for the complete picture. It is agreed that such small-scale nonlinear structures could be handled by Newtonian gravity often based on numerical simulations. The current paradigm of large-scale structure generation and evolution processes is based on an underlying assumption that linear processes dominated until nonlinear processes take over on subhorizon scales in the Newtonian regime. Thus, it seems this paradigm of our understanding of the origin and evolution of the large-scale structures is rather satisfactory within the linear theory concerning the regimes where the relativistic gravity theory is needed. It is well known that in the linear theory there can be no structure formation. In fact, this "no structure formation" in the scenario is precisely why we were successful in describing the structure generation and evolution processes in a simple manner as we describe below. In the standard scenario, the initial conditions (seed fluctuations) generated from quantum fluctuations are imprinted into ripples in the spacetime, and its spatial structures are preserved as the raw large-scale structures (i.e. are as yet unaffected by nonlinear processes). This is a trait which can be traced to the linearity assumption we adopt. It gives a simple, but fictitious system. We could, perhaps, fairly describe the current situation as: the linear paradigm is not inconsistent with observations, especially with the low level of observed anisotropy in the CMB [3]. However, we should remember that the actual equations we are dealing with, both in the gravity and the quantum, are highly nonlinear. It forms an intrinsically complex system.

We can decompose the perturbations into three different types: the scalar-type (associated with the density condensation), the vector-type (rotation) and the tensor-type (gravitational wave) perturbations. To the linear-order perturbation, due to the high symmetry of the background space, these three types of perturbations decouple from each other and evolve independently. Both (the linearity and the homogeneous-isotropic background) conditions are necessary to have natural descriptions of the three types of perturbations independently. We will see how couplings occur to the second-order perturbations; for the couplings in a simplest spatially homogeneous but anisotropic spacetime, see [5].

Another aspect of the simple nature of the linear processes in relativistic gravity theories is characterized by conservations (in expanding phase) of certain amplitudes in the super-horizon scale where we naively anticipate the independent evolution of causally disconnected regions. On superhorizon scales this conserved character is presented by an equation of the form

$$\Phi(\mathbf{x},t) = C(\mathbf{x}),\tag{1}$$

which applies for both the scalar- and tensor-type perturbations; for the vector-type perturbation the angular momentum is conserved in all scales. $C(\mathbf{x})$ is an integration constant of the integral form general solutions available in the large-scale limit, see eqs. (311,332,333). The coefficient $C(\mathbf{x})$ contains information about spatial structure which will eventually grow to the large-scale structure and the gravitational wave background. It can be considered as an initial condition for each perturbation variable which is preserved during the linear evolution. Whether a similar conserved variable can exist even in a nonlinear analysis is apparently an interesting question: for the presence of such variables to the second-order in perturbation see \S VII D-VII F. In the analyses of the large-scale structures in the linear stage, the simple behavior of the conserved variables is practically important. In fact, if we know C the behavior of all the other variables can be determined through linear algebra. Using the conserved quantity one can trivially relate the currently observable (or deducible) linear structure directly to the initial state of the structure at the early universe; probably, just after the scale effectively becomes the large-scale during the hypothetical early acceleration (inflation) stage. Of course, the underlying assumption for all of these results is the applicability of the linear analysis. As long as this assumption is valid, the initial condition is imprinted onto the large-scale structure and is preserved until the nonlinear effects become important. Although this is a big advantage, in a sense this is very consistent with the fact that no structure formation occurs in the linearized system.

The linear perturbation theory is currently well developed; see [6–11]. Although the observations do not desparately demand to go beyond the linear theory, the second-order perturbation theory is a natural next step in the theoretical investigations. The second-order perturbation theory, if well developed, will have important implications on our understanding of the large-scale structure formation processes. Not only the structures we discover are nonlinear, according to the gravitational instability there should occur (perhaps smooth) transitions from the linear to the nonlinear ones. Even in the theoretical point of view, in order to know the limit of the linear perturbation theory we need the behavior of perturbations beyond the linear theory. It is not possible to know the limit of linear theory within the context of the linear theory. It is yet unclear whether the second-order 'perturbation theory' will provide an answer to such a question, but we expect it could provide a better perspective on the problem than the simple linear theory. There will be more practical applications as well, like investigating the non-Gaussian signature in the inflation generated seed fluctuations which could have left a detectable signature in the CMB anisotropies and the large-scale structures. Other possible situations where the trans-linear analyses might be useful are summarized in $\frac{8}{100}$

Now, we discuss the gauge issue present in the relativistic perturbation theory briefly. Since the unperturbed background spacetime is spatially homogeneous and isotropic, to the linear-order the ambiguity caused by spatial gauge (coordinate transformation) freedom does not play a role [7]. Thus, to the linear-order it is appropriate to

write the perturbed set of equations in terms of natural combinations of variables which are invariant under spatial gauge transformation. However, the temporal gauge freedom can be conveniently used in analyzing various aspects of perturbation problem in the Friedmann background. There are infinitely many different ways of taking the temporal gauge (choosing the spatial hypersurface) conditions, and we can identify several fundamental gauge choices [7,12]. Except for a widely used temporal synchronous gauge fixing condition ($\delta g_{00} \equiv 0$), each of the other gauge conditions fixes completely the temporal gauge freedom. Thus each having its own corresponding gauge invariant formulation. In our study of the linear theory we found that some particular gauge invariant variables show the correct Newtonian behaviors. A perturbed density variable in the comoving gauge and a perturbed potential variable and a perturbed velocity variable in the zero-shear gauge most closely resemble the behaviors of the corresponding Newtonian variables [13]. Also, the scalar field perturbation in the uniform-curvature gauge most closely resembles the scalar field equation in the quantum field in curved space [14,15]. Since the gauge conditions mentioned allow no room for remaining gauge mode the variable in a given such gauges has the unique corresponding gauge invariant combination of variables. Thus, the variable in such a gauge is equivalent to the corresponding gauge invariant combination.

In the gauge theory it is well known that proper choice of the gauge condition is often necessary for proper handling of the problem. Either by fixing certain gauge conditions or by choosing certain gauge-invariant combinations in the early calculation stage we are likely to lose possible advantages available in the other gauge conditions. In order to use the various gauge conditions as advantage in handling cosmological perturbations we have proposed a *gaugeready* method which allows the flexible use of the various fundamental gauge conditions. The strategy is that, in order to use the various available temporal gauge conditions as advantage, we had better present the basic equations without choosing any temporal gauge condition, and arrange the equations so that we could choose the fundamental gauge conditions. Our gauge-ready strategy, together with our notation for indicating the gauge-invariant combinations, allows us to use the gauge freedom as the advantage in analysing various given problems. We follow the wisdom suggested by Bardeen in 1988 "the moral is that one should work in the gauge that is mathematically most convenient for the problem at hand" [7].

As long as we are taking perturbative approach the gauge issue in the higher-orders can be resolved similarly as in the linear theory. To the second-order we will identify two variables which can be used to fix the spatial gauge freedom. One gauge condition completely removes the spatial gauge mode, whereas the other condition does not; *i.e.*, in the latter case even after imposing the gauge condition there still remains a degree of freedom which is a gauge mode (a coordinate artifact). We call this incomplete gauge condition the *B*-gauge ($\tilde{g}_{0\alpha} \equiv 0$ where α is a spatial index) whereas the other complete condition is called the *C*-gauge. To the second-order we can identify the same several temporal gauge fixing conditions. Again, except for the synchronous gauge each of the other gauge conditions removes completely the temporal gauge modes.

It is amusing to note that the classic study by Lifshitz [1] adopted the synchronous gauge condition which is a combination of the temporal synchronous gauge condition and the spatial *B*-gauge condition, thus failing to fix both the temporal and the spatial gauge modes completely. This has caused some prevalent errors in the literature based on the synchronous gauge: see the Appendix of [13]. However, we note that these errors are simple algebraic ones probably caused by slightly more complicated algebra due to the presence of the gauge mode after the synchronous gauge fixing. We would like to emphasize that the gauge condition should be appropriately used according to the character of each problem at hand. We have such a freedom because Einstein's gravity theory might be regarded as a gauge theory [16]. In this sense, although the temporal synchronous and the spatial *B*-gauge conditions do not remove completely the gauge modes, often even these conditions could possibly turn out to be convenient in certain problems. Since physically measurable quantities should be gauge invariant we propose to use the gauge conditions in this pragmatic sense.

In a classic study of CMB anisotropy in 1967 Sachs and Wolfe have mentioned that "the linear perturbations are so surprisingly simple that a perturbation analysis accurate to second order may be feasible ..." [17]. In this work, we will present the basic formulation of the second-order perturbation of the Friedmann world model in details. We will present the basic equations needed to investigate the second-order perturbation in a rather general context. We will consider the most general Friedmann background with K and Λ . We will consider the most general imperfect fluid situation. This includes multiple imperfect fluids with general interactions among them. We will also include minimally coupled scalar fields, a class of generalized gravity theories, the electromagnetic fields, the null geodesic, and the relativistic Boltzmann equation. In order to use the gauge fixing conditions optimally we will present the complete sets of perturbed equations in the gauge-ready form. In this manner, as in the linear theory, we could easily apply the equations to any gauge conditions which make the mathematical analyses of given problems simplest. Our formulation will be suitable to handle the nonlinear evolutions in the perturbative manner. Ours will be a useful complement to the other methods suggested in the literature to investigate the trans-linear regimes. In the Discussions we summarize the related studies, the different methods we could employ for further applications, and the cosmological situations where our formulation could be applied fruitfully. Although we will present some trivial applications in the later part, the main applications are left for future studies.

In §II we summarize the basic equations of the Einstein's gravity expressed in the ADM (3+1) formulation and in the covariant (1+3) formulation. In §III we introduce our definition of the metric and the energy-momentum tensor to the second-order in perturbation and present some useful quantities appearing in the ADM and the covariant formulations. In §IV we present the complete sets of perturbed equations up to the second-order in the Friedmann world model. We consider the general spatial curvature and the cosmological constant in the background. We consider systems with the completely general imperfect fluids. Such a general formulation can be reinterpreted to include the cases of the minimally coupled scalar fields, the electro-magnetic field, and even the generalized gravity theories. We present the complete sets of equations for these systems as well. We also present the case of null geodesic equations, and the one based on the relativistic Boltzmann equation. In §V we introduce decomposition of perturbations to three different types and show how these couple to each other to the second-order. All equations up to this point are presented without introducing any gauge condition. Thus, the equations are presented in the most general forms, and any suitable gauge conditions can be easily deployed to these equations. In this sense, our set of equations is in a gauge-ready form. In §VI we address the gauge issue, and show that the gauge issue can be resolved to each perturbation order, just like the case in the linear perturbation. We implement our gauge-ready strategy to the second-order in perturbations. In §VII we make several applications. In §VIII we summarize the main results and outline future applications of our work.

As a unit we set $c \equiv 1$.

II. BASIC EQUATIONS

A. ADM (3+1) equations

The ADM (Arnowitt-Deser-Misner) equations [18] are based on splitting the spacetime into the spatial and the temporal parts using a normal vector field \tilde{n}_a . The metric is written as (we put tilde on the covariant variables):

$$\tilde{g}_{00} \equiv -N^2 + N^{\alpha} N_{\alpha}, \quad \tilde{g}_{0\alpha} \equiv N_{\alpha}, \quad \tilde{g}_{\alpha\beta} \equiv h_{\alpha\beta}, \\ \tilde{g}^{00} = -N^{-2}, \quad \tilde{g}^{0\alpha} = N^{-2} N^{\alpha}, \quad \tilde{g}^{\alpha\beta} = h^{\alpha\beta} - N^{-2} N^{\alpha} N^{\beta},$$
(2)

where N_{α} is based on $h_{\alpha\beta}$ as the metric and $h^{\alpha\beta}$ is an inverse metric of $h_{\alpha\beta}$. Indices a, b, \ldots indicate the spacetime indices, and α, β, \ldots indicate the spatial ones. The normal vector \tilde{n}_a is introduced as:

$$\tilde{n}_0 \equiv -N, \quad \tilde{n}_\alpha \equiv 0, \quad \tilde{n}^0 = N^{-1}, \quad \tilde{n}^\alpha = -N^{-1}N^\alpha.$$
(3)

The fluid quantities are defined as:

$$E \equiv \tilde{n}_a \tilde{n}_b \tilde{T}^{ab}, \quad J_\alpha \equiv -\tilde{n}_b \tilde{T}^b_\alpha, \quad S_{\alpha\beta} \equiv \tilde{T}_{\alpha\beta}, \quad S \equiv h^{\alpha\beta} S_{\alpha\beta}, \quad \bar{S}_{\alpha\beta} \equiv S_{\alpha\beta} - \frac{1}{3} h_{\alpha\beta} S, \tag{4}$$

where J_{α} and $S_{\alpha\beta}$ are based on $h_{\alpha\beta}$. The extrinsic curvature is introduced as

$$K_{\alpha\beta} \equiv \frac{1}{2N} \left(N_{\alpha:\beta} + N_{\beta:\alpha} - h_{\alpha\beta,0} \right), \quad K \equiv h^{\alpha\beta} K_{\alpha\beta}, \quad \bar{K}_{\alpha\beta} \equiv K_{\alpha\beta} - \frac{1}{3} h_{\alpha\beta} K, \tag{5}$$

where $K_{\alpha\beta}$ is based on $h_{\alpha\beta}$. A colon ':' denotes a covariant derivative based on $h_{\alpha\beta}$. The connections become:

$$\tilde{\Gamma}_{00}^{0} = \frac{1}{N} \left(N_{,0} + N_{,\alpha} N^{\alpha} - K_{\alpha\beta} N^{\alpha} N^{\beta} \right), \quad \tilde{\Gamma}_{0\alpha}^{0} = \frac{1}{N} \left(N_{,\alpha} - K_{\alpha\beta} N^{\beta} \right), \quad \tilde{\Gamma}_{\alpha\beta}^{0} = -\frac{1}{N} K_{\alpha\beta},$$

$$\tilde{\Gamma}_{00}^{\alpha} = \frac{1}{N} N^{\alpha} \left(-N_{,0} - N_{,\beta} N^{\beta} + K_{\beta\gamma} N^{\beta} N^{\gamma} \right) + N N^{,\alpha} + N^{\alpha}_{,0} - 2N K^{\alpha\beta} N_{\beta} + N^{\alpha:\beta} N_{\beta},$$

$$\tilde{\Gamma}_{0\beta}^{\alpha} = -\frac{1}{N} N_{,\beta} N^{\alpha} - N K^{\alpha}_{\beta} + N^{\alpha}_{;\beta} + \frac{1}{N} N^{\alpha} N^{\gamma} K_{\beta\gamma}, \quad \tilde{\Gamma}_{\beta\gamma}^{\alpha} = \Gamma^{(h)\alpha}_{\ \beta\gamma} + \frac{1}{N} N^{\alpha} K_{\beta\gamma}, \quad (6)$$

where $\Gamma^{(h)\alpha}_{\beta\gamma}$ is the connection based on $h_{\alpha\beta}$ as the metric, $\Gamma^{(h)\alpha}_{\beta\gamma} \equiv \frac{1}{2}h^{\alpha\delta}(h_{\beta\delta,\gamma} + h_{\delta\gamma,\beta} - h_{\beta\gamma,\delta})$. The intrinsic curvatures are based on $h_{\alpha\beta}$ as the metric:

$$R^{(h)\alpha}_{\ \beta\gamma\delta} \equiv \Gamma^{(h)\alpha}_{\ \beta\delta,\gamma} - \Gamma^{(h)\alpha}_{\ \beta\gamma,\delta} + \Gamma^{(h)\epsilon}_{\ \beta\delta}\Gamma^{(h)\alpha}_{\ \gamma\epsilon} - \Gamma^{(h)\epsilon}_{\ \beta\gamma}\Gamma^{(h)\alpha}_{\ \delta\epsilon},$$

$$R^{(h)}_{\alpha\beta} \equiv R^{(h)\gamma}_{\ \alpha\gamma\beta}, \quad R^{(h)} \equiv h^{\alpha\beta}R^{(h)}_{\alpha\beta}, \quad \bar{R}^{(h)}_{\alpha\beta} \equiv R^{(h)}_{\alpha\beta} - \frac{1}{3}h_{\alpha\beta}R^{(h)}.$$
(7)

A complete set of the ADM equations is the following [6]. Energy constraint equation:

$$R^{(h)} = \bar{K}^{\alpha\beta}\bar{K}_{\alpha\beta} - \frac{2}{3}K^2 + 16\pi GE + 2\Lambda,$$
(8)

where Λ is the cosmological constant. Momentum constraint equation:

$$\bar{K}^{\beta}_{\alpha:\beta} - \frac{2}{3}K_{,\alpha} = 8\pi G J_{\alpha}.$$
(9)

Trace of ADM propagation equation:

$$K_{,0}N^{-1} - K_{,\alpha}N^{\alpha}N^{-1} + N^{\alpha}{}_{\alpha}N^{-1} - \bar{K}^{\alpha\beta}\bar{K}_{\alpha\beta} - \frac{1}{3}K^2 - 4\pi G\left(E+S\right) + \Lambda = 0.$$
(10)

Tracefree ADM propagation equation:

$$\bar{K}^{\alpha}_{\beta,0}N^{-1} - \bar{K}^{\alpha}_{\beta;\gamma}N^{\gamma}N^{-1} + \bar{K}_{\beta\gamma}N^{\alpha;\gamma}N^{-1} - \bar{K}^{\alpha}_{\gamma}N^{\gamma}{}_{;\beta}N^{-1} \\
= K\bar{K}^{\alpha}_{\beta} - \left(N^{;\alpha}{}_{\beta} - \frac{1}{3}\delta^{\alpha}_{\beta}N^{;\gamma}{}_{\gamma}\right)N^{-1} + \bar{R}^{(h)\alpha}{}_{\beta} - 8\pi G\bar{S}^{\alpha}_{\beta}.$$
(11)

Energy conservation equation:

$$E_{,0}N^{-1} - E_{,\alpha}N^{\alpha}N^{-1} - K\left(E + \frac{1}{3}S\right) - \bar{S}^{\alpha\beta}\bar{K}_{\alpha\beta} + N^{-2}\left(N^2J^{\alpha}\right)_{:\alpha} = 0.$$
 (12)

Momentum conservation equation:

$$J_{\alpha,0}N^{-1} - J_{\alpha;\beta}N^{\beta}N^{-1} - J_{\beta}N^{\beta}{}_{;\alpha}N^{-1} - KJ_{\alpha} + EN_{,\alpha}N^{-1} + S^{\beta}_{\alpha;\beta} + S^{\beta}_{\alpha}N_{,\beta}N^{-1} = 0.$$
(13)

B. Covariant (1+3) equations

The covariant formulation of Einstein gravity was investigated in [19,20]. The 1 + 3 covariant decomposition is based on the time-like normalized ($\tilde{u}^a \tilde{u}_a \equiv -1$) four-vector field \tilde{u}_a introduced in all spacetime points. The expansion ($\tilde{\theta}$), the acceleration (\tilde{a}_a), the rotation ($\tilde{\omega}_{ab}$), and the shear ($\tilde{\sigma}_{ab}$) are kinematic quantities of the projected covariant derivative of flow vector \tilde{u}_a introduced as

$$\tilde{h}_{a}^{c}\tilde{h}_{b}^{d}\tilde{u}_{c;d} = \tilde{h}_{[a}^{c}\tilde{h}_{b]}^{d}\tilde{u}_{c;d} + \tilde{h}_{(a}^{c}\tilde{h}_{b]}^{d}\tilde{u}_{c;d} \equiv \tilde{\omega}_{ab} + \tilde{\theta}_{ab} = \tilde{u}_{a;b} + \tilde{a}_{a}\tilde{u}_{b},$$

$$\tilde{\sigma}_{ab} \equiv \tilde{\theta}_{ab} - \frac{1}{3}\tilde{\theta}\tilde{h}_{ab}, \quad \tilde{\theta} \equiv \tilde{u}^{a}{}_{;a}, \quad \tilde{a}_{a} \equiv \tilde{\tilde{u}}_{a} \equiv \tilde{u}_{a;b}\tilde{u}^{b},$$
(14)

where $\tilde{h}_{ab} \equiv \tilde{g}_{ab} + \tilde{u}_a \tilde{u}_b$ is the projection tensor with $\tilde{h}_{ab} \tilde{u}^b = 0$ and $\tilde{h}^a_a = 3$. An overdot with tilde \tilde{i} indicates a covariant derivative along \tilde{u}^a . We have

$$\tilde{u}_{a;b} = \tilde{\omega}_{ab} + \tilde{\sigma}_{ab} + \frac{1}{3}\tilde{\theta}\tilde{h}_{ab} - \tilde{a}_a\tilde{u}_b.$$
(15)

We introduce

$$\tilde{\omega}^{a} \equiv \frac{1}{2} \tilde{\eta}^{abcd} \tilde{u}_{b} \tilde{\omega}_{cd}, \quad \tilde{\omega}_{ab} = \tilde{\eta}_{abcd} \tilde{\omega}^{c} \tilde{u}^{d}, \quad \tilde{\omega}^{2} \equiv \frac{1}{2} \tilde{\omega}^{ab} \tilde{\omega}_{ab} = \tilde{\omega}^{a} \tilde{\omega}_{a}, \quad \tilde{\sigma}^{2} \equiv \frac{1}{2} \tilde{\sigma}^{ab} \tilde{\sigma}_{ab}, \tag{16}$$

where $\tilde{\omega}^a$ is a vorticity vector which has the same information as the vorticity tensor $\tilde{\omega}_{ab}$. We have $\tilde{\eta}^{abcd} = \tilde{\eta}^{[abcd]}$ with $\tilde{\eta}^{1234} = 1/\sqrt{-\tilde{g}}$; indices surrounded by () and [] are the symmetrization and anti-symmetrization symbols, respectively. Our convention of the Riemann curvature and Einstein's equation are:

fur convention of the memanic curvature and Emistern's equation are.

$$\tilde{u}_{a;bc} - \tilde{u}_{a;cb} = \tilde{u}_d R^d_{\ abc},\tag{17}$$

$$\tilde{R}_{ab} - \frac{1}{2}\tilde{R}\tilde{g}_{ab} = 8\pi G\tilde{T}_{ab} - \Lambda\tilde{g}_{ab}.$$
(18)

The Weyl (conformal) curvature is introduced as

$$\tilde{C}_{abcd} \equiv \tilde{R}_{abcd} - \frac{1}{2} \left(\tilde{g}_{ac} \tilde{R}_{bd} + \tilde{g}_{bd} \tilde{R}_{ac} - \tilde{g}_{bc} \tilde{R}_{ad} - \tilde{g}_{ad} \tilde{R}_{bc} \right) + \frac{\dot{R}}{6} \left(\tilde{g}_{ac} \tilde{g}_{bd} - \tilde{g}_{ad} \tilde{g}_{bc} \right).$$

$$\tag{19}$$

The electric and magnetic parts of the Weyl curvature are introduced as:

$$\tilde{E}_{ab} \equiv \tilde{C}_{acbd} \tilde{u}^c \tilde{u}^d, \quad \tilde{H}_{ab} \equiv \frac{1}{2} \tilde{\eta}_{ac}{}^{ef} \tilde{C}_{efbd} \tilde{u}^c \tilde{u}^d.$$
⁽²⁰⁾

The energy-momentum tensor is decomposed into fluid quantities based on the four-vector field \tilde{u}^a as

$$\tilde{T}_{ab} \equiv \tilde{\mu}\tilde{u}_a\tilde{u}_b + \tilde{p}\left(\tilde{g}_{ab} + \tilde{u}_a\tilde{u}_b\right) + \tilde{q}_a\tilde{u}_b + \tilde{q}_b\tilde{u}_a + \tilde{\pi}_{ab},\tag{21}$$

where

$$\tilde{u}^a \tilde{q}_a = 0 = \tilde{u}^a \tilde{\pi}_{ab}, \quad \tilde{\pi}_{ab} = \tilde{\pi}_{ba}, \quad \tilde{\pi}^a_a = 0.$$
(22)

The variables $\tilde{\mu}$, \tilde{p} , \tilde{q}_a and $\tilde{\pi}_{ab}$ are the energy density, the isotropic pressure (including the entropic one), the energy flux and the anisotropic pressure based on \tilde{u}_a -frame, respectively. We have

$$\tilde{\mu} \equiv \tilde{T}_{ab}\tilde{u}^a\tilde{u}^b, \quad \tilde{p} \equiv \frac{1}{3}\tilde{T}_{ab}\tilde{h}^{ab}, \quad \tilde{q}_a \equiv -\tilde{T}_{cd}\tilde{u}^c\tilde{h}_a^d, \quad \tilde{\pi}_{ab} \equiv \tilde{T}_{cd}\tilde{h}_a^c\tilde{h}_b^d - \tilde{p}\tilde{h}_{ab}.$$
(23)

The specific entropy \tilde{S} can be introduced by $\tilde{T}d\tilde{S} = d\tilde{\varepsilon} + \tilde{p}_T d\tilde{v}$ where $\tilde{\varepsilon}$ is specific internal energy density with $\tilde{\mu} = \tilde{\varrho}(1+\tilde{\varepsilon}), \ \tilde{p}_T$ the thermodynamic pressure, $\tilde{v} \equiv 1/\tilde{\varrho}$ the specific volume, and \tilde{T} the temperature. We have the isotropic pressure $\tilde{p} = \tilde{p}_T + \tilde{\pi}$ where $\tilde{\pi}$ is the entropic pressure. Using eq. (26) below we can show

$$\tilde{\varrho}\tilde{T}\tilde{\tilde{S}} = -\left(\tilde{\pi}\tilde{\theta} + \tilde{\pi}^{ab}\tilde{\sigma}_{ab} + \tilde{q}^{a}{}_{;a} + \tilde{q}^{a}\tilde{a}_{a}\right).$$
(24)

Thus, we notice that $\tilde{\pi}$, $\tilde{\pi}^{ab}$ and \tilde{q}^{a} generate the entropy. Using a four-vector $\tilde{S}^{a} \equiv \tilde{\varrho}\tilde{u}^{a}\tilde{S} + \frac{1}{\tilde{T}}\tilde{q}^{a}$ which is termed the entropy flow density [19] we can derive

$$\tilde{S}^{a}_{;a} = -\frac{1}{\tilde{T}^{2}} \left(\tilde{T}_{,a} + \tilde{T}\tilde{a}_{a} \right) \tilde{q}^{a} - \frac{1}{\tilde{T}} \left(\tilde{\pi}\tilde{\theta} + \tilde{\pi}^{ab}\tilde{\sigma}_{ab} \right).$$
⁽²⁵⁾

The covariant formulation provides a useful complement to the ADM formulation. We summarize the covariant (1+3) set of equations in the following. For details, see [19,20] and the Appendix in [21]. The energy and the momentum conservation equations:

$$\tilde{\tilde{\mu}} + (\tilde{\mu} + \tilde{p})\,\tilde{\theta} + \tilde{\pi}^{ab}\tilde{\sigma}_{ab} + \tilde{q}^a_{\ ;a} + \tilde{q}^a\tilde{a}_a = 0,$$
(26)

$$(\tilde{\mu} + \tilde{p})\,\tilde{a}_a + \tilde{h}^b_a \left(\tilde{p}_{,b} + \tilde{\pi}^c_{b;c} + \tilde{\tilde{q}}_b\right) + \left(\tilde{\omega}_{ab} + \tilde{\sigma}_{ab} + \frac{4}{3}\tilde{\theta}\tilde{h}_{ab}\right)\tilde{q}^b = 0.$$

$$(27)$$

Raychaudhuri equation:

$$\tilde{\tilde{\theta}} + \frac{1}{3}\tilde{\theta}^2 - \tilde{a}^a{}_{;a} + 2\left(\tilde{\sigma}^2 - \tilde{\omega}^2\right) + 4\pi G\left(\tilde{\mu} + 3\tilde{p}\right) - \Lambda = 0.$$
⁽²⁸⁾

Vorticity propagation:

$$\tilde{h}_{b}^{a}\tilde{\tilde{\omega}}^{b} + \frac{2}{3}\tilde{\theta}\tilde{\omega}^{a} = \tilde{\sigma}_{b}^{a}\tilde{\omega}^{b} + \frac{1}{2}\tilde{\eta}^{abcd}\tilde{u}_{b}\tilde{a}_{c;d}.$$
(29)

Shear propagation:

$$\tilde{h}_{a}^{c}\tilde{h}_{b}^{d}\left(\tilde{\tilde{\sigma}}_{cd}-\tilde{a}_{(c;d)}\right)-\tilde{a}_{a}\tilde{a}_{b}+\tilde{\omega}_{a}\tilde{\omega}_{b}+\tilde{\sigma}_{ac}\tilde{\sigma}_{b}^{c}+\frac{2}{3}\tilde{\theta}\tilde{\sigma}_{ab}-\frac{1}{3}\tilde{h}_{ab}\left(\tilde{\omega}^{2}+2\tilde{\sigma}^{2}-\tilde{a}_{;c}^{c}\right)+\tilde{E}_{ab}-4\pi G\tilde{\pi}_{ab}=0.$$
(30)

Three constraint equations:

$$\tilde{h}_{ab}\left(\tilde{\omega}^{bc}_{\ ;c} - \tilde{\sigma}^{bc}_{\ ;c} + \frac{2}{3}\tilde{\theta}^{;b}\right) + \left(\tilde{\omega}_{ab} + \tilde{\sigma}_{ab}\right)\tilde{a}^b = 8\pi G\tilde{q}_a,\tag{31}$$

$$\tilde{\omega}^{a}{}_{;a} = 2\tilde{\omega}^{b}\tilde{a}_{b},\tag{32}$$

$$\tilde{H}_{ab} = 2\tilde{a}_{(a}\tilde{\omega}_{b)} - \tilde{h}_{a}^{c}\tilde{h}_{b}^{d} \left(\tilde{\omega}_{(c}^{\ e;f} + \tilde{\sigma}_{(c}^{\ e;f} \right) \tilde{\eta}_{d)gef} \tilde{u}^{g}.$$

$$(33)$$

Four quasi-Maxwellian equations:

$$\tilde{h}^a_b \tilde{h}^c_d \tilde{E}^{bd}_{;c} - \tilde{\eta}^{abcd} \tilde{u}_b \tilde{\sigma}^e_c \tilde{H}_{de} + 3\tilde{H}^a_b \tilde{\omega}^b = 4\pi G \left(\frac{2}{3} \tilde{h}^{ab} \tilde{\mu}_{,b} - \tilde{h}^a_b \tilde{\pi}^{bc}_{;c} - 3\tilde{\omega}^a_{\ b} \tilde{q}^b + \tilde{\sigma}^a_b \tilde{q}^b + \tilde{\pi}^a_b \tilde{a}^b - \frac{2}{3} \tilde{\theta} \tilde{q}^a \right),$$
(34)

$$\tilde{h}^{a}_{b}\tilde{h}^{c}_{d}\tilde{H}^{bd}_{;c} + \tilde{\eta}^{abcd}\tilde{u}_{b}\tilde{\sigma}^{e}_{c}\tilde{E}_{de} - 3\tilde{E}^{a}_{b}\tilde{\omega}^{b} = 4\pi G \left\{ 2\left(\tilde{\mu} + \tilde{p}\right)\tilde{\omega}^{a} + \tilde{\eta}^{abcd}\tilde{u}_{b}\left[\tilde{q}_{c;d} + \tilde{\pi}_{ce}\left(\tilde{\omega}^{e}_{\ d} + \tilde{\sigma}^{e}_{\ d}\right)\right] \right\},\tag{35}$$

$$\tilde{h}_{c}^{a}\tilde{h}_{d}^{b}\tilde{E}^{cd} + \left(\tilde{H}_{d;e}^{f}\tilde{h}_{f}^{(a} - 2\tilde{a}_{d}\tilde{H}_{e}^{(a)}\right)\tilde{\eta}^{b)cde}\tilde{u}_{c} + \tilde{h}^{ab}\tilde{\sigma}^{cd}\tilde{E}_{cd} + \tilde{\theta}\tilde{E}^{ab} - \tilde{E}_{c}^{(a)}\left(3\tilde{\sigma}^{b)c} + \tilde{\omega}^{b)c}\right) = 4\pi G \left[-\left(\tilde{\mu} + \tilde{p}\right)\tilde{\sigma}^{ab} - 2\tilde{a}^{(a}\tilde{q}^{b)} - \tilde{h}_{c}^{(a}\tilde{h}_{d}^{b)}\left(\tilde{q}^{c;d} + \tilde{\tilde{\pi}}^{cd}\right) - \left(\tilde{\omega}_{c}^{(a)} + \tilde{\sigma}_{c}^{(a)}\right)\tilde{\pi}^{b)c} - \frac{1}{\tilde{\pi}}\tilde{\theta}\tilde{\pi}^{ab} + \frac{1}{\tilde{\pi}}\left(\tilde{q}^{c}_{,c} + \tilde{a}_{c}\tilde{q}^{c} + \tilde{\pi}^{cd}\tilde{\sigma}_{cd}\right)\tilde{h}^{ab}\right],$$
(36)

$$\tilde{h}_{c}^{a}\tilde{h}_{d}^{b}\overset{\tilde{i}}{\dot{H}}^{cd} - \left(\tilde{E}_{d;e}^{f}\tilde{h}_{f}^{(a} - 2\tilde{a}_{d}\tilde{E}_{e}^{(a)}\right)\tilde{\eta}^{b)cde}\tilde{u}_{c} + \tilde{h}^{ab}\tilde{\sigma}^{cd}\tilde{H}_{cd} + \tilde{\theta}\tilde{H}^{ab} - \tilde{H}_{c}^{(a}\left(3\tilde{\sigma}^{b)c} + \tilde{\omega}^{b)c}\right)$$

$$= 4\pi G \left[\left(\tilde{q}_{e}\tilde{\sigma}_{d}^{(a)} - \tilde{\pi}_{d;e}^{f}\tilde{h}_{f}^{(a)}\right)\tilde{\eta}^{b)cde}\tilde{u}_{c} + \tilde{h}^{ab}\tilde{\omega}_{c}\tilde{q}^{c} - 3\tilde{\omega}^{(a}\tilde{q}^{b)} \right].$$

$$(37)$$

Evaluated in the normal-frame eqs. (26,27,28,30,31) reproduce eqs. (12,13,10,11,9) in the ADM formulation.

Now, we take the normal-frame vector, thus $\tilde{u}_a = \tilde{n}_a$ with $\tilde{n}_{\alpha} \equiv 0$, thus $\tilde{\omega}_{ab} = 0$. The trace and tracefree parts of the Gauss equation give [21]:

$$\tilde{R}^{(3)} = 2\left(-\frac{1}{3}\tilde{\theta}^2 + \tilde{\sigma}^2 + 8\pi G\tilde{\mu} + \Lambda\right),\tag{38}$$

$$\tilde{R}_{ab}^{(3)} - \frac{1}{3}\tilde{R}^{(3)}\tilde{h}_{ab} = \tilde{h}_{a}^{c}\tilde{h}_{b}^{d}\left(-\tilde{\tilde{\sigma}}_{cd} - \tilde{\theta}\tilde{\sigma}_{cd} + \tilde{a}_{(c;d)}\right) + \tilde{a}_{a}\tilde{a}_{b} - \frac{1}{3}\tilde{h}_{ab}\tilde{a}^{c}_{;c} + 8\pi G\tilde{\pi}_{ab},\tag{39}$$

where $\tilde{R}_{ab}^{(3)}$ and $\tilde{R}^{(3)}$ are the Ricci and scalar curvatures of the hypersurface normal to \tilde{n}_a ; for an arbitrary vector \tilde{V}_a we have

$$\tilde{R}^{(3)}_{abcd}\tilde{V}^b \equiv 2\tilde{\nabla}^{(3)}_{[c}\tilde{\nabla}^{(3)}_{d]}\tilde{V}_a \equiv 2\tilde{h}^e_c\tilde{h}^f_d\tilde{h}^g_a\tilde{\nabla}_{[e}\left(\tilde{h}^h_f]\tilde{h}^i_g\tilde{\nabla}_h\tilde{V}_i\right), \quad \tilde{R}^{(3)}_{ab} \equiv \tilde{h}^{cd}\tilde{R}^{(3)}_{cadb}, \quad \tilde{R}^{(3)} \equiv \tilde{h}^{ab}\tilde{R}^{(3)}_{ab}.$$
(40)

From this we have

$$\tilde{R}^{(3)}_{abcd} = \tilde{h}^e_a \tilde{h}^f_b \tilde{h}^g_c \tilde{h}^h_d \tilde{R}_{efgh} - \tilde{\theta}_{ca} \tilde{\theta}_{db} + \tilde{\theta}_{bc} \tilde{\theta}_{ad}, \tag{41}$$

which is the Gauss equation. We can show that $\tilde{R}^{(3)}_{\alpha\beta\gamma\delta} = R^{(h)}_{\alpha\beta\gamma\delta}$. Equation (39) follows from eq. (30) evaluated in the normal-frame. Using eqs. (19,20) we can show that eq. (39) reproduces eq. (11) in the ADM formulation. Equation (38) gives eq. (8) in the ADM formulation.

Compared with the ADM equations in (8-13) part of the covariant equations in (29,32,33,34-37) look new. In the normal-frame eqs. (29,32) are identically satisfied; using eqs. (14,6,3) we can show

$$\tilde{a}_{\alpha} = (\ln N)_{,\alpha} \,, \tag{42}$$

thus $\tilde{a}_{[\beta;\gamma]} = 0$. Still, eqs. (8-13) provide a complete set. These additional equations in the covariant forms should be regarded as complementary equations which could possibly show certain aspects of the system better. In our perturbation analyses we will use parts of these equations as complement ones. Although the covariant set of equations is based on the general frame vector, this does not add any new physics which is not covered by the normal-frame taken in the ADM formulation, see §III E.

The covariant equations for the scalar fields, generalized gravity, electromagnetic field, null geodesic, and Boltzmann equation will be introduced individually in the corresponding sections later.

C. Multi-component situation

In the multi-component situation we have

$$\tilde{T}_{ab} = \sum_{l} \tilde{T}_{(l)ab}; \quad \tilde{T}_{(i)a;b}^{\ \ b} \equiv \tilde{I}_{(i)a}, \quad \sum_{l} \tilde{I}_{(l)a} = 0.$$
(43)

Based on the normal-frame vector, we have

$$\tilde{\mu} = \sum_{l} \tilde{\mu}_{(l)}, \quad \tilde{p} = \sum_{l} \tilde{p}_{(l)}, \quad \tilde{q}_a = \sum_{l} \tilde{q}_{(l)a}, \quad \tilde{\pi}_{ab} = \sum_{l} \tilde{\pi}_{(l)ab}.$$
(44)

The ADM formulation is based on the normal-frame vector $\tilde{u}_a = \tilde{n}_a$. The ADM fluid quantities in eq. (4) correspond to the fluid quantities based on the normal-frame vector as

$$E = \tilde{\mu}, \quad S = 3\tilde{p}, \quad J_{\alpha} = \tilde{q}_{\alpha}, \quad \bar{S}_{\alpha\beta} = \tilde{\pi}_{\alpha\beta}.$$
 (45)

From eq. (4) or eq. (45) we have

$$E = \sum_{l} E_{(l)}, \quad S = \sum_{l} S_{(l)}, \quad J_{\alpha} = \sum_{l} J_{(l)\alpha}, \quad S_{\alpha\beta} = \sum_{l} S_{(l)\alpha\beta}.$$
(46)

Equation (43) gives

$$E_{(i),0}N^{-1} - E_{(i),\alpha}N^{\alpha}N^{-1} - K\left(E_{(i)} + \frac{1}{3}S_{(i)}\right) - \bar{S}_{(i)}^{\alpha\beta}\bar{K}_{\alpha\beta} + N^{-2}\left(N^2J_{(i)}^{\alpha}\right)_{:\alpha} = -\frac{1}{N}\left(\tilde{I}_{(i)0} - \tilde{I}_{(i)\alpha}N^{\alpha}\right), \quad (47)$$

$$J_{(i)\alpha,0}N^{-1} - J_{(i)\alpha;\beta}N^{\beta}N^{-1} - J_{(i)\beta}N^{\beta}{}_{;\alpha}N^{-1} - KJ_{(i)\alpha} + E_{(i)}N_{,\alpha}N^{-1} + S^{\ \beta}_{(i)\alpha;\beta} + S^{\ \beta}_{(i)\alpha}N_{,\beta}N^{-1} = \tilde{I}_{(i)\alpha}.$$
(48)

The ADM equations in eqs. (8-13) remain valid, with the above additional equations of motion for the individual component. Thus, in the multi-component situation eqs. (8-13,46-48) provide a complete set.

III. PERTURBED QUANTITIES

A. Metric and connections

We use the following convention for the metric variables:

$$\tilde{g}_{00} \equiv -a^2 \left(1 + 2A\right), \quad \tilde{g}_{0\alpha} \equiv -a^2 B_\alpha, \quad \tilde{g}_{\alpha\beta} \equiv a^2 \left(g^{(3)}_{\alpha\beta} + 2C_{\alpha\beta}\right), \tag{49}$$

where A, B_{α} and $C_{\alpha\beta}$ are perturbed order variables and are *assumed* to be based on $g_{\alpha\beta}^{(3)}$ as the metric. To the second-order, we can write the perturbation variables explicitly as:

$$A \equiv A^{(1)} + A^{(2)}, \quad B_{\alpha} \equiv B_{\alpha}^{(1)} + B_{\alpha}^{(2)}, \quad C_{\alpha\beta} \equiv C_{\alpha\beta}^{(1)} + C_{\alpha\beta}^{(2)}.$$
 (50)

As we are interested in the perturbation to the second-order, as our ansatz, we include up to second-order (quadratic) terms in the deviation from the Friedmann background. This can be extended to any higher-order perturbation as long as we take the perturbative approach where the lower-order solutions drive (work as sources for) the next higher-order variables. Thus, in this work we ignore the terms higher than quadratic (second-order) combination of the perturbed metric $(A, B_{\alpha}, C_{\alpha\beta})$, the perturbed fluid quantities $(\delta\mu, \delta p, Q_{\alpha}, \Pi_{\alpha\beta})$ to be introduced in eq. (72), the perturbed field $(\delta\phi)$ to be introduced in eq. (111), etc.

The inverse metric expanded to the second-order in perturbation variables is:

$$\tilde{g}^{00} = \frac{1}{a^2} \left(-1 + 2A - 4A^2 + B_{\alpha} B^{\alpha} \right),$$

$$\tilde{g}^{0\alpha} = \frac{1}{a^2} \left(-B^{\alpha} + 2AB^{\alpha} + 2B_{\beta} C^{\alpha\beta} \right),$$

$$\tilde{g}^{\alpha\beta} = \frac{1}{a^2} \left(g^{(3)\alpha\beta} - 2C^{\alpha\beta} - B^{\alpha} B^{\beta} + 4C^{\alpha}_{\gamma} C^{\beta\gamma} \right).$$
(51)

The connections are:

$$\begin{split} \tilde{\Gamma}_{00}^{0} &= \frac{a'}{a} + A' - 2AA' - A_{,\alpha}B^{\alpha} + B_{\alpha} \left(B^{\alpha\prime} + \frac{a'}{a}B^{\alpha} \right), \\ \tilde{\Gamma}_{0\alpha}^{0} &= A_{,\alpha} - \frac{a'}{a}B_{\alpha} - 2AA_{,\alpha} + 2\frac{a'}{a}AB_{\alpha} - B_{\beta}C_{\alpha}^{\beta\prime} + B^{\beta}B_{[\beta|\alpha]}, \\ \tilde{\Gamma}_{00}^{0} &= A^{|\alpha} - B^{\alpha\prime} - \frac{a'}{a}B^{\alpha} + A'B^{\alpha} - 2A_{,\beta}C^{\alpha\beta} + 2C_{\beta}^{\alpha} \left(B^{\beta\prime} + \frac{a'}{a}B^{\beta} \right), \\ \tilde{\Gamma}_{\alpha\beta}^{0} &= \frac{a'}{a}g_{\alpha\beta}^{(3)} - 2\frac{a'}{a}g_{\alpha\beta}^{(3)}A + B_{(\alpha|\beta)} + C'_{\alpha\beta} + 2\frac{a'}{a}C_{\alpha\beta} \\ &\quad + \frac{a'}{a}g_{\alpha\beta}^{(3)} \left(4A^{2} - B_{\gamma}B^{\gamma} \right) - 2A \left(B_{(\alpha|\beta)} + C'_{\alpha\beta} + 2\frac{a'}{a}C_{\alpha\beta} \right) - B_{\gamma} \left(2C_{(\alpha|\beta)}^{\gamma} - C_{\alpha\beta}^{\ |\gamma} \right), \\ \tilde{\Gamma}_{0\beta}^{\alpha} &= \frac{a'}{a}\delta_{\beta}^{\alpha} + \frac{1}{2} \left(B_{\beta}^{\ |\alpha} - B^{\alpha}_{\ |\beta} \right) + C_{\beta}^{\alpha\prime} + B^{\alpha} \left(A_{,\beta} - \frac{a'}{a}B_{\beta} \right) + 2C^{\alpha\gamma} \left(B_{[\gamma|\beta]} - C'_{\gamma\beta} \right), \\ \tilde{\Gamma}_{\beta\gamma}^{\alpha} &= \Gamma_{\beta\gamma}^{(3)\alpha} + \frac{a'}{a}g_{\beta\gamma}^{(3)}B^{\alpha} + 2C_{(\beta|\gamma)}^{\alpha} - C_{\beta\gamma}^{\ |\alpha} - 2C_{\delta}^{\alpha} \left(2C_{(\beta|\gamma)}^{\delta} - C_{\beta\gamma}^{\ |\delta} \right) \\ - 2\frac{a'}{a}g_{\gamma\beta}^{(3)} \left(AB^{\alpha} + B^{\delta}C_{\delta}^{\alpha} \right) + B^{\alpha} \left(B_{(\beta|\gamma)} + C'_{\beta\gamma} + 2\frac{a'}{a}C_{\beta\gamma} \right), \end{split}$$

$$(52)$$

where a vertical bar indicates a covariant derivative based on $g_{\alpha\beta}^{(3)}$. An index 0 indicates the conformal time η , and a prime indicates a time derivative with respect to η . The components of the frame four-vector \tilde{u}_a are introduced as:

$$\tilde{u}^{0} \equiv \frac{1}{a} \left(1 - A + \frac{3}{2}A^{2} + \frac{1}{2}V^{\alpha}V_{\alpha} - V^{\alpha}B_{\alpha} \right), \quad \tilde{u}^{\alpha} \equiv \frac{1}{a}V^{\alpha},$$
$$\tilde{u}_{0} = -a \left(1 + A - \frac{1}{2}A^{2} + \frac{1}{2}V^{\alpha}V_{\alpha} \right), \quad \tilde{u}_{\alpha} = a \left(V_{\alpha} - B_{\alpha} + AB_{\alpha} + 2V^{\beta}C_{\alpha\beta} \right), \tag{53}$$

where V^{α} is based on $g^{(3)}_{\alpha\beta}$.

B. Normal-frame quantities

The normal-frame vector \tilde{n}_a has a property $\tilde{n}_{\alpha} \equiv 0$. Thus we have

$$\tilde{n}^{0} \equiv \frac{1}{a} \left(1 - A + \frac{3}{2} A^{2} - \frac{1}{2} B^{\alpha} B_{\alpha} \right), \quad \tilde{n}^{\alpha} \equiv \frac{1}{a} \left(B^{\alpha} - A B^{\alpha} - 2 B^{\beta} C^{\alpha}_{\beta} \right),$$
$$\tilde{n}_{0} = -a \left(1 + A - \frac{1}{2} A^{2} + \frac{1}{2} B^{\alpha} B_{\alpha} \right), \quad \tilde{n}_{\alpha} = 0.$$
(54)

Using eqs. (2,3) the ADM metric variables become:

$$N = a \left(1 + A - \frac{1}{2} A^2 + \frac{1}{2} B^{\alpha} B_{\alpha} \right), \quad N_{\alpha} = -a^2 B_{\alpha}, \quad N^{\alpha} = -B^{\alpha} + 2B^{\beta} C^{\alpha}_{\beta},$$
$$h_{\alpha\beta} = a^2 \left(g^{(3)}_{\alpha\beta} + 2C_{\alpha\beta} \right), \quad h^{\alpha\beta} = \frac{1}{a^2} \left(g^{(3)\alpha\beta} - 2C^{\alpha\beta} + 4C^{\alpha}_{\gamma} C^{\beta\gamma} \right). \tag{55}$$

The connection becomes

$$\Gamma^{(h)\gamma}_{\ \alpha\beta} = \Gamma^{(3)\gamma}_{\ \alpha\beta} + \left(g^{(3)\gamma\delta} - 2C^{\gamma\delta}\right) \left(C_{\delta\alpha|\beta} + C_{\delta\beta|\alpha} - C_{\alpha\beta|\delta}\right).$$
(56)

The extrinsic curvature in eq. (5) gives:

$$K_{\alpha\beta} = -a \left[\left(\frac{a'}{a} g_{\alpha\beta}^{(3)} + B_{(\alpha|\beta)} + C'_{\alpha\beta} + 2\frac{a'}{a} C_{\alpha\beta} \right) (1-A) + \frac{1}{2} \frac{a'}{a} g_{\alpha\beta}^{(3)} \left(3A^2 - B_{\gamma}B^{\gamma} \right) - B_{\gamma} \left(2C^{\gamma}_{(\alpha|\beta)} - C^{\beta}_{\alpha\beta} \right) \right],$$

$$K = -\frac{1}{a} \left[\left(3\frac{a'}{a} + B^{\alpha}_{\ |\alpha} + C^{\alpha'}_{\alpha} \right) (1-A) + \frac{3}{2} \frac{a'}{a} \left(3A^2 - B^{\alpha}B_{\alpha} \right) - B^{\beta} \left(2C^{\alpha}_{\beta|\alpha} - C^{\alpha}_{\alpha|\beta} \right) - 2C^{\alpha\beta} \left(C'_{\alpha\beta} + B_{\alpha|\beta} \right) \right],$$

$$\bar{K}_{\alpha\beta} = -a \Biggl\{ \left(B_{(\alpha|\beta)} + C_{\alpha\beta}' \right) (1-A) - B_{\gamma} \left(2C_{(\alpha|\beta)}^{\gamma} - C_{\alpha\beta}^{\ |\gamma} \right) - \frac{2}{3} C_{\alpha\beta} \left(B_{\ |\gamma}^{\gamma} + C_{\gamma}^{\gamma\prime} \right) - \frac{1}{3} g_{\alpha\beta}^{(3)} \left[\left(B_{\ |\gamma}^{\gamma} + C_{\gamma}^{\gamma\prime} \right) (1-A) - B^{\gamma} \left(2C_{\gamma|\delta}^{\delta} - C_{\delta|\gamma}^{\delta} \right) - 2C^{\gamma\delta} \left(C_{\gamma\delta}' + B_{\gamma|\delta} \right) \Biggr] \Biggr\}.$$

$$(57)$$

The intrinsic curvature in eq. (7) becomes:

$$R_{\alpha\beta}^{(h)} = R_{\alpha\beta}^{(3)} + \left(g^{(3)\gamma\delta} - 2C^{\gamma\delta}\right) \left(C_{\delta\alpha|\beta\gamma} + C_{\delta\beta|\alpha\gamma} - C_{\alpha\beta|\delta\gamma} - C_{\delta\gamma|\alpha\beta}\right) + 2C^{\gamma\delta}{}_{|\beta}C_{\gamma\delta|\alpha} \\ - \left(2C_{\delta|\gamma}^{\gamma} - C_{\gamma|\delta}^{\gamma}\right) \left(C_{\alpha|\beta}^{\delta} + C_{\beta|\alpha}^{\delta} - C_{\alpha\beta}{}^{|\delta}\right) - \left(C_{\alpha|\gamma}^{\delta} + C_{\gamma|\alpha}^{\delta} - C_{\alpha\gamma}{}^{|\delta}\right) \left(C_{\beta|\delta}^{\gamma} + C_{\delta|\beta}^{\gamma} - C_{\beta\delta}{}^{|\gamma}\right), \\ R^{(h)} = \frac{1}{a^{2}} \left[R^{(3)} - 2C^{\alpha\beta}R_{\alpha\beta}^{(3)} + 2C_{\alpha\beta}{}^{\beta|\alpha} - 2C_{\alpha\beta}{}^{\alpha|\beta} + 4C_{\gamma}^{\alpha}C^{\beta\gamma}R_{\alpha\beta}^{(3)} + 4C^{\alpha\beta}\left(-C_{\alpha|\beta\gamma}^{\gamma} - C_{\alpha|\gamma\beta}^{\gamma} + C_{\alpha\beta}{}^{|\gamma} + C_{\gamma|\alpha\beta}^{\gamma}\right) \\ - \left(2C_{\beta|\gamma}^{\gamma} - C_{\gamma|\beta}^{\gamma}\right) \left(2C_{\beta\alpha}{}^{\alpha\beta} - C_{\alpha}{}^{\alpha\beta}\right) + C^{\alpha\beta|\gamma}\left(3C_{\alpha\beta|\gamma} - 2C_{\alpha\gamma|\beta}\right)\right],$$
(58)

where

$$R^{(3)\alpha}{}_{\beta\gamma\delta} = \frac{1}{6} R^{(3)} \left(\delta^{\alpha}_{\gamma} g^{(3)}_{\beta\delta} - \delta^{\alpha}_{\delta} g^{(3)}_{\beta\gamma} \right), \quad R^{(3)}_{\alpha\beta} = \frac{1}{3} R^{(3)} g^{(3)}_{\alpha\beta}, \quad R^{(3)} = 6K, \tag{59}$$

with a normalized $K(=0,\pm 1)$, the sign of the background three-space curvature. Thus,

$$\bar{R}^{(h)\alpha}{}_{\beta} = \frac{1}{a^2} \Biggl\{ C^{\alpha\gamma}{}_{|\beta\gamma} + C^{\gamma|\alpha}_{\beta}{}_{\gamma} - C^{\alpha|\gamma}_{\beta}{}_{\gamma} - C^{\gamma|\alpha}_{\gamma}{}_{\beta} - \frac{2}{3} R^{(3)} C^{\alpha}_{\beta} \\
-2C^{\gamma\delta} \left(C^{\alpha}_{\delta|\beta\gamma} + C_{\delta\beta}{}^{|\alpha}{}_{\gamma} - C^{\alpha}_{\beta|\delta\gamma} - C_{\delta\gamma}{}^{|\alpha}{}_{\beta} \right) - 2C^{\alpha\gamma} \left(C^{\delta}_{\gamma|\beta\delta} + C^{\delta}_{\beta|\gamma\delta} - C_{\beta\gamma}{}^{|\delta}{}_{\delta} - C^{\delta}_{\delta|\gamma\beta} \right) + \frac{4}{3} R^{(3)} C^{\alpha}_{\gamma} C^{\gamma}_{\beta} \\
- \left(2C^{\gamma}_{\delta|\gamma} - C^{\gamma}_{\gamma|\delta} \right) \left(C^{\alpha\delta}{}_{|\beta} + C^{\delta|\alpha}_{\beta} - C^{\alpha|\delta}_{\beta} \right) + C_{\gamma\delta|\beta} C^{\gamma\delta|\alpha} + 2C^{\alpha\gamma|\delta} \left(C_{\beta\gamma|\delta} - C_{\beta\delta|\gamma} \right) \\
- \frac{1}{3} \delta^{\alpha}_{\beta} \Biggl[- \frac{2}{3} R^{(3)} C^{\gamma}_{\gamma} + 2C^{\delta|\gamma}_{\gamma} - 2C^{\gamma|\delta}_{\gamma} + 4C^{\gamma\delta} \left(-C^{\epsilon}_{\gamma|\delta\epsilon} - C^{\epsilon}_{\gamma|\epsilon\delta} + C_{\gamma\delta}{}^{|\epsilon}_{\epsilon} + C^{\epsilon}_{\epsilon|\gamma\delta} \right) \\
+ \frac{4}{3} R^{(3)} C^{\delta}_{\gamma} C^{\gamma}_{\delta} - \left(2C^{\epsilon}_{\delta|\epsilon} - C^{\epsilon}_{\epsilon|\delta} \right) \left(2C^{\gamma\delta}_{\ |\gamma} - C^{\gamma|\delta}_{\gamma} \right) + C^{\gamma\delta|\epsilon} \left(3C_{\gamma\delta|\epsilon} - 2C_{\gamma\epsilon|\delta} \right) \Biggr] \Biggr\}.$$
(60)

It is convenient to have

$$B^{\alpha}{}_{|\beta\gamma} = B^{\alpha}{}_{|\gamma\beta} - R^{(3)\alpha}{}_{\delta\beta\gamma}B^{\delta}, \quad B_{\alpha|\beta\gamma} = B_{\alpha|\gamma\beta} + R^{(3)\delta}{}_{\alpha\beta\gamma}B_{\delta}.$$
(61)

C. General (\tilde{u}_a) -frame quantities

To the second-order perturbation, using eqs. (49,51,53), the kinematic quantities in eq. (14) become:

$$\tilde{h}_{\alpha\beta} = a^{2} \left[g_{\alpha\beta}^{(3)} + 2C_{\alpha\beta} + (V_{\alpha} - B_{\alpha})(V_{\beta} - B_{\beta}) \right], \quad \tilde{h}_{0\alpha} = -a^{2} \left(V_{\alpha} + AV_{\alpha} + 2C_{\alpha\beta}V^{\beta} \right), \quad \tilde{h}_{00} = a^{2}V^{\alpha}V_{\alpha}, \tag{62}$$

$$\tilde{\theta} = a^{-1} \left[3\frac{a'}{a}(1-A) + V^{\alpha}{}_{|\alpha} + C^{\alpha'}_{\alpha} + \frac{9}{2}\frac{a'}{a}A^{2} + B^{\alpha}B'_{\alpha} - 2C^{\alpha\beta}C'_{\alpha\beta} - AC^{\alpha'}_{\alpha} - V^{\alpha'}B_{\alpha} + V^{\alpha} \left(V_{\alpha}' + \frac{3}{2}\frac{a'}{a}V_{\alpha} - B'_{\alpha} - 3\frac{a'}{a}B_{\alpha} + A_{,\alpha} + C^{\beta}_{\beta|\alpha} \right) \right], \tag{63}$$

$$\tilde{a}_{\alpha} = A_{,\alpha} + V_{\alpha}' - B'_{\alpha} + \frac{a'}{a} \left(V_{\alpha} - B_{\alpha} \right) + A'B_{\alpha} + A \left(-2A_{,\alpha} + 2B'_{\alpha} + 2\frac{a'}{a}B_{\alpha} - V'_{\alpha} - \frac{a'}{a}V_{\alpha} \right)$$

$$+V^{\beta}\left(V_{\alpha|\beta}+B_{\beta|\alpha}-B_{\alpha|\beta}\right)+2C_{\alpha\beta}\left(V^{\beta\prime}+\frac{a'}{a}V^{\beta}\right)+2C_{\alpha\beta}'V^{\beta},\quad\tilde{a}_{0}=-V^{\alpha}\tilde{a}_{\alpha},\tag{64}$$

$$\tilde{\omega}_{\alpha\beta} = a \Big(V_{[\alpha|\beta]} - B_{[\alpha|\beta]} + AB_{[\alpha|\beta]} - V_{[\alpha}A_{,\beta]} + 2B_{[\alpha}A_{,\beta]} - B_{[\alpha}B'_{\beta]} \\ - V_{[\alpha}V'_{\beta]} + B_{[\alpha}V'_{\beta]} + V_{[\alpha}B'_{\beta]} + 2V_{\gamma}C^{\gamma}_{[\alpha|\beta]} + 2C_{\gamma[\alpha}V^{\gamma}{}_{|\beta]} \Big), \quad \tilde{\omega}_{0\alpha} = V^{\beta}\tilde{\omega}_{\alpha\beta}, \quad \tilde{\omega}_{00} = 0,$$

$$\tilde{\sigma}_{\alpha\beta} = a \Big[V_{(\alpha|\beta)} + C'_{\alpha\beta} - \frac{1}{2}a^{(3)}_{\alpha\beta} \left(V^{\gamma}{}_{|\alpha|} + C^{\gamma'}_{\alpha} \right) + V_{(\alpha}V'_{\alpha}) - V_{(\alpha}B'_{\alpha)} - V'_{\alpha}B_{\alpha} + B_{(\alpha}B'_{\alpha)} \Big]$$
(65)

$$= \left[V^{(\alpha|\beta)} + C_{\alpha\beta} - \frac{3}{3} g_{\alpha\beta} \left(V^{\gamma}|\gamma + C_{\gamma} \right) + V^{\alpha} A_{\beta} \right] - V^{\alpha} B_{\beta} - V^{\alpha} B_{\beta} + D^{\alpha} B_{\beta} + D^{\alpha} B_{\beta} + D^{\alpha} B_{\beta} + V^{\alpha} A_{\beta} + V_{\alpha} B_{\beta} + V^{\alpha} A_{\beta} + C_{\alpha} B_{\beta} + D^{\alpha} B_{\beta} + D$$

In the normal-frame we have $\tilde{u}_{\alpha} \equiv 0$, thus $V_{\alpha} = B_{\alpha} - AB_{\alpha} - 2B^{\beta}C_{\alpha\beta}$. In this frame we have

$$\tilde{h}_{\alpha\beta} = a^2 \left(g^{(3)}_{\alpha\beta} + 2C_{\alpha\beta} \right), \quad \tilde{h}_{0\alpha} = -a^2 B_\alpha, \quad \tilde{h}_{00} = a^2 B^\alpha B_\alpha, \tag{67}$$

$$\tilde{\theta} = -K,\tag{68}$$

$$\tilde{a}_{\alpha} = (\ln N)_{,\alpha} = \left(A - A^2 + \frac{1}{2}B^{\beta}B_{\beta}\right)_{,\alpha}, \quad \tilde{a}_0 = -B^{\alpha}A_{,\alpha},$$
(69)

$$\tilde{\sigma}_{\alpha\beta} = -\bar{K}_{\alpha\beta}, \quad \tilde{\sigma}_{0\alpha} = B^{\beta}\bar{K}_{\alpha\beta}, \quad \tilde{\sigma}_{00} = 0, \tag{70}$$

$$\tilde{\omega}_{ab} = 0. \tag{71}$$

In this frame we have $\tilde{\theta} = -K$ and $\tilde{\sigma}_{\alpha\beta} = -\bar{K}_{\alpha\beta}$. These are natural because K and $\bar{K}_{\alpha\beta}$ are the same as negatives of the expansion scalar and the shear, respectively, of the normal-frame vector field.

D. Fluid quantities

To the perturbed order we decompose the fluid quantities as:

$$\widetilde{\mu} \equiv \mu + \delta \mu, \quad \widetilde{p} \equiv p + \delta p, \quad \widetilde{q}_{\alpha} \equiv a Q_{\alpha}, \quad \widetilde{\pi}_{\alpha\beta} \equiv a^2 \Pi_{\alpha\beta},$$
(72)

where Q_{α} and $\Pi_{\alpha\beta}$ are based on $g_{\alpha\beta}^{(3)}$. In the Friedmann world model we have $\tilde{\mu} = \mu$ and $\tilde{p} = p$ and zeros for the other fluid quantities. We have

$$\Pi^{\alpha}_{\alpha} - 2C^{\alpha\beta}\Pi_{\alpha\beta} = 0, \tag{73}$$

which follows from $\tilde{\pi}_a^a = 0$ or $\bar{S}_{\alpha}^{\alpha} = 0$. The perturbed order fluid quantities $\delta \mu$, δp , Q_{α} and $\Pi_{\alpha\beta}$ in eq. (72) can be expanded similarly as in equation (50)

$$\delta\mu = \delta\mu^{(1)} + \delta\mu^{(2)}, \quad \delta p = \delta p^{(1)} + \delta p^{(2)}, \quad Q_{\alpha} = Q_{\alpha}^{(1)} + Q_{\alpha}^{(2)}, \quad \Pi_{\alpha\beta} = \Pi_{\alpha\beta}^{(1)} + \Pi_{\alpha\beta}^{(2)}.$$
(74)

In the multi-component situation, from eqs. (44,72) we set

$$\mu = \sum_{l} \mu_{(l)}, \quad p = \sum_{l} p_{(l)},$$

$$\delta \mu = \sum_{l} \delta \mu_{(l)}, \quad \delta p = \sum_{l} \delta p_{(l)}, \quad Q_{\alpha} = \sum_{l} Q_{(l)\alpha}, \quad \Pi_{\alpha\beta} = \sum_{l} \Pi_{(l)\alpha\beta}.$$
(75)

Thus, from eq. (45) the ADM fluid variables become:

$$E \equiv \mu + \delta \mu,$$

$$S \equiv 3(p + \delta p),$$

$$J_{\alpha} \equiv aQ_{\alpha}, \quad J^{\alpha} = \frac{1}{a} \left(Q^{\alpha} - 2C^{\alpha\beta} Q_{\beta} \right),$$

$$\bar{S}_{\alpha\beta} \equiv a^{2} \Pi_{\alpha\beta}, \quad \bar{S}_{\beta}^{\alpha} = \Pi_{\beta}^{\alpha} - 2C^{\alpha\gamma} \Pi_{\beta\gamma}, \quad \bar{S}^{\alpha\beta} = \frac{1}{a^{2}} \left(\Pi^{\alpha\beta} - 4C^{\gamma(\alpha} \Pi_{\gamma}^{\beta)} \right).$$
(76)

From eqs. (22,72) we have

$$\tilde{q}_{\alpha} \equiv aQ_{\alpha}, \quad \tilde{q}_{0} = -aQ_{\alpha}B^{\alpha}, \\ \tilde{\pi}_{\alpha\beta} \equiv a^{2}\Pi_{\alpha\beta}, \quad \tilde{\pi}_{0\alpha} = -a^{2}\Pi_{\alpha\beta}B^{\beta}, \quad \tilde{\pi}_{00} = 0.$$
(77)

From eqs. (21,72,54) we have

$$\tilde{T}_{00} = a^{2} \left[\mu + \delta \mu + 2\mu A + 2\delta \mu A + (\mu + p) B^{\alpha} B_{\alpha} + 2Q_{\alpha} B^{\alpha} \right],$$

$$\tilde{T}_{0\alpha} = -a^{2} \left(Q_{\alpha} + pB_{\alpha} + \delta pB_{\alpha} + AQ_{\alpha} + \Pi_{\alpha\beta} B^{\beta} \right),$$

$$\tilde{T}_{\alpha\beta} = a^{2} \left(pg_{\alpha\beta}^{(3)} + \delta pg_{\alpha\beta}^{(3)} + \Pi_{\alpha\beta} + 2pC_{\alpha\beta} + 2\delta pC_{\alpha\beta} \right).$$
(78)

From eqs. (4,76) we have

$$\mu + \delta\mu = \tilde{T}_{ab}\tilde{n}^{a}\tilde{n}^{b}, \quad p + \delta p = \frac{1}{3}h^{\alpha\beta}\tilde{T}_{\alpha\beta}, \quad Q_{\alpha} = -\frac{1}{a}\tilde{T}_{\alpha b}\tilde{n}^{b}, \quad \Pi_{\alpha\beta} = \frac{1}{a^{2}}\left(\tilde{T}_{\alpha\beta} - h_{\alpha\beta}\tilde{p}\right). \tag{79}$$

Equation (24) gives

$$\tilde{\varrho}\tilde{T}\tilde{\tilde{S}} = \tilde{\pi}K + \frac{1}{a^2}\Pi^{\alpha\beta}\bar{K}_{\alpha\beta} - 2\frac{1}{a}Q^{\alpha}A_{,\alpha} - \frac{1}{a}Q^{\alpha}{}_{|\alpha} + 2\frac{1}{a}\left(C^{\alpha\beta}Q_{\beta}\right)_{|\alpha} - \frac{1}{a}C^{\alpha}{}_{\alpha|\beta}Q^{\beta}.$$
(80)

For the interaction terms in eq. (43) we set

$$\tilde{I}_{(i)0} \equiv I_{(i)0} + \delta I_{(i)0}, \quad \tilde{I}_{(i)\alpha} \equiv \delta I_{(i)\alpha}, \tag{81}$$

where $\delta I_{(i)\alpha}$ is based in $g_{\alpha\beta}^{(3)}$.

E. Frame choice

The energy-momentum tensor in the general (\tilde{u}_a) frame follows from eqs. (21,72,53):

$$\tilde{T}^{0}_{0} = -\mu - \delta\mu - (\mu + p) V^{\alpha} (V_{\alpha} - B_{\alpha}) - Q^{\alpha} (2V_{\alpha} - B_{\alpha}),$$

$$\tilde{T}^{0}_{\alpha} = (1 - A) [Q_{\alpha} + (\mu + p) (V_{\alpha} - B_{\alpha})] + (\mu + p) (AB_{\alpha} + 2V^{\beta}C_{\alpha\beta}) + (\delta\mu + \delta p) (V_{\alpha} - B_{\alpha}) + (V^{\beta} - B^{\beta}) \Pi_{\alpha\beta},$$

$$\tilde{T}^{\alpha}_{\beta} = (p + \delta p) \delta^{\alpha}_{\beta} + \Pi^{\alpha}_{\beta} + V^{\alpha} [Q_{\beta} + (\mu + p) (V_{\beta} - B_{\beta})] + Q^{\alpha} (V_{\beta} - B_{\beta}) - 2C^{\alpha\gamma}\Pi_{\beta\gamma}.$$
(82)

In the energy-frame we set $Q_{\alpha} \equiv 0$, thus $\tilde{q}_a = 0$. In the normal-frame we have $\tilde{u}_{\alpha} \equiv 0$, thus from eq. (53) we have

Energy – frame :
$$Q_{\alpha} \equiv 0$$
,
Normal – frame : $V_{\alpha} - B_{\alpha} + AB_{\alpha} + 2B^{\beta}C_{\alpha\beta} \equiv 0.$ (83)

Although we can take infinitely many different combination of the two frames, the energy- and the normal-frames are the ones often used in the literature [22]. By choosing a frame (which is a decision about Q_{α} and V_{α}) we lose no generality. This is because we have 10 independent informations in \tilde{T}_{ab} which can be allocated to the energy density $\tilde{\mu}$ (one), the pressure \tilde{p} (one), the anisotropic stress $\Pi_{\alpha\beta}$ (five, because it is tracefree). The remaining (three) informations can be assigned to either the velocity V_{α} (three) or the flux Q_{α} (three); or some combinations of V_{α} and Q_{α} with total three informations.

Thus, in the normal-frame (indicated by a superscript N) we have

$$\tilde{T}_{0}^{0} = -\mu - \delta\mu^{N} - Q^{N\alpha}B_{\alpha},
\tilde{T}_{\alpha}^{0} = (1 - A)Q_{\alpha}^{N},
\tilde{T}_{\beta}^{\alpha} = \left(p + \delta p^{N}\right)\delta_{\beta}^{\alpha} + \Pi^{N\alpha}{}_{\beta} + B^{\alpha}Q_{\beta}^{N} - 2C^{\alpha\gamma}\Pi_{\beta\gamma}^{N},$$
(84)

which is consistent with eq. (78). To the linear-order we notice that $\delta\mu$, δp , $\Pi_{\alpha\beta}$ are independent of the frame choice, and $Q_{\alpha} + (\mu + p)(V_{\alpha} - B_{\alpha})$ is a frame-invariant combination [21]. However, to the second-order we no longer have such a luxury. As the fluid quantities are defined based on the frame vector as in eq. (23) the values of $\delta\mu$, δp and $\Pi_{\alpha\beta}$ are dependent on the frame.

By comparing the energy-momentum tensor in the normal-frame in eq. (84) with the one in the general frame in eq. (82), we find that by replacing the normal-frame fluid quantities to

$$\begin{split} \delta\mu^{N} &\equiv \delta\mu + \left(V^{\alpha} - B^{\alpha}\right) \left[\left(\mu + p\right)\left(V_{\alpha} - B_{\alpha}\right) + 2Q_{\alpha}\right], \\ \deltap^{N} &\equiv \delta p + \frac{1}{3} \left(V^{\alpha} - B^{\alpha}\right) \left[\left(\mu + p\right)\left(V_{\alpha} - B_{\alpha}\right) + 2Q_{\alpha}\right], \\ Q^{N}_{\alpha} &\equiv Q_{\alpha} + \left(\mu + p\right)\left(V_{\alpha} - B_{\alpha}\right) + \left(\mu + p\right)\left(AB_{\alpha} + 2V^{\beta}C_{\alpha\beta}\right) + \left(\delta\mu + \delta p\right)\left(V_{\alpha} - B_{\alpha}\right) + \left(V^{\beta} - B^{\beta}\right)\Pi_{\alpha\beta}, \\ \Pi^{N}_{\alpha\beta} &\equiv \Pi_{\alpha\beta} + \left(\mu + p\right)\left(V_{\alpha} - B_{\alpha}\right)\left(V_{\beta} - B_{\beta}\right) + 2Q_{(\alpha}\left(V_{\beta)} - B_{\beta}\right)\right) - \frac{1}{3}g^{(3)}_{\alpha\beta}\left(V^{\gamma} - B^{\gamma}\right)\left[\left(\mu + p\right)\left(V_{\gamma} - B_{\gamma}\right) + 2Q_{\gamma}\right], \end{split}$$
(85)

we recover the general frame energy-momentum tensor. Thus, by imposing the energy-frame condition $(Q_{\alpha} = 0)$ we recover the fluid quantities in the energy-frame.

Now, similarly, by comparing eq. (82) with the same one evaluated in the energy-frame, we find that by replacing the energy-frame fluid quantities (indicated by a superscript E) to

$$\delta\mu^{E} \equiv \delta\mu - \frac{1}{\mu + p}Q^{\alpha}Q_{\alpha},$$

$$\delta p^{E} \equiv \delta p - \frac{1}{3}\frac{1}{\mu + p}Q^{\alpha}Q_{\alpha},$$

$$(\mu + p)\left(V_{\alpha}^{E} - B_{\alpha}\right) \equiv (\mu + p)\left(V_{\alpha} - B_{\alpha}\right) + Q_{\alpha} - 2Q^{\beta}C_{\alpha\beta} - \frac{\delta\mu + \delta p}{\mu + p}Q_{\alpha} - Q^{\beta}\frac{\Pi_{\alpha\beta}}{\mu + p},$$

$$\Pi_{\alpha\beta}^{E} \equiv \Pi_{\alpha\beta} - \frac{1}{\mu + p}\left(Q_{\alpha}Q_{\beta} - \frac{1}{3}g_{\alpha\beta}^{(3)}Q^{\gamma}Q_{\gamma}\right),$$
(86)

we recover the general frame energy-momentum tensor. By imposing the normal-frame condition in eq. (83) we recover the fluid quantities in the normal-frame. Thus, using eqs. (85,86) we can transform the fluid quantities in one frame to the other:

$$\delta\mu^{N} = \delta\mu^{E} + (\mu + p) \left(V^{E\alpha} - B^{\alpha} \right) \left(V^{E}_{\alpha} - B_{\alpha} \right),$$

$$\deltap^{N} = \deltap^{E} + \frac{1}{3} \left(\mu + p \right) \left(V^{E\alpha} - B^{\alpha} \right) \left(V^{E}_{\alpha} - B_{\alpha} \right),$$

$$Q^{N}_{\alpha} = (\mu + p) \left(V^{E}_{\alpha} - B_{\alpha} \right) + (\mu + p) \left(AB_{\alpha} + 2V^{E\beta}C_{\alpha\beta} \right) + \left(\delta\mu^{E} + \delta p^{E} \right) \left(V^{E}_{\alpha} - B_{\alpha} \right) + \left(V^{E\beta} - B^{\beta} \right) \Pi^{E}_{\alpha\beta},$$

$$\Pi^{N}_{\alpha\beta} = \Pi^{E}_{\alpha\beta} + (\mu + p) \left(V^{E}_{\alpha} - B_{\alpha} \right) \left(V^{E}_{\beta} - B_{\beta} \right) - \frac{1}{3}g^{(3)}_{\alpha\beta} \left(\mu + p \right) \left(V^{E\gamma} - B^{\gamma} \right) \left(V^{E}_{\gamma} - B_{\gamma} \right);$$

$$\delta\mu^{E} - \delta\mu^{N} - \frac{1}{2} O^{N\alpha}O^{N}$$

(87)

$$\delta\mu^{-} = \delta\mu^{N} - \frac{1}{\mu + p}Q^{N\alpha}Q_{\alpha}^{N},$$

$$\delta p^{E} = \delta p^{N} - \frac{1}{3}\frac{1}{\mu + p}Q^{N\alpha}Q_{\alpha}^{N},$$

$$(\mu + p)\left(V_{\alpha}^{E} - B_{\alpha}\right) = Q_{\alpha}^{N} - 2Q^{N\beta}C_{\alpha\beta} - \frac{\delta\mu^{N} + \delta p^{N}}{\mu + p}Q_{\alpha}^{N} - Q^{N\beta}\frac{\Pi_{\alpha\beta}^{N}}{\mu + p} - (\mu + p)\left(AB_{\alpha} + 2B^{\beta}C_{\alpha\beta}\right),$$

$$\Pi_{\alpha\beta}^{E} = \Pi_{\alpha\beta}^{N} - \frac{1}{\mu + p}\left(Q_{\alpha}^{N}Q_{\beta}^{N} - \frac{1}{3}g_{\alpha\beta}^{(3)}Q^{N\gamma}Q_{\gamma}^{N}\right).$$
(88)

F. Spacetime curvatures to the linear-order

Although straightforward, it is not an easy task to derive the spacetime curvature to the second-order. For our purpose, fortunately, it is not necessary to have the forms. Still, it is convenient to have the curvatures to the

linear-order and we present them in the following. These follow from eqs. (49,51,52). Curvature tensors are:

$$\begin{split} \tilde{R}^{a}{}_{b00} = 0, \quad \tilde{R}^{b}{}_{00\alpha} = -\left(\frac{a'}{a}\right)' B_{\alpha}, \quad \tilde{R}^{b}{}_{0\alpha\beta} = 0, \\ \tilde{R}^{b}{}_{\alpha\alpha\beta} = \left(\frac{a'}{a}\right)' g_{\alpha\beta}^{\alpha\beta} - \left[\frac{a'}{a}A' + 2\left(\frac{a'}{a}\right)'A\right] g_{\alpha\beta}^{\alpha\beta} - A_{\alpha|\beta} + B'_{(\alpha|\beta)} + \frac{a'}{a} B_{(\alpha|\beta)} + C''_{\alpha\beta} + \frac{a'}{a} C'_{\alpha\beta} + 2\left(\frac{a'}{a}\right)' C_{\alpha\beta}, \\ \tilde{R}^{b}{}_{\alpha\beta\gamma} = 2\frac{a'}{a} g_{\alpha|\beta}^{\alpha\beta} A_{\gamma|} - B_{\alpha|\beta\gamma|} + \frac{1}{2} (B_{\gamma|\alpha\beta} - B_{\beta|\alpha\gamma}) - 2C'_{\alpha|\beta|\gamma|}, \\ \tilde{R}^{a}{}_{00\beta} = \left(\frac{a'}{a}\right)' \delta^{\alpha}{}_{\beta} - \frac{a'}{a} A' \delta^{\alpha}{}_{\beta} - A^{|\alpha}{}_{\beta} + \frac{1}{2} \left(B_{\beta}^{|\alpha} + B^{\alpha}{}_{|\beta}\right)' + \frac{1}{2}\frac{a'}{a} \left(B_{\beta}^{|\alpha} + B^{\alpha}{}_{|\beta}\right) + C^{\alpha''}_{\beta''} + \frac{a'}{a} C^{\alpha'}_{\beta'}, \\ \tilde{R}^{a}{}_{0\beta\gamma} = \frac{a'}{a} \delta^{\alpha}{}_{|\beta} A_{\gamma|} - B_{|\beta}^{|\alpha}{}_{\gamma|} + B^{\alpha}{}_{|\beta\gamma|} - 2\left(\frac{a'}{a}\right)^{2} \delta^{\alpha}{}_{|\beta} B_{\gamma|} - 2C^{\alpha'}_{|\beta\gamma|}, \\ \tilde{R}^{\alpha}{}_{\beta0\gamma} = \frac{a'}{a} \left(g_{\beta\gamma}^{\alpha} A^{,\alpha} - \delta^{\alpha}_{\gamma} A_{,\beta}\right) + \left(\frac{a'}{a}\right)' g_{\beta\gamma}^{(3)} B^{\alpha} - \left(\frac{a'}{a}\right)^{2} \left(g_{\beta\gamma}^{(3)} B^{\alpha} - \delta^{\alpha}_{\gamma} B_{\beta}\right) - \frac{1}{2} \left(B_{\beta}^{|\alpha} - B^{\alpha}{}_{|\beta}\right)_{|\gamma} + C^{\alpha''}_{\gamma|\beta} - C'_{\beta\gamma}^{|\alpha}, \\ \tilde{R}^{\alpha}{}_{\beta\gamma\gamma} = R^{(3)\alpha}{}_{\beta\gamma\delta} + \left(\frac{a'}{a}\right)^{2} \left(\delta^{\alpha}_{\gamma} g_{\beta\beta}^{(3)} - \delta^{\beta}_{\delta} g_{\beta\gamma}^{(3)}\right) (1 - 2A) \\ & + \frac{1}{2}\frac{a'}{a} \left[g_{\beta\delta}^{(3)} \left(B_{\gamma}^{|\alpha} + B^{\alpha}{}_{|\gamma}\right) - g_{\beta\gamma}^{(3)} \left(B_{\delta}^{|\alpha} + B^{\alpha}{}_{|\delta}\right) + 2\delta^{\alpha}_{\gamma} B_{|\beta|\delta} - 2\delta^{\alpha}_{\delta} B_{|\beta|\gamma}\right] \\ + 2C^{\alpha}_{|\beta|\beta\gamma} - 2C^{\alpha}_{(\beta|\gamma)} + \delta^{\alpha}_{\gamma} C'_{\beta\delta} - \delta^{\alpha}_{\delta} C'_{\beta\gamma} + 2\frac{a'}{a} \left(\delta^{\alpha}_{\gamma} C_{\beta\delta} - \delta^{\alpha}_{\alpha} C_{\beta\gamma}\right) \right] \\ & + 2C^{\alpha}_{|\beta|\beta\gamma} - 2C^{\alpha}_{|\beta|\gamma} + \delta^{\alpha}_{|\alpha} A' + \Delta A - B^{\alpha}_{|\alpha} - \frac{a'}{a} B^{\alpha}_{|\alpha} - C^{\alpha''}_{\alpha} - \frac{a'}{a} C^{\beta'}_{\alpha}, \\ \bar{R}_{0\alpha} = 2\frac{a'}{a} A_{,\alpha} - \left(\frac{a'}{a}\right)' B_{\alpha} - 2\left(\frac{a'}{a}\right)^{2} B_{\alpha} + \frac{1}{2} \Delta B_{\alpha} - \frac{1}{2} B^{\beta}_{|\alpha\beta} - C^{\beta'}_{|\alpha\beta} - C^{\beta'}_{|\beta\beta} + C^{\beta'}_{|\beta\beta} + C^{\alpha'}_{|\beta\beta}, \\ \bar{R}_{\alpha\beta} = 2Kg_{\alpha\beta}^{\alpha\beta} + \left[\left(\frac{a'}{a}\right)' + 2\left(\frac{a'}{a}\right)^{2}\right\right] g_{\alpha\beta}^{\alpha\beta} (1 - 2A) - \frac{a'}{a} A'^{\alpha}_{\alpha\beta}^{\alpha\beta} - A_{\alpha\beta}^{\beta} - A_{\alpha\beta}^{\beta} + C^{\alpha'}_{|\beta\beta} + C^{\alpha'}_{|\beta\beta} + \frac{a'}{a} B_{|\alpha|\beta\beta} + 2\frac{a'}{a} B_{|\alpha|\beta\beta} + 2\frac{a'}{a} B_{|\alpha|\beta\beta} + 2\frac{a'}{a} B_{|\alpha|\beta\beta} + \frac{a'}{a} B_{|\alpha|\beta\beta} + 2\frac{a'}{a} B_{|\alpha|\beta\beta} + \frac{a'}{a} B_{|\alpha|\beta\beta}$$

The nonvanishing components of the electric and magnetic parts of the Weyl curvature in eq. (20) are:

$$\begin{split} \tilde{E}_{\alpha\beta} &= -\tilde{C}^{0}_{\ \alpha0\beta} \\ &= \frac{1}{2}A_{,\alpha|\beta} - \frac{1}{2}B'_{(\alpha|\beta)} - \frac{1}{2}C''_{\alpha\beta} - \frac{1}{2}\Delta C_{\alpha\beta} - 2KC_{\alpha\beta} + C^{\gamma}_{(\alpha|\beta)\gamma} - \frac{1}{2}C^{\gamma}_{\gamma|\alpha\beta} \\ &\quad -\frac{1}{3}g^{(3)}_{\alpha\beta} \left(\frac{1}{2}\Delta A - \frac{1}{2}B^{\gamma'}_{\ |\gamma} - \frac{1}{2}C^{\gamma''}_{\gamma} - \Delta C^{\gamma}_{\gamma} - 2KC^{\gamma}_{\gamma} + C^{\gamma\delta}_{\ |\gamma\delta}\right) \\ &= \frac{1}{2}\left(\nabla_{\alpha}\nabla_{\beta} - \frac{1}{3}g^{(3)}_{\alpha\beta}\Delta\right) \left(\alpha - \varphi - \frac{1}{a}\chi' + \frac{a'}{a^{2}}\chi\right) - \frac{1}{2}\Psi^{(\nu)}_{(\alpha|\beta)} - \frac{1}{2}\left[C^{(t)\prime\prime}_{\alpha\beta} + (\Delta - 2K)C^{(t)}_{\alpha\beta}\right], \end{split}$$
(92)
$$\tilde{H}_{\alpha\beta} &= -\frac{1}{2}\tilde{\eta}_{0(\alpha}{}^{\gamma\delta}\tilde{C}^{0}_{\ \beta)\gamma\delta} \end{split}$$

$$= -\eta_{(\alpha}^{\gamma\delta} \left(\frac{1}{2} B_{\gamma|\beta)\delta} + C'_{\beta)\gamma|\delta}\right) = -\eta_{(\alpha}^{\gamma\delta} \left(\frac{1}{2} \Psi^{(v)}_{\gamma|\beta)\delta} + C^{(t)\prime}_{\beta)\gamma|\delta}\right),$$
(93)

where the symmetrization is only over α and β indices. The last steps are evaluated in decomposed forms which will be introduced in §V. We introduced $\eta^{\alpha\beta\gamma}$ which is based on $g^{(3)}_{\alpha\beta}$ with $\eta^{\alpha\beta\gamma} \equiv \eta^{[\alpha\beta\gamma]}$ and $\eta^{123} \equiv 1/\sqrt{g^{(3)}}$. We have

$$\tilde{\eta}^{0\alpha\beta\gamma} = \frac{1}{a^4\sqrt{1+D}}\eta^{\alpha\beta\gamma},$$

$$\tilde{g} = -a^8(1+D)g^{(3)}, \quad D \equiv 2A + 2C^{\alpha}_{\alpha} + 4AC^{\alpha}_{\alpha} + B_{\alpha}B^{\alpha} + 2C^{\alpha}_{\alpha}C^{\beta}_{\beta} - 2C^{\alpha}_{\beta}C^{\beta}_{\alpha},$$
(94)

which is valid to the second-order in perturbation.

Deriving the electric and magnetic parts of Weyl tensor to the second-order using eqs. (19,20) requires quite lengthy algebra. Instead, using eqs. (30,33) we can derive them easily. Evaluated in the normal-frame we have

$$\begin{split} \tilde{E}_{\alpha\beta} &= \frac{1}{a} \left(1 - A \right) \left(\bar{K}_{\alpha\beta}' - 2\frac{a'}{a} \bar{K}_{\alpha\beta} \right) - \frac{1}{a} \left(B_{(\alpha}^{\ |\gamma} - B^{\gamma}_{\ |(\alpha} + 2C_{(\alpha)}^{\gamma\prime}) \bar{K}_{\beta)\gamma} + \frac{1}{a} \bar{K}_{\alpha\beta|\gamma} B^{\gamma} - \frac{1}{a^2} g^{(3)\gamma\delta} \bar{K}_{\alpha\gamma} \bar{K}_{\beta\delta} - \frac{2}{3} K \bar{K}_{\alpha\beta} \right. \\ &+ \left(A - A^2 + \frac{1}{2} B^{\gamma} B_{\gamma} \right)_{,\alpha|\beta} - \left(2C_{(\alpha|\beta)}^{\gamma} - C_{\alpha\beta}^{\ |\gamma} \right) A_{,\gamma} + A_{,\alpha} A_{,\beta} - \frac{2}{3} C_{\alpha\beta} \Delta A + 4\pi G a^2 \Pi_{\alpha\beta} \\ &- \frac{1}{3} g_{\alpha\beta}^{(3)} \left[-\frac{1}{a^2} g^{(3)\alpha\gamma} g^{(3)\beta\delta} \bar{K}_{\alpha\beta} \bar{K}_{\gamma\delta} + A_{,\gamma} \left(A^{,\gamma} - 2C^{\gamma\delta}_{\ |\delta} + C^{\delta|\gamma}_{\delta} \right) + \Delta \left(A - A^2 + \frac{1}{2} B^{\gamma} B_{\gamma} \right) - 2C^{\gamma\delta} A_{,\gamma|\delta} \right], \end{split}$$

$$\tag{95}$$

$$\tilde{H}_{\alpha\beta} = \left\{ \left[g_{\delta(\beta)}^{(3)} \left(1 - C_{\nu}^{\nu} \right) + 2C_{\delta(\beta)} \right] \bar{K}_{\alpha)\gamma|\mu} - \left(C_{(\alpha|\mu}^{\nu} + C_{\mu|\alpha}^{\nu} - C_{\mu(\alpha)}^{\ |\nu|} \right) g_{\beta)\delta}^{(3)} \bar{K}_{\gamma\nu} \right\} \frac{1}{a} \eta^{\delta\gamma\mu}, \tag{96}$$

where K and $\bar{K}_{\alpha\beta}$ are given in eq. (57); the symmetrization is only over α and β indices. We have $\tilde{E}_{ab}\tilde{u}^b = 0 = \tilde{H}_{ab}\tilde{u}^b$, thus $\tilde{E}_{\alpha0} = -\tilde{E}_{\alpha\beta}B^{\beta}$, $\tilde{E}_{00} = 0$, and similarly for \tilde{H}_{ab} .

For later use, it is convenient to have the spacetime scalar curvature expanded to the second-order. In terms of the ADM notation, using eq. (6), we have

$$\tilde{R} = R^{(h)} + K^{\alpha\beta}K_{\alpha\beta} + K^2 + \frac{2}{N}\left(-K_{,0} + K_{,\alpha}N^{\alpha} - N^{:\alpha}_{\ \alpha}\right).$$
(97)

To the second-order in perturbation, using quantities in §III B we have

$$\begin{split} \tilde{R} &= R + \delta R \\ &= 6 \left(\frac{K}{a^2} + \dot{H} + 2H^2 \right) \\ &- 6H\dot{A} - 12 \left(\dot{H} + 2H^2 \right) A - 2 \frac{\Delta}{a^2} A + 2 \left(\frac{1}{a} B^{\alpha}{}_{|\alpha} + \dot{C}^{\alpha}_{\alpha} \right) + 8H \left(\frac{1}{a} B^{\alpha}{}_{|\alpha} + \dot{C}^{\alpha}_{\alpha} \right) + 2 \frac{1}{a^2} \left[C^{\beta|\alpha}{}_{\alpha\beta} - (\Delta + 2K) C^{\alpha}_{\alpha} \right] \\ &+ 24HA\dot{A} - 4A \left(\frac{1}{a} B^{\alpha}{}_{|\alpha} + \dot{C}^{\alpha}_{\alpha} \right) - 2 \left(\dot{A} + 8HA \right) \left(\frac{1}{a} B^{\alpha}{}_{|\alpha} + \dot{C}^{\alpha}_{\alpha} \right) + 24 \left(\dot{H} + 2H^2 \right) A^2 + 4A \frac{\Delta}{a^2} A + 2 \frac{1}{a^2} A^{\alpha} A_{,\alpha} \\ &- 6H \frac{1}{a} A_{,\alpha} B^{\alpha} + 4 \frac{1}{a^2} A_{,\alpha|\beta} C^{\alpha\beta} + 2 \frac{1}{a^2} A_{,\beta} \left(2C^{\alpha\beta}{}_{|\alpha} - C^{\alpha|\beta}_{\alpha} \right) + \frac{1}{a^2} \left[B_{(\alpha|\beta)} B^{\alpha|\beta} + B^{\alpha}{}_{|\alpha} B^{\beta}{}_{|\beta} - 2 \left(B_{\alpha} B^{\alpha|\beta} \right)_{|\beta} \right] \\ &- 6H B^{\alpha} \dot{B}_{\alpha} - 6 \left(\dot{H} + 2H^2 \right) B^{\alpha} B_{\alpha} - 2 \frac{1}{a} B^{\alpha} \left(2C^{\beta}{}_{\alpha|\beta} - C^{\beta}{}_{\beta|\alpha} \right) + 2 \frac{1}{a} B^{\alpha}{}_{|\alpha} \dot{C}^{\beta}_{\beta} \\ &- 2 \frac{1}{a} \left(2C^{\beta}_{\alpha|\beta} - C^{\beta}_{\beta|\alpha} \right) \left(\dot{B}^{\alpha} + 3HB^{\alpha} \right) - 4 \frac{1}{a} C^{\alpha\beta} \dot{B}_{\alpha|\beta} - 2 \frac{1}{a} \left(\dot{C}^{\alpha\beta} + 6HC^{\alpha\beta} \right) B_{\alpha|\beta} + 2 \frac{1}{a} B^{\alpha} \left(\frac{1}{a} B^{\beta}{}_{|\beta} + \dot{C}^{\beta}_{\beta} \right)_{|\alpha} \\ &+ \dot{C}^{\alpha}_{\alpha} \dot{C}^{\beta}_{\beta} - 3\dot{C}^{\alpha\beta} \dot{C}_{\alpha\beta} - 4C^{\alpha\beta} \left(\ddot{C}_{\alpha\beta} + 4H\dot{C}_{\alpha\beta} - 2\frac{K}{a^2} C_{\alpha\beta} \right) + 4 \frac{1}{a^2} C^{\alpha\beta} \left(-C^{\gamma}_{\alpha|\beta\gamma} - C^{\gamma}_{\alpha|\gamma\beta} + \Delta C_{\alpha\beta} + C^{\gamma}_{\gamma|\alpha\beta} \right) \\ &- \frac{1}{a^2} \left(2C^{\gamma}_{\beta|\gamma} - C^{\gamma}_{\gamma|\beta} \right) \left(2C^{\alpha\beta}{}_{|\alpha} - C^{\alpha|\beta}_{\alpha} \right) + \frac{1}{a^2} C^{\alpha\beta|\gamma} \left(3C_{\alpha\beta|\gamma} - 2C_{\alpha\gamma|\beta} \right).$$
(98)

An overdot indicates a time derivative with respect to t, with $dt = ad\eta$.

IV. PERTURBED EQUATIONS

A. Basic equations with general fluids

In the following we present complete sets of equations valid up to second-order in the perturbation without fixing the gauge conditions. As the basic set we consider eqs. (5,8-13,47,48) in the ADM formulation. Definition of δK :

$$\bar{K} + 3H
+\delta K - 3HA + \dot{C}^{\alpha}_{\alpha} + \frac{1}{a}B^{\alpha}_{\ |\alpha}
= -A\left(\frac{9}{2}HA - \dot{C}^{\alpha}_{\alpha} - \frac{1}{a}B^{\alpha}_{\ |\alpha}\right) + \frac{3}{2}HB^{\alpha}B_{\alpha} + \frac{1}{a}B^{\alpha}\left(2C^{\beta}_{\alpha|\beta} - C^{\beta}_{\beta|\alpha}\right) + 2C^{\alpha\beta}\left(\dot{C}_{\alpha\beta} + \frac{1}{a}B_{\alpha|\beta}\right)
\equiv N_{0},$$
(99)

where $K \equiv \overline{K} + \delta K$ and K is read from eq. (57). Energy constraint equation:

$$16\pi G\mu + 2\Lambda - 6H^{2} - \frac{1}{a^{2}}R^{(3)}$$

$$+16\pi G\delta\mu + 4H\delta K - \frac{1}{a^{2}}\left(2C_{\alpha}^{\beta|\alpha}{}_{\beta} - 2C_{\alpha}^{\alpha|\beta}{}_{\beta} - \frac{2}{3}R^{(3)}C_{\alpha}^{\alpha}\right)$$

$$= \frac{2}{3}\delta K^{2} - \left(\dot{C}_{\alpha\beta} + \frac{1}{a}B_{(\alpha|\beta)}\right)\left(\dot{C}^{\alpha\beta} + \frac{1}{a}B^{\alpha|\beta}\right) + \frac{1}{3}\left(\dot{C}_{\alpha}^{\alpha} + \frac{1}{a}B^{\alpha}{}_{|\alpha}\right)^{2}$$

$$+ \frac{1}{a^{2}}\left[4C^{\alpha\beta}\left(-C_{\alpha|\beta\gamma}^{\gamma} - C_{\alpha|\gamma\beta}^{\gamma} + C_{\alpha\beta}^{\gamma}{}_{\gamma} + C_{\gamma|\alpha\beta}^{\gamma}\right) + \frac{4}{3}R^{(3)}C_{\gamma}^{\alpha}C_{\alpha}^{\gamma}$$

$$- \left(2C_{\beta|\gamma}^{\gamma} - C_{\gamma|\beta}^{\gamma}\right)\left(2C^{\alpha\beta}{}_{|\alpha} - C_{\alpha}^{\alpha|\beta}\right) + C^{\alpha\beta|\gamma}\left(3C_{\alpha\beta|\gamma} - 2C_{\alpha\gamma|\beta}\right)\right]$$

$$\equiv N_{1}.$$
(100)

Momentum constraint equation:

$$\begin{split} \left[\dot{C}_{\alpha}^{\beta} + \frac{1}{2a} \left(B_{|\alpha}^{\beta} + B_{\alpha}^{|\beta}\right)\right]_{|\beta} &- \frac{1}{3} \left(\dot{C}_{\gamma}^{\gamma} + \frac{1}{a} B_{|\gamma}^{\gamma}\right)_{,\alpha} + \frac{2}{3} \delta K_{,\alpha} + 8\pi Ga Q_{\alpha} \\ &= A \left(-\frac{2}{3} \delta K_{,\alpha} - 8\pi Ga Q_{\alpha}\right) + A_{,\beta} \left[\dot{C}_{\alpha}^{\beta} + \frac{1}{2a} \left(B_{|\alpha}^{\beta} + B_{\alpha}^{|\beta}\right)\right] \\ &+ \left(2C_{\gamma|\beta}^{\beta} - C_{\beta|\gamma}^{\beta}\right) \left[\dot{C}_{\alpha}^{\gamma} + \frac{1}{2a} \left(B_{\alpha}^{|\gamma} + B_{|\alpha}^{\gamma}\right)\right] + 2C^{\beta\gamma} \left(\dot{C}_{\alpha\gamma} + \frac{1}{a} B_{(\alpha|\gamma)}\right)_{|\beta} \\ &+ \frac{1}{a} \left[B_{\gamma} \left(C^{\beta\gamma}_{|\alpha} + C_{\alpha}^{\gamma|\beta} - C_{\alpha}^{\beta|\gamma}\right)\right]_{|\beta} + \frac{1}{3}C_{\beta|\alpha}^{\gamma} \left(\dot{C}_{\gamma}^{\beta} + \frac{1}{a} B_{|\gamma}^{\beta}\right) \\ &- \frac{1}{3} \left\{A_{,\alpha} \left(\dot{C}_{\gamma}^{\gamma} + \frac{1}{a} B_{|\gamma}^{\gamma}\right) + 2C^{\gamma\delta} \left(\dot{C}_{\gamma\delta} + \frac{1}{a} B_{\gamma|\delta}\right)_{|\alpha} + \frac{1}{a} \left[B^{\delta} \left(2C_{\delta|\gamma}^{\gamma} - C_{\gamma|\delta}^{\gamma}\right)\right]_{|\alpha}\right\} \\ &\equiv N_{2\alpha}. \end{split}$$

$$\tag{101}$$

Trace of the ADM propagation equation:

$$-\left[3\dot{H} + 3H^2 + 4\pi G\left(\mu + 3p\right) - \Lambda\right]$$
$$+\delta\dot{K} + 2H\delta K - 4\pi G\left(\delta\mu + 3\delta p\right) + \frac{1}{a^2}A^{|\alpha}{}_{\alpha} + 3\dot{H}A$$
$$= A\delta\dot{K} - \frac{1}{a}\delta K_{,\alpha}B^{\alpha} + \frac{1}{3}\delta K^2 + \frac{3}{2}\dot{H}\left(3A^2 - B^{\alpha}B_{\alpha}\right)$$

$$+\frac{1}{a^{2}}\left[2AA^{|\alpha}{}_{\alpha}+A_{,\alpha}A^{,\alpha}-B^{\beta}B_{\beta}{}^{|\alpha}{}_{\alpha}-B^{\beta|\alpha}B_{\beta|\alpha}+A^{,\alpha}\left(2C^{\beta}_{\alpha|\beta}-C^{\beta}_{\beta|\alpha}\right)+2C^{\alpha\beta}A_{,\alpha|\beta}\right]$$
$$+\left(\dot{C}_{\alpha\beta}+\frac{1}{a}B_{(\alpha|\beta)}\right)\left(\dot{C}^{\alpha\beta}+\frac{1}{a}B^{\alpha|\beta}\right)-\frac{1}{3}\left(\dot{C}^{\alpha}_{\alpha}+\frac{1}{a}B^{\alpha}{}_{|\alpha}\right)^{2}$$
$$\equiv N_{3}.$$
(102)

Tracefree ADM propagation equation:

$$\begin{split} \left[\dot{C}_{\beta}^{\alpha} + \frac{1}{2a} \left(B^{\alpha}{}_{|\beta} + B_{\beta}{}^{|\alpha}\right)\right]^{-} + 3H \left[\dot{C}_{\gamma}^{\alpha} + \frac{1}{2a} \left(B^{\alpha}{}_{|\beta} + B_{\beta}{}^{|\alpha}\right)\right] - \frac{1}{a^{2}} A^{l\alpha}{}_{\beta} \\ & - \frac{1}{3} \delta_{\beta}^{\alpha} \left[\left(\dot{C}_{\gamma}^{\gamma} + \frac{1}{a} B^{\gamma}{}_{|\gamma}\right)^{-} + 3H \left(\dot{C}_{\gamma}^{\gamma} + \frac{1}{a} B^{\gamma}{}_{|\gamma}\right) - \frac{1}{a^{2}} A^{l\gamma}{}_{\gamma}\right] \\ & + \frac{1}{a^{2}} \left[C^{\alpha\gamma}{}_{|\beta\gamma} + C_{\gamma}^{\beta|\alpha}{}_{\gamma} - C_{\beta}^{\alpha|\gamma}{}_{\gamma} - C_{\gamma}^{\alpha|\alpha}{}_{\beta} - \frac{2}{3} R^{(3)} C_{\beta}^{\alpha} - \frac{1}{3} \delta_{\beta}^{\beta} \left(2C_{\gamma}^{\beta|\gamma}{}_{\delta} - 2C_{\gamma}^{\gamma|\delta}{}_{\delta} - \frac{2}{3} R^{(3)} C_{\gamma}^{\gamma}\right)\right] - 8\pi G \Pi_{\beta}^{\alpha} \\ & = \left\{ \left[\dot{C}_{\beta}^{\alpha} + \frac{1}{2a} \left(B^{\alpha}{}_{|\beta} + B_{\beta}{}^{|\alpha}\right)\right] A + 2C^{\alpha\gamma} \left(\dot{C}_{\beta\gamma} + \frac{1}{a} B_{(\beta|\gamma)}\right) + \frac{1}{a} B_{\gamma} \left(C^{\alpha\gamma}{}_{|\beta} + C_{\beta}^{\beta|\alpha} - C_{\beta}^{\alpha|\gamma}\right)\right)^{\prime} \\ & + 3H \left\{ \left[\dot{C}_{\beta}^{\alpha} + \frac{1}{2a} \left(B^{\alpha}{}_{|\beta} + B_{\beta}{}^{|\alpha}\right)\right]^{-} A - \frac{1}{a} \left[\dot{C}_{\beta}^{\alpha} + \frac{1}{2a} \left(B^{\alpha}{}_{|\beta} + B_{\beta}{}^{|\alpha}\right)\right]_{|\gamma} B^{\gamma} + \delta K \left[\dot{C}_{\beta}^{\alpha} + \frac{1}{2a} \left(B^{\alpha}{}_{|\beta} + B_{\beta}{}^{|\alpha}\right)\right] \\ & + \left[\dot{C}_{\beta}^{\alpha} + \frac{1}{2a} \left(B^{\alpha}{}_{|\beta} + B_{\beta}{}^{|\alpha}\right)\right]^{-} A - \frac{1}{a} \left[\dot{C}_{\gamma}^{\alpha} + \frac{1}{2a} \left(B^{\alpha}{}_{|\beta} + B_{\beta}{}^{|\alpha}\right)\right]_{|\gamma} B^{\gamma} + \delta K \left[\dot{C}_{\beta}^{\alpha} + \frac{1}{2a} \left(B^{\alpha}{}_{|\beta} + B_{\beta}{}^{|\alpha}\right)\right] \\ & + \frac{1}{a^{2}} \left[-AA^{|\alpha}{}_{\beta} + \frac{1}{2} \left(-A^{2} + B^{\gamma} B_{\gamma}\right) A + 2C^{\gamma\delta} \left(\dot{C}_{\gamma\delta} + \frac{1}{a} B_{\gamma|\delta}\right) + \frac{1}{a} B^{\delta} \left(2C_{\delta|\gamma}^{\gamma} - C_{\gamma|\delta}^{\gamma}\right)\right]^{-} \\ & + 3H \left[\left(\dot{C}_{\gamma}^{\gamma} + \frac{1}{a} B^{\gamma}{}_{|\gamma}\right) A + 2C^{\gamma\delta} \left(\dot{C}_{\gamma\delta} + \frac{1}{a} B_{\gamma|\delta}\right) + \frac{1}{a} B^{\delta} \left(2C_{\delta|\gamma}^{\gamma} - C_{\gamma|\delta}^{\gamma}\right)\right] \\ & + \left(\dot{C}_{\gamma}^{\gamma} + \frac{1}{a} B^{\gamma}{}_{|\gamma}\right) A - 2C^{\gamma\delta} \left(\dot{C}_{\gamma\delta} + \frac{1}{a} B_{\gamma|\delta}\right) + \frac{1}{a} B^{\delta} \left(2C_{\delta|\gamma}^{\gamma} - C_{\gamma|\delta}^{\gamma}\right)\right] \\ & + \left(\dot{C}_{\gamma}^{\gamma} + \frac{1}{a} B^{\gamma}{}_{|\gamma}\right) A - 2C^{\gamma\delta} \left(\dot{C}_{\gamma\delta} + \frac{1}{a} B_{\gamma|\delta}\right) + \frac{1}{a} B^{\delta} \left(2C_{\delta|\gamma}^{\gamma} - C_{\gamma|\delta}^{\gamma}\right)\right] \\ & + \frac{1}{a^{2}} \left[-AA^{|\gamma} + \frac{1}{2} \left(-A^{2} + B^{\delta} B_{\delta}\right)^{|\gamma} - 2C^{\gamma\delta} A_{\gamma|\delta}\right) + \frac{1}{a} B^{\delta} \left(2C_{\delta|\gamma}^{\gamma} - C_{\gamma|\delta}^{\gamma}\right)\right] \\ & + \frac{1}{a^{2}} \left[-AA^{|\gamma} + \frac{1}{2} \left(-A^{2} + B^{\delta} B_{\delta}\right)^{|\gamma} - 2C^{\gamma\delta} A_{\gamma|\delta} - \left(2C^{\gamma\delta} + \frac{1}{a} B^{\beta} + C_{\delta|\gamma}^{\gamma}\right) - \frac{1}{a} B^{\gamma} \left(2C^{\gamma$$

Energy conservation equation:

$$\begin{split} \dot{\mu} + 3H\left(\mu + p\right) \\ + \delta\dot{\mu} + 3H\left(\delta\mu + \delta p\right) - \left(\mu + p\right)\left(\delta K - 3HA\right) + \frac{1}{a}Q^{\alpha}{}_{|\alpha} \end{split}$$

$$= -\frac{1}{a}\delta\mu_{,\alpha}B^{\alpha} + (\delta\mu + \delta p)\left(\delta K - 3HA\right) + (\mu + p)\left[A\delta K + \frac{3}{2}H\left(A^{2} - B^{\alpha}B_{\alpha}\right)\right]$$
$$-\frac{1}{a}\left[AQ^{\alpha}{}_{|\alpha} + Q^{\alpha}\left(2A_{,\alpha} + C^{\beta}_{\beta|\alpha} - 2C^{\beta}_{\alpha|\beta}\right) - 2C^{\alpha\beta}Q_{\alpha|\beta}\right] - \Pi^{\alpha\beta}\left(\dot{C}_{\alpha\beta} + \frac{1}{a}B_{\alpha|\beta}\right)$$
$$\equiv N_{5}.$$
(104)

Momentum conservation equation:

$$\begin{split} \dot{Q}_{\alpha} + 4HQ_{\alpha} + \frac{1}{a} \left[(\mu + p) A_{,\alpha} + \delta p_{,\alpha} + \Pi^{\beta}_{\alpha|\beta} \right] \\ &= Q_{\alpha} \left(\delta K - 3HA \right) + \frac{1}{a} \Biggl\{ -Q_{\alpha|\beta} B^{\beta} - Q_{\beta} B^{\beta}_{\ |\alpha} - (\delta \mu + \delta p) A_{,\alpha} + A \left[(\mu + p) A_{,\alpha} - \delta p_{,\alpha} - \Pi^{\beta}_{\alpha|\beta} \right] \\ &- (\mu + p) B^{\beta} B_{\beta|\alpha} + 2 \left(C^{\gamma\beta} \Pi_{\alpha\gamma} \right)_{|\beta} - \Pi^{\gamma}_{\alpha} C^{\beta}_{\beta|\gamma} + \Pi^{\beta}_{\gamma} C^{\gamma}_{\beta|\alpha} - A_{,\beta} \Pi^{\beta}_{\alpha} \Biggr\} \\ &\equiv N_{6\alpha}. \end{split}$$
(105)

In the multi-component situation we additionally have the energy and the momentum conservation of individual component in eqs. (47,48).

Energy conservation equation for the i-th component:

$$\begin{split} \dot{\mu}_{(i)} + 3H\left(\mu_{(i)} + p_{(i)}\right) + \frac{1}{a}I_{(i)0} \\ + \delta\dot{\mu}_{(i)} + 3H\left(\delta\mu_{(i)} + \delta p_{(i)}\right) - \left(\mu_{(i)} + p_{(i)}\right)\left(\delta K - 3HA\right) + \frac{1}{a}Q_{(i)|\alpha}^{\alpha} + \frac{1}{a}\delta I_{(i)0} \\ &= -\frac{1}{a}\delta\mu_{(i),\alpha}B^{\alpha} + \left(\delta\mu_{(i)} + \delta p_{(i)}\right)\left(\delta K - 3HA\right) + \left(\mu_{(i)} + p_{(i)}\right)A\delta K + \frac{3}{2}H\left(\mu_{(i)} + p_{(i)}\right)\left(A^{2} - B^{\alpha}B_{\alpha}\right) \\ &+ \frac{1}{a}\left[-Q_{(i)|\alpha}^{\alpha}A + 2\left(C^{\alpha\beta}Q_{(i)\beta}\right)_{|\alpha} - C_{\alpha}^{\alpha|\beta}Q_{(i)\beta} - 2A_{,\alpha}Q_{(i)}^{\alpha}\right] - \Pi_{(i)}^{\alpha\beta}\left(\frac{1}{a}B_{\alpha|\beta} + \dot{C}_{\alpha\beta}\right) - \frac{1}{a}\delta I_{(i)\alpha}B^{\alpha} \\ &\equiv N_{(i)5}. \end{split}$$
(106)

Momentum conservation equation for the i-th component:

$$\dot{Q}_{(i)\alpha} + 4HQ_{(i)\alpha} + \frac{1}{a} \left(\mu_{(i)} + p_{(i)} \right) A_{,\alpha} + \frac{1}{a} \left(\delta p_{(i),\alpha} + \Pi_{(i)\alpha|\beta}^{\ \beta} - \delta I_{(i)\alpha} \right)$$

$$= \frac{1}{a} \Biggl\{ - \left(\delta p_{(i),\alpha} + \Pi_{(i)\alpha|\beta}^{\ \beta} - \delta I_{(i)\alpha} \right) A - \left(\delta \mu_{(i)} + \delta p_{(i)} \right) A_{,\alpha} + \left(\mu_{(i)} + p_{(i)} \right) \left(AA_{,\alpha} - B^{\beta} B_{\beta|\alpha} \right)$$

$$- Q_{(i)\alpha|\beta} B^{\beta} - Q_{(i)\beta} B^{\beta}_{\ |\alpha} + a \left(\delta K - 3HA \right) Q_{(i)\alpha} + 2 \left(C^{\beta\gamma} \Pi_{(i)\alpha\gamma} \right)_{|\beta} - C^{\beta}_{\beta|\gamma} \Pi_{(i)\alpha}^{\ \gamma} + C^{\gamma}_{\beta|\alpha} \Pi_{(i)\gamma}^{\ \beta} - A_{,\beta} \Pi_{(i)\alpha}^{\ \beta} \Biggr\}$$

$$\equiv N_{(i)6\alpha}.$$

$$(107)$$

The collective fluid quantities are given in eq. (75). The equations are presented with the quadratic combination of the linear order terms located in the right-hand-side. Still, notice that the equations are presented to the second-order without separating the background order part.

Equations (99-107) provide a complete set valid for the Einstein's gravity with an imperfect fluid, thus the most general form of energy-momentum tensor. We have not imposed any condition like the gauge condition. In the following subsections we will consider the cases of minimally coupled scalar fields, an electromagnetic field, and a broad class of generalized gravity theories. We emphasize that even in these additional fields or generalized gravity the above equations *remain valid* with the fluid quantities reinterpreted to absorb the contributions from the fields and the generalized gravity.

B. Scalar field

1. Covariant equations

The action for a minimally coupled scalar field is given as

$$S = \int \sqrt{-\tilde{g}} \left[\frac{1}{16\pi G} \tilde{R} - \frac{1}{2} \tilde{\phi}^{,c} \tilde{\phi}_{,c} - \tilde{V}(\tilde{\phi}) \right] d^4x.$$

$$\tag{108}$$

The equation of motion follows from the variation in $\tilde{\phi}$

$$\tilde{\phi}^{;c}{}_{c} - \tilde{V}_{,\tilde{\phi}} = 0, \tag{109}$$

where $\tilde{V}_{,\tilde{\phi}} \equiv \frac{\partial \tilde{V}}{\partial \tilde{\phi}}$. From $\delta_{\tilde{g}_{ab}} \mathcal{L}_M \equiv \frac{1}{2} \sqrt{-\tilde{g}} \tilde{T}^{ab} \delta \tilde{g}_{ab}$ we have the energy-momentum tensor

$$\tilde{T}_{ab}^{(\phi)} = \tilde{\phi}_{,a}\tilde{\phi}_{,b} - \frac{1}{2}\tilde{g}_{ab}\tilde{\phi}^{,c}\tilde{\phi}_{,c} - \tilde{g}_{ab}\tilde{V}(\tilde{\phi}).$$
(110)

2. Perturbations

We decompose

$$\tilde{\phi} \equiv \phi + \delta\phi. \tag{111}$$

The equation of motion becomes

$$\begin{split} \ddot{\phi} + 3H\dot{\phi} + V_{,\phi} \\ + \delta\ddot{\phi} + 3H\delta\dot{\phi} - \frac{\Delta}{a^{2}}\delta\phi + V_{,\phi\phi}\delta\phi - 2A\ddot{\phi} - \dot{\phi}\left(\dot{A} + 6HA - \frac{1}{a}B^{\alpha}{}_{|\alpha} - \dot{C}^{\alpha}_{\alpha}\right) \\ &= 2A\left[\delta\ddot{\phi} + 3H\delta\dot{\phi} - 2A\ddot{\phi} - \dot{\phi}\left(2\dot{A} + 6HA - \frac{1}{a}B^{\alpha}{}_{|\alpha} - \dot{C}^{\alpha}_{\alpha}\right)\right] + \delta\dot{\phi}\left(\dot{A} - \frac{1}{a}B^{\alpha}{}_{|\alpha} - \dot{C}^{\alpha}_{\alpha}\right) - 2\frac{1}{a}B^{\alpha}\delta\dot{\phi}_{,\alpha} \\ &\quad + \frac{1}{a}\delta\phi_{,\alpha}\left(\frac{1}{a}A^{,\alpha} - \dot{B}^{\alpha} - 2HB^{\alpha} - 2\frac{1}{a}C^{\alpha\beta}{}_{|\beta} + \frac{1}{a}C^{\beta}{}_{\beta}{}^{|\alpha}\right) - 2\frac{1}{a^{2}}\delta\phi_{,\alpha|\beta}C^{\alpha\beta} - \frac{1}{2}V_{,\phi\phi\phi}\delta\phi^{2} \\ &\quad + \ddot{\phi}B_{\alpha}B^{\alpha} + \dot{\phi}\left[\frac{1}{a}A_{,\alpha}B^{\alpha} + B_{\alpha}\dot{B}^{\alpha} + 3HB^{\alpha}B_{\alpha} + \frac{1}{a}B_{\beta}\left(2C^{\alpha\beta}{}_{|\alpha} - C^{\alpha|\beta}_{\alpha}\right) + 2C^{\alpha\beta}\left(\frac{1}{a}B_{\alpha|\beta} + \dot{C}_{\alpha\beta}\right)\right] \\ &\equiv N_{\phi}. \end{split}$$

$$\tag{112}$$

The energy-momentum tensor gives:

$$\begin{split} \tilde{T}_{00}^{(\phi)} &= \frac{1}{2} \phi'^2 + a^2 V + \phi' \delta \phi' + a^2 \left(V_{,\phi} \delta \phi + 2VA \right) \\ &\quad + \frac{1}{2} \delta \phi'^2 + \frac{1}{2} \delta \phi_{,\alpha} \delta \phi^{,\alpha} + \frac{1}{2} a^2 V_{,\phi\phi} \delta \phi^2 + 2a^2 V_{,\phi} \delta \phi A - \phi' \delta \phi_{,\alpha} B^{\alpha} + \frac{1}{2} \phi'^2 B_{\alpha} B^{\alpha}, \\ \tilde{T}_{0\alpha}^{(\phi)} &= \phi' \delta \phi_{,\alpha} - \left(\frac{1}{2} \phi'^2 - a^2 V \right) B_{\alpha} + \delta \phi' \delta \phi_{,\alpha} + \left(-\phi' \delta \phi' + a^2 V_{,\phi} \delta \phi + \phi'^2 A \right) B_{\alpha}, \\ \tilde{T}_{\alpha\beta}^{(\phi)} &= g_{\alpha\beta}^{(3)} \left(\frac{1}{2} \phi'^2 - a^2 V \right) + g_{\alpha\beta}^{(3)} \left(\phi' \delta \phi' - a^2 V_{,\phi} \delta \phi - \phi'^2 A \right) + \left(\phi'^2 - 2a^2 V \right) C_{\alpha\beta} \\ &\quad + \delta \phi_{,\alpha} \delta \phi_{,\beta} + 2 \left[\phi' \delta \phi' - a^2 V_{,\phi} \delta \phi - \phi'^2 A \right] C_{\alpha\beta} \\ &\quad - \frac{1}{2} g_{\alpha\beta}^{(3)} \left[-\delta \phi'^2 + \delta \phi_{,\gamma} \delta \phi^{,\gamma} + a^2 V_{,\phi\phi} \delta \phi^2 + 4 \phi' \delta \phi' A - 2 \phi' \delta \phi_{,\gamma} B^{\gamma} + \phi'^2 \left(-4A^2 + B_{\gamma} B^{\gamma} \right) \right]. \end{split}$$
(113)

Fluid quantities can be read from eq. (79) as:

$$\begin{split} \mu^{(\phi)} + \delta\mu^{(\phi)} &= \frac{1}{2}\dot{\phi}^2 + V + \dot{\phi}\dot{\delta\phi} - \dot{\phi}^2A + V_{,\phi}\delta\phi \\ &\quad + \frac{1}{2}\delta\dot{\phi}^2 + \frac{1}{2a^2}\delta\phi_{,\alpha}\delta\phi^{,\alpha} + \frac{1}{2}V_{,\phi\phi}\delta\phi^2 - 2\dot{\phi}\dot{\delta\phi}A + \frac{1}{a}\dot{\phi}\delta\phi_{,\alpha}B^{\alpha} + 2\dot{\phi}^2A^2 - \frac{1}{2}\dot{\phi}^2B^{\alpha}B_{\alpha}, \\ p^{(\phi)} + \delta p^{(\phi)} &= \frac{1}{2}\dot{\phi}^2 - V + \dot{\phi}\dot{\delta\phi} - \dot{\phi}^2A - V_{,\phi}\delta\phi \\ &\quad + \frac{1}{2}\delta\dot{\phi}^2 - \frac{1}{6a^2}\delta\phi_{,\alpha}\delta\phi^{,\alpha} - \frac{1}{2}V_{,\phi\phi}\delta\phi^2 - 2\dot{\phi}\dot{\delta\phi}A + \frac{1}{a}\dot{\phi}\delta\phi_{,\alpha}B^{\alpha} + 2\dot{\phi}^2A^2 - \frac{1}{2}\dot{\phi}^2B^{\alpha}B_{\alpha}, \end{split}$$

$$Q_{\alpha}^{(\phi)} = -\frac{1}{a} \left[\dot{\phi} \delta\phi_{,\alpha} + \delta\phi_{,\alpha} \left(\delta\dot{\phi} - \dot{\phi}A \right) \right],$$

$$\Pi_{\alpha\beta}^{(\phi)} = \frac{1}{a^2} \left(\delta\phi_{,\alpha} \delta\phi_{,\beta} - \frac{1}{3} g_{\alpha\beta}^{(3)} \delta\phi_{,\gamma} \delta\phi^{,\gamma} \right).$$
(114)

We indicate the quadratic parts (the quadratic combinations of two linear-order terms) as $\delta\mu^{(q)}$, $\delta p^{(q)}$, $Q^{(q)}_{\alpha}$, and $\Pi^{(q)}_{\alpha\beta}$.

C. Scalar fields

1. Covariant equations

The action for multiple components of minimally coupled scalar fields is

$$S = \int \sqrt{-\tilde{g}} \left[\frac{1}{16\pi G} \tilde{R} - \frac{1}{2} \sum_{k} \tilde{\phi}_{(k)}{}^{c} \tilde{\phi}_{(k),c} - \tilde{V}(\tilde{\phi}_{(l)}) \right] d^{4}x, \tag{115}$$

where $i, j, \ldots = 1, 2 \ldots n$ indicate the *n* scalar fields. The equation of motion for *i*-th component is

$$\tilde{\phi}_{(i)\ c}^{\ ;c} - \tilde{V}_{,\tilde{\phi}_{(i)}} = 0.$$
(116)

The energy-momentum tensor is

$$\tilde{T}_{ab}^{(\phi)} = \sum_{k} \left(\tilde{\phi}_{(k),a} \tilde{\phi}_{(k),b} - \frac{1}{2} \tilde{g}_{ab} \tilde{\phi}_{(k)}^{\ c} \tilde{\phi}_{(k),c} \right) - \tilde{g}_{ab} \tilde{V}(\tilde{\phi}_{(l)}).$$
(117)

2. Perturbations

We introduce

$$\tilde{\phi}_{(i)} \equiv \phi_{(i)} + \delta \phi_{(i)}. \tag{118}$$

The equation of motion for i-th component becomes

$$\begin{split} \ddot{\phi}_{(i)} + 3H\dot{\phi}_{(i)} + V_{,\phi_{(i)}} \\ + \delta\ddot{\phi}_{(i)} + 3H\delta\dot{\phi}_{(i)} - \frac{\Delta}{a^{2}}\delta\phi_{(i)} + \sum_{k}V_{,\phi_{(i)}\phi_{(k)}}\delta\phi_{(k)} - 2A\ddot{\phi}_{(i)} - \dot{\phi}_{(i)}\left(\dot{A} + 6HA - \frac{1}{a}B^{\alpha}_{\ |\alpha} - \dot{C}^{\alpha}_{\alpha}\right) \\ &= 2A\left[\delta\ddot{\phi}_{(i)} + 3H\delta\dot{\phi}_{(i)} - 2A\ddot{\phi}_{(i)} - \dot{\phi}_{(i)}\left(2\dot{A} + 6HA - \frac{1}{a}B^{\alpha}_{\ |\alpha} - \dot{C}^{\alpha}_{\alpha}\right)\right] + \delta\dot{\phi}_{(i)}\left(\dot{A} - \frac{1}{a}B^{\alpha}_{\ |\alpha} - \dot{C}^{\alpha}_{\alpha}\right) - 2\frac{1}{a}B^{\alpha}\delta\dot{\phi}_{(i),\alpha} \\ &+ \frac{1}{a}\delta\phi_{(i),\alpha}\left(\frac{1}{a}A^{,\alpha} - \dot{B}^{\alpha} - 2HB^{\alpha} - 2\frac{1}{a}C^{\alpha\beta}_{\ |\beta} + \frac{1}{a}C^{\beta|\alpha}_{\beta}\right) - 2\frac{1}{a^{2}}\delta\phi_{(i),\alpha|\beta}C^{\alpha\beta} - \frac{1}{2}\sum_{k,l}V_{,\phi_{(i)}\phi_{(k)}\phi_{(l)}}\delta\phi_{(k)}\delta\phi_{(l)} \\ &+ \ddot{\phi}_{(i)}B_{\alpha}B^{\alpha} + \dot{\phi}_{(i)}\left[\frac{1}{a}A_{,\alpha}B^{\alpha} + B_{\alpha}\dot{B}^{\alpha} + 3HB^{\alpha}B_{\alpha} + \frac{1}{a}B_{\beta}\left(2C^{\alpha\beta}_{\ |\alpha} - C^{\alpha|\beta}_{\alpha}\right) + 2C^{\alpha\beta}\left(\frac{1}{a}B_{\alpha|\beta} + \dot{C}_{\alpha\beta}\right)\right] \\ &\equiv N_{\phi_{(i)}}. \end{split}$$
(119)

The energy-momentum tensor gives:

$$\begin{split} \tilde{T}_{00}^{(\phi)} &= \frac{1}{2} \sum_{k} \phi_{(k)}^{\prime 2} + a^{2}V + \sum_{k} \left(\phi_{(k)}^{\prime} \delta \phi_{(k)}^{\prime} + a^{2} V_{,\phi_{(k)}} \delta \phi_{(k)} \right) + 2a^{2} V A + \sum_{k} \left(\frac{1}{2} \delta \phi_{(k)}^{\prime 2} + \frac{1}{2} \delta \phi_{(k),\alpha} \delta \phi_{(k)}^{,\alpha} + \frac{1}{2} a^{2} \sum_{l} V_{,\phi_{(k)}\phi_{(l)}} \delta \phi_{(k)} \delta \phi_{(l)} + 2a^{2} V_{,\phi_{(k)}} \delta \phi_{(k)} A - \phi_{(k)}^{\prime} \delta \phi_{(k),\alpha} B^{\alpha} + \frac{1}{2} \phi_{(k)}^{\prime 2} B_{\alpha} B^{\alpha} \right), \end{split}$$

$$\begin{split} \tilde{T}_{0\alpha}^{(\phi)} &= \sum_{k} \phi_{(k)}^{\prime} \delta\phi_{(k),\alpha} - \left(\frac{1}{2} \sum_{k} \phi_{(k)}^{\prime 2} - a^{2}V\right) B_{\alpha} + \sum_{k} \left[\delta\phi_{(k)}^{\prime} \delta\phi_{(k),\alpha} + \left(-\phi_{(k)}^{\prime} \delta\phi_{(k)}^{\prime} + a^{2}V_{,\phi_{(k)}} \delta\phi_{(k)} + \phi_{(k)}^{\prime 2}A\right) B_{\alpha}\right], \\ \tilde{T}_{\alpha\beta}^{(\phi)} &= g_{\alpha\beta}^{(3)} \left(\frac{1}{2} \sum_{k} \phi_{(k)}^{\prime 2} - a^{2}V\right) + g_{\alpha\beta}^{(3)} \sum_{k} \left(\phi_{(k)}^{\prime} \delta\phi_{(k)}^{\prime} - a^{2}V_{,\phi_{(k)}} \delta\phi_{(k)} - \phi_{(k)}^{\prime 2}A\right) + \left(\sum_{k} \phi_{(k)}^{\prime 2} - 2a^{2}V\right) C_{\alpha\beta} \\ &+ \sum_{k} \left\{\delta\phi_{(k),\alpha} \delta\phi_{(k),\beta} + 2 \left[\phi_{(k)}^{\prime} \delta\phi_{(k)}^{\prime} - a^{2}V_{,\phi_{(k)}} \delta\phi_{(k)} - \phi_{(k)}^{\prime 2}A\right] C_{\alpha\beta} - \frac{1}{2}g_{\alpha\beta}^{(3)} \left[-\delta\phi_{(k)}^{\prime 2} + \delta\phi_{(k),\gamma} \delta\phi_{(k)}^{,\gamma} + a^{2}\sum_{l} V_{,\phi_{(k)}\phi_{(l)}} \delta\phi_{(k)} \delta\phi_{(l)} + 4\phi_{(k)}^{\prime} \delta\phi_{(k)}^{\prime} A - 2\phi_{(k)}^{\prime} \delta\phi_{(k),\gamma} B^{\gamma} + \phi_{(k)}^{\prime 2} \left(-4A^{2} + B_{\gamma} B^{\gamma}\right)\right] \right\}.$$

$$(120)$$

Fluid quantities can be read from eq. (79) as:

$$\begin{split} \mu^{(\phi)} + \delta\mu^{(\phi)} &= \frac{1}{2} \sum_{k} \dot{\phi}_{(k)}^{2} + V + \sum_{k} \left(\dot{\phi}_{(k)} \delta\dot{\phi}_{(k)} - \dot{\phi}_{(k)}^{2} A + V_{,\phi_{(k)}} \delta\phi_{(k)} \right) + \sum_{k} \left[\frac{1}{2} \delta\dot{\phi}_{(k)}^{2} + \frac{1}{2a^{2}} \delta\phi_{(k),\alpha} \delta\phi_{(k)}^{,\alpha} + \frac{1}{2} \sum_{l} V_{,\phi_{(k)}\phi_{(l)}} \delta\phi_{(k)} \delta\phi_{(l)} - 2\dot{\phi}_{(k)} \delta\dot{\phi}_{(k)} A + \frac{1}{a} \dot{\phi}_{(k)} \delta\phi_{(k),\alpha} B^{\alpha} + \left(2A^{2} - \frac{1}{2}B^{\alpha}B_{\alpha} \right) \dot{\phi}_{(k)}^{2} \right], \\ p^{(\phi)} + \delta p^{(\phi)} &= \sum_{k} \frac{1}{2} \dot{\phi}_{(k)}^{2} - V + \sum_{k} \left(\dot{\phi}_{(k)} \delta\dot{\phi}_{(k)} - \dot{\phi}_{(k)}^{2} A - V_{,\phi_{(k)}} \delta\phi_{(k)} \right) + \sum_{k} \left[\frac{1}{2} \delta\dot{\phi}_{(k)}^{2} - \frac{1}{6a^{2}} \delta\phi_{(k),\alpha} \delta\phi_{(k)}^{,\alpha} - \frac{1}{2} \sum_{k} V_{,\phi_{(k)}\phi_{(l)}} \delta\phi_{(k)} \delta\phi_{(l)} - 2\dot{\phi}_{(k)} \delta\dot{\phi}_{(k)} A + \frac{1}{a} \dot{\phi}_{(k)} \delta\phi_{(k),\alpha} B^{\alpha} + \left(2A^{2} - \frac{1}{2}B^{\alpha}B_{\alpha} \right) \dot{\phi}_{(k)}^{2} \right], \\ Q^{(\phi)}_{\alpha} &= -\frac{1}{a} \sum_{k} \left[\dot{\phi}_{(k)} \delta\phi_{(k),\alpha} + \left(\delta\dot{\phi}_{(k)} - \dot{\phi}_{(k)} A \right) \delta\phi_{(k),\alpha} \right], \\ \Pi^{(\phi)}_{\alpha\beta} &= \frac{1}{a^{2}} \sum_{k} \left(\delta\phi_{(k),\alpha} \delta\phi_{(k),\beta} - \frac{1}{3}g^{(3)}_{\alpha\beta} \delta\phi_{(k),\gamma} \delta\phi_{(k)}^{,\gamma} \right). \end{split}$$
(121)

We indicate the quadratic parts as $\delta \mu^{(q)}$, $\delta p^{(q)}$, $Q^{(q)}_{\alpha}$, and $\Pi^{(q)}_{\alpha\beta}$.

D. Generalized gravity theories

1. Covariant equations

As the action for a class of generalized gravity theories we consider

$$S = \int \sqrt{-\tilde{g}} \left[\frac{1}{2} \tilde{f}(\tilde{\phi}^K, \tilde{R}) - \frac{1}{2} \tilde{g}_{IJ}(\tilde{\phi}^K) \tilde{\phi}^{I,c} \tilde{\phi}^J_{,c} - \tilde{V}(\tilde{\phi}^K) + \tilde{L}_m \right] d^4x.$$
(122)

 $\tilde{\phi}^{I}$ is the *I*-th component of *N* scalar fields. The capital indices $I, J, K, \ldots = 1, 2, 3, \ldots, N$ indicate the scalar fields, and the summation convention is used for repeated indices. $\tilde{f}(\tilde{\phi}^{K}, \tilde{R})$ is a general algebraic function of \tilde{R} and the scalar fields $\tilde{\phi}^{I}$, and $\tilde{g}_{IJ}(\tilde{\phi}^{K})$ and $\tilde{V}(\tilde{\phi}^{K})$ are general algebraic functions of the scalar fields. We include a nonlinear sigma type kinetic term where the kinetic matrix \tilde{g}_{IJ} is considered as a Riemannian metric on the manifold with the coordinates $\tilde{\phi}^{I}$. The matter part Lagrangian \tilde{L}_{m} includes the fluids, the kinetic components, and the interaction with the fields, as well. We have introduced the general action in eq. (122) in [23,12] as a toy model which allows quite general handling of various different generalized gravity theories in a unified manner, see [24]. Our generalized gravity includes as subset: f(R) gravity which includes R^{2} gravity, the scalar-tensor theory which includes the Jordan-Brans-Dicke theory, the non-minimally coupled scalar field, the induced gravity, the low-energy effective action of string theory, etc, and various combinations of such gravity theories with additional multiple fields and fluids. It does not, however, include higher-derivative theories with terms like $R^{ab}R_{ab}$, see [25] for its role.

The gravitational field equation and the equation of motion become:

$$\tilde{G}_{ab} = \frac{1}{\tilde{F}} \left[\tilde{T}_{ab} + \tilde{g}_{IJ} \left(\tilde{\phi}^{I}{}_{,a} \tilde{\phi}^{J}{}_{,b} - \frac{1}{2} \tilde{g}_{ab} \tilde{\phi}^{I,c} \tilde{\phi}^{J}{}_{,c} \right) + \frac{1}{2} \left(\tilde{f} - \tilde{R}\tilde{F} - 2\tilde{V} \right) \tilde{g}_{ab} + \tilde{F}_{,a;b} - \tilde{g}_{ab} \tilde{F}^{;c}{}_{c} \right] \equiv 8\pi G \tilde{T}^{(\text{eff})}_{ab}, \quad (123)$$

$$\tilde{\phi}^{I;c}{}_{c} + \frac{1}{2} \left(\tilde{f} - 2\tilde{V} \right)^{,I} + \tilde{\Gamma}^{I}_{JK} \tilde{\phi}^{J,c} \tilde{\phi}^{K}{}_{,c} = -\tilde{L}_{m}{}^{,I} \equiv \tilde{\Gamma}^{I},$$
(124)

$$\tilde{T}^b_{a;b} = \tilde{L}_{m,J} \tilde{\phi}^J_{\ ,a},\tag{125}$$

where $\tilde{F} \equiv \partial \tilde{f} / \partial \tilde{R}$; \tilde{g}^{IJ} is the inverse metric of \tilde{g}_{IJ} , $\tilde{\Gamma}^{I}_{JK} \equiv \frac{1}{2} \tilde{g}^{IL} (\tilde{g}_{LJ,K} + \tilde{g}_{LK,J} - \tilde{g}_{JK,L})$, and $\tilde{V}_{,\tilde{I}} \equiv \partial \tilde{V} / (\partial \tilde{\phi}^{I})$. Introduction of the effective energy-momentum tensor $\tilde{T}_{ab}^{(\text{eff})}$ provides a useful trick to derive and handle the perturbed set of equations [23]. It allows the equations derived in Einstein gravity remain valid with the energy-momentum parts replaced by the effective ones.

We note that the gravity theory in eq. (122) can be transformed to Einstein's gravity through a conformal rescaling of the metric and rescaling of one of the fields. As the result we have Einstein's gravity sector with only complications appearing in the modified form of the field potential; the nonlinear sigma type couplings in the kinetic part also remain. We have studied the conformal transformation properties to the linear order perturbation in [23,26], and in most general form in the Appendix A of [29]. Extention to the second-order perturbation is trivial.

2. Perturbed equations

The perturbed set of equations can be derived similarly as in the previous sections on the scalar fields. We set

$$\tilde{F} \equiv F + \delta F, \quad \tilde{\Gamma}^I \equiv \Gamma^I + \delta \Gamma^I.$$
 (126)

Thus,

$$\delta F = F_{,I}\delta\phi^{I} + F_{,R}\delta R + \frac{1}{2}F_{,IJ}\delta\phi^{I}\delta\phi^{J} + F_{,IR}\delta\phi^{I}\delta R + \frac{1}{2}F_{,RR}\delta R^{2}.$$
(127)

The equation of motion in eq. (124) gives

$$\begin{split} \dot{\phi}^{I} + 3H\dot{\phi}^{I} - \frac{1}{2}g^{IJ}\left(f_{,J} - 2V_{,J}\right) + \Gamma_{JK}^{I}\dot{\phi}^{J}\dot{\phi}^{K} + \Gamma^{I} \\ + \delta\ddot{\phi}^{I} + 3H\delta\dot{\phi}^{I} - \frac{\Delta}{a^{2}}\delta\phi^{I} - 2A\ddot{\phi}^{I} + \dot{\phi}^{I}\left(-\dot{A} - 6HA + \frac{1}{a}B^{\alpha}{}_{|\alpha} + \dot{C}^{\alpha}{}_{\alpha}\right) - \frac{1}{2}g^{IJ}\left[F_{,J}\delta R + (f_{,LJ} - 2V_{,LJ})\delta\phi^{L}\right] \\ - \frac{1}{2}g^{IJ}{}_{,L}\delta\phi^{L}\left(f_{,J} - 2V_{,J}\right) + 2\Gamma_{JK}^{I}\left(\dot{\phi}^{J}\delta\dot{\phi}^{K} - A\dot{\phi}^{J}\dot{\phi}^{K}\right) + \Gamma_{JK,L}^{J}\delta\phi^{L}\dot{\phi}^{J}\dot{\phi}^{K} + \delta\Gamma^{I} \\ = 2A\left[\delta\ddot{\phi}^{I} + 3H\delta\dot{\phi}^{I} - 2A\ddot{\phi}^{I} - \dot{\phi}^{I}\left(2\dot{A} + 6HA - \frac{1}{a}B^{\alpha}{}_{|\alpha} - \dot{C}^{\alpha}{}_{\alpha}\right) + 2\Gamma_{JK}^{I}\left(\dot{\phi}^{J}\delta\dot{\phi}^{K} - A\dot{\phi}^{J}\dot{\phi}^{K}\right) + \Gamma_{JK,L}^{I}\delta\phi^{L}\dot{\phi}^{J}\dot{\phi}^{K} \\ + B^{\alpha}B_{\alpha}\left(\ddot{\phi}^{I} + 3H\dot{\phi}^{I} + \Gamma_{JK}^{I}\dot{\phi}^{J}\dot{\phi}^{K}\right) + \left(\dot{A} - \frac{1}{a}B^{\alpha}{}_{|\alpha} - \dot{C}^{\alpha}{}_{\alpha}\right)\delta\dot{\phi}^{I} - 2\frac{1}{a}B^{\alpha}\dot{\delta}\dot{\phi}^{I}{}_{,\alpha} - 2\frac{1}{a^{2}}C^{\alpha\beta}\delta\phi^{I}{}_{,\alpha|\beta} \\ + \frac{1}{a}\delta\phi^{I}{}_{,\alpha}\left(\frac{1}{a}A^{,\alpha} - \dot{B}^{\alpha} - 2HB^{\alpha} - 2\frac{1}{a}C^{\alpha\beta}{}_{|\beta} + \frac{1}{a}C^{\beta|\alpha}{}_{\beta}\right) \\ + \dot{\phi}^{I}\left[\frac{1}{a}A_{,\alpha}B^{\alpha} + B^{\alpha}\dot{B}_{\alpha} + \frac{1}{a}B_{\alpha}\left(2C^{\alpha\beta}{}_{|\beta} - C^{\beta|\alpha}{}_{\beta}\right) + 2C^{\alpha\beta}\left(\frac{1}{a}B_{\alpha|\beta} + \dot{C}_{\alpha\beta}\right)\right] \\ + \frac{1}{4}g^{IJ}\left[F_{,RJ}\delta R^{2} + 2F_{,LJ}\delta R\delta\phi^{L} + (f_{,LMJ} - 2V_{,LMJ})\delta\phi^{L}\delta\phi^{M}\right] \\ + \frac{1}{2}g^{IJ}{}_{,L}\delta\phi^{L}\left[F_{,J}\delta R + (f_{,LJ} - 2V_{,LJ})\delta\phi^{L}\right] + \frac{1}{4}g^{IJ}{}_{,LM}\delta\phi^{L}\delta\phi^{M}\left(f_{,J} - 2V_{,J}\right) \\ + \Gamma_{JK}^{I}\left(-\delta\dot{\phi}^{J}\delta\dot{\phi}^{K} - 2\frac{1}{a}\dot{\phi}^{J}\delta\phi^{K}{}_{,\alpha}B^{\alpha} + \frac{1}{a^{2}}\delta\phi^{J|\alpha}\delta\phi^{K}{}_{,\alpha}\right) - 2\Gamma_{JK,L}^{I}\delta\phi^{L}\dot{\phi}^{J}\delta\dot{\phi}^{K} - \frac{1}{2}\Gamma_{JK,LM}^{I}\delta\phi^{L}\delta\phi^{M}\dot{\phi}^{J}\dot{\phi}^{K} \\ \equiv N_{g}. \end{split}$$
(128)

 δR can be read from eq. (98). From eq. (123) the effective energy momentum tensor gives:

$$-2A\delta F'' - \left(A' + 2\frac{a'}{a}A - B^{\gamma}_{|\gamma} - C^{\gamma\prime}_{\gamma}\right)\delta F' + 2B^{\gamma}\delta F'_{,\gamma} + \left(-A^{,\gamma} + B^{\gamma\prime} + \frac{a'}{a}B^{\gamma} + 2C^{\gamma\delta}_{|\delta} - C^{\delta|\gamma}_{\delta}\right)\delta F_{,\gamma} + 2C^{\gamma\delta}\delta F_{,\gamma|\delta} + \left(4A^2 - B^{\gamma}B_{\gamma}\right)\left(F'' + \frac{a'}{a}F'\right) + \left[4AA' - A_{,\gamma}B^{\gamma} - B^{\gamma}B'_{\gamma} - 2A\left(B^{\gamma}_{|\gamma} + C^{\gamma\prime}_{\gamma}\right) - B_{\gamma}\left(2C^{\gamma\delta}_{|\delta} - C^{\delta|\gamma}_{\delta}\right) - 2C^{\gamma\delta}\left(B_{\gamma|\delta} + C'_{\gamma\delta}\right)\right]F'\right]\right\}.$$
(129)

The fluid quantities follow from eq. (79):

$$\begin{split} \mu^{(\text{eff})} + \delta\mu^{(\text{eff})} &= \frac{1}{8\pi G \tilde{F}} \Biggl\{ \mu + \frac{1}{2} g_{IJ} \dot{\phi}^{I} \dot{\phi}^{J} - \frac{1}{2} \left(f - RF - 2V \right) - 3H \dot{F} \\ &+ \delta\mu + g_{IJ} \left(\dot{\phi}^{I} \delta \dot{\phi}^{J} - A \dot{\phi}^{I} \dot{\phi}^{J} \right) + \frac{1}{2} g_{IJ,L} \delta \phi^{L} \dot{\phi}^{I} \dot{\phi}^{J} - \frac{1}{2} \left(f_{,L} - 2V_{,L} \right) \delta \phi^{L} \\ &- 3H \delta \dot{F} + \left(\frac{1}{2} R + \frac{\Delta}{a^{2}} \right) \delta F - \left(-6HA + \frac{1}{a} B^{\alpha}{}_{|\alpha} + \dot{C}^{\alpha}{}_{\alpha} \right) \dot{F} \\ &+ \frac{1}{2} g_{IJ} \left[\delta \dot{\phi}^{I} \delta \dot{\phi}^{J} + \frac{1}{a^{2}} \delta \phi^{I,\alpha} \delta \phi^{J}{}_{,\alpha} - 4A \dot{\phi}^{I} \delta \dot{\phi}^{J} + 2\frac{1}{a} B^{\alpha} \dot{\phi}^{I} \delta \phi^{J}{}_{,\alpha} + \left(4A^{2} - B^{\alpha}B_{\alpha} \right) \dot{\phi}^{I} \dot{\phi}^{J} \right] \\ &+ g_{IJ,L} \delta \phi^{L} \left(\dot{\phi}^{I} \delta \dot{\phi}^{J} - A \dot{\phi}^{I} \dot{\phi}^{J} \right) + \frac{1}{4} g_{IJ,LM} \delta \phi^{L} \delta \phi^{M} \dot{\phi}^{I} \dot{\phi}^{J} \\ &+ \frac{1}{4} \left[F_{,R} \delta R^{2} - \left(f_{,LM} - 2V_{,LM} \right) \delta \phi^{L} \delta \phi^{M} \right] - \left(-6HA + \frac{1}{a} B^{\alpha}{}_{|\alpha} + \dot{C}^{\alpha}{}_{\alpha} \right) \left(\delta \dot{F} - 2A \dot{F} \right) \\ &- \left[3H B^{\alpha} + \frac{1}{a} \left(2C^{\alpha\beta}{}_{|\beta} - C^{\beta|\alpha}{}_{\beta} \right) \right] \left(\frac{1}{a} \delta F_{,\alpha} - B_{\alpha} \dot{F} \right) - 2\frac{1}{a^{2}} C^{\alpha\beta} \delta F_{,\alpha|\beta} + 2\dot{F} C^{\alpha\beta} \left(\frac{1}{a} B_{\alpha|\beta} + \dot{C}_{\alpha\beta} \right) \right\}, \\ p^{(\text{eff})} + \delta p^{(\text{eff})} = \frac{1}{8\pi G \tilde{F}} \Biggl\{ p + \frac{1}{2} g_{IJ} \dot{\phi}^{I} \dot{\phi}^{J} + \frac{1}{2} \left(f - RF - 2V \right) + \ddot{F} + 2H \dot{F} \end{aligned}$$

$$\begin{split} \delta\pi GF\left(\begin{array}{c} 2 \\ +\delta p + g_{IJ}\left(\dot{\phi}^{I}\delta\dot{\phi}^{J} - A\dot{\phi}^{I}\dot{\phi}^{J}\right) + \frac{1}{2}g_{IJ,L}\delta\phi^{L}\dot{\phi}^{I}\dot{\phi}^{J} + \frac{1}{2}\left(f_{,L} - 2V_{,L}\right)\delta\phi^{L} \\ +\delta\ddot{F} + 2H\delta\dot{F} - \left(\frac{1}{2}R + \frac{2}{3}\frac{\Delta}{a^{2}}\right)\deltaF - 2A\ddot{F} - \left[\dot{A} + 4HA - \frac{2}{3}\left(\frac{1}{a}B^{\alpha}{}_{|\alpha} + \dot{C}^{\alpha}_{\alpha}\right)\right]\dot{F} \\ +g_{IJ}\left[\frac{1}{2}\delta\phi^{I}\delta\dot{\phi}^{J} - 2A\dot{\phi}^{I}\delta\dot{\phi}^{J} + \frac{1}{2}\left(4A^{2} - B^{\alpha}B_{\alpha}\right)\dot{\phi}^{I}\dot{\phi}^{J} + \frac{1}{a}B^{\alpha}\dot{\phi}^{I}\delta\phi^{J}, \alpha - \frac{1}{6a^{2}}\delta\phi^{I,\alpha}\delta\phi^{J}, \alpha\right] \\ +g_{IJ,L}\delta\phi^{L}\left(\dot{\phi}^{I}\delta\dot{\phi}^{J} - A\dot{\phi}^{I}\dot{\phi}^{J}\right) + \frac{1}{4}g_{IJ,LM}\delta\phi^{L}\delta\phi^{M}\phi^{I}\dot{\phi}^{J} - \frac{1}{4}\left[F_{,R}\delta R^{2} - \left(f_{,LM} - 2V_{,LM}\right)\delta\phi^{L}\delta\phi^{M}\right] \\ -2A\delta\ddot{F} - \left[\dot{A} + 4HA - \frac{2}{3}\left(\frac{1}{a}B^{\alpha}{}_{|\alpha} + \dot{C}^{\alpha}_{\alpha}\right)\right]\left(\delta\dot{F} - 2A\dot{F}\right) + 2\frac{1}{a}B^{\alpha}\delta\dot{F}, \alpha \\ + \left[-\frac{1}{a}A^{,\alpha} + \dot{B}^{\alpha} + HB^{\alpha} + \frac{2}{3a}\left(2C^{\alpha\beta}{}_{|\beta} - C^{\beta|\alpha}_{\beta}\right)\right]\left(\frac{1}{a}\delta F_{,\alpha} - B_{\alpha}\dot{F}\right) + \frac{4}{3a^{2}}C^{\alpha\beta}\delta F_{,\alpha|\beta} \\ + \left(4A^{2} - B^{\alpha}B_{\alpha}\right)\ddot{F} + \left[2A\dot{A} - \frac{2}{a}A_{,\alpha}B^{\alpha} - HB^{\alpha}B_{\alpha} - \frac{4}{3}C^{\alpha\beta}\left(\frac{1}{a}B_{\alpha|\beta} + \dot{C}_{\alpha\beta}\right)\right]\dot{F}\right], \\ Q_{\alpha}^{(\text{eff})} = \frac{1}{8\pi G} \left\{Q_{\alpha} - \frac{1}{a}g_{IJ}\dot{\phi}^{I}\delta\phi^{J}, \alpha + \frac{1}{a}\left(-\delta\dot{F}, \alpha + H\delta F, \alpha\right) + \frac{1}{a}A_{,\alpha}\dot{F} \\ - \frac{1}{a}g_{IJ}\left(\delta\dot{\phi}^{I} - A\dot{\phi}^{I}\right)\delta\phi^{J}, \alpha - \frac{1}{a}g_{IJ,L}\delta\phi^{L}\dot{\phi}^{I}\delta\phi^{J}, \alpha + A\left[-3\frac{1}{a}\dot{F}A_{,\alpha} + \frac{1}{a}\left(\delta\dot{F}, \alpha - H\delta F, \alpha\right)\right] \\ + \frac{1}{a}A_{,\alpha}\delta\dot{F} + \frac{1}{a}\left[\frac{1}{2a}\left(B_{\alpha}{}^{|\beta} - B^{\beta}{}_{|\alpha}\right) + \dot{C}^{\beta}_{\alpha}\right]\delta F_{,\beta} - \frac{1}{a^{2}}B^{\beta}\delta F_{,\alpha|\beta} + \frac{1}{a}B^{\beta}B_{\alpha|\beta}\dot{F}\right\}, \\ \Pi_{\alpha\beta}^{(\text{eff})} = \frac{1}{8\pi G}\left[\frac{1}{a}\left[1_{\alpha\beta} + \frac{1}{a^{2}}\delta F_{,\alpha|\beta} - \left(\frac{1}{a}B_{(\alpha|\beta)} + \dot{C}_{\alpha\beta}\right)\dot{F} - \frac{1}{3}g_{\alpha\beta}^{(3)}\left[\frac{\Delta}{a^{2}}\delta F - \left(\frac{1}{a}B^{\gamma}_{|\gamma} + \dot{C}^{\gamma}_{\gamma}\right)\dot{F}\right] \right] \end{cases}$$

$$+\frac{1}{a^{2}}g_{IJ}\delta\phi^{I}{}_{,\alpha}\delta\phi^{J}{}_{,\beta} - \frac{2}{3}C_{\alpha\beta}\left[\frac{\Delta}{a^{2}}\delta F - \left(\frac{1}{a}B^{\gamma}{}_{|\gamma} + \dot{C}^{\gamma}_{\gamma}\right)\dot{F}\right] \\ - \left(\frac{1}{a}B_{(\alpha|\beta)} + \dot{C}_{\alpha\beta}\right)\left(\delta\dot{F} - 2A\dot{F}\right) - \frac{1}{a}\left(2C^{\gamma}{}_{(\alpha|\beta)} - C_{\alpha\beta}{}^{|\gamma}\right)\left(\frac{1}{a}\delta F_{,\gamma} - B_{\gamma}\dot{F}\right) \\ - \frac{1}{3}g^{(3)}_{\alpha\beta}\left[\frac{1}{a^{2}}g_{IJ}\delta\phi^{I,\gamma}\delta\phi^{J}{}_{,\gamma} + 2C^{\gamma\delta}\left(\frac{1}{a}B_{\gamma|\delta} + \dot{C}_{\gamma\delta}\right)\dot{F} - \frac{2}{a^{2}}C^{\gamma\delta}\delta F_{,\gamma|\delta} \\ - \left(\frac{1}{a}B^{\gamma}{}_{|\gamma} + \dot{C}^{\gamma}_{\gamma}\right)\left(\delta\dot{F} - 2A\dot{F}\right) - \frac{1}{a}\left(2C^{\gamma\delta}{}_{|\delta} - C^{\delta|\gamma}_{\delta}\right)\left(\frac{1}{a}\delta F_{,\gamma} - B_{\gamma}\dot{F}\right)\right]\right\}.$$
(130)

We have used eq. (78) for the energy-momentum tensor. We indicate the quadratic parts as $\delta\mu^{(\text{eff},q)}$, $\delta p^{(\text{eff},q)}$, $Q_{\alpha}^{(\text{eff},q)}$, and $\Pi_{\alpha\beta}^{(\text{eff},q)}$. We note again that in this generalized gravity the basic equations in §IVA *remain valid* with the fluid quantities replaced by the effective ones.

E. Electromagnetic field

1. Covariant equations

The Lagrangian of electromagnetic field is given as

$$\mathcal{L}_{\rm em} = -\frac{1}{4}\sqrt{-\tilde{g}}\tilde{F}^{ab}\tilde{F}_{ab},\tag{131}$$

where $\tilde{F}_{ab} \equiv \tilde{A}_{a,b} - \tilde{A}_{b,a}$. The energy-momentum tensor becomes

$$\tilde{T}_{ab}^{(\mathrm{em})} = \tilde{F}_{ac}\tilde{F}_{b}^{\ c} - \frac{1}{4}\tilde{g}_{ab}\tilde{F}_{cd}\tilde{F}^{cd}.$$
(132)

We introduce [20]

$$\tilde{F}_{ab} = \tilde{u}_a \tilde{E}_b - \tilde{u}_b \tilde{E}_a - \tilde{\eta}_{abcd} \tilde{u}^c \tilde{H}^d,
\tilde{E}_a \equiv \tilde{F}_{ab} \tilde{u}^b, \quad \tilde{H}_a \equiv \frac{1}{2} \tilde{\eta}_{abcd} \tilde{u}^b \tilde{F}^{cd}, \quad \tilde{E}^2 \equiv \tilde{E}^a \tilde{E}_a, \quad \tilde{H}^2 \equiv \tilde{H}^a \tilde{H}_a,
\tilde{q} \equiv -\tilde{j}^a \tilde{u}_a, \quad \tilde{J}^a \equiv \tilde{h}_b^a \tilde{j}^b.$$
(133)

Then we have

$$\tilde{T}_{ab}^{(em)} = \frac{1}{2} \tilde{u}_{a} \tilde{u}_{b} \left(\tilde{E}^{2} + \tilde{H}^{2} \right) + 2 \tilde{u}_{(a} \tilde{\eta}_{b)cgd} \tilde{u}^{c} \tilde{E}^{g} \tilde{H}^{d} - \tilde{E}_{a} \tilde{E}_{b} - \tilde{H}_{a} \tilde{H}_{b} + \frac{1}{2} \tilde{h}_{ab} \left(\tilde{E}^{2} + \tilde{H}^{2} \right),$$

$$\tilde{\mu}^{(em)} = \frac{1}{2} \left(\tilde{E}^{2} + \tilde{H}^{2} \right), \quad \tilde{p}^{(em)} = \frac{1}{6} \left(\tilde{E}^{2} + \tilde{H}^{2} \right),$$

$$\tilde{q}_{a}^{(em)} = \tilde{\eta}_{acgd} \tilde{u}^{c} \tilde{E}^{g} \tilde{H}^{d}, \quad \tilde{\pi}_{ab}^{(em)} = -\tilde{E}_{a} \tilde{E}_{b} - \tilde{H}_{a} \tilde{H}_{b} + \frac{1}{3} \tilde{h}_{ab} \left(\tilde{E}^{2} + \tilde{H}^{2} \right).$$
(134)

From the Maxwell equations and the conservation equations

$$\tilde{F}^{ab}_{\ ;b} = \tilde{j}^{a}, \quad \tilde{F}_{[ab;c]} = 0, \quad \tilde{j}^{a}_{\ ;a} = 0,$$
(135)

we can derive the covariant forms of relativistic Maxwell's equations [20]:

$$\tilde{E}^a{}_{;b}\tilde{h}^b_a + 2\tilde{H}_a\tilde{\omega}^a = \tilde{q},\tag{136}$$

$$\tilde{H}^a{}_{;b}\tilde{h}^b_a - 2\tilde{E}_a\tilde{\omega}^a = 0, \tag{137}$$

$$\tilde{h}^{a}_{b}\tilde{E}^{b}_{;c}\tilde{u}^{c} = \tilde{E}^{b}\left(\tilde{\omega}^{a}_{\ b} + \tilde{\sigma}^{a}_{\ b} - \frac{2}{3}\tilde{\theta}\tilde{h}^{a}_{b}\right) + \tilde{\eta}^{abcd}u_{b}\left(\tilde{a}_{c}\tilde{H}_{d} - \tilde{H}_{c;d}\right) - \tilde{J}^{a},\tag{138}$$

$$\tilde{h}^{a}_{b}\tilde{H}^{b}_{;c}\tilde{u}^{c} = \tilde{H}^{b}\left(\tilde{\omega}^{a}_{\ b} + \tilde{\sigma}^{a}_{\ b} - \frac{2}{3}\tilde{\theta}\tilde{h}^{a}_{b}\right) + \tilde{\eta}^{abcd}\tilde{u}_{b}\left(\tilde{a}_{c}\tilde{E}_{d} - \tilde{E}_{c;d}\right) - \tilde{J}^{a},\tag{139}$$

$$\tilde{q}_{,a}\tilde{u}^a + \tilde{\theta}\tilde{q} + \tilde{h}^a_b \tilde{J}^b_{;a} + \tilde{J}^a \tilde{a}_a = 0.$$
(140)

2. Perturbations

We take the normal-frame, thus $\tilde{u}_a = \tilde{n}_a$, thus $\tilde{\omega}_{ab} = 0$. Due to the high symmetry the Friedmann background does not support the electric or magnetic field. Thus, \tilde{E}_a and \tilde{H}_a are already perturbed order. We set

$$\tilde{E}_{\alpha} \equiv E_{\alpha}, \quad \tilde{H}_{\alpha} \equiv H_{\alpha},$$
(141)

where E_{α} and H_{α} are based on $g_{\alpha\beta}^{(3)}$. Thus, $\tilde{E}_0 = -E_{\alpha}B^{\alpha}$ (which follows from $\tilde{E}_a \tilde{n}^a = 0$), etc. For η^{abcd} see eq. (94). Equations (136-140) become:

$$E^{\alpha}_{\ |\alpha} - a^2 \delta q = 2 \left(C^{\alpha\beta} E_{\beta} \right)_{|\alpha} - C^{\alpha}_{\alpha|\beta} E^{\beta}, \tag{142}$$

$$H^{\alpha}_{\ |\alpha} = 2\left(C^{\alpha\beta}H_{\beta}\right)_{|\alpha} - C^{\alpha}_{\alpha|\beta}H^{\beta},\tag{143}$$

$$\dot{E}^{\alpha} + HE^{\alpha} + \frac{1}{a}\eta^{\alpha\beta\gamma}H_{\beta|\gamma} + J^{\alpha} = A\left(\dot{E}^{\alpha} + HE^{\alpha}\right) - \frac{1}{a}E^{\alpha}{}_{|\beta}B^{\beta} + E^{\beta}\left(\frac{1}{a}B^{\alpha}{}_{|\beta} + 2\dot{C}^{\alpha}_{\beta}\right) -E^{\alpha}\left(\frac{1}{a}B^{\beta}{}_{|\beta} + \dot{C}^{\beta}_{\beta}\right) + \frac{1}{a}\eta^{\alpha\beta\gamma}\left(H_{\gamma}A_{,\beta} - H_{\beta|\gamma}C^{\delta}_{\delta}\right) - \frac{2}{a}\eta^{\beta\gamma\delta}C^{\alpha}_{\beta}H_{\gamma|\delta},$$
(144)

$$(H_{\alpha} \Leftrightarrow E_{\alpha}), \tag{145}$$

$$\delta \dot{q} + 3H\delta q = -3HA\delta q - \frac{1}{a}\delta q_{,\alpha}B^{\alpha} + \delta K\delta q - \frac{1}{a^2} \left[\left(J^{\alpha} - 2C^{\alpha\beta}J_{\beta} \right)_{|\alpha} + J^{\alpha} \left(C^{\beta}_{\beta|\alpha} + A_{,\alpha} \right) \right],\tag{146}$$

where we set

$$\tilde{q} \equiv q + \delta q, \quad \tilde{J}_{\alpha} \equiv J_{\alpha},$$
(147)

with J_{α} based on $g_{\alpha\beta}^{(3)}$; J_{α} in this subsection differs from the flux term in ADM notation used in the other sections. We have q = 0.

The energy-momentum tensor becomes:

$$\tilde{T}_{00}^{(\text{em})} = \frac{1}{2} \left(E^{\alpha} E_{\alpha} + H^{\alpha} H_{\alpha} \right),$$

$$\tilde{T}_{0\alpha}^{(\text{em})} = -\eta_{\alpha\beta\gamma} E^{\beta} H^{\gamma},$$

$$\tilde{T}_{\alpha\beta}^{(\text{em})} = -E_{\alpha} E_{\beta} - H_{\alpha} H_{\beta} + \frac{1}{2} g_{\alpha\beta}^{(3)} \left(E^{\gamma} E_{\gamma} + H^{\gamma} H_{\gamma} \right).$$
(148)

Fluid quantities can be read from eq. (79) as:

$$\delta\mu^{(\mathrm{em})} = 3\delta p^{(\mathrm{em})} = \frac{1}{2a^2} \left(E^{\alpha} E_{\alpha} + H^{\alpha} H_{\alpha} \right),$$

$$Q^{(\mathrm{em})}_{\alpha} = \frac{1}{a^2} \eta_{\alpha\beta\gamma} E^{\beta} H^{\gamma},$$

$$\Pi^{(\mathrm{em})}_{\alpha\beta} = -\frac{1}{a^2} \left[E_{\alpha} E_{\beta} + H_{\alpha} H_{\beta} - \frac{1}{3} g^{(3)}_{\alpha\beta} \left(E^{\gamma} E_{\gamma} + H^{\gamma} H_{\gamma} \right) \right].$$
(149)

We have $\mu^{(em)} = 0 = p^{(em)}$.

F. Null-geodesic and temperature anisotropy

We introduce the photon four-velocity as:

$$\tilde{k}^{0} \equiv \frac{1}{a} \left(\nu + \delta \nu \right), \quad \tilde{k}^{\alpha} \equiv -\frac{\nu}{a} \left(e^{\alpha} + \delta e^{\alpha} \right);$$

$$\tilde{k}_{0} = -a\nu \left(1 + \frac{\delta\nu}{\nu} + 2A - B_{\alpha}e^{\alpha} + 2A\frac{\delta\nu}{\nu} - B_{\alpha}\delta e^{\alpha} \right),$$

$$\tilde{k}_{\alpha} = -a\nu \left(e_{\alpha} + \delta e_{\alpha} + B_{\alpha} + 2C_{\alpha\beta}e^{\beta} + B_{\alpha}\frac{\delta\nu}{\nu} + 2C_{\alpha\beta}\delta e^{\beta} \right), \quad (150)$$

where e^{α} and δe^{α} are based on $g^{(3)}_{\alpha\beta}$, and ν and e^{α} are assumed to be the background order. We have

$$\frac{d}{d\lambda} = \frac{\partial x^a}{\partial \lambda} \frac{\partial}{\partial x^a} = \tilde{k}^a \partial_a = \frac{\nu}{a} \left(\partial_0 - e^\alpha \partial_\alpha + \frac{\delta \nu}{\nu} \partial_0 - \delta e^\alpha \partial_\alpha \right).$$
(151)

Thus,

$$\frac{d}{dy} \equiv \partial_0 - e^\alpha \partial_\alpha,\tag{152}$$

can be considered as a derivative along the background photon four-velocity. The null equation, $\tilde{k}^a \tilde{k}_a = 0$, gives

$$\tilde{k}^{a}\tilde{k}_{a} = \nu^{2} \left[e^{\alpha}e_{\alpha} - 1 + 2\left(e^{\alpha}\delta e_{\alpha} - \frac{\delta\nu}{\nu} - A + B_{\alpha}e^{\alpha} + C_{\alpha\beta}e^{\alpha}e^{\beta}\right) + \delta e^{\alpha}\delta e_{\alpha} - \frac{\delta\nu^{2}}{\nu^{2}} - 2\frac{\delta\nu}{\nu}\left(2A - B_{\alpha}e^{\alpha}\right) + 2\left(B_{\alpha} + 2C_{\alpha\beta}e^{\beta}\right)\delta e^{\alpha} \right] = 0.$$
(153)

The geodesic equation, $\tilde{k}^a_{\ ;b}\tilde{k}^b=0$, using eq. (52), gives

$$\begin{split} \tilde{k}^{0}_{,b}\tilde{k}^{b} &= \frac{\nu^{2}}{a^{2}} \Biggl\{ \frac{(a\nu)'}{a\nu} + \left(\frac{\delta\nu}{\nu}\right)' + 2\frac{\nu'}{\nu}\frac{\delta\nu}{\nu} - \frac{\delta\nu_{,\alpha}}{\nu}e^{\alpha} + 2\frac{a'}{a}e^{\alpha}\delta e_{\alpha} + A' - 2\frac{a'}{a}A \\ &+ \left(B_{\alpha|\beta} + C'_{\alpha\beta} + 2\frac{a'}{a}C_{\alpha\beta}\right)e^{\alpha}e^{\beta} - 2\left(A_{,\alpha} - \frac{a'}{a}B_{\alpha}\right)e^{\alpha} \\ &+ \frac{\delta\nu}{\nu}\frac{\delta\nu'}{\nu} - \frac{\delta\nu_{,\alpha}}{\nu}\delta e^{\alpha} + 2\frac{\delta\nu}{\nu}A' - 2\frac{\delta\nu}{\nu}\left(A_{,\alpha} - \frac{a'}{a}B_{\alpha}\right)e^{\alpha} + \frac{a'}{a}\delta e^{\alpha}\delta e_{\alpha} \\ &- 2\delta e^{\alpha}\left(A_{,\alpha} - \frac{a'}{a}B_{\alpha}\right) - 4\frac{a'}{a}e^{\alpha}\delta e_{\alpha}A + 2e^{\alpha}\delta e^{\beta}\left(C'_{\alpha\beta} + 2\frac{a'}{a}C_{\alpha\beta} + B_{(\alpha|\beta)}\right) \\ &- \left[A\left(2B_{\alpha|\beta} + 2C'_{\alpha\beta} + 4\frac{a'}{a}C_{\alpha\beta}\right) + B_{\gamma}\left(2C^{\gamma}_{\alpha|\beta} - C_{\alpha\beta}^{-|\gamma|}\right)\right]e^{\alpha}e^{\beta} + \frac{a'}{a}\left(4A^{2} - B^{\alpha}B_{\alpha}\right) \\ &+ 2\left(2AA_{,\alpha} - 2\frac{a'}{a}AB_{\alpha} + B_{\beta}C^{\beta\prime}_{\alpha} + B^{\beta}B_{[\alpha|\beta]}\right)e^{\alpha} - 2AA' - A_{,\alpha}B^{\alpha} + B^{\alpha}\left(B'_{\alpha} + \frac{a'}{a}B_{\alpha}\right)\right\} = 0, \quad (154) \\ \tilde{k}^{\alpha}_{,b}\tilde{k}^{b} = \frac{\nu^{2}}{a^{2}}\left\{-e^{\alpha\prime} + e^{\beta}e^{\alpha}_{|\beta} - \delta e^{\alpha\prime} - \frac{\delta\nu}{\nu}e^{\alpha\prime} + \delta e^{\alpha}_{|\beta}e^{\beta} + \delta e^{\beta}e^{\alpha}_{|\beta} \\ &+ \left(2C^{\alpha}_{\beta|\gamma} - C_{\beta\gamma}\right)^{|\alpha|}\right)e^{\beta}e^{\gamma} - \left(B_{\beta}^{-|\alpha} - B^{\alpha}_{|\beta} + 2C^{\alpha\prime}_{\beta}\right)e^{\beta} + A^{,\alpha} - B^{\alpha\prime} \\ &- \frac{\delta\nu}{\nu}\delta e^{\alpha\prime} + 2\frac{\delta\nu}{\nu}\left(A^{,\alpha} - B^{\alpha\prime}\right) - \left(\delta e^{\beta} + \frac{\delta\nu}{\nu}e^{\beta}\right)\left(B_{\beta}^{-|\alpha} - B^{\alpha}_{|\beta} + 2C^{\alpha\prime}_{\beta}\right) + \delta e^{\beta}\delta e^{\alpha}_{-|\beta} \\ &+ 2e^{\beta}\delta e^{\gamma}\left(2C^{\alpha}_{\beta|\gamma} - C_{\beta\gamma}^{-|\alpha}\right) + A'B^{\alpha} - 2A_{,\beta}C^{\alpha\beta} + 2C^{\alpha}_{\beta}B^{\beta\prime} - 2B^{\alpha}A_{,\beta}e^{\beta} + 4C^{\alpha\gamma}\left(B_{[\beta|\gamma]} + C'_{\beta\gamma}\right)e^{\beta} \\ &- 2C^{\alpha}_{\delta}\left(2C^{\delta}_{\beta|\gamma} - C^{\gamma}_{\beta\gamma}^{-|\delta}\right)e^{\beta}e^{\gamma} + B^{\alpha}\left(B_{\beta|\gamma} + C'_{\beta\gamma}\right)e^{\beta}e^{\gamma}\right\} = 0, \quad (155)$$

where we used the null equation in eq. (153). To the background order eqs. (153-155) give

$$e^{\alpha}e_{\alpha} = 1, \quad \nu \propto a^{-1}, \quad e^{\alpha\prime} = e^{\beta}e^{\alpha}{}_{|\beta}.$$
 (156)

Using eqs. (152,153), eq. (154) becomes

$$\frac{d}{dy}\left(\frac{\delta\nu}{\nu}+A\right) - A_{,\alpha}e^{\alpha} + \left(B_{\alpha|\beta}+C_{\alpha\beta}'\right)e^{\alpha}e^{\beta}$$
$$= -\frac{\delta\nu}{\nu}\frac{\delta\nu'}{\nu} - \frac{a'}{a}\frac{\delta\nu^2}{\nu^2} - 2\frac{\delta\nu}{\nu}\left(A'+2\frac{a'}{a}A\right) + \frac{\delta\nu_{,\alpha}}{\nu}\delta e^{\alpha} + 2\frac{\delta\nu}{\nu}A_{,\alpha}e^{\alpha}$$

$$+2\delta e^{\alpha}A_{,\alpha} + 4\frac{a'}{a}e^{\alpha}\delta e_{\alpha}A - 2e^{\alpha}\delta e^{\beta}\left(B_{(\alpha|\beta)} + C'_{\alpha\beta}\right) \\ + \left[2A\left(B_{\alpha|\beta} + C'_{\alpha\beta} + 2\frac{a'}{a}C_{\alpha\beta}\right) + B_{\gamma}\left(2C^{\gamma}_{\alpha|\beta} - C_{\alpha\beta}^{\ |\gamma}\right)\right]e^{\alpha}e^{\beta} - \frac{a'}{a}\left(4A^{2} - B^{\alpha}B_{\alpha}\right) \\ -2\left(2AA_{,\alpha} - 2\frac{a'}{a}AB_{\alpha} + B_{\beta}C^{\beta\prime}_{\alpha} + B^{\beta}B_{[\alpha|\beta]}\right)e^{\alpha} + 2AA' + A_{,\alpha}B^{\alpha} - B^{\alpha}\left(B'_{\alpha} + \frac{a'}{a}B_{\alpha}\right) \\ \equiv N_{\nu}.$$

$$(157)$$

Thus, we have

$$\left(\frac{\delta\nu}{\nu} + A\right) \Big|_{E}^{O} = \int_{E}^{O} \left[A_{,\alpha}e^{\alpha} - \left(B_{\alpha|\beta} + C_{\alpha\beta}' \right)e^{\alpha}e^{\beta} + N_{\nu} \right] dy, \tag{158}$$

where the integral is along the ray's null-geodesic path from E the emitted event at the intersection of the ray and the last scattering surface to O the observed event here and now.

The temperatures of the CMB at two different points (O and E) along a single null-geodesic ray in a given observational direction are [27,17],

$$\frac{\tilde{T}_O}{\tilde{T}_E} \equiv \frac{1}{1+\tilde{z}} \equiv \frac{(\tilde{k}^a \tilde{u}_a)_O}{(\tilde{k}^b \tilde{u}_b)_E},\tag{159}$$

where \tilde{u}_a at O and E are the local four-velocities of the observer and the emitter, respectively. Thus, \tilde{u}_a should be considered as the one based on the energy-frame which sets $\tilde{q}_a \equiv 0$; or equivalently in a general frame vector which absorbs the flux term to the frame vector to the second-order.

Using eqs. (53,150) we have

$$\tilde{k}^{a}\tilde{u}_{a} = -\nu\left[1 + \frac{\delta\nu}{\nu} + A + \left(V_{\alpha}^{E} - B_{\alpha}\right)e^{\alpha} + \frac{\delta\nu}{\nu}A + \delta e^{\alpha}\left(V_{\alpha}^{E} - B_{\alpha}\right) + \left(AB_{\alpha} + 2C_{\alpha\beta}V^{E\beta}\right)e^{\alpha} + \frac{1}{2}V^{E\alpha}V_{\alpha}^{E} - \frac{1}{2}A^{2}\right].$$
(160)

We have denoted the energy-frame nature by replacing V_{α} to V_{α}^{E} ; if we consider eq. (86), eq. (160) in this form is valid in the general frame. Since the calculations in the rest of this paper are based on the normal-frame vector we use eq. (88) to derive the result in the normal-frame. We have

$$\tilde{k}^{a}\tilde{u}_{a} = -\nu \left[1 + \frac{\delta\nu}{\nu} + A + \frac{1}{\mu + p}Q_{\alpha}e^{\alpha} - \frac{\left(\delta\mu + \delta p\right)Q_{\alpha} + \Pi_{\alpha\beta}Q^{\beta}}{(\mu + p)^{2}}e^{\alpha} + \frac{\delta\nu}{\nu}A + \delta e^{\alpha}\frac{Q_{\alpha}}{\mu + p} + \frac{1}{2}\left(B_{\alpha} + \frac{Q_{\alpha}}{\mu + p}\right)\left(B^{\alpha} + \frac{Q^{\alpha}}{\mu + p}\right) - \frac{1}{2}A^{2} \right].$$
(161)

Using eq. (159) we have

$$\left(1 - \frac{T_E}{T_O} \frac{\nu_O}{\nu_E}\right) \left(1 + \frac{\delta T_E}{T_E}\right)$$

$$+ \frac{\delta T}{T} \Big|_E^O = \frac{T_E}{T_O} \frac{\nu_O}{\nu_E} \left[\frac{\delta \nu}{\nu} + A + \frac{1}{\mu + p} Q_\alpha e^\alpha - \frac{(\delta \mu + \delta p) Q_\alpha + \Pi_{\alpha\beta} Q^\beta}{(\mu + p)^2} e^\alpha + \frac{\delta \nu}{\nu} A + \delta e^\alpha \frac{Q_\alpha}{\mu + p} \right]$$

$$+ \frac{1}{2} \left(B_\alpha + \frac{Q_\alpha}{\mu + p}\right) \left(B^\alpha + \frac{Q^\alpha}{\mu + p}\right) - \frac{1}{2} A^2 \Big|_E^O \times \left[1 - \left(\frac{\delta \nu}{\nu} + A + \frac{1}{\mu + p} Q_\gamma e^\gamma\right) + \frac{\delta T}{T}\right]\Big|_E$$

$$= \frac{T_E}{T_O} \frac{\nu_O}{\nu_E} \left(\frac{\delta \nu}{\nu} + A + \frac{1}{\mu + p} Q_\alpha e^\alpha\right)\Big|_E^O + N_T,$$

$$(162)$$

where $\frac{\delta T}{T}|_{E}^{O} \equiv \frac{\delta T}{T}|_{O} - \frac{\delta T}{T}|_{E}$ and $\frac{\delta T}{T}|_{E} \equiv \frac{\delta T}{T}$ at E. Thus, if we take $T_{O}/T_{E} = \nu_{O}/\nu_{E}$, eqs. (162,158) give

$$\frac{\delta T}{T}\Big|_{O} = \frac{\delta T}{T}\Big|_{E} + \frac{1}{\mu + p}Q_{\alpha}e^{\alpha}\Big|_{E}^{O} + \int_{E}^{O}\left[A_{,\alpha}e^{\alpha} - \left(B_{\alpha|\beta} + C_{\alpha\beta}'\right)e^{\alpha}e^{\beta} + N_{\nu}\right]dy + N_{T}.$$
(163)

In the large angular scale we are considering (larger than the horizon size at the last scattering era) the detailed dynamics at last scattering is not important. The physical processes of last scattering are important in the small angular scale where we need to solve the Boltzmann equation for the photon distribution function, see §IV G.

G. Boltzmann equation

1. Covariant equations

The relativistic Boltzmann equation is [28]

$$\frac{d}{d\lambda}\tilde{f} = \frac{dx^a}{d\lambda}\frac{\partial\tilde{f}}{\partial x^a} + \frac{d\tilde{p}^a}{d\lambda}\frac{\partial\tilde{f}}{\partial\tilde{p}^a} = \tilde{p}^a\frac{\partial\tilde{f}}{\partial x^a} - \tilde{\Gamma}^a_{bc}\tilde{p}^b\tilde{p}^c\frac{\partial\tilde{f}}{\partial\tilde{p}^a} = \tilde{C}[\tilde{f}],\tag{164}$$

where $\tilde{f}(x^a, \tilde{p}^b)$ is a distribution function with the phase space variables x^a and $\tilde{p}^a \equiv dx^a/d\lambda$, and $\tilde{C}[\tilde{f}]$ is the collision term. The energy-momentum tensor of the collisionless (or collisional) component is

$$\tilde{T}_{ab}^{(c)} = \int 2\theta(\tilde{p}^0)\delta(\tilde{p}^c\tilde{p}_c + m^2)\tilde{p}_a\tilde{p}_b\tilde{f}\frac{d^4\tilde{p}_{0123}}{\sqrt{-\tilde{g}}} = \int 2\theta(\tilde{p}^0)\delta(\tilde{p}^c\tilde{p}_c + m^2)\tilde{p}_a\tilde{p}_b\tilde{f}\sqrt{-\tilde{g}}d^4\tilde{p}^{0123},$$
(165)

where

$$\delta(\tilde{p}^c \tilde{p}_c + m^2) = \delta(\tilde{g}_{00} \tilde{p}^0 \tilde{p}^0 + 2\tilde{g}_{0\alpha} \tilde{p}^0 \tilde{p}^\alpha + \tilde{g}_{\alpha\beta} \tilde{p}^\alpha \tilde{p}^\beta + m^2) = \frac{\delta(\text{mass shell})}{|2\tilde{g}_{00} \tilde{p}^0 + 2\tilde{g}_{0\alpha} \tilde{p}^\alpha|} = \frac{\delta(\text{mass shell})}{2|\tilde{p}_0|}.$$
 (166)

Thus, after integrating over \tilde{p}^0 , we have

$$\tilde{T}_{ab}^{(c)} = \int \frac{\sqrt{-g} d^3 \tilde{p}^{123}}{|\tilde{p}_0|} \tilde{p}_a \tilde{p}_b \tilde{f},$$
(167)

with the mass-shell condition $\tilde{p}^a \tilde{p}_a + m^2 = 0$.

2. Perturbed equations

Under our metric, using \tilde{p}^a as the phase space variable, we have

$$\begin{split} \tilde{p}^{0}\tilde{f}' + \tilde{p}^{\alpha}\tilde{f}_{,\alpha} - \left\{ \frac{a'}{a} \left(\tilde{p}^{0}\tilde{p}^{0} + g^{(3)}_{\alpha\beta}\tilde{p}^{\alpha}\tilde{p}^{\beta} \right) + A'\tilde{p}^{0}\tilde{p}^{0} + 2\left(A_{,\alpha} - \frac{a'}{a}B_{\alpha}\right)\tilde{p}^{0}\tilde{p}^{\alpha} \\ + \left(-2\frac{a'}{a}g^{(3)}_{\alpha\beta}A + B_{\alpha|\beta} + C'_{\alpha\beta} + 2\frac{a'}{a}C_{\alpha\beta}\right)\tilde{p}^{\alpha}\tilde{p}^{\beta} \\ + \left(-2AA' - A_{,\alpha}B^{\alpha} + B^{\alpha}B'_{\alpha} + \frac{a'}{a}B^{\alpha}B_{\alpha}\right)\tilde{p}^{0}\tilde{p}^{0} + 2\left(-2AA_{,\alpha} + 2\frac{a'}{a}AB_{\alpha} - B_{\beta}C^{\beta'}_{\alpha} + B^{\beta}B_{[\beta|\alpha]}\right)\tilde{p}^{0}\tilde{p}^{\alpha} \\ + \left[\frac{a'}{a}g^{(3)}_{\alpha\beta}\left(4A^{2} - B^{\gamma}B_{\gamma}\right) - 2A\left(B_{\alpha|\beta} + C'_{\alpha\beta} + 2\frac{a'}{a}C_{\alpha\beta}\right) - B_{\gamma}\left(2C^{\gamma}_{\alpha|\beta} - C_{\alpha\beta}^{-\beta}\right)\right]\tilde{p}^{\alpha}\tilde{p}^{\beta} \right\}\frac{\partial\tilde{f}}{\partial\tilde{p}^{0}} \\ - \left[\left(2\frac{a'}{a}\tilde{p}^{0}\tilde{p}^{\alpha} + \Gamma^{(3)\alpha}_{\ \beta\gamma}\tilde{p}^{\beta}\tilde{p}^{\gamma} \right) + \left(A^{,\alpha} - B^{\alpha'} - \frac{a'}{a}B^{\alpha}\right)\tilde{p}^{0}\tilde{p}^{0} + \left(B_{\beta}^{-\alpha} - B^{\alpha}_{\ -\beta} + 2C^{\alpha'}_{\beta}\right)\tilde{p}^{0}\tilde{p}^{\beta} \\ + \left(\frac{a'}{a}g^{(3)}_{\beta\gamma}B^{\alpha} + 2C^{\alpha}_{\beta|\gamma} - C_{\beta\gamma}^{-\alpha}\right)\tilde{p}^{\beta}\tilde{p}^{\gamma} \right]\frac{\partial\tilde{f}}{\partial\tilde{p}^{\alpha}} = \tilde{C}[\tilde{f}]. \end{split}$$

$$(168)$$

As the phase space variable it is convenient to use $(q,\,\gamma^\alpha)$ introduced as

$$q \equiv a\sqrt{a^{2}(\tilde{p}^{0})^{2} - m^{2}} \left[1 + \frac{a^{2}(\tilde{p}^{0})^{2}}{a^{2}(\tilde{p}^{0})^{2} - m^{2}} \left(A + \frac{1}{2}B^{\alpha}B_{\alpha} \right) - \frac{1}{2} \frac{a^{4}(\tilde{p}^{0})^{4}}{[a^{2}(\tilde{p}^{0})^{2} - m^{2}]^{2}} A^{2} \right],$$

$$\gamma^{\alpha} \equiv \frac{a}{\sqrt{a^{2}(\tilde{p}^{0})^{2} - m^{2}}} \left[\tilde{p}^{\alpha} - \frac{a^{2}(\tilde{p}^{0})^{2}}{a^{2}(\tilde{p}^{0})^{2} - m^{2}} A \tilde{p}^{\alpha} - B^{\alpha} \tilde{p}^{0} + C^{\alpha}_{\beta} \tilde{p}^{\beta} + \frac{3}{2} \frac{a^{4}(\tilde{p}^{0})^{4}}{[a^{2}(\tilde{p}^{0})^{2} - m^{2}]^{2}} A^{2} \tilde{p}^{\alpha} + \frac{a^{2}(\tilde{p}^{0})^{2}}{a^{2}(\tilde{p}^{0})^{2} - m^{2}} \left(AB^{\alpha} \tilde{p}^{0} - AC^{\alpha}_{\beta} \tilde{p}^{\beta} - \frac{1}{2} B^{\beta} B_{\beta} \tilde{p}^{\alpha} \right) + C^{\alpha}_{\beta} B^{\beta} \tilde{p}^{0} - \frac{1}{2} C^{\gamma}_{\beta} C^{\alpha}_{\gamma} \tilde{p}^{\beta} \right],$$

$$\tilde{p}^{0} = \frac{1}{a^{2}} \sqrt{q^{2} + m^{2}a^{2}} \left(1 - A + \frac{3}{2} A^{2} - \frac{1}{2} B^{\alpha} B_{\alpha} \right),$$

$$\tilde{p}^{\alpha} = \frac{1}{a^{2}} \left[q\gamma^{\alpha} + \sqrt{q^{2} + m^{2}a^{2}} B^{\alpha} - qC^{\alpha}_{\beta} \gamma^{\beta} - \sqrt{q^{2} + m^{2}a^{2}} \left(AB^{\alpha} + 2C^{\alpha}_{\beta} B^{\beta} \right) + \frac{3}{2} qC^{\gamma}_{\beta} C^{\alpha}_{\gamma} \gamma^{\beta} \right],$$
(169)

where γ^{α} is based on $g_{\alpha\beta}^{(3)}$. The mass-shell condition gives

$$(\tilde{p}^{0})^{2} - g^{(3)}_{\alpha\beta}\tilde{p}^{\alpha}\tilde{p}^{\beta} - m^{2}/a^{2} + 2A(\tilde{p}^{0})^{2} + 2B_{\alpha}\tilde{p}^{\alpha}\tilde{p}^{0} - 2C_{\alpha\beta}\tilde{p}^{\alpha}\tilde{p}^{\beta} = 0,$$
(170)

and we can show $\gamma^{\alpha}\gamma_{\alpha} = 1$. The Boltzmann equation becomes

$$\begin{split} \tilde{f}' + \frac{q}{\sqrt{q^2 + m^2 a^2}} \left(\gamma^{\alpha} \delta f_{,\alpha} - \Gamma^{(3)\alpha}_{\ \beta\gamma} \gamma^{\beta} \gamma^{\gamma} \frac{\partial \delta f}{\partial \gamma^{\alpha}} \right) - \left[\frac{\sqrt{q^2 + m^2 a^2}}{q} \gamma^{\alpha} A_{,\alpha} + \left(B_{\alpha|\beta} + C'_{\alpha\beta} \right) \gamma^{\alpha} \gamma^{\beta} \right] q \frac{\partial \tilde{f}}{\partial q} \\ &= -\frac{1}{\sqrt{q^2 + m^2 a^2}} \left(q A \gamma^{\alpha} + \sqrt{q^2 + m^2 a^2} B^{\alpha} - q C^{\alpha}_{\beta} \gamma^{\beta} \right) \delta f_{,\alpha} \\ &- \left\{ \frac{\sqrt{q^2 + m^2 a^2}}{q} \left[A_{,\alpha} \left(A \gamma^{\alpha} + C^{\alpha}_{\beta} \gamma^{\beta} \right) - B^{\beta} B_{\beta|\alpha} \gamma^{\alpha} \right] + 2 C^{\beta}_{\gamma} \left(C'_{\alpha\beta} + B_{(\alpha|\beta)} \right) \gamma^{\alpha} \gamma^{\gamma} \\ &+ B_{\gamma} \left(2 C^{\gamma}_{\alpha|\beta} - C_{\alpha\beta}^{\ |\gamma} \right) \gamma^{\alpha} \gamma^{\beta} \right\} q \frac{\partial f}{\partial q} \\ &+ \left[\Gamma^{(3)\alpha}_{\ \beta\gamma} \left(\frac{q}{\sqrt{q^2 + m^2 a^2}} A \gamma^{\beta} \gamma^{\gamma} + B^{\beta} \gamma^{\gamma} - \frac{q}{\sqrt{q^2 + m^2 a^2}} C^{\gamma}_{\delta} \gamma^{\beta} \gamma^{\delta} \right) + \frac{\sqrt{q^2 + m^2 a^2}}{q} \left(A^{,\alpha} - A_{,\beta} \gamma^{\beta} \gamma^{\alpha} \right) \\ &+ \left(B_{\beta}^{\ |\alpha} + C^{\alpha}_{\beta} \right) \gamma^{\beta} - \left(B_{\beta|\gamma} + C'_{\beta\gamma} \right) \gamma^{\beta} \gamma^{\gamma} \gamma^{\alpha} + \frac{q}{\sqrt{q^2 + m^2 a^2}} \left(C^{\alpha}_{\beta|\gamma} - C^{\ |\alpha|}_{\beta\gamma} \right) \gamma^{\beta} \gamma^{\gamma} \right] \frac{\partial \delta f}{\partial \gamma^{\alpha}} \\ &+ \frac{a^2}{\sqrt{q^2 + m^2 a^2}} \left(1 + A - \frac{1}{2} A^2 + \frac{1}{2} B^{\alpha} B_{\alpha} \right) \tilde{C}[\tilde{f}] \\ &\equiv N_c. \end{split}$$

$$\tag{171}$$

For convenience we located the collision term in N_c . The energy-momentum tensor becomes

$$\tilde{T}_{ab}^{(c)} = \frac{1}{a^2} \int \tilde{p}_a \tilde{p}_b \tilde{f} \frac{q^2 dq d\Omega_q}{\sqrt{q^2 + m^2 a^2}}.$$
(172)

Thus, using $\int \gamma_{\alpha} \gamma_{\beta} d\Omega_q = \frac{1}{3} g_{\alpha\beta}^{(3)}$, we have

$$\tilde{T}_{00}^{(c)} = \frac{1}{a^2} \int \sqrt{q^2 + m^2 a^2} q^2 dq d\Omega_q \left[f(1+2A) + \delta f(1+2A) + \left(1 + \frac{1}{3} \frac{q^2}{q^2 + m^2 a^2}\right) B^\alpha B_\alpha f \right. \\ \left. + \frac{2q}{\sqrt{q^2 + m^2 a^2}} B_\alpha \gamma^\alpha \delta f \right], \\ \tilde{T}_{\alpha 0}^{(c)} = -\frac{1}{a^2} \int \frac{q^4 dq d\Omega_q}{\sqrt{q^2 + m^2 a^2}} \left\{ \frac{1}{3} B_\alpha f + \left[\frac{\sqrt{q^2 + m^2 a^2}}{q} \left(\gamma_\alpha + A\gamma_\alpha + C_{\alpha\beta} \gamma^\beta\right) + B_\beta \gamma^\beta \gamma_\alpha \right] \delta f \right\}, \\ \tilde{T}_{\alpha\beta}^{(c)} = \frac{1}{a^2} \int \frac{q^4 dq d\Omega_q}{\sqrt{q^2 + m^2 a^2}} \left(\frac{1}{3} g_{\alpha\beta}^{(3)} f + \delta f \gamma_\alpha \gamma_\beta + \frac{2}{3} f C_{\alpha\beta} + 2\delta f \gamma_{(\alpha} C_{\beta)\gamma} \gamma^\gamma \right).$$
(173)

From eq. (79) the fluid quantities become:

$$\mu^{(c)} + \delta\mu^{(c)} = \frac{1}{a^4} \int \sqrt{q^2 + m^2 a^2} q^2 dq d\Omega_q \left(f + \delta f\right),$$

$$p^{(c)} + \delta p^{(c)} = \frac{1}{3a^4} \int \frac{q^4 dq d\Omega_q}{\sqrt{q^2 + m^2 a^2}} \left(f + \delta f\right),$$

$$Q^{(c)}_{\alpha} = \frac{1}{a^4} \int q^3 dq d\Omega_q \left(\gamma_{\alpha} + C_{\alpha\beta}\gamma^{\beta}\right) \delta f,$$

$$\Pi^{(c)}_{\alpha\beta} = \frac{1}{a^4} \int \frac{q^4 dq d\Omega_q}{\sqrt{q^2 + m^2 a^2}} \left(\gamma_{\alpha}\gamma_{\beta} - \frac{1}{3}g^{(3)}_{\alpha\beta} + 2\gamma_{(\alpha}C_{\beta)\gamma}\gamma^{\gamma} - \frac{2}{3}C_{\alpha\beta}\right) \delta f.$$
(174)

If we have multiple components each described by the Boltzmann equation, all equations in this subsection remain valid for any component with \tilde{f} replaced by $\tilde{f}_{(i)}$, etc., and the total (collective) fluid quantities as the sum of the individual ones.

V. DECOMPOSITION

A. Three perturbation types

We decompose the perturbation variables as follow:

$$A \equiv \alpha,$$

$$B_{\alpha} \equiv \beta_{,\alpha} + B_{\alpha}^{(v)},$$

$$C_{\alpha\beta} \equiv \varphi g_{\alpha\beta}^{(3)} + \gamma_{,\alpha|\beta} + C_{(\alpha|\beta)}^{(v)} + C_{\alpha\beta}^{(t)},$$

$$Q_{\alpha} \equiv Q_{,\alpha} + Q_{\alpha}^{(v)} \equiv (\mu + p) \left(-v_{,\alpha} + v_{\alpha}^{(v)} \right),$$

$$\Pi_{\alpha\beta} \equiv \frac{1}{a^{2}} \left(\Pi_{,\alpha|\beta} - \frac{1}{3} g_{\alpha\beta}^{(3)} \Delta \Pi \right) + \frac{1}{a} \Pi_{(\alpha|\beta)}^{(v)} + \Pi_{\alpha\beta}^{(t)},$$
(175)

with the properties:

$$B^{(v)\alpha}_{\ |\alpha} \equiv 0, \quad C^{(v)\alpha}_{\ |\alpha} \equiv 0, \quad v^{(v)\alpha}_{\ |\alpha} \equiv 0, \quad \Pi^{(v)\alpha}_{\ |\alpha} \equiv 0,$$

$$C^{(t)\alpha}_{\ \alpha} \equiv 0, \quad \Pi^{(t)\alpha}_{\ \alpha} \equiv 0, \quad C^{(t)\beta}_{\ \alpha|\beta} \equiv 0, \quad \Pi^{(t)\beta}_{\ \alpha|\beta} \equiv 0.$$
(176)

 $\mu + p$ appearing in the decomposition of Q_{α} is assumed to be the background order quantity. The vector- and the tensor-type perturbations are denoted by superscripts (v) and (t), respectively. We assume all these variables are based on $g_{\alpha\beta}^{(3)}$. The decomposed variables also can be expressed in terms of the original variables. For example, we have $\beta = \Delta^{-1} \nabla^{\alpha} B_{\alpha}$ and $B_{\alpha}^{(v)} = B_{\alpha} - \nabla_{\alpha} \Delta^{-1} \nabla^{\beta} B_{\beta}$, etc, where ∇_{α} means $\nabla_{\alpha}^{(3)}$. For the fluid quantities we have

$$Q = \Delta^{-1} \nabla^{\alpha} Q_{\alpha},$$

$$Q_{\alpha}^{(v)} = Q_{\alpha} - \nabla_{\alpha} \Delta^{-1} \nabla^{\beta} Q_{\beta},$$

$$\Pi = \frac{3}{2} a^{2} \left(\Delta + \frac{1}{2} R^{(3)} \right)^{-1} \Delta^{-1} \nabla^{\alpha} \nabla^{\beta} \Pi_{\alpha\beta},$$

$$\Pi_{\alpha}^{(v)} = 2a \left(\Delta + \frac{1}{3} R^{(3)} \right)^{-1} \left(\nabla^{\beta} \Pi_{\alpha\beta} - \nabla_{\alpha} \Delta^{-1} \nabla^{\beta} \nabla^{\gamma} \Pi_{\beta\gamma} \right),$$

$$\Pi_{\alpha\beta}^{(t)} = \Pi_{\alpha\beta} - \frac{3}{2} \left(\nabla_{\alpha} \nabla_{\beta} - \frac{1}{3} g_{\alpha\beta}^{(3)} \Delta \right) \left(\Delta + \frac{1}{2} R^{(3)} \right)^{-1} \Delta^{-1} \nabla^{\gamma} \nabla^{\delta} \Pi_{\gamma\delta}$$

$$-2 \nabla_{(\alpha} \left(\Delta + \frac{1}{3} R^{(3)} \right)^{-1} \left(\nabla^{\gamma} \Pi_{\beta\gamma} - \nabla_{\beta} \Delta^{-1} \nabla^{\gamma} \nabla^{\delta} \Pi_{\gamma\delta} \right).$$
(177)

We introduce

$$\chi \equiv a \left(\beta + a \dot{\gamma}\right), \quad \Psi_{\alpha}^{(v)} \equiv B_{\alpha}^{(v)} + a \dot{C}_{\alpha}^{(v)}, \tag{178}$$

and let

$$\kappa \equiv \delta K.$$
 (179)

In the multi-component situation we have eq. (75). For the individual component we have

$$Q_{(i)\alpha} \equiv (\mu_{(i)} + p_{(i)}) \left(-v_{(i),\alpha} + v_{(i)\alpha}^{(v)} \right),$$

$$\Pi_{(i)\alpha\beta} \equiv \frac{1}{a^2} \left(\Pi_{(i),\alpha|\beta} - \frac{1}{3} g^{(3)}_{\alpha\beta} \Delta \Pi_{(i)} \right) + \frac{1}{a} \Pi^{(v)}_{(i)(\alpha|\beta)} + \Pi^{(t)}_{(i)\alpha\beta},$$
(180)

with

$$v_{(i)}^{(v)\alpha}{}_{|\alpha} \equiv 0, \quad \Pi_{(i)}^{(v)\alpha}{}_{|\alpha} \equiv 0, \quad \Pi_{(i)\alpha}^{(t)\alpha} \equiv 0, \quad \Pi_{(i)\alpha|\beta}^{(t)\beta} \equiv 0,$$
 (181)

and

$$\delta I_{(i)\alpha} \equiv \delta I_{(i),\alpha} + \delta I_{(i)\alpha}^{(v)}, \quad \delta I_{(i)|\alpha}^{(v)\alpha} \equiv 0.$$
(182)

The definitions for the scalar-type perturbation variables are introduced to match with our notation used in the linear analysis [7,12,29]; compared with our previous definitions in the linear theory our v and $v_{(i)}$ correspond to v/k and $v_{(i)}/k$ in [29] where k is the wave number. These are the notations introduced by Bardeen in 1988 [7]. A complete set of equations written separately for the three perturbation types will be presented in equations (195)-(210) where the quadratic combinations of the linear-order variables contribute to the second-order perturbations. Thus, to the second-order the three perturbation types couple each other through quadratic combinations of the linear-order terms. If needed we may decompose the perturbed order quantities explicitly as in eq. (50)

$$\alpha \equiv \alpha^{(1)} + \alpha^{(2)}, \quad \varphi \equiv \varphi^{(1)} + \varphi^{(2)}, \tag{183}$$

etc.

B. Background equations

To the background order, eqs. (100, 102, 104, 112) give

$$H^{2} = \frac{8\pi G}{3}\mu - \frac{K}{a^{2}} + \frac{\Lambda}{3},$$
(184)

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \left(\mu + 3p\right) + \frac{\Lambda}{3},\tag{185}$$

$$\dot{\mu} + 3H(\mu + p) = 0, \tag{186}$$

$$\phi + 3H\phi + V_{,\phi} = 0. \tag{187}$$

In the multi-component situations from eqs. (106, 119) we have

$$\dot{\mu}_{(i)} + 3H\left(\mu_{(i)} + p_{(i)}\right) = -\frac{1}{a}I_{(i)0},\tag{188}$$

$$\ddot{\phi}_{(i)} + 3H\dot{\phi}_{(i)} + V_{,\phi_{(i)}} = 0, \tag{189}$$

with

$$\mu_{(\phi)} = \frac{1}{2} \sum_{k} \dot{\phi}_{(k)}^{2} + V, \quad p_{(\phi)} = \frac{1}{2} \sum_{k} \dot{\phi}_{(k)}^{2} - V, \tag{190}$$

which follow from eq. (121).

In the generalized gravity considered in §IV D, eqs. (184,185) remain valid by replacing the fluid quantities to the effective one in eq. (130):

$$\mu^{\text{(eff)}} = \frac{1}{8\pi GF} \left[\mu + \frac{1}{2} g_{IJ} \dot{\phi}^{I} \dot{\phi}^{J} - \frac{1}{2} \left(f - RF - 2V \right) - 3H\dot{F} \right],$$

$$p^{\text{(eff)}} = \frac{1}{8\pi GF} \left[p + \frac{1}{2} g_{IJ} \dot{\phi}^{I} \dot{\phi}^{J} + \frac{1}{2} \left(f - RF - 2V \right) + \ddot{F} + 2H\dot{F} \right].$$
 (191)

For the equation of motion, eq. (128) gives

$$\ddot{\phi}^{I} + 3H\dot{\phi}^{I} - \frac{1}{2}g^{IJ}\left(f_{,J} - 2V_{,J}\right) + \Gamma^{I}_{JK}\dot{\phi}^{J}\dot{\phi}^{K} + \Gamma^{I} = 0.$$
(192)

The null-geodesic equations are presented in eq. (156). The Boltzmann equation in eq. (171) gives

$$f' = \frac{a^2}{\sqrt{q^2 + m^2 a^2}} C[f],$$
(193)

and from eq. (174) we have

$$\mu^{(c)} = \frac{1}{a^4} \int \sqrt{q^2 + m^2 a^2} q^2 dq d\Omega_q f, \quad p^{(c)} = \frac{1}{3a^4} \int \frac{q^4 dq d\Omega_q}{\sqrt{q^2 + m^2 a^2}} f.$$
(194)

C. Decomposed equations

We summarize a complete set of equations necessary to analyse each perturbation type. We decompose the perturbation variables according to equation (175). Algebraic manipulations are made which can be recognized by examining the right hand side of the following equations.

The scalar-type perturbation:

$$\kappa - 3H\alpha + 3\dot{\varphi} + \frac{\Delta}{a^2}\chi = N_0,\tag{195}$$

$$4\pi G\delta\mu + H\kappa + \frac{\Delta + 3K}{a^2}\varphi = \frac{1}{4}N_1,\tag{196}$$

$$\kappa + \frac{\Delta + 3K}{a^2} \chi - 12\pi G(\mu + p)av = \frac{3}{2} \Delta^{-1} \nabla^{\alpha} N_{2\alpha} \equiv N_2^{(s)},$$
(197)

$$\dot{\kappa} + 2H\kappa - 4\pi G \left(\delta\mu + 3\delta p\right) + \left(3\dot{H} + \frac{\Delta}{a^2}\right)\alpha = N_3,\tag{198}$$

$$\dot{\chi} + H\chi - \varphi - \alpha - 8\pi G\Pi = \frac{3}{2}a^2 \left(\Delta + 3K\right)^{-1} \Delta^{-1} \nabla^{\alpha} \nabla_{\beta} N_{4\alpha}^{\ \beta} \equiv N_4^{(s)},\tag{199}$$

$$\delta\dot{\mu} + 3H\left(\delta\mu + \delta p\right) - \left(\mu + p\right)\left(\kappa - 3H\alpha + \frac{1}{a}\Delta v\right) = N_5,\tag{200}$$

$$\frac{[a^4(\mu+p)v]}{a^4(\mu+p)} - \frac{1}{a}\alpha - \frac{1}{a(\mu+p)}\left(\delta p + \frac{2}{3}\frac{\Delta+3K}{a^2}\Pi\right) = -\frac{1}{\mu+p}\Delta^{-1}\nabla^{\alpha}N_{6\alpha} \equiv N_6^{(s)},$$
(201)

$$\delta\dot{\mu}_{(i)} + 3H\left(\delta\mu_{(i)} + \delta p_{(i)}\right) - \left(\mu_{(i)} + p_{(i)}\right)\left(\kappa - 3H\alpha + \frac{1}{a}\Delta v_{(i)}\right) + \frac{1}{a}\delta I_{(i)0} = N_{5(i)},\tag{202}$$

$$\frac{\left[a^{4}(\mu_{(i)}+p_{(i)})v_{(i)}\right]^{\cdot}}{a^{4}(\mu_{(i)}+p_{(i)})} - \frac{1}{a}\alpha - \frac{1}{a(\mu_{(i)}+p_{(i)})}\left(\delta p_{(i)} + \frac{2}{3}\frac{\Delta+3K}{a^{2}}\Pi_{(i)} - \delta I_{(i)}\right) \\
= -\frac{1}{\mu_{(i)}+p_{(i)}}\Delta^{-1}\nabla^{\alpha}N_{6(i)\alpha} \equiv N_{6(i)}^{(s)},$$
(203)

$$\delta\ddot{\phi} + 3H\delta\dot{\phi} - \frac{\Delta}{a^2}\delta\phi + V_{,\phi\phi}\delta\phi - \dot{\phi}\left(\kappa + \dot{\alpha}\right) - \left(2\ddot{\phi} + 3H\dot{\phi}\right)\alpha = N_{\phi} - \dot{\phi}N_0,\tag{204}$$

$$\delta\ddot{\phi}_{(i)} + 3H\delta\dot{\phi}_{(i)} - \frac{\Delta}{a^2}\delta\phi_{(i)} + \sum_k V_{,\phi_{(i)}\phi_{(k)}}\delta\phi_{(k)} - \dot{\phi}_{(i)}(\kappa + \dot{\alpha}) - \left(2\ddot{\phi}_{(i)} + 3H\dot{\phi}_{(i)}\right)\alpha = N_{\phi_{(i)}} - \dot{\phi}_{(i)}N_0.$$
(205)

The vector-type perturbation:

$$\frac{\Delta + 2K}{2a^2}\Psi_{\alpha}^{(v)} + 8\pi G(\mu + p)v_{\alpha}^{(v)} = \frac{1}{a}\left(N_{2\alpha} - \nabla_{\alpha}\Delta^{-1}\nabla^{\beta}N_{2\beta}\right) \equiv N_{2\alpha}^{(v)},\tag{206}$$

$$\dot{\Psi}_{\alpha}^{(v)} + 2H\Psi_{\alpha}^{(v)} - 8\pi G\Pi_{\alpha}^{(v)} = 2a\left(\Delta + 2K\right)^{-1} \left(\nabla_{\beta}N_{4\alpha}^{\ \beta} - \nabla_{\alpha}\Delta^{-1}\nabla^{\gamma}\nabla_{\beta}N_{4\gamma}^{\ \beta}\right) \equiv N_{4\alpha}^{(v)},\tag{207}$$

$$\frac{[a^4(\mu+p)v_{\alpha}^{(v)}]}{a^4(\mu+p)} + \frac{\Delta+2K}{2a^2}\frac{\Pi_{\alpha}^{(v)}}{\mu+p} = \frac{1}{\mu+p}\left(N_{6\alpha} - \nabla_{\alpha}\Delta^{-1}\nabla^{\beta}N_{6\beta}\right) \equiv N_{6\alpha}^{(v)},$$
(208)

$$\frac{\left[a^{4}(\mu_{(i)}+p_{(i)})v_{(i)\alpha}^{(v)}\right]^{\cdot}}{a^{4}(\mu_{(i)}+p_{(i)})} + \frac{\Delta+2K}{2a^{2}}\frac{\Pi_{(i)\alpha}^{(v)}}{\mu_{(i)}+p_{(i)}} - \frac{1}{a}\frac{\delta I_{(i)\alpha}^{(v)}}{\mu_{(i)}+p_{(i)}} \\
= \frac{1}{\mu_{(i)}+p_{(i)}}\left(N_{6(i)\alpha}-\nabla_{\alpha}\Delta^{-1}\nabla^{\beta}N_{6(i)\beta}\right) \equiv N_{6(i)\alpha}^{(v)}.$$
(209)

The tensor-type perturbation

$$\ddot{C}^{(t)}_{\alpha\beta} + 3H\dot{C}^{(t)}_{\alpha\beta} - \frac{\Delta - 2K}{a^2}C^{(t)}_{\alpha\beta} - 8\pi G\Pi^{(t)}_{\alpha\beta}$$

$$= N_{4\alpha\beta} - \frac{3}{2}\left(\nabla_{\alpha}\nabla_{\beta} - \frac{1}{3}g^{(3)}_{\alpha\beta}\Delta\right)(\Delta + 3K)^{-1}\Delta^{-1}\nabla^{\gamma}\nabla_{\delta}N^{\delta}_{4\gamma}$$

$$-2\nabla_{(\alpha}\left(\Delta + 2K\right)^{-1}\left(\nabla^{\gamma}N_{4\beta\gamma\gamma} - \nabla_{\beta}\Delta^{-1}\nabla^{\gamma}\nabla_{\delta}N^{\delta}_{4\gamma}\right) \equiv N^{(t)}_{4\alpha\beta}.$$
(210)

In order to derive eqs. (199,207,210) it is convenient to show

$$\frac{1}{a^{2}} \left(\nabla_{\alpha} \nabla_{\beta} - \frac{1}{3} g_{\alpha\beta}^{(3)} \Delta \right) \left(\dot{\chi} + H \chi - \varphi - \alpha - 8\pi G \Pi \right) + \frac{1}{a^{3}} \left(a^{2} \Psi_{(\alpha|\beta)}^{(v)} \right)^{\cdot} - 8\pi G \frac{1}{a} \Pi_{(\alpha|\beta)}^{(v)} \\
+ \ddot{C}_{\alpha\beta}^{(t)} + 3H \dot{C}_{\alpha\beta}^{(t)} - \frac{\Delta - 2K}{a^{2}} C_{\alpha\beta}^{(t)} - 8\pi G \Pi_{\alpha\beta}^{(t)} = N_{4\alpha\beta},$$
(211)

which follows from eq. (103). In our perturbative approach, the second-order perturbations are sourced by the quadratic combinations of all three-types of linear-order terms.

For the scalar field, from eq. (121), we have:

$$\delta\mu^{(\phi)} = \sum_{k} \left(\dot{\phi}_{(k)} \delta \dot{\phi}_{(k)} - \dot{\phi}_{(k)}^{2} \alpha + V_{,\phi_{(k)}} \delta \phi_{(k)} \right) + \delta\mu^{(q)},$$

$$\delta p^{(\phi)} = \sum_{k} \left(\dot{\phi}_{(k)} \delta \dot{\phi}_{(k)} - \dot{\phi}_{(k)}^{2} \alpha - V_{,\phi_{(k)}} \delta \phi_{(k)} \right) + \delta p^{(q)},$$

$$Q^{(\phi)} = -(\mu^{(\phi)} + p^{(\phi)}) v^{(\phi)} = -\frac{1}{a} \sum_{k} \dot{\phi}_{(k)} \delta \phi_{(k)} + \Delta^{-1} \nabla^{\alpha} Q^{(q)}_{\alpha},$$

$$Q^{(\phi,v)}_{\alpha} = (\mu^{(\phi)} + p^{(\phi)}) v^{(\phi,v)}_{\alpha} = Q^{(q)}_{\alpha} - \nabla_{\alpha} \Delta^{-1} \nabla^{\beta} Q^{(q)}_{\beta}.$$
(212)

The anisotropic pressure follows from eqs. (121,177).

In the generalized gravity theory in §IV D, eq. (128) gives

$$\begin{split} \delta\ddot{\phi}^{I} + 3H\delta\dot{\phi}^{I} - \frac{\Delta}{a^{2}}\delta\phi^{I} - 2\alpha\ddot{\phi}^{I} + \dot{\phi}^{I}\left(3\dot{\varphi} - \dot{\alpha} - 6H\alpha + \frac{\Delta}{a^{2}}\chi\right) \\ - \frac{1}{2}g^{IJ}\left[F_{,J}\delta R + (f_{,LJ} - 2V_{,LJ})\,\delta\phi^{L}\right] - \frac{1}{2}g^{IJ}_{,L}\delta\phi^{L}\left(f_{,J} - 2V_{,J}\right) \\ + 2\Gamma^{I}_{JK}\left(\dot{\phi}^{J}\delta\dot{\phi}^{K} - A\dot{\phi}^{J}\dot{\phi}^{K}\right) + \Gamma^{I}_{JK,L}\delta\phi^{L}\dot{\phi}^{J}\dot{\phi}^{K} + \delta\Gamma^{I} = N_{g}. \end{split}$$
(213)

Equations (195-201,206-208,210) remain valid even in the generalized gravity by replacing the fluid quantities to the effective ones. The decomposed effective fluid quantities follow from eqs. (130,177) as:

$$\delta\mu^{(\text{eff})} = \frac{1}{8\pi GF} \left[\delta\mu + g_{IJ} \left(\dot{\phi}^I \delta \dot{\phi}^J - \alpha \dot{\phi}^I \dot{\phi}^J \right) + \frac{1}{2} g_{IJ,L} \delta\phi^L \dot{\phi}^I \dot{\phi}^J - \frac{1}{2} \left(f_{,L} - 2V_{,L} \right) \delta\phi^L - 3H\delta \dot{F} + \left(\frac{1}{2}R + \frac{\Delta}{a^2} \right) \delta F + \left(6H\alpha - \frac{\Delta}{a^2} \chi - 3\dot{\varphi} \right) \dot{F} - 8\pi G \mu^{(\text{eff})} \delta F \right] + \delta\mu^{(\text{eff},q)},$$

$$\begin{split} \delta p^{(\text{eff})} &= \frac{1}{8\pi GF} \left[\delta p + g_{IJ} \left(\dot{\phi}^{I} \delta \dot{\phi}^{J} - \alpha \dot{\phi}^{I} \dot{\phi}^{J} \right) + \frac{1}{2} g_{IJ,L} \delta \phi^{L} \dot{\phi}^{I} \dot{\phi}^{J} + \frac{1}{2} \left(f_{,L} - 2V_{,L} \right) \delta \phi^{L} + \delta \ddot{F} + 2H \delta \dot{F} \\ &- \left(\frac{1}{2}R + \frac{2}{3} \frac{\Delta}{a^{2}} \right) \delta F - 2\alpha \ddot{F} - \left(\dot{\alpha} + 4H\alpha - \frac{2}{3} \frac{\Delta}{a^{2}} \chi - 2\dot{\varphi} \right) \dot{F} - 8\pi G p^{(\text{eff})} \delta F \right] + \delta p^{(\text{eff},q)}, \\ Q^{(\text{eff})} &= \frac{1}{8\pi GF} \left[Q - \frac{1}{a} g_{IJ} \dot{\phi}^{I} \delta \phi^{J} + \frac{1}{a} \left(-\delta \dot{F} + H \delta F \right) + \frac{1}{a} A \dot{F} \right] + Q^{(\text{eff},q)}, \\ Q^{(\text{eff})}_{\alpha} &= \frac{1}{8\pi GF} Q^{(v)}_{\alpha} + Q^{(\text{eff},v,q)}_{\alpha}, \\ \Pi^{(\text{eff})} &= \frac{1}{8\pi GF} \left(\Pi + \delta F - \chi \dot{F} \right) + \Pi^{(\text{eff},q)}, \\ \Pi^{(\text{eff},v)}_{\alpha} &= \frac{1}{8\pi GF} \left(\Pi^{(v)}_{\alpha} - \Psi^{(v)}_{\alpha} \dot{F} \right) + \Pi^{(\text{eff},v,q)}_{\alpha}, \\ \Pi^{(\text{eff},t)}_{\alpha\beta} &= \frac{1}{8\pi GF} \left(\Pi^{(t)}_{\alpha\beta} - \dot{C}^{(t)}_{\alpha\beta} \dot{F} \right) + \Pi^{(\text{eff},t,q)}_{\alpha\beta}, \end{split}$$
(214)

where the quadratic parts follow from eq. (177). As an example, from eqs. (210,214), the gravitational wave equation in generalized gravity becomes

$$\ddot{C}_{\alpha\beta}^{(t)} + \left(3H + \frac{\dot{F}}{F}\right)\dot{C}_{\alpha\beta}^{(t)} - \frac{\Delta - 2K}{a^2}C_{\alpha\beta}^{(t)} = \frac{1}{F}\Pi_{\alpha\beta}^{(t)} + 8\pi G\Pi_{\alpha\beta}^{(\text{eff},t,q)} + N_{4\alpha\beta}^{(t)}.$$
(215)

For the electromagnetic field we can decompose:

$$E_{\alpha} \equiv E_{,\alpha}^{(\mathrm{em})} + E_{\alpha}^{(v)}, \quad H_{\alpha} \equiv H_{,\alpha}^{(\mathrm{em})} + H_{\alpha}^{(v)}; \quad E_{\alpha}^{(v)|\alpha} \equiv 0 \equiv H_{\alpha}^{(v)|\alpha}.$$
(216)

The decomposed forms of fluid quantities can be read from eqs. (149,177). Similarly, for the null-geodesic equations we decompose

$$\delta e_{\alpha} \equiv \delta e_{,\alpha} + \delta e_{\alpha}^{(v)}; \quad \delta e_{\alpha}^{(v)|\alpha} \equiv 0.$$
(217)

For the temperature anisotropy, eq. (163) gives

$$\frac{\delta T}{T}\Big|_{O} = \frac{\delta T}{T}\Big|_{E} - v_{,\alpha}e^{\alpha}\Big|_{E}^{O} + \int_{E}^{O}\Big(-\varphi' + \alpha_{,\alpha}e^{\alpha} - \frac{1}{a}\chi_{,\alpha|\beta}e^{\alpha}e^{\beta}\Big)dy + v_{\alpha}^{(v)}e^{\alpha}\Big|_{E}^{O} - \int_{E}^{O}\Psi_{\alpha|\beta}^{(v)}e^{\alpha}e^{\beta}dy - \int_{E}^{O}C_{\alpha\beta}^{(t)'}e^{\alpha}e^{\beta}dy + \int_{E}^{O}N_{\nu}dy + N_{T}.$$
(218)

To the linear-order this result was first presented by Sachs and Wolfe [17]; for further analyses using our notation, see [30].

For the Boltzmann equation, eq. (171) becomes

$$\tilde{f}' + \frac{q}{\sqrt{q^2 + m^2 a^2}} \left(\gamma^{\alpha} \delta f_{,\alpha} - \Gamma^{(3)\alpha}_{\ \beta\gamma} \gamma^{\beta} \gamma^{\gamma} \frac{\partial \delta f}{\partial \gamma^{\alpha}} \right) - \left[\varphi' + \frac{\sqrt{q^2 + m^2 a^2}}{q} \gamma^{\alpha} \alpha_{,\alpha} + \left(\frac{1}{a} \chi_{,\alpha|\beta} + \Psi^{(v)}_{\alpha|\beta} + C^{(t)\prime}_{\alpha\beta} \right) \gamma^{\alpha} \gamma^{\beta} \right] q \frac{\partial \tilde{f}}{\partial q} = N_c.$$
(219)

The fluid quantities can be read from eqs. (174,177).

We emphasize that all the equations up to this point are presented without fixing the gauge conditions. In order to solve the equations in a given situation, we can choose *any* allowed gauge conditions suitable for the situation. In this sense, the equations are presented in a *gauge-ready* form.

VI. GAUGE ISSUE

A. Gauge transformation

We consider the following transformation between two coordinates x^a and \hat{x}^a

$$\hat{x}^a \equiv x^a + \tilde{\xi}^a(x^e) \equiv x^a + \tilde{\zeta}^a + \frac{1}{2}\tilde{\zeta}^a{}_{,b}\tilde{\zeta}^b.$$
(220)

The variables $\tilde{\xi}^a$ and $\tilde{\zeta}^a$ are perturbed order quantities. To the second-order we may have

$$\tilde{\xi}^a \equiv \tilde{\xi}^{(1)a} + \tilde{\xi}^{(2)a},\tag{221}$$

and similarly for $\tilde{\zeta}^a$. For any tensor quantity we use the tensor transformation property between x^a and \hat{x}^a spacetimes.

$$\tilde{\phi}(x^e) = \hat{\tilde{\phi}}(\hat{x}^e), \quad \tilde{v}_a(x^e) = \frac{\partial \hat{x}^b}{\partial x^a} \hat{\tilde{v}}_b(\hat{x}^e), \quad \tilde{t}_{ab}(x^e) = \frac{\partial \hat{x}^c}{\partial x^a} \frac{\partial \hat{x}^d}{\partial x^b} \hat{\tilde{t}}_{cd}(\hat{x}^e).$$
(222)

Comparing the tensor quantities at the same spacetime point, x^a , we can derive the gauge transformation property of the tensor quantity. We can show that a tensor quantity **t** transforms as [31]

$$\hat{\mathbf{t}}(x^c) = \mathbf{t}(x^c) - \pounds_{\tilde{\zeta}} \mathbf{t} + \frac{1}{2} \pounds_{\tilde{\zeta}}^2 \mathbf{t}, \qquad (223)$$

where $\pounds_{\tilde{\zeta}}$ is a Lie derivative along $\tilde{\zeta}^a.$ We have

$$\hat{\tilde{\phi}}(x^e) = \tilde{\phi}(x^e) - \tilde{\phi}_{,c}\tilde{\xi}^c + \tilde{\phi}_{,b}\tilde{\xi}^b_{,c}\tilde{\xi}^c + \frac{1}{2}\tilde{\phi}_{,bc}\tilde{\xi}^b\tilde{\xi}^c,$$
(224)

$$\hat{\tilde{v}}_{a}(x^{e}) = \tilde{v}_{a}(x^{e}) - \tilde{v}_{a,b}\tilde{\xi}^{b} - \tilde{v}_{b}\tilde{\xi}^{b}{}_{,a} + \frac{1}{2}\tilde{v}_{a,bc}\tilde{\xi}^{b}\tilde{\xi}^{c} + \tilde{v}_{a,b}\tilde{\xi}^{b}{}_{,c}\tilde{\xi}^{c} + \tilde{v}_{b,c}\tilde{\xi}^{b}{}_{,a}\tilde{\xi}^{c} + \tilde{v}_{b}\tilde{\xi}^{b}{}_{,ac}\tilde{\xi}^{c} + \tilde{v}_{c}\tilde{\xi}^{c}{}_{,b}\tilde{\xi}^{b}{}_{,a},$$
(225)

$$\tilde{\tilde{t}}_{ab}(x^e) = \tilde{t}_{ab}(x^e) - 2\tilde{t}_{c(a}\tilde{\xi}^c{}_{,b)} - \tilde{t}_{ab,c}\tilde{\xi}^c$$

$$+2\tilde{t}_{c(a}\tilde{\xi}^{d}{}_{,b)}\tilde{\xi}^{c}{}_{,d}+\tilde{t}_{cd}\tilde{\xi}^{c}{}_{,a}\tilde{\xi}^{d}{}_{,b}+\tilde{\xi}^{d}\left(2\tilde{\xi}^{c}{}_{,(b}\tilde{t}_{a)c,d}+2\tilde{t}_{c(a}\tilde{\xi}^{c}{}_{,b)d}+\frac{1}{2}\tilde{t}_{ab,cd}\tilde{\xi}^{c}+\tilde{t}_{ab,c}\tilde{\xi}^{c}{}_{,d}\right).$$

$$(226)$$

We define

$$\tilde{\xi}^0 \equiv \xi^0, \quad \tilde{\xi}^\alpha \equiv \xi^\alpha,$$
(227)

where ξ^{α} is based on $g_{\alpha\beta}^{(3)}$. In terms of $\tilde{\zeta}^{a}$ we set $\tilde{\zeta}^{0} \equiv \zeta^{0}$ and $\tilde{\zeta}^{\alpha} \equiv \zeta^{\alpha}$ where ζ^{α} is based on $g_{\alpha\beta}^{(3)}$. Thus, we have

$$\xi^{0} = \zeta^{0} + \frac{1}{2}\zeta^{0'}\zeta^{0} + \frac{1}{2}\zeta^{0}{}_{,\alpha}\zeta^{\alpha}, \quad \xi^{\alpha} = \zeta^{\alpha} + \frac{1}{2}\zeta^{\alpha'}\zeta^{0} + \frac{1}{2}\zeta^{\alpha}{}_{,\beta}\zeta^{\beta}.$$
(228)

From the gauge transformation property of \tilde{g}_{ab} and the definitions of our perturbation variables we can derive:

$$\hat{A} = A - \left(\xi^{0'} + \frac{a'}{a}\xi^{0}\right) - A'\xi^{0} - 2A\left(\xi^{0'} + \frac{a'}{a}\xi^{0}\right) - A_{,\alpha}\xi^{\alpha} - B_{\alpha}\xi^{\alpha'}
+ \frac{3}{2}\xi^{0'}\xi^{0'} + \xi^{0}_{,\alpha}\xi^{\alpha'} + \xi^{\alpha}\left(\xi^{0'}_{,\alpha} + \frac{a'}{a}\xi^{0}_{,\alpha}\right) + \xi^{0}\left[\xi^{0''} + 3\frac{a'}{a}\xi^{0'} + \frac{1}{2}\left(\frac{a''}{a} + \frac{a'^{2}}{a^{2}}\right)\xi^{0}\right] - \frac{1}{2}\xi^{\alpha'}\xi'_{\alpha},$$
(229)
$$\hat{B}_{\alpha} = B_{\alpha} - \xi^{0}_{,\alpha} + \xi'_{\alpha} - 2A\xi^{0}_{,\alpha} - \left(B'_{\alpha} + 2\frac{a'}{a}B_{\alpha}\right)\xi^{0} - B_{\alpha}\xi^{0'} - B_{\alpha,\beta}\xi^{\beta} - B_{\beta}\xi^{\beta}_{,\alpha} + 2C_{\alpha\beta}\xi^{\beta'}
-\xi'_{\alpha}\xi^{0'} + 2\xi^{0'}\xi^{0}_{,\alpha} + \xi^{0}_{,\beta}\xi^{\beta}_{,\alpha} + \xi^{\gamma}\xi^{0}_{,\alpha\gamma} - \xi^{0}\left(\xi''_{\alpha} + 2\frac{a'}{a}\xi'_{\alpha} - \xi^{0'}_{,\alpha} - 2\frac{a'}{a}\xi^{0}_{,\alpha}\right)
-\xi^{\beta}_{,\alpha}\xi'_{\beta} - g^{(3)}_{\alpha\beta}\xi^{\beta}_{,\gamma}\xi^{\gamma'} - \xi^{\gamma}\left(g^{(3)}_{\alpha\beta,\gamma}\xi^{\beta'} + g^{(3)}_{\alpha\beta}\xi^{\beta'}_{,\gamma}\right),$$
(230)

$$\hat{C}_{\alpha\beta} = C_{\alpha\beta} - \frac{a'}{a} \xi^{0} g^{(3)}_{\alpha\beta} - \frac{1}{2} g^{(3)}_{\alpha\beta,\gamma} \xi^{\gamma} - g^{(3)}_{\gamma(\alpha} \xi^{\gamma}_{,\beta)}
+ B_{(\alpha} \xi^{0}_{,\beta)} - \left(C'_{\alpha\beta} + 2\frac{a'}{a} C_{\alpha\beta} \right) \xi^{0} - C_{\alpha\beta,\gamma} \xi^{\gamma} - 2C_{\gamma(\alpha} \xi^{\gamma}_{,\beta)} + \xi'_{(\alpha} \xi^{0}_{,\beta)} - \frac{1}{2} \xi^{0}_{,\alpha} \xi^{0}_{,\beta} + \frac{a'}{a} g^{(3)}_{\alpha\beta} \xi^{\gamma} \xi^{0}_{,\gamma}
+ \xi^{0} \left[\frac{a'}{a} g^{(3)}_{\alpha\beta} \xi^{0'} + \frac{1}{2} \left(\frac{a''}{a} + \frac{a'^{2}}{a^{2}} \right) g^{(3)}_{\alpha\beta} \xi^{0} + \left(\frac{1}{2} \xi^{\gamma'} + \frac{a'}{a} \xi^{\gamma} \right) g^{(3)}_{\alpha\beta,\gamma} + 2\frac{a'}{a} g^{(3)}_{\gamma(\alpha} \xi^{\gamma}_{,\beta)} + g^{(3)}_{\gamma(\alpha} \xi^{\gamma'}_{,\beta)} \right]
+ \xi^{\delta}_{,(\beta} g^{(3)}_{\alpha\gamma\gamma} \xi^{\gamma}_{,\delta} + \frac{1}{2} g^{(3)}_{\gamma\delta} \xi^{\gamma}_{,\alpha} \xi^{\delta}_{,\beta} + \xi^{\delta} \left(\frac{1}{2} g^{(3)}_{\alpha\beta,\gamma} \xi^{\gamma}_{,\delta} + \xi^{\gamma}_{,(\beta} g^{(3)}_{\alpha)\gamma,\delta} + \frac{1}{4} g^{(3)}_{\alpha\beta,\gamma\delta} \xi^{\gamma} + g^{(3)}_{\gamma(\alpha} \xi^{\gamma}_{,\beta)\delta} \right).$$
(231)

From the gauge transformation property of \tilde{T}_{ab} and using the definitions of our perturbed fluid variables in the normal-frame we can derive:

$$\delta\hat{\mu} = \delta\mu - (\mu' + \delta\mu')\xi^0 - \delta\mu_{,\alpha}\xi^{\alpha} + \mu'\left(\xi^0\xi^{0\prime} + \xi^0_{,\alpha}\xi^{\alpha}\right) + \frac{1}{2}\mu''\xi^0\xi^0 + \left[2Q^{\alpha} + (\mu + p)\xi^{0,\alpha}\right]\xi^0_{,\alpha},\tag{232}$$

$$\delta \hat{p} = \delta p - (p' + \delta p') \xi^{0} - \delta p_{,\alpha} \xi^{\alpha} + p' \left(\xi^{0} \xi^{0'} + \xi^{0}_{,\alpha} \xi^{\alpha}\right) + \frac{1}{2} p'' \xi^{0} \xi^{0} + \frac{1}{3} \left[2Q^{\alpha} + (\mu + p)\xi^{0,\alpha}\right] \xi^{0}_{,\alpha}, \tag{233}$$

$$\hat{Q}_{\alpha} = Q_{\alpha} + (\mu + p)\xi^{0}{}_{,\alpha} - Q_{\beta}\xi^{\beta}{}_{,\alpha} - Q_{\alpha,\beta}\xi^{\beta} - \left(Q'_{\alpha} + \frac{a'}{a}Q_{\alpha}\right)\xi^{0} + (\mu + p)A\xi^{0}{}_{,\alpha} + \left[\delta\mu + \delta p - (\mu' + p')\xi^{0}\right]\xi^{0}{}_{,\alpha} + \Pi^{\beta}_{\alpha}\xi^{0}{}_{,\beta} - (\mu + p)\left[\xi^{0}{}_{,\alpha}\xi^{0\prime} + \xi^{\beta}{}_{,\alpha}\xi^{0}{}_{,\beta} + \xi^{0}\left(\xi^{0\prime}{}_{,\alpha} + \frac{a'}{a}\xi^{0}{}_{,\alpha}\right) + \xi^{\beta}\xi^{0}{}_{,\alpha\beta}\right],$$
(234)

$$\hat{\Pi}_{\alpha\beta} = \Pi_{\alpha\beta} - 2\Pi_{\gamma(\alpha}\xi^{\gamma}{}_{,\beta)} - \left(\Pi'_{\alpha\beta} + 2\frac{a'}{a}\Pi_{\alpha\beta}\right)\xi^{0} - \Pi_{\alpha\beta,\gamma}\xi^{\gamma} + \left[2Q_{(\alpha} + (\mu+p)\xi^{0}{}_{,(\alpha)}\right]\xi^{0}{}_{,\beta)} - \frac{1}{3}g^{(3)}_{\alpha\beta}\left[2Q^{\gamma} + (\mu+p)\xi^{0,\gamma}\right]\xi^{0}{}_{,\gamma}.$$
(235)

Under the gauge transformation the individual fluid quantities $\delta \mu_{(i)}$, $\delta p_{(i)}$, $Q_{(i)\alpha}$, $\Pi_{(i)\alpha\beta}$, and $\delta \phi_{(i)}$ transform just like the corresponding collective fluid quantities in eqs. (232-239) with all the fluid quantities changed into the ones for the individual one. Using the vector nature of $\tilde{I}_{(i)\alpha}$ we have

$$\delta \hat{I}_{(i)0} = \delta I_{(i)0} - (I_{(i)0}\xi^{0})' - (\delta I_{(i)0}\xi^{0})' - \delta I_{(i)0,\alpha}\xi^{\alpha} + \delta I_{(i)\alpha}\xi^{\alpha'} + I'_{(i)0}\xi^{0'}\xi^{0} + [I_{(i)0}(\xi^{0'}\xi^{0} + \xi^{0}_{,\alpha}\xi^{\alpha})]', \qquad (236)$$

$$\delta \hat{I}_{(i)\alpha} = \delta I_{(i)\alpha} - I_{(i)0}\xi^{0}_{,\alpha} - \delta I_{(i)0}\xi^{0}_{,\alpha} - \delta I'_{(i)\alpha}\xi^{0} - \delta I_{(i)\alpha,\beta}\xi^{\beta} - \delta I_{(i)\beta}\xi^{\beta}_{,\alpha} + I'_{(i)0}\xi^{0}_{,\alpha}\xi^{0} + I_{(i)0}\xi^{0}_{,\alpha}\xi^{0} + I_{(i)0}[(\xi^{0}_{,\alpha}\xi^{0})' + (\xi^{0}_{,\beta}\xi^{\beta})_{,\alpha}]. \qquad (237)$$

The fluid quantities we use in this work are based on the normal-frame four-vector where $\tilde{n}_{\alpha} = 0$, see eq. (83). It is convenient to have the gauge-transformation properties of the fluid quantities in the energy-frame where we set $Q_{\alpha} = 0$. These can be derived either by applying the frame-transformation rule presented in eqs. (87,88) or directly from the gauge transformation property of the energy-momentum tensor in eq. (82) with $Q_{\alpha} = 0$ in the energy-frame. We have

$$\begin{split} \delta\hat{\mu}^{E} &= \delta\mu^{E} - \left(\mu' + \delta\mu^{E'}\right)\xi^{0} - \delta\mu^{E}_{,\alpha}\xi^{\alpha} + \frac{1}{2}\mu''\xi^{0}\xi^{0} + \mu'\left(\xi^{0}\xi^{0'} + \xi^{\alpha}\xi^{0}_{,\alpha}\right), \\ \delta\hat{p}^{E} &= \delta p^{E} - \left(p' + \delta p^{E'}\right)\xi^{0} - \delta p^{E}_{,\alpha}\xi^{\alpha} + \frac{1}{2}p''\xi^{0}\xi^{0} + p'\left(\xi^{0}\xi^{0'} + \xi^{\alpha}\xi^{0}_{,\alpha}\right), \\ \hat{V}^{E}_{\alpha} - \hat{B}_{\alpha} &= V^{E}_{\alpha} - B_{\alpha} + \xi^{0}_{,\alpha} - \left(V^{E}_{\alpha} - B_{\alpha}\right)'\xi^{0} + \frac{a'}{a}\left(V^{E}_{\alpha} - B_{\alpha}\right)\xi^{0} - \left(V^{E}_{\beta} - B_{\beta}\right)\xi^{\beta}_{,\alpha} - \left(V^{E}_{\alpha} - B_{\alpha}\right)_{,\beta}\xi^{\beta} \\ &+ \left(V^{E\beta} + \xi^{\beta'}\right)\left(g^{(3)}_{\alpha\beta,\gamma}\xi^{\gamma} + 2g^{(3)}_{\gamma(\alpha}\xi^{\gamma}_{,\beta)}\right) + \left(A - \xi^{0'} - \frac{a'}{a}\xi^{0}\right)\left(2\xi^{0}_{,\alpha} - \xi'_{\alpha}\right) + B_{\alpha}\left(\xi^{0'} + 3\frac{a'}{a}\xi^{0}\right) - 2C_{\alpha\beta}\xi^{\beta'} \\ &- \xi^{\beta}_{,\alpha}\xi^{0}_{,\beta} - \xi^{0}\xi^{0'}_{,\alpha} - \xi^{\beta}\xi^{0}_{,\alpha\beta} + 2\frac{a'}{a}\xi^{0}\xi'_{\alpha}, \\ \hat{\Pi}^{E}_{\alpha\beta} &= \Pi^{E}_{\alpha\beta} - \left(\Pi^{E'}_{\alpha\beta} + 2\frac{a'}{a}\Pi^{E}_{\alpha\beta}\right)\xi^{0} - \Pi^{E}_{\alpha\beta,\gamma}\xi^{\gamma} - 2\Pi^{E}_{\gamma(\alpha}\xi^{\gamma}_{,\beta)}. \end{split}$$

From the gauge transformation of $\tilde{\phi}$ we have

$$\delta\hat{\phi} = \delta\phi - (\phi' + \delta\phi')\xi^{0} - \delta\phi_{,\alpha}\xi^{\alpha} + \phi'\left(\xi^{0'}\xi^{0} + \xi^{0}_{,\alpha}\xi^{\alpha}\right) + \frac{1}{2}\phi''\xi^{0}\xi^{0}.$$
(239)

Using the gauge-transformation property of a vector quantity \tilde{k}^a similar to eqs. (222,225), and using the definition of \tilde{k}^a in eq. (150) we can derive

$$\begin{split} \frac{\delta\hat{\nu}}{\nu} &= \frac{\delta\nu}{\nu} + \xi^{0\prime} + 2\frac{a'}{a}\xi^0 - \xi^0{}_{,\alpha}e^{\alpha} - \frac{\delta\nu'}{\nu}\xi^0 + \frac{\delta\nu}{\nu}\left(\xi^{0\prime} + \frac{a'}{a}\xi^0\right) - \frac{\delta\nu{}_{,\alpha}}{\nu}\xi^{\alpha} - \xi^0{}_{,\alpha}\delta e^{\alpha} \\ &+ \xi^0\left[-\xi^{0\prime\prime} - \left(\frac{a''}{a} - 3\frac{a'^2}{a^2}\right)\xi^0 + \xi^{0\prime}{}_{,\alpha}e^{\alpha} + \xi^0{}_{,\alpha}\left(e^{\alpha\prime} - 2\frac{a'}{a}e^{\alpha}\right)\right] \end{split}$$

$$+\xi^{\alpha}\left(-\xi^{0}_{,\alpha}-2\frac{a'}{a}\xi^{0}_{,\alpha}+\xi^{0}_{,\alpha\beta}e^{\beta}+\xi^{0}_{,\beta}e^{\beta}_{,\alpha}\right),$$

$$(240)$$

$$\delta\hat{e}^{\alpha}=\delta e^{\alpha}-\xi^{0}\left(e^{\alpha\prime}-2\frac{a'}{a}e^{\alpha}\right)-\xi^{\alpha\prime}+\xi^{\alpha}_{,\beta}e^{\beta}-e^{\alpha}_{,\beta}\xi^{\beta}-\xi^{0}\left(\delta e^{\alpha\prime}-2\frac{a'}{a}\delta e^{\alpha}\right)-\xi^{\alpha\prime}\frac{\delta\nu}{\nu}+\xi^{\alpha}_{,\beta}\delta e^{\beta}-\delta e^{\alpha}_{,\beta}\xi^{\beta}$$

$$+\xi^{0}\left\{\xi^{0\prime}\left(e^{\alpha\prime}-2\frac{a'}{a}e^{\alpha}\right)+\xi^{0}\left[\frac{1}{2}e^{\alpha\prime\prime}-2\frac{a'}{a}e^{\alpha\prime}-\left(\frac{a^{\prime\prime}}{a}-3\frac{a^{\prime2}}{a^{2}}\right)e^{\alpha}\right]$$

$$+\xi^{\alpha\prime\prime}-2\frac{a'}{a}\xi^{\alpha\prime}-\xi^{\alpha\prime}_{,\beta}e^{\beta}+e^{\alpha}_{,\beta}\xi^{\beta\prime}+\xi^{\beta}\left(e^{\alpha\prime}_{,\beta}-2\frac{a'}{a}e^{\alpha}_{,\beta}\right)-\xi^{\alpha}_{,\beta}\left(e^{\beta\prime}-2\frac{a'}{a}e^{\beta}\right)\right\}$$

$$+\xi^{\beta}\left[\frac{1}{2}e^{\alpha}_{,\beta\gamma}\xi^{\gamma}-\xi^{\alpha}_{,\gamma}e^{\gamma}_{,\beta}+e^{\alpha}_{,\gamma}\xi^{\gamma}_{,\beta}+\xi^{\alpha\prime}_{,\beta}-\xi^{\alpha}_{,\beta\gamma}e^{\gamma}+\xi^{0}_{,\beta}\left(e^{\alpha\prime}-2\frac{a'}{a}e^{\alpha}\right)\right].$$

$$(241)$$

Using the scalar nature of the temperature \tilde{T} and eq. (224) we can show

$$\delta \hat{T}(x^e) = \delta T(x^e) - (T' + \delta T')\xi^0 - \delta T_{,\alpha}\xi^{\alpha} + T'\left(\xi^{0\prime}\xi^0 + \xi^0_{,\alpha}\xi^{\alpha}\right) + \frac{1}{2}T''\xi^0\xi^0.$$
(242)

Using the vector nature of the electric and magnetic vectors and eq. (225) we can show

$$\hat{E}_{\alpha}(x^{e}) = E_{\alpha}(x^{e}) - E_{\alpha}'\xi^{0} - E_{\alpha,\beta}\xi^{\beta} - E_{\beta}\xi^{\beta}_{,\alpha}, \qquad (243)$$

and similarly for H_{α} . Thus, E_{α} and H_{α} are gauge-invariant to the linear-order. Since $\tilde{p}^a \equiv dx^a/d\lambda$, under the gauge transformation we have $\hat{\tilde{p}}^a = \tilde{p}^a + \tilde{\xi}^a{}_{,b}\tilde{p}^b$. Using the definitions of q and γ^{α} in Eq. (169) we can derive

$$\hat{q} = q \left\{ 1 + \frac{a'}{a} \xi^{0} + \frac{\sqrt{q^{2} + m^{2}a^{2}}}{q} \xi^{0}{}_{,\alpha} \gamma^{\alpha} + \frac{\sqrt{q^{2} + m^{2}a^{2}}}{q} \xi^{0}{}_{,\alpha} \left[\gamma^{\alpha} \left(A - \xi^{0'} + \frac{a'}{a} \xi^{0} \right) - C^{\alpha}_{\beta} \gamma^{\beta} \right] \right. \\ \left. + \frac{q^{2} + m^{2}a^{2}}{q^{2}} \left[-A'\xi^{0} - A_{,\alpha}\xi^{\alpha} + \frac{1}{2} \xi^{0}{}_{,\alpha}\xi^{0,\alpha} + \xi^{\alpha} \left(\xi^{0'}{}_{,\alpha} + \frac{a'}{a} \xi^{0}{}_{,\alpha} \right) + \xi^{0}\xi^{0''} + \frac{a'}{a} \xi^{0}\xi^{0'} \right. \\ \left. + \left(\frac{1}{2} \frac{a''}{a} \frac{3q^{2} + 2m^{2}a^{2}}{q^{2} + m^{2}a^{2}} - \frac{a'^{2}}{a^{2}} \right) \xi^{0}\xi^{0} \right] - \frac{1}{2} \frac{m^{2}a^{2}}{q^{2}} \xi^{0}{}_{,\alpha}\xi^{0}{}_{,\beta}\gamma^{\alpha}\gamma^{\beta} \right\},$$

$$(244)$$

$$\hat{\gamma}^{\alpha} = \gamma^{\alpha} + \frac{\sqrt{q^2 + m^2 a^2}}{q} \left(\xi^{0,\alpha} - \xi^{0}{}_{,\beta}\gamma^{\beta}\gamma^{\alpha}\right) + \left(\xi^{\alpha}{}_{,\beta} - \frac{1}{2}\xi^{\alpha}{}_{|\beta} - \frac{1}{2}\xi^{|\alpha}{}_{\beta}\right)\gamma^{\beta}.$$
(245)

As $\hat{\gamma}^{\alpha}$ always appears together with perturbed order terms multiplied, it is evaluated only to the linear-order. From the scalar nature of \tilde{f} we have

$$\tilde{f}(x^e, q, \gamma^{\epsilon}) = \hat{\tilde{f}}(\hat{x}^e, \hat{q}, \hat{\gamma}^{\epsilon}) = \hat{\tilde{f}}(x^e + \tilde{\xi}^e, q + \delta q, \gamma^{\epsilon} + \delta \gamma^{\epsilon}).$$
(246)

At the same momentum space and the spacetime point, we have

$$\delta \hat{f} = \delta f - \tilde{f}' \xi^0 - \tilde{f}_{,q} \delta q + f' \xi^0 \xi^{0\prime} + \frac{1}{2} f'' \xi^0 \xi^0 + 2f'_{,q} \xi^0 \delta q + f_{,q} \delta q_{,q} \delta q + \frac{1}{2} f_{,qq} \delta q^2 - \delta f_{,\alpha} \xi^\alpha - \delta f_{,\gamma^\alpha} \delta \gamma^\alpha.$$
(247)

Using eqs. (244, 245, 229) we have

$$\begin{split} \delta \hat{f} &= \delta f - \tilde{f}' \xi^0 - q \tilde{f}_{,q} \left(\frac{a'}{a} \xi^0 + \frac{\sqrt{q^2 + m^2 a^2}}{q} \xi^0_{,\alpha} \gamma^\alpha \right) \\ &- q f_{,q} \Biggl\{ \frac{\sqrt{q^2 + m^2 a^2}}{q} \xi^0_{,\alpha} \left[\gamma^\alpha \left(A - \xi^{0'} \right) - C^\alpha_\beta \gamma^\beta \right] + \frac{q^2 + m^2 a^2}{q^2} \Biggl[-A' \xi^0 - A_{,\alpha} \xi^\alpha + \xi^\alpha \left(\xi^{0'}_{,\alpha} + \frac{a'}{a} \xi^0_{,\alpha} \right) + \xi^0 \xi^{0''} + \frac{1}{2} \frac{a''}{a} \frac{3q^2 + 2m^2 a^2}{q^2 + m^2 a^2} \xi^0 \xi^0 + \frac{a'}{a} \xi^0 \xi^{0'} + \frac{1}{2} \xi^0_{,\alpha} \xi^{0,\alpha} \Biggr] \end{split}$$

$$-\frac{1}{2}\frac{2q^{2}+m^{2}a^{2}}{q^{2}}\xi^{0}{}_{,\alpha}\xi^{0}{}_{,\beta}\gamma^{\alpha}\gamma^{\beta} - \frac{2q^{2}+m^{2}a^{2}}{q^{2}}\frac{a'^{2}}{a^{2}}\xi^{0}\xi^{0} - \frac{q}{\sqrt{q^{2}+m^{2}a^{2}}}\frac{a'}{a}\xi^{0}\xi^{0}{}_{,\alpha}\gamma^{\alpha}\right\}$$

$$+\frac{1}{2}q^{2}f_{,qq}\left(\frac{a'^{2}}{a^{2}}\xi^{0}\xi^{0} + 2\frac{\sqrt{q^{2}+m^{2}a^{2}}}{q}\frac{a'}{a}\xi^{0}\xi^{0}{}_{,\alpha}\gamma^{\alpha} + \frac{q^{2}+m^{2}a^{2}}{q^{2}}\xi^{0}{}_{,\alpha}\xi^{0}{}_{,\beta}\gamma^{\alpha}\gamma^{\beta}\right)$$

$$+2qf'_{,q}\xi^{0}\left(\frac{a'}{a}\xi^{0} + \frac{\sqrt{q^{2}+m^{2}a^{2}}}{q}\xi^{0}{}_{,\alpha}\gamma^{\alpha}\right) + f'\xi^{0}\xi^{0'} + \frac{1}{2}f''\xi^{0}\xi^{0}$$

$$-\delta f_{,\alpha}\xi^{\alpha} - \delta f_{,\gamma^{\alpha}}\left[\frac{\sqrt{q^{2}+m^{2}a^{2}}}{q}\left(\xi^{0,\alpha}-\xi^{0}{}_{,\beta}\gamma^{\beta}\gamma^{\alpha}\right) + \left(\xi^{\alpha}{}_{,\beta}-\frac{1}{2}\xi^{\alpha}{}_{|\beta}-\frac{1}{2}\xi^{|\alpha}{}_{,\beta}\right)\gamma^{\beta}\right].$$
(248)

Notice that with our phase space variables introduced in Eq. (169) the distribution function f is spatially gaugeinvariant to the linear-order. We can check that the gauge transformation property of δf in Eq. (248) is consistent with the gauge transformation properties of the fluid quantities identified in Eq. (174).

We further decompose ξ_{α} (and similarly for ζ_{α}) into the scalar- and vector-types as

$$\xi_{\alpha} \equiv \frac{1}{a} \xi_{,\alpha} + \xi_{\alpha}^{(v)}, \tag{249}$$

with $\xi^{(v)\alpha}{}_{|\alpha} \equiv 0$. In order to fix the gauge we can impose three conditions on three variables such that these conditions can fix ξ^0 , ξ and $\xi^{(v)}_{\alpha}$. We call these conditions fixing ξ^0 , ξ and $\xi^{(v)}_{\alpha}$ as the temporal, the spatial and the rotational gauge fixing conditions, respectively.

The decomposed variables in eq. (175) and others transform as

$$\begin{split} \hat{\alpha} &= \alpha - \frac{1}{a} \left(a\xi^0 \right)' + A_{\xi}, \\ \hat{\beta} &= \beta - \xi^0 + \left(\frac{1}{a} \xi \right)' + \Delta^{-1} \nabla^{\alpha} B_{\xi \alpha}, \\ \hat{B}_{\alpha}^{(v)} &= B_{\alpha}^{(v)} + \xi_{\alpha}^{(v)\prime} + B_{\xi \alpha} - \nabla_{\alpha} \Delta^{-1} \nabla^{\beta} B_{\xi \beta}, \\ \hat{\gamma} &= \gamma - \frac{1}{a} \xi + \frac{1}{2} \left(\Delta + \frac{1}{2} R^{(3)} \right)^{-1} \left(3\Delta^{-1} \nabla^{\alpha} \nabla^{\beta} C_{\xi \alpha \beta} - C_{\xi \alpha}^{\alpha} \right), \\ \hat{\varphi} &= \varphi - \frac{a'}{a} \xi^0 + \frac{1}{3} C_{\xi \alpha}^{\alpha} - \frac{1}{6} \Delta \left(\Delta + \frac{1}{2} R^{(3)} \right)^{-1} \left(3\Delta^{-1} \nabla^{\alpha} \nabla^{\beta} C_{\xi \alpha \beta} - C_{\xi \alpha}^{\alpha} \right), \\ \hat{C}_{\alpha}^{(v)} &= C_{\alpha}^{(v)} - \xi_{\alpha}^{(v)} + 2 \left(\Delta + \frac{1}{3} R^{(3)} \right)^{-1} \left(\nabla^{\beta} C_{\xi \alpha \beta} - \nabla_{\alpha} \Delta^{-1} \nabla^{\gamma} \nabla^{\beta} C_{\xi \gamma \beta} \right), \\ \hat{C}_{\alpha\beta}^{(t)} &= C_{\alpha\beta}^{(t)} - C_{\xi \alpha \beta} - \frac{1}{3} C_{\xi \gamma}^{\gamma} g_{\alpha\beta}^{(3)} - \frac{1}{2} \left(\nabla_{\alpha} \nabla_{\beta} - \frac{1}{3} g_{\alpha\beta}^{(3)} \Delta \right) \left(\Delta + \frac{1}{2} R^{(3)} \right)^{-1} \left(3\Delta^{-1} \nabla^{\gamma} \nabla^{\delta} C_{\xi \gamma \delta} - C_{\xi \gamma}^{\gamma} \right) \\ - \nabla_{(\alpha} \left(\Delta + \frac{1}{3} R^{(3)} \right)^{-1} \left(\nabla^{\gamma} C_{\xi \beta) \gamma} - \nabla_{\beta} \right) \Delta^{-1} \nabla^{\gamma} \nabla^{\delta} C_{\xi \gamma \delta} \right), \\ \delta \hat{\mu} &= \delta \mu - \mu' \xi^0 + \delta \mu_{\xi}, \\ \delta \hat{p} &= \delta p - p' \xi^0 + \delta p_{\xi}, \\ \hat{v} &= v - \xi^0 - \frac{1}{\mu + p} \Delta^{-1} \nabla^{\alpha} Q_{\xi \alpha}, \\ \hat{v}_{\alpha}^{(v)} &= v_{\alpha}^{(v)} + \frac{1}{\mu + p} \left(Q_{\xi \alpha} - \nabla_{\alpha} \Delta^{-1} \nabla^{\beta} \beta \Pi_{\xi \alpha \beta}, \right) \\ \hat{\Pi}_{\alpha\beta}^{(v)} &= \Pi_{\alpha\beta}^{(v)} + 2a \left(\Delta + \frac{1}{3} R^{(3)} \right)^{-1} \left(\nabla^{\beta} \Pi_{\xi \alpha \beta} - \nabla_{\alpha} \Delta^{-1} \nabla^{\beta} \nabla^{\gamma} \Pi_{\xi \beta \gamma} \right), \\ \hat{\Pi}_{\alpha\beta}^{(t)} &= \Pi_{\alpha\beta}^{(t)} - \Pi_{\xi \alpha\beta} - \frac{3}{2} \left(\nabla_{\alpha} \nabla_{\beta} - \frac{1}{3} g_{\alpha\beta}^{(3)} \Delta \right) \left(\Delta + \frac{1}{2} R^{(3)} \right)^{-1} \Delta^{-1} \nabla^{\gamma} \nabla^{\delta} \Pi_{\xi \gamma \delta} \end{split}$$

$$-\nabla_{(\alpha} \left(\Delta + \frac{1}{3} R^{(3)} \right)^{-1} \left(\nabla^{\gamma} \Pi_{\xi\beta)\gamma} - \nabla_{\beta} \Delta^{-1} \nabla^{\gamma} \nabla^{\delta} \Pi_{\xi\gamma\delta} \right),$$

$$\delta \hat{\phi} = \delta \phi - \phi' \xi^{0} + \delta \phi_{\xi},$$

$$\frac{\delta \hat{\nu}}{\nu} = \frac{\delta \nu}{\nu} + \xi^{0'} + 2\frac{a'}{a} \xi^{0} - \xi^{0}{}_{,\alpha} e^{\alpha} + \frac{\delta \nu_{\xi}}{\nu},$$

$$\delta \hat{e} = \delta e - \frac{1}{a} \xi' + \frac{a'}{a^{2}} \xi - \Delta^{-1} \nabla_{\alpha} \left[\xi^{0} \left(e^{\alpha'} - 2\frac{a'}{a} e^{\alpha} \right) \right] + \Delta^{-1} \left[\frac{1}{a} \Delta(\xi, \alpha) e^{\alpha} + 2K \xi^{(\nu)}_{\alpha} e^{\alpha} - e^{\alpha}{}_{|\beta\alpha} \left(\frac{1}{a} \xi^{,\beta} + \xi^{(\nu)\beta} \right) + \nabla_{\alpha} \delta e^{\alpha}_{\xi} \right],$$

$$\delta \hat{e}^{(\nu)}_{\alpha} = \delta e_{\alpha} - \delta e_{,\alpha},$$

$$\delta \hat{T} = \delta T - T' \xi^{0} + \delta T_{\xi},$$

$$\hat{E}^{(\text{em})} = E^{(\text{em})} + \Delta^{-1} \nabla^{\alpha} E_{\alpha\xi},$$

$$\hat{E}^{(\text{em})} = E^{(\text{em})} + \Delta^{-1} \nabla^{\alpha} E_{\alpha\xi},$$

$$\delta \tilde{f} = \delta f - q \frac{\partial f}{\partial q} \left(\frac{a'}{a} \xi^{0} + \frac{\sqrt{q^{2} + m^{2}a^{2}}}{q} \xi^{0}{}_{,\alpha} \gamma^{\alpha} \right) + \delta f_{\xi},$$
(250)

where A_{ξ} indicates the quadratic parts of eq. (229) and similarly for other variables. For δf we have used f' = 0 which follows from eq. (171) for C[f] = 0 to the background order.

Using t instead of η (indicated as 0) as the time variable, from the definition $dt \equiv ad\eta$ we can show

$$\xi^{0} = \frac{1}{a}\xi^{t} \left(1 - \frac{1}{2}H\xi^{t}\right).$$
(251)

B. Linear-order

From eq. (250) we find that the decomposed metric and matter variables transform to the linear-order as:

$$\hat{\alpha} = \alpha - \dot{\xi}^t, \quad \hat{\beta} = \beta - \frac{1}{a}\xi^t + a\left(\frac{\xi}{a}\right)^t, \quad \hat{\gamma} = \gamma - \frac{1}{a}\xi, \quad \hat{\varphi} = \varphi - H\xi^t, \quad \hat{\chi} = \chi - \xi^t, \quad \hat{\kappa} = \kappa + \left(3\dot{H} + \frac{\Delta}{a^2}\right)\xi^t,$$

$$\delta\hat{\mu} = \delta\mu - \dot{\mu}\xi^t, \quad \delta\hat{p} = \delta p - \dot{p}\xi^t, \quad \hat{v} = v - \frac{1}{a}\xi^t, \quad \hat{\Pi} = \Pi, \quad \delta\hat{\phi} = \delta\phi - \dot{\phi}\xi^t,$$

$$\hat{B}^{(v)}_{\alpha} = B^{(v)}_{\alpha} + a\dot{\xi}^{(v)}_{\alpha}, \quad \hat{C}^{(v)}_{\alpha} = C^{(v)}_{\alpha} - \xi^{(v)}_{\alpha}, \quad \hat{\Psi}^{(v)}_{\alpha} = \Psi^{(v)}_{\alpha}, \quad \hat{v}^{(v)}_{\alpha} = v^{(v)}_{\alpha}, \quad \hat{\Pi}^{(v)}_{\alpha} = \Pi^{(v)}_{\alpha},$$

$$\hat{C}^{(t)}_{\alpha\beta} = C^{(t)}_{\alpha\beta}, \quad \hat{\Pi}^{(t)}_{\alpha\beta} = \Pi^{(t)}_{\alpha\beta}.$$

$$(252)$$

1. Temporal gauge conditions

Temporal gauge fixing condition, fixing ξ^t , applies only to the scalar-type perturbation. To the linear-order, we can impose any one of the following temporal gauge conditions to be valid at any spacetime point:

synchronous gauge :	$\alpha \equiv 0$	\rightarrow	$\xi^{\iota}(\mathbf{x}),$	
comoving gauge :	$v \equiv 0$	\rightarrow	$\xi^t = 0,$	
zero-shear gauge :	$\chi \equiv 0$	\rightarrow	$\xi^t = 0,$	
uniform-expansion gauge :	$\kappa\equiv 0$	\rightarrow	$\xi^t = 0,$	
uniform-curvature gauge :	$\varphi \equiv 0$	\rightarrow	$\xi^t = 0,$	
uniform-density gauge :	$\delta\mu\equiv 0$	\rightarrow	$\xi^t = 0,$	
uniform—pressure gauge :	$\delta p \equiv 0$	\rightarrow	$\xi^t = 0,$	
uniform-field gauge:	$\delta\phi\equiv 0$	\rightarrow	$\xi^t = 0.$	(253)

Except for the synchronous gauge condition, each of the other temporal gauge fixing conditions completely removes the temporal gauge mode. In the multi-component situations in addition we can choose one of the following conditions as the proper temporal gauge condition which also removes the temporal gauge-mode completely:

$$\delta\mu_{(i)} \equiv 0, \quad \delta p_{(i)} \equiv 0, \quad v_{(i)} \equiv 0, \quad \delta\phi_{(i)} \equiv 0.$$
(254)

All these variables which can be used to fix the temporal gauge freedom in fact do not depend on the spatial gauge transformation, ξ , thus are naturally spatially gauge-invariant.

The followings are some examples of combinations of variables which are temporally gauge-invariant:

$$\delta\mu_v \equiv \delta\mu - \dot{\mu}av, \quad \varphi_\chi \equiv \varphi - H\chi, \quad v_\chi \equiv v - \frac{1}{a}\chi, \quad \varphi_v \equiv \varphi - aHv, \quad \varphi_{\delta\phi} \equiv \varphi - \frac{H}{\dot{\phi}}\delta\phi \equiv -\frac{H}{\dot{\phi}}\delta\phi_\varphi. \tag{255}$$

These are completely (i.e., both spatially and temporally) gauge-invariant to the linear-order. Any variable under any gauge condition in eqs. (253,254) (except for the synchronous gauge), has a unique equivalent gauge-invariant combination. For example, we have

$$\varphi_{\chi} = \varphi|_{\chi \equiv 0}.\tag{256}$$

Thus, φ_{χ} is the same as φ variable in the zero-shear gauge where we set $\chi \equiv 0$.

All the equations in \S IV and V are presented without imposing any gauge condition. The equations are arranged using the above variables in eqs. (253,254) which can be used in fixing the temporal gauge condition. This allows us to use the various temporal gauge conditions optimally depending on the situation, thus the equations are presented in a sort of *gauge-ready* manner. Usually we do not know the most suitable gauge condition *a priori*. In order to take the advantage of gauge choice in the most optimal way it is desirable to use the gauge-ready form equations presented in this paper. Our set of equations is arranged so that we can easily impose various fundamental gauge conditions in eqs. (253,254), and their suitable combinations as well. As we have so many different ways of fixing the temporal gauge conditions it is convenient to denote the gauge condition, or equivalently, gauge-invariant combination, we are using. Our notation for gauge-invariant combinations proposed in eq. (255) is convenient for such a purpose in the spirit of our gauge strategy [12,29]. The notation is also practically convenient for connecting solutions in different gauge conditions as well as tracing the associated gauge conditions easily. Compared with the notations for gauge-invariant variables which were introduced by Bardeen [6,7] we have

$$\epsilon_m = \delta_v \equiv \delta \mu_v / \mu, \quad \Psi_H = \varphi_\chi, \quad v_s^{(0)} = k v_\chi, \quad p \pi_L^{(0)} = \delta p, \quad p \pi_T^{(0)} = -\frac{\Delta}{a^2} \Pi, \quad \zeta \equiv \varphi_\delta, \tag{257}$$

etc.; we ignored the harmonic functions used in [6]. The perturbed curvature variable in the comoving gauge \mathcal{R} often used in the literature is the same as our φ_v which is the same as $\varphi_{\delta\phi}$ in the scalar field.

2. Spatial gauge conditions

The spatial gauge transformations ξ and $\xi_{\alpha}^{(v)}$ affect the scalar- and the vector-type perturbations, respectively. Due to spatial homogeneity of the background we have natural spatial gauge fixing conditions to choose [7]. We have two natural spatial gauge fixing conditions. From eq. (252) we can see that:

B-gauge:
$$\beta \equiv 0, \quad B_{\alpha}^{(v)} \equiv 0 \quad \rightarrow \quad \xi(\mathbf{x}, t) \propto a, \quad \xi_{\alpha}^{(v)}(\mathbf{x}),$$
 (258)

$$C-gauge: \quad \gamma \equiv 0, \quad C_{\alpha}^{(v)} \equiv 0 \quad \to \quad \xi = 0, \quad \xi_{\alpha}^{(v)} = 0.$$
(259)

For β we have considered a situation where the temporal gauge condition already completely removed ξ^t . We call the spatial gauge fixing conditions in eqs. (258,259) the B-gauge and the C-gauge, respectively [5]. These gauge conditions are imposed so that we have

$$B-\text{gauge}: \quad B_{\alpha} \equiv 0,$$

$$C-\text{gauge}: \quad C_{\alpha\beta} \equiv \varphi g_{\alpha\beta}^{(3)} + C_{\alpha\beta}^{(t)}.$$
(260)

Apparently, the *B*-gauge conditions fail to fix the spatial and the rotational gauge modes completely, thus, even after imposing the gauge conditions we still have the remaining gauge modes. Whereas, the *C*-gauge conditions successfully remove the gauge modes. To the linear-order, the variables χ and $\Psi_{\alpha}^{(v)}$ introduced in eq. (178) are natural and unique spatially gauge-invariant combinations. Notice that in the *C*-gauge χ is the same as $a\beta$, and $\Psi^{(v)}$ is the same as $B_{\alpha}^{(v)}$. Thus, β and $B_{\alpha}^{(v)}$ variables in the *C*-gauge conditions are equivalent to the corresponding (spatially and rotationally) gauge-invariant combinations χ/a and $\Psi_{\alpha}^{(v)}$, respectively.

C. Second-order

1. Gauge conditions

If we use any of the gauge conditions which completely fixes both the temporal and the spatial gauge modes to the linear-order, the gauge transformation properties of the second-order variables, say $\varphi^{(2)}$ in eq. (183), follow *exactly* the same forms as their linear counterparts. Using the transformation of $\delta \phi$ in eq. (239) as an example, to the linear order we have

$$\delta\hat{\phi}^{(1)} = \delta\phi^{(1)} - \phi'\xi^{0(1)}.$$
(261)

If we take gauge conditions which remove (fix) ξ^0 and ξ^{α} completely to the linear order we have $\xi^{0(1)} = 0 = \xi^{\alpha(1)}$. Thus, from eq. (239) we have

$$\delta\hat{\phi}^{(2)} = \delta\phi^{(2)} - \phi'\xi^{0(2)},\tag{262}$$

which shows exactly the same form as in eq. (261). Thus, the gauge conditions in eqs. (252,254) apply to secondorder perturbation variables as well, and we can impose similar gauge conditions even to the second-order. For example, in the zero-shear gauge we impose $\chi = 0$ as the gauge condition to the second-order, thus $\chi^{(1)} = 0 = \chi^{(2)}$; otherwise mentioned, we always take the *C*-gauge for the spatial and rotational ones. In this gauge condition the gauge transforamtion properties are completely fixed, and the gauge-modes do not appear. Thus, we anticipate that each variable in that gauge condition has unique corresponding gauge-invariant combination of variables. Thus, using φ , we have $\varphi|_{\chi=0,C-\text{gauge}}$ is free of the gauge-modes. We denote the corresponding gauge-invariant combination as

$$\varphi_{\chi}$$
 (263)

with the C-gauge condition assumed always. To the linear-order we have $\varphi_{\chi} \equiv \varphi - H\chi$, but to the second-order we need correction terms to make φ_{χ} gauge-invariant. Construction of such a gauge-invariant combination will be shown below.

From eqs. (57,58), assuming pure scalar mode in the C-gauge we can show

$$\bar{K}_{\alpha\beta} = -(1-\alpha)\chi_{,\alpha|\beta} + 2\chi_{,(\alpha}\varphi_{,\beta)} - \frac{1}{3}g^{(3)}_{\alpha\beta}\left[-(1-\alpha)\Delta\chi + 2\chi^{,\gamma}\varphi_{,\gamma}\right],$$
(264)

$$R^{(h)} = \frac{1}{a^2} \left[R^{(3)} - 4\left(\Delta + \frac{1}{2}R^{(3)}\right)\varphi + 16\varphi\left(\Delta + \frac{1}{4}R^{(3)}\right)\varphi + 6\varphi^{,\alpha}\varphi_{,\alpha} \right].$$
 (265)

Thus, the gauge condition $\chi \equiv 0$ implies $\bar{K}_{\alpha\beta} = 0$, justifying its name as the zero-shear gauge to the second-order. Similarly, the gauge condition $\varphi \equiv 0$ implies $R^{(h)} = \frac{1}{a^2}R^{(3)}$ (we also have $R^{(h)}_{\alpha\beta} = \frac{1}{3}R^{(3)}g^{(3)}_{\alpha\beta}$), justifying its name as the uniform-curvature gauge to the second-order. We can show that the names of gauge conditions in eq. (253) remain valid to the second-order.

In the perturbative approach, apparently, this method can be similarly applied to any higher-order perturbations. As long as we work in any of these gauge conditions, the gauge modes are completely removed and the behavior of all the variables is equivalently gauge-invariant. As the variables are free of gauge mode, these can be considered as physically important ones in that particular gauge conditions we choose. We can also choose different gauge conditions in the second-order compared with the ones imposed to the linear-order. Examples will be shown below.

2. Constructing gauge-invariant combinations

Let us explain a method to derive the gauge-invariant combinations using an example. Since the gauge transformation properties of $\delta\mu$ and $\delta\phi$ are available in a convenient forms in eqs. (232,239) we consider the gauge-invariant combinations involving these two variables to the second-order. Thus, we consider the case with a scalar field. To the linear-order we can construct various gauge invariant combinations involving $\delta\mu$, and as examples we consider two cases

$$\delta\mu_{\delta\phi} \equiv \delta\mu - \frac{\mu'}{\phi'}\delta\phi, \quad \delta\mu_{\varphi} \equiv \delta\mu - \phi'\frac{a}{a'}\varphi.$$
(266)

Apparently, the combinations in eqs. (266) are not gauge-invariant to the second-order. In order to construct the gauge-invariant combination in the gauge with $\delta\phi^{(2)} = 0$ we construct $\delta\hat{\mu} - \frac{\mu'}{\phi'}\delta\hat{\phi}$ using eqs. (232,239). Then, in the RHS we have quadratic combination of linear-order terms involving ξ^0 and ξ^{α} . As the spatial and rotational gauge we consider the *C*-gauge conditions which remove the corresponding gauge modes completely. This can be achieved by taking

$$\xi^{\alpha} = -\left(\hat{\gamma}^{,\alpha} + \hat{C}^{(v)\alpha}\right) + \gamma^{,\alpha} + C^{(v)\alpha},\tag{267}$$

which follows from eqs. (252,249), and moving terms with overhat to the LHS. Now, coming to the temporal gauge freedom, if we want to consider the uniform-field gauge we take

$$\xi^0 = -\frac{1}{\phi'} \left(\delta \hat{\phi} - \delta \phi \right), \tag{268}$$

which follows from eq. (252), and move terms with overhat to the LHS. Then we have a gauge-invariant combination

$$\delta\mu_{\delta\phi} \equiv \delta\mu - \frac{\mu'}{\phi'}\delta\phi - \left(\delta\mu - \frac{\mu'}{\phi'}\delta\phi\right)_{,\alpha} \left(\gamma^{,\alpha} + C^{(v)\alpha}\right) - \frac{1}{\phi'}\left(\delta\mu - \frac{\mu'}{\phi'}\delta\phi\right)'\delta\phi - \frac{1}{2}\left(\frac{\mu'}{\phi'}\right)'\frac{1}{\phi'}\delta\phi^2 - \frac{1}{a^2}\delta\phi^{,\alpha}\delta\phi_{,\alpha}$$
$$\equiv \delta\mu - \frac{\mu'}{\phi'}\delta\phi + \delta\mu^{(q)}_{\delta\phi}.$$
(269)

We have $\delta\mu_{\delta\phi} = \delta\mu|_{\delta\phi^{(1)}=0=\delta\phi^{(2)}, C-\text{gauge}}$, thus $\delta\mu_{\delta\phi}$ is the same as $\delta\mu$ under the gauge conditions $\delta\phi^{(1)} = 0 = \delta\phi^{(2)}$ and the *C*-gauges. If we want to take the uniform-curvature gauge to the linear-order we take

$$\xi^0 = -\frac{a}{a'} \left(\hat{\varphi} - \varphi \right) \tag{270}$$

which follows from eq. (252), and move terms with overhat to the LHS. Then we can identify a gauge-invariant combination

$$\delta\mu - \frac{\mu'}{\phi'}\delta\phi - \left(\delta\mu - \frac{\mu'}{\phi'}\delta\phi\right)_{,\alpha} \left(\gamma^{,\alpha} + C^{(v)\alpha}\right) - \frac{a}{a'} \left(\delta\mu - \frac{\mu'}{\phi'}\delta\phi\right)'\varphi - \left(\frac{\mu'}{\phi'}\right)'\frac{a}{a'} \left(\delta\phi - \frac{1}{2}\phi'\frac{a}{a'}\varphi\right)\varphi - 2\frac{1}{a^2}\phi'\frac{a}{a'} \left(\delta\phi - \frac{1}{2}\phi'\frac{a}{a'}\varphi\right)^{,\alpha}\varphi_{,\alpha}.$$
(271)

This combination is equivalent to $\delta\mu$ in the following gauge conditions: $\delta\phi = 0$ in the linear and the pure second-order part [i.e., $\delta\mu^{(2)} - (\mu'/\phi')\delta\phi^{(2)}$], and $\varphi = 0$ in the quadratic parts, and the *C*-gauges. By replacing the linear-order part of eq. (271) with $\delta\mu^{(1)} - \phi'(a/a')\varphi^{(1)}$ we can make another gauge-invariant combination

$$\delta\mu_{\varphi^{(1)},\delta\phi^{(2)}} \equiv \delta\mu^{(1)} - \phi'\frac{a}{a'}\varphi^{(1)} + \delta\mu^{(2)} - \frac{\mu'}{\phi'}\delta\phi^{(2)} - \left(\delta\mu - \frac{\mu'}{\phi'}\delta\phi\right)_{,\alpha}\left(\gamma^{,\alpha} + C^{(v)\alpha}\right) - \frac{a}{a'}\left(\delta\mu - \frac{\mu'}{\phi'}\delta\phi\right)'\varphi - \left(\frac{\mu'}{\phi'}\right)'\frac{a}{a'}\left(\delta\phi - \frac{1}{2}\phi'\frac{a}{a'}\varphi\right)\varphi - 2\frac{1}{a^2}\phi'\frac{a}{a'}\left(\delta\phi - \frac{1}{2}\phi'\frac{a}{a'}\varphi\right)^{,\alpha}\varphi_{,\alpha},$$
(272)

which is the same as $\delta \mu |_{\varphi^{(1)}=0=\delta\phi^{(2)},C-\text{gauge}}$. The calculation becomes simpler if we take the *C*-gauge condition: this sets $\gamma \equiv 0 \equiv C_{\alpha}^{(v)}$, thus we can simply set $\xi_{\alpha} \equiv 0$ ($\xi \equiv 0 \equiv \xi_{\alpha}^{(v)}$). Similarly we can construct diverse combinations of the gauge-invariant variables: several useful gauge-invariant combinations will be presented in the next subsection.

In the following, as in eq. (269), a gauge-invariant notation, say, φ_v indicates a combination which is equivalent to φ in the comoving gauge (v = 0) to all orders, thus, $v^{(1)} = 0 = v^{(2)}$, and in the *C*-gauge. In order to denote gauge-invariant combinations valid to the second-order, we introduce the following notation

$$\varphi_v \equiv \varphi - aHv + \varphi_v^{(q)}, \quad \varphi_\chi \equiv \varphi - H\chi + \varphi_\chi^{(q)}, \quad \delta\mu_v \equiv \delta\mu - \dot{\mu}av + \delta\mu_v^{(q)}, \quad v_\chi \equiv v - \frac{1}{a}\chi + v_\chi^{(q)}, \tag{273}$$

etc., where the upper (q) index indicates the quadratic combinations of the linear-order terms. In the following we always take the spatial *C*-gauge. We note that $\delta \mu_v^{(q)}$ is the quadratic correction terms to make $\delta \mu_v$ a gauge-invariant

combination to the second-order; thus it differs from, say, $\delta \mu^{(q)} - \dot{\mu} a v^{(q)}$. As φ_v is the same as φ in the v = 0 gauge, we have $\varphi_v^{(q)}$ vanishes under the v = 0 gauge, i.e.,

$$\varphi_v^{(q)}|_v = \varphi_v^{(q)}|_{v=0} = 0.$$
(274)

Using the definition of our gauge-invariant combinations we can show, for example,

$$\varphi_v = \varphi - aHv + \varphi_v^{(q)} = \varphi_\chi - aHv_\chi + \varphi_v^{(q)}|_\chi, \tag{275}$$

where in the second step we have evaluated the first step in the zero-shear gauge. Thus

$$\varphi_{v}^{(q)}|_{\chi} = \varphi_{v} - (\varphi_{\chi} - aHv_{\chi}) = \varphi_{v}^{(q)} - \left(\varphi_{\chi}^{(q)} - aHv_{\chi}^{(q)}\right),$$
(276)

and similarly for other correction terms.

3. Gauge-invariant variables

Now, we present several useful gauge-invariant combinations explicitly. We assume $R^{(3)} = 0$ and pure scalar-type perturbations. We take the spatial *C*-gauge, $\gamma \equiv 0$. As long as we take the temporal gauge which fixes ξ^0 completely, we can set $\xi_{\alpha} \equiv 0$. The metric becomes

$$A = \alpha, \quad B_{\alpha} = \frac{1}{a} \chi_{,\alpha}, \quad C_{\alpha\beta} = \varphi g_{\alpha\beta}^{(3)}.$$
(277)

From eqs. (229-235,239) we have

$$\begin{split} \hat{\alpha} &= \alpha - \frac{1}{a} \left(a\xi^{0} \right)' - \alpha'\xi^{0} - 2\alpha \left(\xi^{0'} + \frac{a'}{a}\xi^{0} \right) + \frac{3}{2}\xi^{0'}\xi^{0'} + \xi^{0} \left[\xi^{0''} + 3\frac{a'}{a}\xi^{0'} + \frac{1}{2} \left(\frac{a''}{a} + \frac{a'^{2}}{a^{2}} \right) \xi^{0} \right] \\ &+ \frac{1}{2} \left(\frac{1}{a}\chi^{,\alpha}\xi^{0} + \xi^{0} \left[-\varphi' - 2\frac{a'}{a}\varphi + \frac{a'}{a}\xi^{0'} + \frac{1}{2} \left(\frac{a''}{a} + \frac{a'^{2}}{a^{2}} \right) \xi^{0} \right] \\ &+ \frac{1}{2} \left(\frac{1}{a}\chi^{,\alpha}\xi^{0} , \alpha - \frac{1}{2}\xi^{0,\alpha}\xi^{0} , \alpha \right) - \frac{1}{2}\Delta^{-1}\nabla^{\alpha}\nabla^{\beta} \left(\frac{1}{a}\chi_{,\alpha}\xi^{0} , \beta - \frac{1}{2}\xi^{0} , \alpha\xi^{0} , \beta \right) , \\ \hat{\chi} &= \chi - a\xi^{0} + a\xi^{0} \left(\xi^{0'} + \frac{a'}{a}\xi^{0} \right) + a\Delta^{-1}\nabla^{\alpha} \left[-2\alpha\xi^{0} , \alpha - \frac{1}{a} \left(\chi' + \frac{a'}{a}\chi \right) _{,\alpha} \xi^{0} - \frac{1}{a}\chi_{,\alpha}\xi^{0'} + \xi^{0'}\xi^{0} , \alpha \right] \\ &- \frac{a}{2}\Delta^{-1} \left[\frac{1}{a}\chi^{,\alpha}\xi^{0} , \alpha - \frac{1}{2}\xi^{0,\alpha}\xi^{0} , \alpha - 3\Delta^{-1}\nabla^{\alpha}\nabla^{\beta} \left(\frac{1}{a}\chi_{,\alpha}\xi^{0} , \beta - \frac{1}{2}\xi^{0} , \alpha\xi^{0} , \beta \right) \right]' , \\ \hat{\kappa} &= \kappa + \left(3\frac{a''}{a^{2}} - 6\frac{a'^{2}}{a^{3}} + \frac{\Delta}{a} \right)\xi^{0} + \text{quadratic terms}, \\ \delta\hat{\mu} &= \delta\mu - \mu'\xi^{0} - \delta\mu'\xi^{0} + \mu'\xi^{0'}\xi^{0} + \frac{1}{2}\mu''\xi^{0}\xi^{0} + (\mu + p) \left(-2v + \xi^{0} \right)^{,\alpha}\xi^{0} , \alpha, \\ \hat{v} &= v - \xi^{0} + \xi^{0}\xi^{0'} + \frac{1}{2} \left(\frac{a'}{a} + \frac{\mu' + p'}{\mu + p} \right)\xi^{0}\xi^{0} - \left[v' + \left(4\frac{a'}{a} + \frac{\mu' + p'}{\mu + p} \right) v \right]\xi^{0} \\ &- \Delta^{-1}\nabla^{\alpha} \left[-3\frac{a'}{a}v_{,\alpha}\xi^{0} + \frac{\delta\mu}{\mu + p}\xi^{0} , \alpha + \frac{1}{a^{2}}\frac{1}{\mu + p} \left(\Pi^{,\beta}_{,\alpha} - \delta^{\beta}_{,\alpha}\Delta\Pi \right)\xi^{0} , \beta \right] , \\ \delta\hat{\phi} &= \delta\phi - \phi'\xi^{0} - \delta\phi'\xi^{0} + \phi'\xi^{0'}\xi^{0} + \frac{1}{2}\phi''\xi^{0}\xi^{0}. \end{split}$$

$$(278)$$

In the transformation of v we have used eq. (201). We have ignored writing the quadratic terms in the transformation of κ ; this can be read from the definition of κ

$$\kappa \equiv -\frac{1}{a} \left[3\varphi' - 3\frac{a'}{a}\alpha + \frac{\Delta\chi}{a} - (\alpha + 2\varphi) \left(3\varphi' + \frac{\Delta\chi}{a} \right) + \frac{3a'}{2a} \left(3\alpha^2 - \frac{1}{a^2}\chi^{,\alpha}\chi_{,\alpha} \right) - \frac{1}{a}\chi^{,\alpha}\varphi_{,\alpha} \right], \tag{279}$$

which follows from eqs. (179,57).

Following the prescription in the previous subsection, from eq. (278) we can construct the following gauge-invariant combinations

$$\varphi_{\chi} \equiv \varphi - H\chi - (\dot{\varphi}_{\chi} + 2H\varphi_{\chi})\chi - \frac{1}{2}\left(\dot{H} + H^{2}\right)\chi^{2} + \frac{1}{4a^{2}}\left[\chi^{,\alpha}\chi_{,\alpha} - \Delta^{-1}\nabla^{\alpha}\nabla^{\beta}\left(\chi_{,\alpha}\chi_{,\beta}\right)\right] + H\Delta^{-1}\nabla^{\alpha}\left[2\alpha_{\chi}\chi_{,\alpha} + (\dot{\chi} - H\chi)\chi_{,\alpha}\right] + \frac{1}{4}a^{2}H\Delta^{-1}\left[\frac{1}{a^{2}}\chi^{,\alpha}\chi_{,\alpha} - 3\frac{1}{a^{2}}\Delta^{-1}\nabla^{\alpha}\nabla^{\beta}\left(\chi_{,\alpha}\chi_{,\beta}\right)\right],$$
(280)

$$\varphi_{v} \equiv \varphi - aHv - (\dot{\varphi}_{v} + 2H\varphi_{v}) av - \frac{1}{2} \left(\dot{H} + 2H^{2} - H\frac{\dot{\mu} + \dot{p}}{\mu + p} \right) a^{2}v^{2} + \frac{1}{4a} \left(2\chi - av \right)^{,\alpha} v_{,\alpha} - \frac{1}{4a} \Delta^{-1} \nabla^{\alpha} \nabla^{\beta} \left[(2\chi - av)_{,\alpha} v_{,\beta} \right] + aH\Delta^{-1} \nabla^{\alpha} \left[\frac{\delta\mu_{v}}{\mu + p} v_{,\alpha} + \frac{1}{a^{2}} \frac{1}{\mu + p} \left(\Pi^{,\beta}{}_{\alpha} - \delta^{\beta}_{\alpha} \Delta \Pi \right) v_{,\beta} \right],$$
(281)

$$\delta_{v} \equiv \delta - \frac{\dot{\mu}}{\mu} av - \frac{\delta \dot{\mu}_{v}}{\mu} av - \frac{1}{2} \frac{\dot{\mu}}{\mu} \frac{\dot{H}}{H} a^{2} v^{2} - \frac{\mu + p}{\mu} v^{,\alpha} v_{,\alpha} + \frac{\dot{\mu}}{\mu} a \Delta^{-1} \nabla^{\alpha} \left[\frac{\delta \mu_{v}}{\mu + p} v_{,\alpha} + \frac{1}{a^{2}} \frac{1}{\mu + p} \left(\Pi^{,\beta}{}_{\alpha} - \delta^{\beta}_{\alpha} \Delta \Pi \right) v_{,\beta} \right],$$

$$(282)$$

$$1 \qquad \left[\int \left(- \frac{\dot{\mu} + \dot{\mu}}{\mu} \right) \right] = 1 \qquad \left(- \frac{\dot{\mu} + \dot{\mu}}{\mu} \right) - 2$$

$$v_{\chi} \equiv v - \frac{1}{a}\chi - \left[\dot{v}_{\chi} + \left(H + \frac{\mu + p}{\mu + p}\right)v_{\chi}\right]\chi + \frac{1}{2a}\left(H - \frac{\mu + p}{\mu + p}\right)\chi^{2} + \frac{1}{a}\Delta^{-1}\nabla^{\alpha}\left[2\alpha_{\chi}\chi_{,\alpha} + \chi_{,\alpha}\left(\dot{\chi} - H\chi\right) - \frac{\delta\mu_{v}}{\mu + p}\chi_{,\alpha} - \frac{1}{a^{2}}\frac{1}{\mu + p}\left(\Pi^{,\beta}{}_{\alpha} - \delta^{\beta}_{\alpha}\Delta\Pi\right)\chi_{,\beta}\right] + \frac{a}{4}\Delta^{-1}\left[\frac{1}{a^{2}}\chi^{,\alpha}\chi_{,\alpha} - \frac{3}{a^{2}}\Delta^{-1}\nabla^{\alpha}\nabla^{\beta}\left(\chi_{,\alpha}\chi_{,\beta}\right)\right],$$
(283)

$$\chi_{v} \equiv \chi - av - \frac{1}{2}a^{2} \left(H - \frac{\dot{\mu} + \dot{p}}{\mu + p}\right)v^{2} + a\Delta^{-1}\nabla^{\alpha} \left[-2\alpha_{v}v_{,\alpha} - (\dot{\chi}_{v} + H\chi_{v})_{,\alpha}v - \chi_{,\alpha}\dot{v} + \frac{\delta\mu_{v}}{\mu + p}v_{,\alpha} + \frac{1}{a^{2}}\frac{1}{\mu + p}\left(\Pi^{,\beta}{}_{\alpha} - \delta^{\beta}_{\alpha}\Delta\Pi\right)v_{,\beta}\right] + \frac{1}{2}a^{2}\Delta^{-1} \left[-\left(\frac{1}{a}\chi^{,\alpha}v_{,\alpha} - \frac{1}{2}v^{,\alpha}v_{,\alpha}\right) + 3\Delta^{-1}\nabla^{\alpha}\nabla^{\beta}\left(\frac{1}{a}\chi_{,\alpha}v_{,\beta} - \frac{1}{2}v_{,\alpha}v_{,\beta}\right)\right]^{'}.$$
(284)

To the linear-order, δ_v (equivalently, δ in the comoving gauge) behaves like the Newtonian density perturbation, and v_{χ} and $-\varphi_{\chi}$ (equivalently, v and $-\varphi$ in the zero-shear gauge) behave like the Newtonian velocity and the gravitational potential. Also to the linear-order φ_v is known to be the best conserved quantity in the super-sound-horizon scale. For extensions of these results to the second-order, see SVIIC and VIID, respectively.

In the case of a scalar field we have

$$\varphi_{\delta\phi} \equiv \varphi - \frac{H}{\dot{\phi}} \delta\phi - \frac{1}{a^2} \left(a^2 \varphi_{\delta\phi} \right)^{\cdot} \frac{\delta\phi}{\dot{\phi}} - \frac{1}{2a^2} \left(\frac{a^2 H}{\dot{\phi}} \right)^{\cdot} \frac{\delta\phi^2}{\dot{\phi}} + \frac{1}{2a^2 \dot{\phi}} \left[\chi^{,\alpha} \delta\phi_{,\alpha} - \frac{1}{2\dot{\phi}} \delta\phi^{,\alpha} \delta\phi_{,\alpha} - \Delta^{-1} \nabla^{\alpha} \nabla^{\beta} \left(\chi_{,\alpha} \delta\phi_{,\beta} - \frac{1}{2\dot{\phi}} \delta\phi_{,\alpha} \delta\phi_{,\beta} \right) \right],$$

$$\delta\phi_{\varphi} \equiv \delta\phi - \frac{\dot{\phi}}{H} \varphi - \frac{1}{H} \delta\dot{\phi}_{\varphi} \varphi + \frac{\dot{\phi}^2}{2a^2 H^3} \left(\frac{a^2 H}{\dot{\phi}} \right)^{\cdot} \varphi^2$$

$$(285)$$

$$-\frac{\dot{\phi}}{2a^{2}H^{2}}\left[\chi^{,\alpha}\varphi_{,\alpha}-\frac{1}{2H}\varphi^{,\alpha}\varphi_{,\alpha}-\Delta^{-1}\nabla^{\alpha}\nabla^{\beta}\left(\chi_{,\alpha}\varphi_{,\beta}-\frac{1}{2H}\varphi_{,\alpha}\varphi_{,\beta}\right)\right].$$
(286)

Thus, to the second-order we have $\varphi_{\delta\phi} \neq -\frac{H}{\phi} \delta\phi_{\varphi}$. By evaluating the RHSs of eqs. (285,286) in the $\varphi = 0$ gauge and $\delta\phi = 0$ gauge, respectively, we have the following relations between the two gauge-invariant variables

$$\varphi_{\delta\phi} = -\frac{H}{\dot{\phi}}\delta\phi_{\varphi} + \frac{H}{\dot{\phi}^{2}}\delta\phi_{\varphi}\delta\dot{\phi}_{\varphi} + \frac{1}{2a^{2}\dot{\phi}}\left(\frac{a^{2}H}{\dot{\phi}}\right)^{\cdot}\delta\phi_{\varphi}^{2} + \frac{1}{2a^{2}\dot{\phi}}\left[\chi_{\varphi}{}^{,\alpha}\delta\phi_{\varphi,\alpha} - \frac{1}{2\dot{\phi}}\delta\phi_{\varphi}{}^{,\alpha}\delta\phi_{\varphi,\alpha} - \Delta^{-1}\nabla^{\alpha}\nabla^{\beta}\left(\chi_{\varphi,\alpha}\delta\phi_{\varphi,\beta} - \frac{1}{2\dot{\phi}}\delta\phi_{\varphi,\alpha}\delta\phi_{\varphi,\beta}\right)\right], \qquad (287)$$

$$\delta\phi_{\varphi} = -\frac{\dot{\phi}}{H}\varphi_{\delta\phi} + \frac{\dot{\phi}}{H^{2}}\varphi_{\delta\phi}\dot{\phi}_{\delta\phi} + \frac{1}{2a^{2}H}\left(\frac{a^{2}\dot{\phi}}{H}\right)^{\cdot}\varphi_{\delta\phi}^{2}$$

$$-\frac{\dot{\phi}}{2a^{2}H^{2}}\left[\chi_{\delta\phi}^{\ ,\alpha}\varphi_{\delta\phi,\alpha} - \frac{1}{2H}\varphi_{\delta\phi}^{\ ,\alpha}\varphi_{\delta\phi,\alpha} - \Delta^{-1}\nabla^{\alpha}\nabla^{\beta}\left(\chi_{\delta\phi,\alpha}\varphi_{\delta\phi,\beta} - \frac{1}{2H}\varphi_{\delta\phi,\alpha}\varphi_{\delta\phi,\beta}\right)\right].$$
(288)

To linear-order $\delta \phi_{\varphi}$ (equivalently, $\delta \phi$ in the uniform-curvature gauge) most closely resembles the scalar-field equation in the fixed cosmological background metric [14]. Since $\delta \phi \equiv 0$ implies $Q_{\alpha}^{(\phi)} = 0$ (thus $v^{(\phi)} = 0$), we have $\varphi_{\delta\phi} = \varphi_v$; from eq. (114) we notice that this is valid to the second-order.

4. Spatial gradient variable

A covariant density gradient variable

$$\tilde{\Delta}_a \equiv \frac{1}{\tilde{\mu}} \tilde{h}_a^b \tilde{\mu}_{,b},\tag{289}$$

is gauge-invariant to the linear-order [32]. In [32] the energy-frame is taken. Using the \tilde{u}_a -frame in eq. (53) we have

$$\Delta_{\alpha} = \frac{1}{1+\delta} \left\{ \delta^{E}_{,\alpha} + \frac{\mu'}{\mu} \left(V^{E}_{\alpha} - B_{\alpha} + AB_{\alpha} + 2V^{E\beta}C_{\alpha\beta} \right) + \left[\delta' + \frac{\mu'}{\mu} \left(\delta - A \right) \right] \left(V^{E}_{\alpha} - B_{\alpha} \right) \right\},\tag{290}$$

where we set $\tilde{\Delta}_{\alpha} \equiv \Delta_{\alpha}$ with Δ_{α} based on $g_{\alpha\beta}^{(3)}$; from $\tilde{u}^a \tilde{\Delta}_a = 0$ we have $\tilde{\Delta}_0 = -V^{\alpha} \Delta_{\alpha}$. From eqs. (85,86) we notice that δ is frame invariant to the linear-order, and we ignore superscript E in such cases. Using the prescription in eq. (88) we can express Δ_{α} in the normal-frame as

$$\Delta_{\alpha} = \frac{1}{1+\delta} \left\{ \delta_{,\alpha}^{N} + \frac{\mu'}{\mu} \frac{Q_{\alpha}^{N}}{\mu+p} + \left[\delta' + \frac{\mu'}{\mu} \left(\delta - A \right) \right] \frac{Q_{\alpha}^{N}}{\mu+p} - \frac{(Q^{N\beta}Q_{\beta}^{N})_{,\alpha}}{\mu(\mu+p)} - \frac{\mu'}{\mu} \frac{1}{(\mu+p)^{2}} \left[\left(\delta\mu + \delta p \right) Q_{\alpha}^{N} + \Pi_{\alpha\beta} Q^{N\beta} \right] \right\}.$$
(291)

Thus, under the condition $Q_{\alpha} = 0$ we have $\Delta_{\alpha} = \delta_{,\alpha}/(1+\delta)$; $Q_{\alpha} = 0$ can be achieved by the comoving gauge condition $(Q_{\alpha}^{(v)} \equiv 0)$ and the irrotational condition $(Q_{\alpha}^{(v)} \equiv 0)$; see [33] and the note added in proof of [21]. Under the gauge transformation, either using eqs. (232-235) for eq. (291) or using eq. (238) for eq. (290) we can show

$$\hat{\Delta}_{\alpha} = \Delta_{\alpha} - \Delta_{\alpha}' \xi^{0} - \left(\Delta_{\beta} \xi^{\beta}\right)_{,\alpha} - 2\Delta_{[\alpha,\beta]} \xi^{\beta}.$$
(292)

Thus, Δ_{α} is not gauge-invariant to the second-order. To the linear-order, the scalar-type part becomes $\Delta_{\alpha} = \delta_{v,\alpha}$ where the gauge-invariant combination δ_v is the same as δ in the comoving gauge.

VII. APPLICATIONS

A. Closed form equations

From eqs. (196,197), eqs. (197,200,201), eqs. (199), eqs. (199,201) and eqs. (195,197,199) we can derive, respectively:

$$\frac{\Delta + 3K}{a^2}\varphi_{\chi} + 4\pi G\delta\mu_v = \frac{\Delta + 3K}{a^2}\varphi_{\chi}^{(q)} + 4\pi G\delta\mu_v^{(q)} + \frac{1}{4}N_1 - HN_2^{(s)},$$
(293)

$$\delta\dot{\mu}_{v} + 3H\delta\mu_{v} - \frac{\Delta + 3K}{a^{2}} \left[a(\mu + p)v_{\chi} + 2H\Pi \right] = \delta\dot{\mu}_{v}^{(q)} + 3H\delta\mu_{v}^{(q)} - \frac{\Delta + 3K}{a^{2}}a(\mu + p)v_{\chi}^{(q)} + N_{5} + (\mu + p)\left(N_{2}^{(s)} + 3aHN_{6}^{(s)}\right),$$
(294)

$$\varphi_{\chi} + \alpha_{\chi} + 8\pi G\Pi = \varphi_{\chi}^{(q)} + \alpha_{\chi}^{(q)} - N_4^{(s)}, \quad \text{or} \quad \varphi_{\chi} + \alpha_{\chi} + 8\pi G\Pi_{\chi} = -N_{4\chi}^{(s)}, \tag{295}$$

$$\dot{v}_{\chi} + Hv_{\chi} - \frac{1}{a} \left(\alpha_{\chi} + \frac{\delta p_{v}}{\mu + p} + \frac{2}{3} \frac{\Delta + 3K}{a^{2}} \frac{\Pi}{\mu + p} \right) = \dot{v}_{\chi}^{(q)} + Hv_{\chi}^{(q)} - \frac{1}{a} \left(\alpha_{\chi}^{(q)} + \frac{\delta p_{v}^{(q)}}{\mu + p} \right) + N_{6}^{(s)},$$
(296)

$$\dot{\varphi}_{\chi} + H\varphi_{\chi} + 4\pi G(\mu + p)av_{\chi} + 8\pi GH\Pi = \dot{\varphi}_{\chi}^{(q)} + H\varphi_{\chi}^{(q)} + 4\pi G(\mu + p)av_{\chi}^{(q)} + \frac{1}{3}\left(N_0 - N_2^{(s)}\right) - HN_4^{(s)}.$$
(297)

These equations are presented using mixed gauge-invariant variables. We note that to the linear-order, $\delta \mu_v$, $-\varphi_{\chi}$, and $kv_{\chi}(\sim \nabla v_{\chi})$ closely resemble the Newtonian density perturbation, perturbed gravitational potential, and the perturbed velocity perturbation, respectively [34,35,13]. To the linear-order these equations were presented by Bardeen in 1980 [6]; see eqs. (4.3,4.8,4.4,4.5,4.7) in [6], and compared with our notation, see eq. (257). Using eq. (297) and eqs. (293,296,297) we can show

$$\Phi \equiv \varphi_v - \frac{K/a^2}{4\pi G(\mu+p)} \varphi_\chi$$
$$= \frac{H^2}{4\pi G(\mu+p)a} \left[\frac{a}{H} \left(\varphi_\chi - \varphi_\chi^{(q)} \right) \right]^{\cdot} + 2H^2 \frac{\Pi}{\mu+p} + \Phi^{(q)} + N_\Phi,$$
(298)

$$\dot{\Phi} = \frac{Hc_s^2 \Delta}{4\pi G(\mu+p)a^2} \left(\varphi_{\chi} - \varphi_{\chi}^{(q)}\right) - \frac{H}{\mu+p} \left(e + \frac{2}{3}\frac{\Delta}{a^2}\Pi\right) + \dot{\Phi}^{(q)} + N_{\dot{\Phi}},\tag{299}$$

where

$$\Phi^{(q)} \equiv \varphi_v^{(q)} - \frac{K/a^2}{4\pi G(\mu+p)} \varphi_\chi^{(q)},$$

$$N_{\Phi} \equiv \frac{H^2}{4\pi G(\mu+p)} \left[N_4^{(s)} + \frac{1}{3H} \left(N_2^{(s)} - N_0 \right) \right],$$

$$N_{\dot{\Phi}} \equiv \frac{1}{3} \left(1 - \frac{K/a^2}{4\pi G(\mu+p)} \right) \left(N_0 - N_2^{(s)} \right) - \frac{Hc_s^2}{4\pi G(\mu+p)} \left(\frac{1}{4} N_1 - HN_2^{(s)} \right) + \frac{K/a^2}{4\pi G(\mu+p)} HN_4^{(s)} - aHN_6^{(s)}.$$
(300)

We have introduced an entropic perturbation e by

$$\delta p \equiv c_s^2 \delta \mu + e, \quad c_s^2 \equiv \frac{\dot{p}}{\dot{\mu}}.$$
(301)

Defined in this way, e is not necessarily gauge-invariant to the second-order. To the linear order we have $e = \tilde{\pi}$ introduced above eq. (24). Combining eqs. (298,299) we can derive

$$\frac{H^{2}c_{s}^{2}}{(\mu+p)a^{3}} \left[\frac{(\mu+p)a^{3}}{H^{2}c_{s}^{2}} \dot{\Phi} \right]^{\cdot} - c_{s}^{2} \frac{\Delta}{a^{2}} \Phi = \frac{Hc_{s}}{a^{3}\sqrt{\mu+p}} \left[v'' - \left(\frac{z''}{z} + c_{s}^{2}\Delta\right) v \right] \\
= \frac{H^{2}c_{s}^{2}}{(\mu+p)a^{3}} \left\{ \frac{(\mu+p)a^{3}}{H^{2}c_{s}^{2}} \left[-\frac{H}{\mu+p} \left(e + \frac{2}{3}\frac{\Delta}{a^{2}}\Pi \right) + \dot{\Phi}^{(q)} + N_{\dot{\Phi}} \right] \right\}^{\cdot} - c_{s}^{2}\frac{\Delta}{a^{2}} \left(2H^{2}\frac{\Pi}{\mu+p} + \Phi^{(q)} + N_{\Phi} \right), \quad (302) \\
\frac{\mu+p}{H} \left[\frac{H^{2}}{(\mu+p)a} \left(\frac{a}{H}\varphi_{\chi} \right)^{\cdot} \right]^{\cdot} - c_{s}^{2}\frac{\Delta}{a^{2}}\varphi_{\chi} = \frac{\sqrt{\mu+p}}{a^{2}} \left[u'' - \left(\frac{(1/\bar{z})''}{1/\bar{z}} + c_{s}^{2}\Delta \right) u \right] \\
= \frac{4\pi G(\mu+p)}{H} \left[-\frac{H}{\mu+p} \left(e + \frac{2}{3}\frac{\Delta}{a^{2}}\Pi \right) - 2 \left(H^{2}\frac{\Pi}{\mu+p} \right)^{\cdot} + N_{\dot{\Phi}} - \dot{N}_{\Phi} \right] \\
+ \frac{\mu+p}{H} \left[\frac{H^{2}}{(\mu+p)a} \left(\frac{a}{H}\varphi_{\chi}^{(q)} \right)^{\cdot} \right]^{\cdot} - c_{s}^{2}\frac{\Delta}{a^{2}}}\varphi_{\chi}^{(q)}, \quad (303)$$

where we used

$$v \equiv z\Phi, \quad u \equiv \frac{1}{\bar{z}}\frac{a}{H}\varphi_{\chi}, \quad c_s z \equiv \frac{a\sqrt{\mu+p}}{H} \equiv \bar{z}.$$
 (304)

[v in eqs. (302-304) differs from the perturbed velocity related variable used in the rest of this paper.] The equation using v in the linear theory was first derived by Field and Shepley in 1968 [36], see also [37,21]. Using eq. (293), eq. (303) gives an equation for δ_v . Using eqs. (293-296) we can derive equation for δ_v in *another* form

$$\frac{\mu + p}{a^{2}\mu H} \left[\frac{H^{2}}{(\mu + p)a} \left(\frac{a^{3}\mu}{H} \delta_{v} \right)^{\cdot} \right]^{\cdot} - c_{s}^{2} \frac{\Delta}{a^{2}} \delta_{v} \\
= \frac{\Delta + 3K}{a^{2}} \left[\frac{e}{\mu} + \frac{2}{3} \frac{\Delta}{a^{2}} \frac{\Pi}{\mu} + 2 \frac{\mu + p}{\mu H} \left(\frac{H^{2}}{\mu + p} \Pi \right)^{\cdot} \right] + \frac{\mu + p}{a^{2}\mu H} \left[\frac{H^{2}}{(\mu + p)a} \left(\frac{a^{3}\mu}{H} \delta_{v}^{(q)} \right)^{\cdot} \right]^{\cdot} - c_{s}^{2} \frac{\Delta}{a^{2}} \delta_{v}^{(q)} \\
+ \frac{\mu + p}{\mu} \left\{ -\frac{1}{4} N_{1} + H N_{2}^{(s)} + \frac{\Delta + 3K}{a^{2}} \left(a N_{6}^{(s)} - N_{4}^{(s)} \right) + \frac{1}{a^{2}} \left[a^{2} \left(N_{2}^{(s)} + \frac{N_{5}}{\mu + p} + 3a H N_{6}^{(s)} \right) \right]^{\cdot} \right\}. \quad (305)$$

The above set of equations is valid for a general imperfect fluid. A minimally coupled scalar field can be regarded as an imperfect fluid with special fluid quantities. We additionally have an equation of motion of the field which is actually included in the energy and momentum conservation equations. In fact, the above set of equations is valid even for multi-component fluids and fields. In such cases, the fluid quantities become collective fluid quantities and we additionally need the energy and momentum conservation equations for individual fluid and the equations of motion for the individual field.

In the single scalar field case, from eqs. (212,190,293,301) we have

$$e = -\frac{1 - c_s^2}{4\pi G} \frac{\Delta + 3K}{a^2} \left(\varphi_{\chi} - \varphi_{\chi}^{(q)}\right) + N_e,$$

$$N_e \equiv \frac{1 - c_s^2}{4\pi G} \left(\frac{1}{4}N_1 - HN_2^{(s)}\right) + \delta p^{(q)} - \delta \mu^{(q)} + 3H(1 - c_s^2)a\Delta^{-1}\nabla^{\alpha}Q_{\alpha}^{(q)}.$$
(306)

Equation (298) remains valid, and eq. (299) becomes

$$\dot{\Phi} = \frac{Hc_A^2 \Delta}{4\pi G(\mu+p)a^2} \left(\varphi_{\chi} - \varphi_{\chi}^{(q)}\right) - \frac{H}{\mu+p} \left(N_e + \frac{2}{3}\frac{\Delta}{a^2}\Pi\right) + \dot{\Phi}^{(q)} + N_{\dot{\Phi}},\tag{307}$$

where

$$c_A^2 \Delta \equiv \Delta + 3(1 - c_s^2) K. \tag{308}$$

Therefore, eqs. (302,303,304) remain valid with c_s and e replaced by c_A and N_e ; in eq. (302) one can show that we can ignore the operator nature of Δ^{-1} in the c_A^2 .

The rotational perturbation and the gravitational wave are described by eqs. (208,210), respectively

$$\frac{[a^4(\mu+p)v_{\alpha}^{(v)}]}{a^4(\mu+p)} = -\frac{\Delta+2K}{2a^2}\frac{\Pi_{\alpha}^{(v)}}{\mu+p} + N_{6\alpha}^{(v)},\tag{309}$$

$$\ddot{C}^{(t)}_{\alpha\beta} + 3H\dot{C}^{(t)}_{\alpha\beta} - \frac{\Delta - 2K}{a^2}C^{(t)}_{\alpha\beta} = \frac{1}{a^3} \left[v^{(t)\prime\prime}_{\alpha\beta} - \left(\frac{a^{\prime\prime}}{a} + \Delta - 2K\right)v^{(t)}_{\alpha\beta} \right] = 8\pi G\Pi^{(t)}_{\alpha\beta} + N^{(t)}_{4\alpha\beta},\tag{310}$$

where $v_{\alpha\beta}^{(t)} \equiv a C_{\alpha\beta}^{(t)}$. We note that all the equations above in this section are valid for general K.

B. Solutions to the linear-order

1. Scalar-type

We consider a single component ideal fluid. Several known solutions in the literature are the followings. (i) In the large-scale limit (the super-sound-horizon scale), i.e., ignoring $c_s^2 \Delta$ term compared with z''/z and $(1/\bar{z})''/(1/\bar{z})$ terms in eqs. (302,303), we have general solutions ¹

$$\Phi(k,t) = C(k) - d(k) \frac{k^2}{4\pi G} \int^t \frac{c_s^2 H^2}{a^3(\mu+p)} dt,$$
(311)

$$\varphi_{\chi}(k,t) = 4\pi GC(k) \frac{H}{a} \int^{t} \frac{a(\mu+p)}{H^2} dt + d(k) \frac{H}{a}.$$
(312)

C(k) and d(k) are the coefficients of the growing and decaying solutions (in an expanding medium), respectively. To the second-order in the large-scale expansion we have

¹**k** is the wave vector with $k \equiv |\mathbf{k}|$. With the wave number k appearing in the equation the variables can be regarded as the Fourier transformed ones. To the *linear-order* each Fourier mode decouple from the other modes and evolves independently. The same equations in configuration space remain valid in Fourier space as well. Thus, we ignore specific symbols distinguishing the variables in the two spaces. Only in this subsection concerning the linear theory we use the Fourier transformation.

$$\Phi = C\left\{1 + k^2 \left[\int^{\eta} \bar{z}^2 \left(\int^{\eta} \frac{d\eta}{z^2}\right) d\eta - \int^{\eta} \bar{z}^2 d\eta \int^{\eta} \frac{d\eta}{z^2}\right]\right\} - d\frac{k^2}{4\pi G} \int^{\eta} \frac{d\eta}{z^2},\tag{313}$$

$$\varphi_{\chi} = 4\pi G C \frac{H}{a} \int^{\eta} \bar{z}^2 d\eta + d\frac{H}{a} \left\{ 1 + k^2 \left[\int^{\eta} \frac{1}{z^2} \left(\int^{\eta} \bar{z}^2 d\eta \right) d\eta - \int^{\eta} \bar{z}^2 d\eta \int^{\eta} \frac{d\eta}{z^2} \right] \right\}.$$
(314)

We emphasize that these solutions are valid for general K and Λ , and time-varying equation of state.

(ii) In the small-scale limits $[c_s^2 k^2 \gg z''/z, (1/\bar{z})''/(1/\bar{z})]$, if we further assume that c_s is constant in time, eqs. (302,303) give the general solutions:

$$v = z\Phi \propto e^{\pm ic_s k\eta}, \quad u = \frac{1}{\sqrt{\mu + p}}\varphi_\chi \propto e^{\pm ic_s k\eta}.$$
 (315)

(iii) For $K = 0 = \Lambda$ and $w \equiv p/\mu = \text{constant}$ we have exact solutions [6,38]. For the background, from eqs. (184,186), we have

$$a \propto t^{\frac{2}{3(1+w)}} \propto \eta^{\frac{2}{1+3w}}, \quad aH\eta = \frac{2}{1+3w}.$$
 (316)

Equation (304) gives $z \propto \bar{z} \propto a$, thus

$$\frac{z''}{z} = \frac{2(1-3w)}{(1+3w)^2} \frac{1}{\eta^2}, \quad \frac{(1/\bar{z})''}{(1/\bar{z})} = \frac{6(1+w)}{(1+3w)^2} \frac{1}{\eta^2}.$$
(317)

In this case, eqs. (302,303) become Bessel's equations with solutions:

$$v = z\Phi \propto \sqrt{\eta} \left(J_{\nu}(x), Y_{\nu}(x) \right), \quad x \equiv c_s k |\eta|, \quad \nu \equiv \frac{3(1-w)}{2(1+3w)},$$
(318)

$$u = \frac{1}{\sqrt{\mu + p}} \varphi_{\chi} \propto \sqrt{\eta} \left(J_{\bar{\nu}}(x), Y_{\bar{\nu}}(x) \right), \quad \bar{\nu} \equiv \frac{5 + 3w}{2(1 + 3w)}.$$
(319)

We have $\bar{\nu} = \nu + 1$. Using eqs. (298,299) we can normalize the solutions as

$$\Phi \equiv c_1(k)\frac{J_{\nu}(x)}{x^{\nu}} + c_2(k)\frac{Y_{\nu}(x)}{x^{\nu}},\tag{320}$$

$$\varphi_{\chi} = \frac{3(1+w)}{1+3w} \left(c_1(k) \frac{J_{\bar{\nu}}(x)}{x^{\bar{\nu}}} + c_2(k) \frac{Y_{\bar{\nu}}(x)}{x^{\bar{\nu}}} \right).$$
(321)

Equation (293) gives

$$\delta_v = \frac{(1+3w)^2}{6w} x^2 \varphi_\chi. \tag{322}$$

In the large-scale limit $(x \ll 1)$ we have

$$\Phi = \frac{c_1}{2^{\nu}\Gamma(\nu+1)} - 2^{\nu}\frac{\Gamma(\nu)}{\pi}x^{-2\nu}c_2, \qquad (323)$$

where for $\nu = 0$ we have additional $2 \ln x$ factor in the c_2 -mode. By matching with the general large-scale solution in eq. (311) we can identify

$$c_1 = 2^{\nu} \Gamma(\nu+1)C, \quad c_2 = -\frac{1}{3(1+w)} \frac{\pi}{2^{\nu} \Gamma(\bar{\nu})} \frac{x^{2\bar{\nu}}}{a^2 \eta} d.$$
(324)

In the large-scale limit $(x \ll 1)$ we have

$$\Phi \quad \propto \quad C, \ da^{-\frac{3}{2}(1-w)}, \tag{325}$$

$$\varphi_{\chi} \propto C, \ da^{-\frac{5+3w}{2}},$$
(326)

$$\delta_v \propto Ca^{1+3w}, \ da^{-\frac{3}{2}(1-w)} \propto Ct^{\frac{2(1+3w)}{3(1+w)}}, \ dt^{-\frac{1-w}{1+w}} \propto C\eta^2, \ d\eta^{-\frac{3(1-w)}{1+3w}}.$$
 (327)

Equation (327) follows from eq. (293) which gives $\delta_v \propto a^{1+3w} \varphi_{\chi} \propto \eta^2 \varphi_{\chi}$ in general. Equation (327) includes the well known solutions in the matter (w = 0) and radiation $(w = \frac{1}{3})$ eras [1]:

mde:
$$\delta_v \propto Ca, da^{-\frac{3}{2}} \propto Ct^{\frac{2}{3}}, dt^{-1} \propto C\eta^2, d\eta^{-3},$$

rde: $\delta_v \propto Ca^2, da^{-1} \propto Ct, dt^{-\frac{1}{2}} \propto C\eta^2, d\eta^{-1}.$ (328)

If we consider only the C-mode which is the relatively growing-mode in an expanding phase, we have

$$\Phi(\mathbf{x},t) = C(\mathbf{x}),\tag{329}$$

$$\varphi_{\chi}(\mathbf{x},t) = \frac{3+3w}{5+3w}C(\mathbf{x}). \tag{330}$$

The nontransient mode of Φ remains constant in the super-sound-horizon scale. Whereas, the one of φ_{χ} jumps as the background equation of state changes. Still, it is $-\varphi_{\chi}$, not $-\varphi_{\nu}$, which closely resembles the perturbed Newtonian gravitational potential [13].

The asymptotic solutions (i) and (ii) remain valid for the scalar field with K = 0; in this case we have c_s^2 replaced by 1. The background solutions for w = constant case considered in (iii), eq. (316), are valid for a scalar field with exponential potential of the following form [39]

$$V = \frac{1 - w}{12\pi G(1 + w)^2} e^{-\sqrt{24\pi G(1 + w)}\phi}, \quad \phi = \sqrt{\frac{1}{6\pi G(1 + w)}\ln t}.$$
(331)

For the perturbation, from eq. (308) and the prescription below it, the same equations of the fluid remain valid with the coefficient of Laplacian term replaced by 1 (for the field) instead of c_s^2 (for the fluid) [14]. Thus, the perturbation solutions in the fluid remain valid for such the scalar field with $x = k|\eta|$, instead of $x = c_s k|\eta|$ for the fluid case.

We emphasize that, if a solution is known in a given gauge condition the rest of the variables in all gauge conditions can be derived through simple algebra; such solutions are presented in tabular forms for an ideal fluid and a scalar field in [38,40,15]. Solutions in the situations of generalized gravity theories considered in §IV D can be found in [24]. The cases with multiple components of the fluids and fields were analysed in [41, 42].

2. Vector-type

The rotational perturbation is described by eq. (309). If we assume $\Pi_{\alpha}^{(v)} = 0$, we have a general solution

$$a \cdot a^3(\mu + p) \cdot v_{\alpha}^{(v)}(\mathbf{x}, t) = L_{\alpha}^{(v)}(\mathbf{x}).$$
(332)

Thus, to the linear-order the rotational perturbation is described by the conservation of angular momentum, and is transient in expanding media. We note that eq. (309) follows from the $T^b_{\alpha;b} = 0$, thus *independent* of the gravitational field equation. Thus the presence of scalar fields or the generalized gravity theories considered in §IV D do not affect the vector-type perturbation of the fluids [23].

3. Tensor-type

Now, we consider the gravitational wave with K = 0 and $\Pi_{\alpha\beta}^{(t)} = 0$. The basic equation is presented in eq. (310). (i) The general large-scale $(k^2 \ll a''/a)$ solution is

$$C_{\alpha\beta}^{(t)}(k,t) = c_{\alpha\beta}^{(t)}(k) + d_{\alpha\beta}^{(t)}(k) \int^t \frac{dt}{a^3}.$$
(333)

Thus, ignoring the transient mode $(d_{\alpha\beta})$ in an expanding phase the tensor-type perturbation is characterized by its conserved amplitude $c_{\alpha\beta}^{(t)}(k)$. (ii) In the small-scale limit $(k^2 \gg a''/a)$ we have a general solution

$$C^{(t)}_{\alpha\beta}(k,t) \propto \frac{1}{a} e^{\pm ik\eta}.$$
(334)

Thus the gravitational wave redshifts away.

(iii) For $K = 0 = \Lambda$ and w = constant, we have an exact solution:

$$v_{\alpha\beta}^{(t)} = aC_{\alpha\beta}^{(t)} \propto \sqrt{\eta} \left(J_{\nu}(x), Y_{\nu}(x) \right), \quad \nu \equiv \frac{3(1-w)}{2(1+3w)}, \quad x \equiv k|\eta|.$$
(335)

Thus, we set

$$C_{\alpha\beta}^{(t)} = c_{1\alpha\beta}^{(t)} \frac{J_{\nu}(x)}{x^{\nu}} + c_{2\alpha\beta}^{(t)} \frac{Y_{\nu}(x)}{x^{\nu}}.$$
(336)

In the large-scale limit $(x \ll 1)$ we have

$$C_{\alpha\beta}^{(t)} = \frac{1}{2^{\nu}\Gamma(\nu+1)}c_{1\alpha\beta}^{(t)} - 2^{\nu}\frac{\Gamma(\nu)}{\pi}x^{-2\nu}c_{2\alpha\beta}^{(t)},$$
(337)

where for $\nu = 0$ we have additional $2 \ln x$ factor in the $c_{2\alpha\beta}^{(t)}$ -mode. By matching with the general large-scale solution in eq. (333) we can identify

$$c_{1\alpha\beta}^{(t)} = 2^{\nu}\Gamma(\nu+1)c_{\alpha\beta}^{(t)}, \quad c_{2\alpha\beta}^{(t)} = \frac{\pi}{2^{\nu+1}\Gamma(\nu+1)} \frac{\eta x^{2\nu}}{a^2} d_{\alpha\beta}^{(t)}.$$
(338)

Thus,

$$C_{\alpha\beta}^{(t)} \propto c_{\alpha\beta}^{(t)}, \ d_{\alpha\beta}^{(t)} a^{-\frac{3}{2}(1-w)} \propto c_{\alpha\beta}^{(t)}, \ d_{\alpha\beta}^{(t)} t^{-\frac{1-w}{1+w}} \propto c_{\alpha\beta}^{(t)}, \ d_{\alpha\beta}^{(t)} \eta^{-\frac{3(1-w)}{1+3w}}.$$
(339)

C. Pressureless irrotational fluid

The equation of δ_v was derived in eq. (305) or can be derived from eqs. (303,296). In the pressureless case, a simpler route is to use the basic equations in eqs. (195-201).

We consider a pressureless fluid, thus $\delta p = 0 = \prod_{\alpha\beta}$, and ignore the vector-type perturbation. For the spatial gauge we take $\gamma \equiv 0$, thus $\beta = \chi/a$. If we take the temporal comoving gauge $(v \equiv 0)$ we have $Q_{\alpha} = 0$. Equation (201) gives $\alpha_v = -aN_{6v}^{(s)} = -\frac{1}{2}\beta_{v,\alpha}\beta_v^{,\alpha}$, thus α vanishes to the linear-order; in the pressureless medium, to the linear-order, the comoving particle follows geodesic, thus v = 0 implies $\alpha = 0$. From eqs. (200,198), first evaluating these in the comoving gauge, we can derive

$$\dot{\delta}_v = \kappa_v + \frac{1}{a} \nabla \cdot \left(\delta_v \nabla v_\chi \right), \tag{340}$$

$$\dot{\kappa}_v + 2H\kappa_v - 4\pi G\mu\delta_v = \frac{1}{a^2} \left[(\nabla v_\chi) \cdot (\nabla v_\chi)^{,\alpha} \right]_{|\alpha} + \dot{C}^{(t)}_{\alpha\beta} \left(\dot{C}^{(t)\alpha\beta} + \frac{2}{a^2} \chi_v^{,\alpha|\beta} \right), \tag{341}$$

where we have used $\chi_v = -av_{\chi}$ following from eq. (255), and $\kappa_v = \frac{1}{a}\Delta v_{\chi}$ following from eq. (197) with K = 0, both to the linear-order.

In order to compare with Newtonian analysis we introduce $\mathbf{u} = -\nabla v_{\chi}$ to the linear-order. By combining eqs. (340,341) we have

$$\ddot{\delta}_v + 2H\dot{\delta}_v - 4\pi G\bar{\mu}\delta_v = -\frac{1}{a^2} \left[a\nabla\cdot(\delta_v\mathbf{u})\right] + \frac{1}{a^2}\nabla\cdot(\mathbf{u}\cdot\nabla\mathbf{u}) + \dot{C}^{(t)}_{\alpha\beta}\left(\frac{2}{a}\nabla^{\alpha}u^{\beta} + \dot{C}^{(t)\alpha\beta}\right).$$
(342)

We note that, to the linear-order, the growing solution of the gravitational wave remains constant in time in the superhorizon scale, whereas it redshifts away $(C_{\alpha\beta}^{(t)} \propto a^{-1})$ in the sub-horizon scale; see §VIIB3. Ignoring the gravitational wave we reproduce correctly the corresponding Newtonian equation

$$\ddot{\delta} + 2H\dot{\delta} - 4\pi G\bar{\varrho}\delta = -\frac{1}{a^2} \left[a\nabla\cdot(\delta\mathbf{u})\right]^{\cdot} + \frac{1}{a^2}\nabla\cdot(\mathbf{u}\cdot\nabla\mathbf{u})\,. \tag{343}$$

We note that our eq. (342) is valid in the super-sound-horizon (Jeans scale) which is negligible in the pressureless medium, thus valid even in the super-(visual)-horizon. In the Newtonian context eq. (343) is valid to all orders in

perturbation, and follows from the mass conservation, the momentum conservation, and the Poissons' equation given as [10]

$$\dot{\delta} + \frac{1}{a} \nabla \cdot \mathbf{u} = -\frac{1}{a} \nabla \cdot (\delta \mathbf{u}), \qquad (344)$$

$$\dot{\mathbf{u}} + H\mathbf{u} + \frac{1}{a}\nabla\delta\Phi = -\frac{1}{a}\mathbf{u}\cdot\nabla\mathbf{u},\tag{345}$$

$$\frac{1}{a^2}\nabla^2\delta\Phi = 4\pi G\bar{\varrho}\delta.$$
(346)

To the linear-order these equations can be compared with the relativistic version in eqs. (294,296,293) with eq. (295). To the second-order, however, although the final result in eq. (342) coincides with the Newtonian one in eq. (343), we notice some difference between eqs. (340,341,294-296) with eq. (295) and eqs. (344-346). From eq. (197), to the second-order, we have

$$\kappa_v - \frac{\Delta}{a} v_\chi = N_{2v}^{(s)} - \frac{\Delta}{a^2} \left(\chi_v^{(q)} + a v_\chi^{(q)} \right).$$
(347)

Since the RHSs of this equation and of eq. (293) do not vanish we cannot directly relate $-\varphi_{\chi}$ and $-\nabla v_{\chi}$ (or $a\Delta^{-1}\nabla\kappa_v$) to the Newtonian counterparts $\delta\Phi$ and **u**, respectively. Still, we emphasize that the final equation identified in eq. (342) coincides exactly with the Newtonian one in eq. (343). For a similar conclusion in the relativistic situation, see [43]. For analyses of eq. (343) in the Newtonian context, see [44].

Using $\beta = 0$ as the spatial gauge condition Kasai [45] has derived a different equation compared with ours in eq. (342). Kasai [45] took both the comoving gauge $v \equiv 0$ and the original synchronous gauge which takes $\alpha = 0 = \beta$. As we have shown above eq. (340) in a pressureless medium, the comoving gauge v = 0 implies $\alpha = -\frac{1}{2}\beta_{,\alpha}\beta^{,\alpha}$, thus vanishes if we take $\beta = 0$ as the spatial gauge condition. However, in that gauge condition (the spatial *B*-gauge) the spatial gauge-mode is incompletely fixed. Thus, comparison with the Newtonian analyses is not transparent in that gauge condition.

1. Nonlinear equation based on 3 + 1 formulation

The general equation of the pressureless and irrotational ideal fluid can be derived from eqs. (10,12,13). The pressureless ideal fluid implies $S_{\alpha\beta} = 0$. We take the temporal comoving gauge condition, v = 0. Together with the irrotational condition we have $Q_{\alpha} = 0$, thus $J_{\alpha} = 0$. Equation (13) gives $N_{,\alpha} = 0$; if we use the normalization in eq. (55), we have N = a. Equations (10,12) give

$$K = \frac{\dot{E}}{E} - \frac{1}{N} \frac{E_{,\alpha}}{E} N^{\alpha}, \tag{348}$$

$$\dot{K} - \frac{1}{N}K_{,\beta}N^{\beta} - \frac{1}{3}K^2 = 4\pi GE - \Lambda + \bar{K}^{\alpha\beta}\bar{K}_{\alpha\beta}.$$
(349)

Apparently, the spatial *B*-gauge condition, $B_{\alpha} \equiv 0$, leads to $N_{\alpha} = 0$, thus simplifying the equations. However, such a gauge condition leaves the spatial gauge-mode removed incompletely. We prefer to take the spatial *C*-gauge condition which fixes the spatial gauge modes completely, in this way the analyses can be equivalently considered as spatially gauge-invariant ones. From eqs. (348,349) we can derive eq. (342) to the second-order. We notice that, in contrast with the Newtonian case, in general we anticipate to have infinite perturbation series expansion, and eq. (342) looks valid only to the second-order. If we have the higher-order terms nonvanishing, these can be regarded as purely relativistic effects.

2. Nonlinear equation based on 1+3 formulation

Assuming pressureless condition, eqs. (26,27,28) in the energy-frame ($\tilde{q}_a = 0$) become:

$$\dot{\tilde{\mu}} + \tilde{\theta}\tilde{\mu} = 0, \tag{350}$$

$$\tilde{a}_a = 0, \tag{351}$$

$$\dot{\tilde{\theta}} + \frac{1}{3}\tilde{\theta}^2 + 4\pi G\tilde{\mu} - \Lambda + 2\left(\tilde{\sigma}^2 - \tilde{\omega}^2\right) = 0.$$
(352)

In the energy-frame the frame-vector follows the possible energy flux, thus, the energy flux term \tilde{q}_a vanishes. Equations (350) and (352) can be combined to give

$$\left(\frac{\tilde{\tilde{\mu}}}{\tilde{\mu}}\right)^{\tilde{\tau}} - \frac{1}{3}\left(\frac{\tilde{\tilde{\mu}}}{\tilde{\mu}}\right)^2 - 4\pi G\tilde{\mu} + \Lambda - 2\left(\tilde{\sigma}^2 - \tilde{\omega}^2\right) = 0.$$
(353)

If we set $\tilde{\mu} \equiv \mu(1+\delta)$, where μ is the background energy density, equation (353) becomes

$$\tilde{\ddot{\delta}} - \frac{2}{3}\frac{\tilde{\mu}}{\mu}\tilde{\delta} - 4\pi G\mu \left(1+\delta\right)\delta - \frac{4}{3}\frac{\tilde{\dot{\delta}}^2}{1+\delta} - 2\left(\tilde{\sigma}^2 - \tilde{\omega}^2\right)\left(1+\delta\right) + \left(1+\delta\right)\left[\left(\frac{\tilde{\mu}}{\mu}\right)^2 - \frac{1}{3}\left(\frac{\tilde{\mu}}{\mu}\right)^2 - 4\pi G\mu + \Lambda\right] = 0.$$
(354)

This is a completely nonlinear equation.

Now, we assume an irrotational fluid. In the energy-frame, the comoving gauge condition leads to $\tilde{u}_{\alpha} \equiv 0$; this is equivalent to taking the normal-frame with vanishing energy flux. The momentum conservation equation in eq. (351) implies that our frame-vector follows geodesic path. In the comoving gauge eqs. (351,69) lead to $A = -\frac{1}{2}B^{\alpha}B_{\alpha}$ to the second-order. Thus, to the linear-order we have A = 0 which coincides with taking the synchronous gauge. We have

$$\tilde{\check{\mu}} = \tilde{\mu}_{,a}\tilde{u}^a = \partial_t\tilde{\mu} + \frac{1}{a}\tilde{\mu}_{,\alpha}B^{\alpha}; \quad \tilde{\check{\mu}} = \partial_t\mu, \quad \tilde{\check{\delta}} = \partial_t\delta + \frac{1}{a}\delta_{,\alpha}B^{\alpha}.$$
(355)

Only in the comoving gauge condition the covariant derivative along \tilde{u}^a simplifies to the second-order as in eq. (355). In this derivation the pressureless condition is used essentially. Thus, in this comoving gauge equation (354) becomes

$$\ddot{\delta} + 2H\dot{\delta} - 4\pi G\mu\delta = \frac{1}{a^2} \left(a^2 \delta \dot{\delta} \right)^{\cdot} + \frac{1}{3} \dot{\delta}^2 - \frac{1}{a^2} \left(a\delta_{,\alpha} B^{\alpha} \right)^{\cdot} - \frac{1}{a} \dot{\delta}_{,\alpha} B^{\alpha} + 2\tilde{\sigma}^2, \tag{356}$$

to the second-order. From this we can derive eq. (342) where $\tilde{\sigma}^2$ follows from eqs. (16,57,71).

3. Gravitational wave as a source

From eq. (342) considering the pure gravitational wave as the source for density perturbation we have

$$\ddot{\delta}_v + 2H\dot{\delta}_v - 4\pi G\mu\delta_v = \dot{C}^{(t)}_{\alpha\beta}\dot{C}^{(t)\alpha\beta} \equiv S.$$
(357)

In the matter dominated era an exact solution is given in eq. (336) with $\nu = \frac{3}{2}$; in the large-scale limit, considering the relatively growing mode, S vanishes, whereas, in the small scale limit it decays proportional to $1/(at)^2 \propto a^{-5}$, see eq. (334). If δ_g and δ_d denote two linear-order solutions, the general solution can be written as

$$\delta_{v}(\mathbf{x},t) = \delta_{g}(\mathbf{x},t) + \delta_{d}(\mathbf{x},t) + \int^{t} S(\mathbf{x},t') \frac{\delta_{g}(\mathbf{x},t')\delta_{d}(\mathbf{x},t) - \delta_{d}(\mathbf{x},t')\delta_{g}(\mathbf{x},t)}{\delta_{g}(\mathbf{x},t')\dot{\delta}_{d}(\mathbf{x},t') - \delta_{d}(\mathbf{x},t')\dot{\delta}_{g}(\mathbf{x},t')} dt'.$$
(358)

The particular solution is proportional to $a^{-5}t^2 \propto a^{-2}$, and decays more rapidly even compared with the decaying mode in the linear theory which behaves as t^{-1} .

D. Pure scalar-type perturbation

Equation (302) can be written as

$$\frac{H^2 c_s^2}{(\mu+p)a^3} \left\{ \frac{(\mu+p)a^3}{H^2 c_s^2} \left[\dot{\Phi} - \dot{\Phi}^{(q)} - N_{\dot{\Phi}} + \frac{H}{\mu+p} \left(e + \frac{2}{3} \frac{\Delta}{a^2} \Pi \right) \right] \right\}^{\cdot} = c_s^2 \frac{\Delta}{a^2} \left(\Phi - \Phi^{(q)} - 2H^2 \frac{\Pi}{\mu+p} - N_{\Phi} \right).$$
(359)

In the large-scale limit, if we could ignore the second-order spatial derivative terms, we have

$$\dot{\Phi} - \dot{\Phi}^{(q)} - N_{\dot{\Phi}} + \frac{H}{\mu + p} e \propto \frac{H^2 c_s^2}{(\mu + p)a^3}.$$
(360)

Now, we consider K = 0 and ideal fluids, thus e = 0. From eq. (298) we have $\Phi = \varphi_v$. Ignoring the second-order spatial derivatives, from eqs. (300,195-201,99-105) we can show that $N_{\dot{\Phi}} = (\varphi_v^2)^{\cdot}$. In the comoving gauge we have $\Phi^{(q)}|_v = \varphi_v^{(q)}|_v = 0$, see eq. (274). Thus, we have

$$\varphi_v - \varphi_v^2 = C(\mathbf{x}) + d(\mathbf{x}) \int^t \frac{H^2 c_s^2}{(\mu + p)a^3} dt.$$
(361)

Therefore, ignoring the transient mode in the expanding phase, we have

$$\varphi_v - \varphi_v^2 = C(\mathbf{x}),\tag{362}$$

which remains constant even to the second-order in perturbations [46,47]. Thus, φ_v is conserved to the second-order in the large-scale (super-sound-horizon) limit.

Equation (359) is valid for $p \neq 0$. For p = 0 we have a simpler form in eq. (299) which gives

$$\dot{\Phi} - \dot{\Phi}^{(q)} - N_{\dot{\Phi}} + \frac{H}{\mu} \left(e + \frac{2}{3} \frac{\Delta}{a^2} \Pi \right) = 0.$$
(363)

Equation (360) includes this as a case in the large-scale (super-sound-horizon) limit. Thus, the above results in eqs. (360-362) remain valid for general p.

From eq. (114) we notice that for a minimally coupled scalar field $\delta \phi = 0$ implies v = 0 to the second-order, thus

$$\varphi_v = \varphi_{\delta\phi},\tag{364}$$

and the uniform-field gauge coincides with the comoving gauge. Thus, the above analyses are valid even for a minimally coupled scalar field.

E. Pure rotation

In the case of pure rotation eqs. (208,105) provide a complete set for a single component fluid; for the multicomponent case see eqs. (209,107). Assuming $\Pi_{\alpha}^{(v)} = 0$ we have

$$\frac{\left[a^{4}(\mu+p)v_{\alpha}^{(v)}\right]^{\cdot}}{a^{4}(\mu+p)} = N_{6\alpha}^{(v)},
N_{6\alpha}^{(v)} = -\frac{1}{a} \left[v_{\alpha|\beta}^{(v)}B^{(v)\beta} + v_{\beta}^{(v)}B^{(v)\beta}_{\ |\alpha} - \nabla_{\alpha}\Delta^{-1}\nabla^{\beta} \left(v_{\beta|\gamma}^{(v)}B^{(v)\gamma} + v_{\gamma}^{(v)}B^{(v)\gamma}_{\ |\beta}\right)\right].$$
(365)

As a simple exercise, using eq. (250), one can check the gauge transformation properties of both sides. In the *C*-gauge condition $(C_{\alpha}^{(v)} \equiv 0)$ we have $B_{\alpha}^{(v)} = \Psi_{\alpha}^{(v)}$ which is gauge-invariant. From eqs. (332,206), to the linear-order we have $\Psi_{\alpha}^{(v)} \propto a^2(\mu + p)v_{\alpha}^{(v)} \propto a^{-2}$, thus from eq. (365) we have

$$\left[a^{4}(\mu+p)v_{\alpha}^{(v)}\right]^{\cdot} = a^{-3}\left[a^{3} \cdot a^{4}(\mu+p)N_{6\alpha}^{(v)}\right] \propto a^{-3}.$$
(366)

Thus, the additional second-order perturbation sourced by the RHS of eq. (365) behaves as

$$a^{4}(\mu+p)v_{\alpha}^{(v)} = L_{\alpha}^{(v)}(\mathbf{x}) + \left[a^{3} \cdot a^{4}(\mu+p)N_{6\alpha}^{(v)}\right] \int^{t} \frac{dt}{a^{3}}.$$
(367)

The time dependent nonlinear solution is proportional to $\int^t dt/a^3$; for w = constant it is proportional to $a^{-3(1-w)/2}$, thus it always decays (in expanding phase) for w < 1. The lower bound of integration which could give a temporally constant nonlinear solution can be absorbed to $L_{\alpha}^{(v)}$.

As we explained in §VIB2, to the linear-order, the *C*-gauge condition removes the rotational gauge-mode completely, whereas the *B*-gauge condition fails to fix it completely. That is, even after imposing the gauge condition we have some modes which are coordinate effect; under the *B*-gauge, from eq. (252) we have $\xi_{\alpha}^{(v)} = \xi_{\alpha}^{(v)}(\mathbf{x})$. Then, in eq. (365) we notice an ironic situation where the *B*-gauge condition gives vanishing quadratic terms, whereas these terms do not vanish in the *C*-gauge condition. That is, although we anticipate the nonlinear solution in the *C*-gauge in eq.

(367) is physical, in the *B*-gauge condition the RHS of eq. (366) vanishes, and we do not have the nonlinear solution in eq. (367). We can check this situation by using the gauge transformation property of $v_{\alpha}^{(v)}$ variable in the two gauge conditions.

Considering pure vector-type perturbation, from eqs. (175,177,234,249) we have

$$\hat{v}_{\alpha}^{(v)} = v_{\alpha}^{(v)} - v_{\beta}^{(v)} \xi^{(v)\beta}_{,\alpha} - v_{\alpha,\beta}^{(v)} \xi^{(v)\beta} + \nabla_{\alpha} \Delta^{-1} \nabla^{\beta} \left(v_{\gamma}^{(v)} \xi^{(v)\gamma}_{,\beta} + v_{\beta,\gamma}^{(v)} \xi^{(v)\gamma} \right).$$
(368)

Now, let the variables with hat and no-hat correspond to the ones in the *B*- and the *C*-gauge conditions, respectively. As the $\xi_{\alpha}^{(v)}$ s appear in quadratic combination, we need it only to the linear-order. From eq. (252), we have $\hat{B}_{\alpha}^{(v)} = B_{\alpha}^{(v)} + a\dot{\xi}_{\alpha}^{(v)}$. Since the hat indicates the *B* gauge, we have $a\dot{\xi}_{\alpha}^{(v)} = -B_{\alpha}^{(v)}$, thus $\xi_{\alpha}^{(v)} = -\int^{t} (B_{\alpha}^{(v)}/a) dt$. Thus, eq. (368) gives

$$a^{4}(\mu+p)\hat{v}_{\alpha}^{(v)}\Big|_{B-\text{gauge}} = \left\{ a^{4}(\mu+p)v_{\alpha}^{(v)} - \left[a^{3} \cdot a^{4}(\mu+p) N_{6\alpha}^{(v)} \right] \int^{t} \frac{dt}{a^{3}} \right\} \Big|_{C-\text{gauge}}.$$
(369)

Therefore, in the *B*-gauge the nonlinear solution in the *C*-gauge in eq. (367) disappears exactly. We note, however, that the solution in eq. (367) is physical (gauge-invariant) one in the *C*-gauge.

F. Pure gravitational wave

In the case of pure gravitational wave eqs. (210,103) provide a complete set. In the large-scale limit, thus ignoring second-order spatial derivative terms, and assuming K = 0 and $\Pi_{\alpha\beta}^{(t)} = 0$, we have

$$\begin{split} \ddot{C}^{(t)\alpha}{}_{\beta} + 3H\dot{C}^{(t)\alpha}{}_{\beta} &= N_{4\beta}^{(t)\alpha}, \\ N_{4\beta}^{(t)\alpha} &= N_{4\beta}^{\alpha} - \Delta^{-1} \left(\nabla^{\alpha} \nabla_{\gamma} N_{4\beta}{}^{\gamma} + \nabla_{\beta} \nabla^{\gamma} N_{4\gamma}{}^{\alpha} \right) + \frac{1}{2} \left(\Delta^{-1} \nabla^{\alpha} \nabla_{\beta} + \delta^{\alpha}_{\beta} \right) \Delta^{-1} \nabla^{\gamma} \nabla_{\delta} N_{4\gamma}{}^{\delta}, \\ N_{4\beta}{}^{\alpha} &= 2 \left(\dot{C}^{(t)\alpha\gamma} \dot{C}^{(t)}_{\beta\gamma} - \frac{1}{3} \delta^{\alpha}_{\beta} \dot{C}^{(t)\gamma\delta} \dot{C}^{(t)}_{\gamma\delta} \right). \end{split}$$
(370)

Notice that in this large-scale limit we have $C_{\alpha\beta}^{(t)} = \text{constant}$ as the relatively growing solution (in expanding phase) even to the second-order perturbation. In this sense, ignoring the transient mode in expanding phase, the amplitude of $C_{\alpha\beta}^{(t)}$ remains constant even to the second-order in perturbations [46,47].

G. Action formulation

We consider the action expanded to the second-order in perturbations which will give equations of motion to the linear-order in perturbation [48,37,9]. We consider the action for a scalar field in eq. (108). The perturbed action can be derived by using eqs. (94,98) and the ADM quantities presented in §III. To the background order, ignoring the surface terms, we have

$$S_{\rm BG} = \frac{1}{16\pi G} \int \sqrt{g^{(3)}} a^3 \left[-6\left(\frac{\dot{a}}{a}\right)^2 + \frac{6K}{a^2} + 16\pi G\left(\frac{1}{2}\dot{\phi}^2 - V\right) \right] dt d^3x.$$
(371)

To the second-order perturbation, ignoring the surface terms, the pure gravitational wave part becomes

$$S_{\rm GW} = \frac{1}{16\pi G} \int \sqrt{g^{(3)}} a^3 \left(\dot{C}^{(t)\alpha\beta} \dot{C}^{(t)}_{\alpha\beta} - \frac{1}{a^2} C^{(t)\alpha\beta|\gamma} C^{(t)}_{\alpha\beta|\gamma} - \frac{2K}{a^2} C^{(t)\alpha\beta} C^{(t)}_{\alpha\beta} \right) dt d^3x.$$
(372)

This action is valid for arbitrary number of scalar fields and fluids with vanishing tensor-type anisotropic stress. Now, we consider the pure scalar-type perturbation. We assume K = 0. To the second-order perturbation, ignoring the surface terms, we have

$$S_{\text{scalar}} = \frac{1}{2} \int a^3 \left\{ \delta \dot{\phi}_{\varphi}^2 - \frac{1}{a^2} \delta \phi_{\varphi}^{,\alpha} \delta \phi_{\varphi,\alpha} + \frac{H}{a^3 \dot{\phi}} \left[a^3 \left(\frac{\dot{\phi}}{H} \right)^* \right]^* \delta \phi_{\varphi}^2 \right\} dt d^3 x$$
$$= \frac{1}{2} \int a^3 \frac{\dot{\phi}^2}{H^2} \left(\dot{\varphi}_{\delta\phi}^2 - \frac{1}{a^2} \varphi_{\delta\phi}^{,\alpha} \varphi_{\delta\phi,\alpha} \right) dt d^3 x.$$
(373)

In this case we used the linear-order equations of motions, thus it is an on-shell action. In the second step we used eq. (255).

Maldacena has considered the perturbed action to the third-order in perturbations which is needed to have equations of motion valid to the second-order [47]. For the temporal gauge he used two gauge conditions, the uniform-field gauge $(\delta \phi \equiv 0)$ and the uniform-curvature gauge ($\varphi \equiv 0$), and the *C*-gauge for the spatial and rotational ones. Compared with our notations we have

$$\zeta_{\text{Maldacena}} = \varphi_{\delta\phi} - \varphi_{\delta\phi}^2, \quad \varphi_{\text{Maldacena}} = \delta\phi_{\varphi}. \tag{374}$$

Thus, $\zeta_{\text{Maldacena}}$ is conserved in the large-scale limit, see eqs. (362,364). To the linear-order, from eq. (255) we have $\varphi_{\delta\phi} = -(H/\dot{\phi})\delta\phi_{\varphi}$, thus $\zeta_{\text{Maldacena}} = -(H/\dot{\phi})\varphi_{\text{Maldacena}}$.

VIII. DISCUSSIONS

We have presented the basic equations to investigate the second-order perturbation of the Friedmann world model. In order to serve as a convenient reference for future studies and applications we have presented some useful relations and quantities needed for the second-order perturbations. The present study is, apparently, not entirely new in this rich field of cosmology and the large-scale structure formation. In the late sixties Tomita has presented a series of work on the subject in the context of a fluid [49]. Studies in the context of the ideal fluid or the minimally coupled field can be found in [31,50,47]. In the case of a pressureless irrotational fluid, see [45,43]. The case with the null-geodesic equations was studied in [51]. And, the case with Boltzmann equation was considered in [52].

Compared with the previous works, perhaps we could emphasize the following as the new points in our work:

(i) We present the complete sets of perturbed equations in the gauge-ready form, so that we could easily apply the equations to any gauge conditions which make the mathematical analyses of given problems simplest.

(ii) We consider the most general Friedmann background with K and Λ . Previous studies have considered the flat Friedmann background only.

(iii) We consider the most general imperfect fluid situation which includes multiple imperfect fluids with general interactions among them. In addition we also include minimally coupled scalar fields, a class of generalized gravity theories, the electromagnetic fields, the null geodesic, and the relativistic Boltzmann equation.

(iv) In §VII A we present closed form equations which are similar to the known ones in linear theory.

(v) In VIIC we show that up to the second-order in perturbations the relativistic pressureless fluid coincides exactly with the Newtonian one. We note that suitable choices and combinations of different gauges (thus gauge-invariant combinations) are important to show the equivalence.

(vi) In VIID we have derived the large-scale (super-sound-horizon) conserved quantity to the second-order, φ_v , directly from the differential equation governing its evolution. This conserved variable was first studied by Salopek and Bond

Our equations are suitable to handle the nonlinear evolutions in the perturbative manner. If we have the solutions to the linear-order (see §VII B for some examples), the evolution of second-order perturbations can be derived using the quadratic combination of the linear variables as sources; our basic sets of equations in §V C and some closed forms in §VII A are presented with such a purpose. As long as we take such a perturbative approach our formulation in this work can be trivially extended to any higher-order perturbation; except for the fact that, of course, the needed algebra would be quite demanding. We also have shown in §VI that the gauge issue can be similarly handled even in such higher-order perturbation.

Our formulation can be applied using several different methods in the following.

(i) Quasilinear analyses using Fourier analyses as often used in the Newtonian case [44]. In this approach the quadratic combination of the linear-order terms will lead to the mode-mode coupling among different scales, as well as among different types of perturbations.

(ii) Nonlinear backreaction. In our approach we have assumed the presence of "fictitious" background metric which is spatially homogeneous and isotropic. As the basic equations of Einstein gravity are nonlinear the nonlinear fluctuations in the metric and the matter can affect the background world model. One anticipates to recover the

background Friedmann world model through averaging the more realistic lumpy world model and finding best fit to the idealized world model [53].

(iii) Fitting and averaging. Our basic equations in §IV are presented without separating the background order quantities from the perturbed-order ones. Thus, the equations are suitable for the operation of averaging. Using our formulation we could apply and check the various different averaging prescriptions suggested in the literature [53,54,45].

Our perturbative formulation would be a useful complement to the following formulations aiming to investigate the nonlinear evolutions of the cosmological structures.

(i) The large-scale (long wavelength) approximation or the spatial gradient expansion studied by Salopek, Tomita, Deruelle and others in [46,55].

(ii) Cosmological post-Newtonian formulation studied by Futamase, Tomita and others in [56].

(iii) Relativistic Zel'dovich approximation studied in [45,43,57].

(iv) General (spatially inhomogeneous and anisotropic) solutions near singularity where the large-scale conditions are well met; in such a situation it was shown that the spatially different points decouple and evolve separately. These were studied by Belinsky, Lifshitz, Khalatnikov, and others in [58].

Our general formulation can be used to study the following situations anticipated during the evolution of our universe:

(i) We can check the limit of linear theory. Current cosmological observations can be successfully explained within the current standard theoretical paradigm. In that paradigm the linear perturbation theory plays significant roles to explain the quantum generation stage in the early universe and the classical evolution processes in the large-scale and in the early era. The linear theory provides a self-consistent explanation of some important aspects of the origin and evolution of large-scale structures. However, the limit of the linear theory *cannot* be estimated within the linear theory. We expect the second-order perturbation theory could provide a meaningful ways to investigate such limits.

(ii) We can investigate the quasilinear process in the relativistic context. In the literature it is commonly assumed that the relativistic linear perturbation theory is sufficient to handle the large-scale structure, and the nonlinear processes occur only in the Newtonian context which are often handled by the numerical simulations. The quasilinear evolution would be useful to investigate the transition regions between the linear and nonlinear evolutions. Our perturbative approach may have its own limit, because if we find the importance of second-order contributions it may naturally follow that higher-order contributions would become important immediately as well. Thus, we anticipate, if successful, the relativistic quasilinear analyses can be developed similarly as the Newtonian cosmological quasilinear analyses studied in [44].

(iii) Fate of fluctuations in the collapsing phase, and possibly through a bounce. The fluctuations of single component medium and the gravitational wave are described by the second-order differential equations. In the linear stage and in the large-scale limit, we have general solutions in eqs. (311,312,333). In an expanding phase, the C-mode is relatively growing and the *d*-mode is decaying, thus transient. If the initial conditions (say, generated from the quantum fluctuations) are imposed in the early expanding phase, the *d*-mode parts disappear in a few e-folding time of the scale factor increase, thus uninteresting. The relatively growing modes both for the scalar- and tensortype perturbations are characterized by the conserved amplitude of certain gauge-invariant variable, φ_v (and the curvature variables in other gauges) and $C_{\alpha\beta}^{(t)}$, in the large-scale limit. In the collapsing phase, however, the roles of growing and decaying modes are switched. In the collapsing phase the *d*-mode, and the vector-mode as well, grows quite rapidly, see eqs. (325,339,332); our solutions in §VII B also cover the collapsing phase by considering $t \to |t|$ with t approaching 0, see [59]. Thus, the linear perturbations grow rapidly and inevitably reach the nonlinear stage [6,60]. Such growths would cause the transition of our simple (spatially homogeneous and isotropic) background world model to the anisotropic and inhomogeneous ones studied in [58]. Although we anticipate the perturbations would become quite nonlinear, we hope we could investigate the transition region based on our second-order perturbation formulation. One simplifying fact is that in the collapsing phase the local range covered by the dynamical time scale $\sim H^{-1}$ shrinks relative to the comoving scale, thus effectively the scales we are interested in satisfy the conditions of large-scale limit². Such large-scale conditions are well met for a given comoving scale during the early evolution stage (near singularity) and as the background model approaches the singularity in the collapsing phase. Investigation of situations in the collapsing and subsequent bouncing background is left for future study; for evolutions under the linear assumption, see [59]. For the general cosmological investigation near singularity, see [58].

(iv) It is well known that the nonlinear effect (either in the quantum generation or in the classical evolution

²Professor James Bardeen and Ewan Stewart have suggested that the large-scale (long wavelength) expansion or the spatial gradient expansion technique [46,55] would be useful to investigate such situations.

processes) could lead to non-Gaussian effects in the observed quantities of the CMB anisotropies and the large-scale galaxy distribution and motion. Maldacena has recently investigated such an effect on the CMB based on the second-order perturbation theory, see [47]. The first year WMAP data shows no positive detection of non-Gaussian nature of the CMB sky maps under a couple of non-Gaussianity tests [61].

(v) Evolutions in the super-horizon scale where the scale is larger than causal domain during the dynamic time scale. See §VIID and VIIF for the conserved quantities to the second-order which were found by Salopek and Bond in [46].

(vi) In §VIIC we have shown that to the second-order a pressureless fluid with pure scalar-type perturbation reproduces Newtonian result. It is likely that the relativistic effect appears in the higher-order perturbation which is left for future investigation.

ACKNOWLEDGMENTS

We thank Ewan Stewart for encouraging us to continue the project and sharing his enthusiasm and opinions on the subject. We also wish to thank Prof. George Efstathiou for inviting us to the Institute of Astronomy where the project was reassumed. JH wishes to thank Toshi Futamase and Masumi Kasai for their hospitality and encouragement in the beginning stage of the work at Hirosaki. JH and HN were supported by grant No. R02-2003-000-10051-0 and No. R04-2003-10004-0, respectively, from the Basic Research Program of the Korea Science and Engineering Foundation.

- [1] E.M. Lifshitz, J. Phys. (USSR) 10, 116 (1946); E.M. Lifshitz and I.M. Khalatnikov, Adv. Phys. 12, 185 (1963).
- [2] A.A. Friedmann, Zeitschrift für Physik **10**, 377 (1922); translated in *Cosmological-constants: papers in modern cosmology*, edited by J. Bernstein and G. Feinberg (Columbia Univ. Press, New York, 1986), p49.
- [3] C.L. Bennett, et al., astro-ph/0302207; T.J. Pearson, et al., astro-ph/0205388; C.L. Kuo, et al., astro-ph/0212289.
- [4] J.R. Bond and G. Efstathiou, Mon. Not. R. Astron. Soc. 226, 655 (1987).
- [5] H. Noh and J. Hwang, Phys. Rev. D 52, 1970, 5643 (1995); H. Noh, *ib id.* 53, 690, 4311 (1996).
- [6] J.M. Bardeen, Phys. Rev. D 22, 1882 (1980).
- [7] J.M. Bardeen, Particle Physics and Cosmology, edited by L. Fang and A. Zee (Gordon and Breach, London, 1988), p1.
- [8] H. Kodama and M. Sasaki, Prog. Theor. Phys. Suppl. 78, 1 (1984).
- [9] V.F. Mukhanov, H.A. Feldman, and R.H. Brandenberger, Phys. Rep. 215, 203 (1992).
- [10] P.J.E. Peebles, The large-scale structure of the universe, (Princeton Univ. Press, Princeton, 1980); Ya.B. Zel'dovich and I.D. Novikov, Relativistic astrophysics, Vol 2, The structure and evolution of the universe, (Univ. Chicago Press, Chicago, 1983).
- [11] T. Padmanabhan, Structure formation in the universe, (Cambridge University Press, Cambridge, 1993); P.J.E. Peebles, Principles of physical cosmology, (Princeton Univ. Press, Princeton, 1993); A.R. Liddle and D.H. Lyth, Cosmological inflation and large-scale structure (Cambridge Univ. Press, Cambridge, 2000).
- [12] J. Hwang, Astrophys. J. **375**, 443 (1991).
- [13] J. Hwang and H. Noh, Gen. Rel. Grav. **31**, 1131 (1999).
- [14] J. Hwang, Phys. Rev. D 48, 3544 (1993); Class. Quant. Grav. 11, 2305 (1994).
- [15] J. Hwang, Astrophys. J. **427**, 542 (1994).
- [16] R. Utiyama, Phys. Rev. 101, 1597 (1958); T.W.B. Kibble, J. Math. Phys. 2, 212 (1961).
- [17] R.K. Sachs and A.M. Wolfe, Astrophys. J. 147, 73 (1967).
- [18] R. Arnowitt, S. Deser, and C.W. Misner, in *Gravitation: an introduction to current research*, edited by L. Witten (Wiley, New York, 1962) p. 227.
- [19] J. Ehlers, Proceedings of the mathematical-natural science of the Mainz academy of science and literature, Nr. 11, 792 (1961), translated in Gen. Rel. Grav. 25, 1225 (1993).
- [20] G.F.R. Ellis, in General relativity and cosmology, Proceedings of the international summer school of physics Enrico Fermi course 47, edited by R. K. Sachs (Academic Press, New York, 1971), 104; in Cargese Lectures in Physics, edited by E. Schatzmann (Gorden and Breach, New York, 1973), 1.
- [21] J. Hwang and E.T. Vishniac, Astrophys. J. 353, 1 (1990).
- [22] W. Israel, Ann. Phys. 100, 310 (1976).
- [23] J. Hwang, Class. Quant. Grav. 7, 1613 (1990).
- [24] J. Hwang, Phys. Rev. D 53, 762 (1996); J. Hwang and H. Noh, *ib id.* 54, 1460 (1996); Class. Quant. Grav. 15, 1387, 1401 (1998).

- [25] H. Noh and J. Hwang, Phys. Rev. D 55, 5222 (1997); 59, 047501 (1999).
- [26] J. Hwang, Class. Quant. Grav. 14, 1981 (1997).
- [27] J. Kristian and R.K. Sachs, Astrophys. J. 143, 379 (1966).
- [28] R.W. Lindquist, Ann. Phys., 37, 487 (1966); J. Ehlers, in General relativity and cosmology, Proceedings of the international summer school of physics Enrico Fermi course 47, edited by R.K. Sachs (Academic Press, New York, 1971), p1.
- [29] J. Hwang and H. Noh, Phys. Rev. D **65**, 023512 (2002).
- [30] J. Hwang and H. Noh, Phys. Rev. D 59, 067302 (1999); J. Hwang, T. Padmanabhan, O, Lahav, and H. Noh, *ib id.* 65, 043005 (2002).
- [31] A.H. Taub, J. Math. Phys. 2, 787 (1961); L.R.W. Abramo, R.H. Brandenberger, and V.F. Mukhanov, Phys. Rev. D 56, 3248 (1997); S. Matarrese, S. Mollerach, and M. Bruni, *ib id.* 58, 043504 (1998).
- [32] G.F.R. Ellis and M. Bruni, Phys. Rev. D 40, 1804 (1989); A. Woszczyna and A. Kułak, Class. Quant. Grav. 6, 1665 (1989).
- [33] G.F.R. Ellis, M. Bruni, and J. Hwang, Phys. Rev. D 42, 1035 (1990).
- [34] E.R. Harrison, Rev. Mod. Phys. **39**, 862 (1967).
- [35] H. Nariai, Prog. Theor. Phys. 41, 686 (1969).
- [36] G.B. Field and L.C. Shepley, Astrophys. Space. Sci. 1, 309 (1968).
- [37] V.N. Lukash, Sov. Phys. JETP Lett. **31**, 596 (1980); Sov. Phys. JETP **52**, 807 (1980); G.V. Chibisov and V.F. Mukhanov, Mon. Not. R. Astron. Soc. **200**, 535 (1982); V.F. Mukhanov, Sov. Phys. JETP **68**, 1297 (1988).
- [38] J. Hwang, Astrophys. J. **415**, 486 (1993).
- [39] F. Lucchin and S. Matarrese, Phys. Rev. D 32, 1316 (1985).
- [40] J. Hwang, Astrophys. J. **427**, 533 (1994).
- [41] H. Kodama and M. Sasaki, Int. J. Mod. Phys. A 1, 265 (1986); 2, 491 (1987).
- [42] J. Hwang and H. Noh, Phys. Lett. B, 495, 277 (2000); Class. Quant. Grav. 19, 527 (2002).
- [43] E. Bertschinger and A.J.S. Hamilton, Astrophys. J. 435, 1 (1994); L. Kofman and D. Pogosyan, *ib id.* 442, 30 (1995); S. Matarrese and D. Terranova, Monthly Not. R. Astron. Soc. 283, 400 (1996).
- [44] E.T. Vishniac, Monthly Not. R. Astron. Soc. 203, 345 (1983); M.H. Goroff, B. Grinstein, S.-J. Rey, and M.B. Wise, Astrophys. J., 311, 6 (1986); N. Makino, M. Sasaki, and Y. Suto, Phys. Rev. D 46, 585 (1992).
- [45] M. Kasai, Phys. Rev. Lett. 69, 2330 (1992); Phys. Rev. D 47, 3214 (1993).
- [46] D.S. Salopek and J.R. Bond, Phys. Rev. D 42, 3936 (1990).
- [47] J. Maldacena, JHEP **0305**, 013 (2003).
- [48] L.H. Ford and L. Parker, Phys. Rev. D 16, 1601 (1977).
- [49] K. Tomita, Prog. Theor. Phys. 37, 831 (1967); 45, 1747 (1971); 47, 416 (1972).
- [50] M. Bruni, S. Matarrese, S. Mollerach, and S. Sonego, Class. Quant. Grav. 14, 2585 (1997); W. Unruh, astro-ph/9802323;
 V. Acquaviva, N. Bartolo, S. Matarrese, and A. Riotto, astro-ph/0209156; G. Rigopoulos, astro-ph/0212141.
- [51] T. Pyne and S.M. Carroll, Phys. Rev. D 53, 2920 (1996); S. Mollerach and S. Matarrese, *ib id.* 56, 4494 (1997).
- [52] W. Hu, D. Scott, J. Silk, Phys. Rev. D 49, 648 (1994); R. Maartens, T. Gebbie, and G.F.R. Ellis, *ib id.* 59, 083506 (1999).
- [53] G.F.R. Ellis, in *General Relativity and Gravitation*, edited by B. Bertotti, et al. (Reidel, Dordrecht, 1984) p215; G.F.R. Ellis and W. Stoeger, Class. Quant. Grav. 4, 1697 (1987).
- [54] M.F. Shirokov and I.Z. Fisher, Sov. Astron. 6, 699 (1963); R.A. Isaacson, Phys. Rev. 166, 1263, 1272 (1968); N.R. Sibgatullin, Sov. Phys. Doklady 16, 697 (1972); A.V. Byalko, Sov. Phys. JETP 38, 421 (1974); H. Nariai, Prog. Theor. Phys. 52, 1539 (1974); 53, 656 (1975); 54, 1356 (1975); L.S. Marochnik, N.V. Pelikhov, and G.M. Vereshkov, Astrophys. Space Sci. 34, 249, 281, (1975); G.M. Vereshkov and A.N. Poltavtsev, Sov. Phys. JETP 44, 1 (1976); A.M. Krymsky, L.S. Marochnik, P.D. Naselsky, and N.V. Pelikhov, Astrophys. Space Sci. 55, 325, (1978); L.S. Marochnik, P.D. Naselsky, and N.V. Pelikhov, Astrophys. Space Sci. 55, 325, (1978); L.S. Marochnik, P.D. Naselsky, and N.V. Pelikhov, *ib id.* 67, 261, (1980); L.S. Marochnik, Sov. Astron. 24, 518, 651, (1980); 25, 8 (1981); T.W. Noonan, Gen. Rel. Grav. 16, 1103 (1984); I.K. Rozgachëva, Sov. Astron. 31, 14 (1987); T. Futamase, Phys. Rev. D 53, 681 (1996); M. Carfora and K. Piotrkowska, *ib id.* 52, 4393 (1995); J.P. Boersma, *ib id.* 57, 798 (1998); W.R. Stoeger, A. Helmi, and D. Torres, gr-qc/9904020; Y. Nambu, Phys. Rev. D 62, 104010 (2000); 63, 044013 (2001); 65, 104013 (2002); T. Buchert and M. Carfora, Class. Quant. Grav. 19, 6109 (2002).
- [55] K. Tomita, Prog. Theor. Phys. 48, 1503 (1972); 54, 730 (1975); D.S. Salopek, Phys. Rev. D 43, 3214 (1991); D.S. Salopek and J.M. Stewart, Class. Quant. Grav. 9, 1943 (1992); Phys. Rev. D 48, 719 (1993); 47, 3235 (1993); K. Tomita, *ib id.* 48, 5634 (1993); G.L. Comer, N. Deruelle, D. Langlois, and J. Parry, *ib id.* 49, 2759 (1994); J. Parry, D.S. Salopek, and J.M. Stewart, *ib id.* 49, 2872 (1994); K. Tomita and N. Deruelle, *ib id.* 50, 7216 (1994); N. Deruelle and D. Langlois, *ib id.* 52, 2007 (1995); Y. Nambu and A. Taruya, Class. Quant. Grav. 13, 705 (1996); G.L. Comer, *ib id.* 14, 407 (1997); G.L. Comer, N. Deruelle, and D. Langlois, Phys. Rev. D 55, 3497 (1997); D.S. Salopek, *ib id.* 56, 2057 (1997); Y. Nambu and Y. Yamaguchi, *ib id.* 60, 104011 (1999); Y. Nambu and S. Ohokata, Class. Quant. Grav. 19, 4263 (2002).
- [56] T. Futamase, Phys. Rev. Lett. **61**, 2175 (1988); K. Tomita, Prog. Theor. Phys. **79**, 258 (1988); T. Futamase, Mon. Not. R. Astron. Soc. **237**, 187 (1989); K. Tomita, Prog. Theor. Phys. **85**, 1041 (1991); T. Futamase, *ib id.* **86**, 389 (1991); **89**, 581 (1993); M. Shibata and H. Asada, *ib id.* **94**, 11 (1995); M. Takada and T. Futamase, Monthly Not. R. Astron. Soc. **306**, 64 (1999).
- [57] S. Matarrese, O. Pantano, and D. Saez, Phys. Rev. D 47, 1331 (1993); Monthly, Not. R. Astron. Soc. 271, 513 (1994);
 K.M. Croudace, J. Parry, D.S. Salopek, and J.M. Stewart, Astrophys. J. 423, 22 (1994); D.S. Salopek, J.M. Stewart, and

K.M. Croudace, Monthly, Not. R. Astron. Soc. **271**, 1005 (1994); M. Kasai, Phys. Rev. D **52**, 5605 (1995); H. Russ, M. Morita, M. Kasai, and G. Börner, *ib id.* **53**, 6881 (1996); M. Morita, K. Nakamura, and M. Kasai, *ib id.* **57**, 6094 (1998).

- [58] V.A. Belinskii, I.M. Khalatnikov, and E.M. Lifshitz, Adv. Phys. **19**, 525 (1970); I.M. Khalatnikov, and E.M. Lifshitz, Phys. Rev. Lett. **24**, 76 (1970); V.A. Belinskii, E.M. Lifshitz, and I.M. Khalatnikov, Sov. Phys. JETP **35**, 838 (1972); D. Eardley, E. Liang, and R. Sachs, J. Math. Phys. **13**, 99 (1972); V.A. Belinskii, I.M. Khalatnikov, and E.M. Lifshitz, Adv. Phys. **31**, 639 (1982); L. Andersson and A.D. Rendall, Commun. Math. Phys. **218**, 479 (2001); B.K. Berger, Living Rev. Relativity 2002-1, gr-qc/0201056.
- [59] J. Hwang and H. Noh, Phys. Rev. D 65, 124010 (2002).
- [60] D.H. Lyth, Phys. Lett. B **526** 171 (2002).
- [61] E. Komatsu, et al., Astrophys. J. Suppl. 148, 119 (2003).