

MESSENGER OF THE EXTREME UNIVERSE

A. V. OLINTO

*Department of Astronomy & Astrophysics, Enrico Fermi Institute,
Center for Cosmological Physics, University of Chicago,
5640 S. Ellis Avenue, Chicago, IL 60637, USA
E-mail: olinto@oddjob.uchicago.edu*

We give a brief overview of the highest energy messengers: very high energy photons, neutrinos, and ultra-high energy cosmic rays. The mysterious workings of the extreme universe should soon be unveiled with new observatories now under construction.

1. Introduction

The highest energy messengers arriving on Earth bring information from the most energetic events occurring in the universe since the Big Bang. These extreme environments cannot be reproduced in terrestrial laboratories. Instead, physics at these extremes is studied using telescopes and detectors that span many orders of magnitude in energy (from 10^9 eV photons to 10^{20} eV cosmic rays). Sources range from Supermassive black holes (masses $\sim 10^9 M_\odot$) in the centers of galaxies (known as active galactic nuclei or AGN) to yet to be discovered sources of the highest energy cosmic rays. These astrophysical environments not only call for applications of physics to extreme cosmic systems but they also challenge fundamental physics beyond what is currently known.

2. High-Energy Gamma-rays

The most important messengers in the history of astronomy and astrophysics are photons. Visible photons started astronomy and photons are still the clearest messengers of most phenomena in the universe since photons are neutral, massless, abundantly produced, and easily detected. They move the fastest and in the absence of strong gravitational fields, they point straight back to their sources. In contrast, cosmic protons and electrons are charged and usually diffuse in cosmic magnetic fields. Electrons and

positrons lose energy fast while unstable particles, such as neutrons, decay on their way to Earth. Neutrinos have some of the advantages of photons, they are neutral and have very small masses, but they only interact through the weak interactions, making their detection especially challenging. Photons are easily detected and easily produced by cosmic events that accelerate charged particles. We observe cosmic photons from radio wavelengths ($\sim 10^{-7}$ eV) to the highest energy photons observed thus far, gamma-rays of tens of TeVs (1 TeV = 10^{12} eV). Here we focus on gamma-rays, i.e. photons above 10^5 eV.

Gamma-ray astronomy was born once rockets and balloons reached above the absorbing atmosphere of the Earth. By 1961, the Explorer-XI satellite had detected a few gamma-rays consistent with a cosmic gamma-ray background. By the 1970s, SAS-2 and COS-B satellites produced maps of the sky in gamma-rays where the Galactic gamma-ray emission, the gamma-ray background, and the presence of a few point sources were identified. The most surprising discovery in early gamma-ray astronomy was made by the defense Vela satellite series. These were designed to monitor compliance with the 1963 Limited Test Ban Treaty by detecting flashes of gamma-rays from man-made nuclear explosions. Instead, Vela 4A and 4B in 1967, followed by Vela 5A and 5B in 1969, discovered many flashes of gamma-rays coming not from the direction of the Earth but from space. These gamma-ray bursts (GRBs) are bright flashes of hard X-rays and gamma rays that last fractions of a second to minutes during which they dominate the gamma-ray sky. Afterglows in lower energy wavelengths have shown that these bursts originate in some highly energetic explosions in galaxies throughout the observable universe.

Great progress in observations of cosmic gamma-ray sources and gamma-ray bursts was achieved with the launch of the Compton Gamma-Ray Observatory (CGRO), on April 4th, 1991. The Burst And Transient Source Experiment (BATSE) aboard CGRO recorded over 2700 gamma-ray bursts that showed a clearly isotropic distribution. The highest energy instrument on CGRO, the Energetic Gamma Ray Experiment Telescope (EGRET), surveyed the whole sky survey from 20 MeV to 30 GeV and detected 66 blazar AGN, 27 lower-confidence AGN, 7 pulsars, many gamma-ray bursts, and 170 unidentified sources (see Grenier¹ in these proceedings for more on gamma-ray sources). In addition, EGRET set a limit on the extragalactic diffuse gamma-ray background that has constrained many models of ultra-high energy phenomena. Compton re-entered the Earth's atmosphere on June 4, 2000.

The next generation of gamma-ray satellites will have improved sensitivity at higher energies. The Gamma-Ray Large Area Space Telescope (GLAST) will have a large area ($\sim 1 \text{ m}^2$) and will observe gamma-rays from 10 MeV to above $\sim 100 \text{ GeV}$ with excellent energy and angular resolutions. GLAST is expected to be launched in 2006. Before GLAST, the Astro-rivelatore Gamma a Immagini Leggero (AGILE) will cover the energy range from 30 MeV to 50 GeV while Astro-E2 should be launched in 2005 with a range from 0.4 keV to 700 keV. AGILE will have similar sensitivity to EGRET with a larger field of view and much better angular resolution.

Cosmic gamma-rays of energies above 100 GeV are hard to detect from space but can be studied on the ground. High-energy gamma-rays entering the atmosphere interact with the atmospheric particles and develop a cascade of particles known as an extensive air-shower. The electromagnetic cascade can be imaged by Atmospheric Cherenkov Telescopes (ACTs) such as the Whipple 10 m telescope in Arizona, the CAT telescope in the Pyrenees, the Cangaroo telescope in Australia, and the HEGRA telescope in the Canary Islands. ACTs detect the Cherenkov radiation generated by the electromagnetic particles in the shower from gamma-ray primaries with energies $\sim 100 \text{ GeV}$ or higher. For lower thresholds, the STACEE and CELESTE experiments utilize large arrays of heliostats at solar power plants to reach just below 100 GeV where overlap with satellite data can occur. For very high energies, the shower is detectable at the ground by Extensive Air Shower (EAS) arrays like CASA and CYGNUS. These arrays sample the showers sparsely and have $\sim 100 \text{ TeV}$ thresholds. To lower the threshold of EAS arrays to $\sim 1 \text{ TeV}$, the Milagro experiment has a large active area employing water as the detecting medium and an array of photomultiplier tubes. The Tibet array can also reach lower thresholds by being located at high altitude. To observe a wide range of gamma-ray energies, a variety of ground detectors need to complement the lower energy satellite observations.

Shower pattern imaging techniques have improved substantially in modern ACTs increasing significantly the ability to reject cosmic ray background showers. Imaging ACTs were the first telescopes to observe TeV sources (see Aharonian² for more on TeV sources). A new generation of ground-based instruments is under development such as HESS, MAGIC, CANGAROO-III, and VERITAS. These ACTs will reach the highest gamma-ray energies and will rival space-based instruments in sensitivity. Satellites provide a platform for all sky surveys and background studies

while ground telescopes are best suited to observe known point sources. These two approaches are highly complementary in energy range and type of source monitoring and together will help understand some of the highest energy environments in the universe.

Gamma-ray bursts show that the universe is quite transparent to 100 MeV gamma-rays, but the universe is likely to be opaque to very high energy gamma-rays. Pair production in extragalactic diffuse backgrounds are likely to degrade gamma-rays above ~ 10 TeV to $\sim 10^{19}$ eV producing a natural limit to the reach of very high energy gamma-ray observatories. It is clear that the CMB strongly suppresses high redshift PeV (10^{15} eV) gamma-rays, but the infrared and radio backgrounds, that influence the TeV and EeV gamma-ray propagation respectively, are less well determined. These less known backgrounds can be studied by contrasting high redshift to low redshift gamma-ray emission. For instance, EGRET observed many blazars up to ~ 30 GeV which have not been detected at \sim TeV energies. In principle, this could be due to a cutoff in the blazar spectrum, but since the few TeV blazars that have been detected are all at relatively low redshifts ($z \sim 0.03$ to 0.1), it is more likely that the lack of observations be due to an energy dependent gamma-ray horizon. The lack of TeV emission from powerful high redshift blazars gives an estimate of the infrared background. Finally, if high redshift gamma-ray sources have spectra that extend well beyond TeV energies, the effect of the diffuse backgrounds is to shift the high energy radiation to the EGRET energy range. Thus, EGRET limits on the extragalactic gamma-ray background are strongly constraining to models where ultra high energy gamma-rays are produced throughout the universe.

3. High-Energy Neutrinos

Neutrino astronomy was born in the mid 1960s with Davis' chlorine experiment detecting neutrinos from the ^8B beta decay in the core of the Sun. The discrepancies between the observed flux and the theoretical predictions led to the well known solar neutrino problem. A number of experiments pointed to the solution of this problem, most recently the Sudbury Neutrino observatory (SNO), which has the ability to detect all three flavors of neutrinos, was able to show that the total flux of the three neutrino flavors matches the theoretical predictions and the deficit of electron neutrinos from the Sun is due to their oscillations most likely into tau neutrinos. Another neutrino flavor oscillation was shown by SuperKamiokande (SuperK)

to explain the atmospheric neutrino problem: a deficit of muon neutrinos relative to electron neutrinos produced by cosmic ray interactions in the Earth's atmosphere. The solution to these problems is now known to be a great new discovery in fundamental physics: neutrinos have masses and oscillate between different flavors. (For the detection of cosmic neutrinos, Davis and Koshiba from SuperK were awarded the 2002 Nobel prize in physics.)

Another great discovery involving neutrinos was the confirmation of the standard scenario of core-collapse supernova by the detection of the first extrasolar neutrinos in 1987. Kamiokande detected 12 neutrinos from Supernova 1987a while IMB detected 9 events. Even though only a few neutrinos were measured from SN1987a, this detection led to great improvement in our understanding of core collapse supernovae. If a future supernova is observed in our own Galaxy, the richness of information carried by the neutrinos would be unparalleled.

Although the observed neutrinos from the SN87a are relatively low energy neutrinos, higher energy neutrinos are also expected to be emitted by supernovae explosions and by most high-energy astrophysical systems in the universe. The advantage of searching for very high-energy neutrinos from astrophysical sources is that, unlike gamma-rays, neutrinos have no horizon due to interactions; they can traverse the observable universe unimpeded. In particular, gamma-ray bursts and AGN should produce observable fluxes of neutrinos from nearby to very high redshifts. The estimated fluxes indicate a minimum scale of km^3 or 1 billion tons of detecting material. There are two approaches being taken for neutrinos around 1 PeV (10^{15} eV) in energy: km^3 water detectors (ANTARES, NEMO, NESTOR) and km^3 ice detectors (IceCube). These detectors use the Earth as a filter of background particles and observe upward going muons generated by neutrinos as they traverse the Earth. High-energy muons emit Cherenkov radiation in water or ice, which is then observed by an array of photomultiplier tubes. Prototype detectors in ice (AMANDA) and water (Lake Baikal) have already observed many events consistent with atmospheric neutrinos. (See Halzen³ in these proceedings for more on high energy neutrinos.)

Another source of even higher energy neutrinos are extragalactic ultra-high energy cosmic rays. As ultra-high energy cosmic rays traverse the universe, they interact with the cosmic microwave background radiation and generate pions. Pions decay generating neutrinos with energies starting $\sim 10^{17}$ eV and higher. The new generation of ultra-high energy cosmic ray detectors, such as the Auger Project and the EUSO satellite experiment,

will be able to study ultra-high energy neutrinos through horizontal and Earth skimming showers. In addition, new techniques for ultra-high energy neutrino observations using radio signals in ice (e.g., RICE and ANITA) or salt mines (e.g., SALSA) are now under development. Over the next decade, new high-energy neutrino detectors will open a new window of high-energy astrophysics. (See Klages⁴, Blasi⁵, and Berezhinsky⁶ in these proceedings for more on high energy neutrinos from ultra-high energy cosmic ray sources.)

4. Cosmic-Rays the highest energy messengers

Observations of extensive air showers have detected the highest energy particles ever observed: cosmic rays with macroscopic energies around tens of Joules or 10^{20} eV. Cosmic rays with energies up to 10^{14} eV can be detected directly with balloon and space experiments while above this energy the flux is too low for space-based detectors and cosmic rays are studied through their air shower development. The cosmic ray spectrum from $\sim 10^9$ eV to 10^{14} eV is well described by a power law of spectral index ~ 2.7 . At higher energies the spectrum steepens into a spectral index ~ 3 and the transition region is called the knee. At the highest energies the spectrum seems to change again but the exact features are still unclear.

Composition studies at low energies show a diffusive history of cosmic ray nuclei through the Galaxy. At lower energies, cosmic rays are dominated by light nuclei (protons and helium) while at the knee the composition seems to become heavier (see Haungs⁷ in these proceedings for recent results on composition from KASCADE). Galactic cosmic ray propagation is diffusive in magnetic fields with a rigidity dependent escape probability. The knee is thought to represent the transition from confined trajectories to trajectories that escape the Galaxy and thus the change in spectral index. The origin and propagation history of cosmic rays are still unclear with the leading theory being shock acceleration in supernova remnant as proposed by Enrico Fermi in 1949. A clear confirmation of this picture is still lacking but can be achieved by gamma-ray detectors sensitive to gamma-rays from pion decay around the acceleration sites.

Fermi acceleration in supernova remnants may be responsible for accelerating cosmic rays below the knee, but more powerful sources seem to be required for the higher energy events. In addition, as the energy of the primary cosmic ray increases, the effect of the Galactic magnetic field in the particle trajectory decreases. As cosmic rays reach 10^{19} eV and above, trajectories should start to point back to cosmic ray sources, i.e., cosmic ray

astronomy should become possible. Instead, observations show an isotropic distribution of arrival directions up to the highest energies observed thus far. With no indication of the Galactic plane or other nearby structures, the isotropy argues for an extra-galactic origin for the highest energy sources of cosmic ray protons or much stronger than expected magnetic fields in the halo or our Galaxy.

Ultra-high energy cosmic rays are detected mainly via two techniques: ground arrays of scintillators or water Cherenkov tanks and fluorescence detectors. Ground arrays sample the extensive air shower as it reaches the ground while fluorescence observatories detect the fluorescence of nitrogen molecules in the atmosphere as the shower develops above the ground. This second technique works best in clear moonless nights and in a stereo mode, when two detectors separated by many kilometers can observe the same shower from two different angles determining the shower axis. The leading experiments using each of these techniques, the Akeno Giant Air Shower Array (AGASA) and the High Resolution Fly's Eye (HiRes), are now probing the predicted Greisen-Zatsepin-Kuzmin (GZK) cutoff⁹ (see Nagano⁸ in these proceedings for more on these experiments). A cutoff should be present if the ultra-high energy particles are extragalactic protons, because protons above 7×10^{19} eV produce pions on their way to Earth through interactions with the cosmic microwave background. The energy lost due to the photo-pion production generates a feature in the spectrum, the GZK feature. Thus, in addition to the extraordinary energy requirements for astrophysical sources to accelerate protons to 10^{20} eV, the photopion threshold reaction was expected to cause a natural high-energy limit to the cosmic ray spectrum known as the GZK cutoff.

The first definitive detection of an ultra high energy cosmic ray was recorded by the Fly's Eye experiment in 1991: a 3.2×10^{20} eV event. This event triggered considerable interest on the origin and nature of these particles that indicate the lack of the predicted GZK cutoff. Following this event, AGASA has reported 11 events above the GZK cutoff which argues for the absence of the cutoff. Together with the isotropy of arrival directions these events pose a challenge to models for the origin of UHECRs. However, the HiRes collaboration reported a suppression of the UHECR flux at high energies based on an analysis of their monocular data. With the presently available low statistics of observed extremely high energy events, the GZK feature associated with the photopion production is still dominated by fluctuations¹⁰ and a new generation of experiments can resolve this issue (see Blasi⁵ and Berezhinsky⁶ in these proceedings).

The hybrid detector of the Auger Project (see Klages⁴) should settle the present disparity between AGASA and HiRes. The Pierre Auger Project is now under construction in Argentina and a second site should be constructed in the northern hemisphere. The Argentinean site will have a 3000 km² array of water Cherenkov tanks together with four fluorescence detectors overlooking the array. The use of both surface array and fluorescence technique will help resolve the apparent inconsistency in the spectrum between AGASA and HiRes by measuring many more events and cross calibrating the two techniques. It can also determine the composition by measuring the depth of shower maximum and the ground footprint of the same shower. The solution to the ultra-high energy cosmic rays mystery will come with the next generation of experiments, which are under construction such as Auger or in the planning stages such as the space-based EUSO and OWL observatories. These space fluorescence detectors can monitor extensive air showers from above the atmosphere on very large volumes.

5. The Future of High energy astrophysics

The future of high-energy astrophysics looks extremely promising. The combination of space missions that span most wavelengths of detectable photons and ground-based detectors that reach thousands of km² will bring the many puzzles involved in the production and propagation of the highest energy gamma-rays, cosmic-rays, and neutrinos into a new era.

After the success of CGRO, the next large gamma-ray space-based observatory now under construction is GLAST. GLAST's new technology includes Si trackers, CsI calorimeters, and anti-coincidence veto and is expected to be launch in 2006. At the higher energies, ground based gamma-ray telescope arrays are now being built such as the Tibet Array, HESS (started in 2002), VERITAS and Cangaroo III (now under construction). At intermediate energies, MAGIC follows STACEE and CELESTE in the overlap with the satellite energy range.

High-energy neutrino observatories will be able to reach the necessary km³ scale in ice (IceCube) and in water following ANTARES, NESTOR and NEMOS. These observatories will open a new window into the workings of the high-energy systems in the universe. At the highest energies, neutrinos should be detectable by the next generation ultra-high energy cosmic ray detectors such as Auger, EUSO, and OWL and the next generation radio detectors such as SuperRICE and ANITA.

In ultra-high energy cosmic rays, the future promises to unveil the origin and nature of these mysterious particles. The present state of observations is particularly puzzling and the necessary experiments to resolve these puzzles will be operating in the very near future. The Pierre Auger project should be fully operational in the Southern site in 2005 and the Northern site shortly after. Once Auger reaches past 10^{20} eV, further exploration into higher energies can be achieved by space observatories, such as the EUSO and OWL missions.

The understanding of the high-energy universe is just beginning with the promise of many new discoveries to come. By reading the messages brought by all the different messengers we will gain important insight into the workings of Nature and the origin and evolution of the Universe.




Figure 1. Spectra of photon backgrounds (based on Ressel and Turner¹¹ modified by Halzen³, cosmic rays, and expected neutrino flux from ultra high energy cosmic ray interactions.

Acknowledgments

We thank the organizers of the XXI Texas Symposium. This work was supported in part by the NSF through grant AST-0071235 and DOE grant DE-FG0291-ER40606 at the University of Chicago.

References

1. I. Grenier, *Proceedings of the XXI Texas Symposium* (2003).
2. F. Aharonian, *Proceedings of the XXI Texas Symposium* (2003).
3. F. Halzen, *Proceedings of the XXI Texas Symposium* (2003).
4. H. Klages, *Proceedings of the XXI Texas Symposium* (2003).
5. P. Blasi, *Proceedings of the XXI Texas Symposium* (2003).
6. V. Berezhinsky, *Proceedings of the XXI Texas Symposium* (2003).
7. A. Haungs, *Proceedings of the XXI Texas Symposium* (2003).
8. M. Nagano, *Proceedings of the XXI Texas Symposium* (2003).
9. K. Greisen, *Phys. Rev. Lett.* 16, 748 (1966); G.T. Zatsepin and V. A. Kuzmin, *Sov. Phys. JETP Lett.* 4, 781966 (1966).
10. D. DeMarco, P. Blasi, A. V. Olinto, *Astroparticle Physics* in press (2003).
11. M. T. Ressel and M. S. Turner, *Bulletin of the American Astronomical Society*, Vol. 22, p.753 (1990).