# New Hipparcos-based Parallaxes for 424 Dim Stars

Andrew Gould and Julio Chanamé

Department of Astronomy, The Ohio State University, 140 W. 18th Ave., Columbus, OH 43210

gould, jchaname@astronomy.ohio-state.edu

### **ABSTRACT**

We present a catalog of 424 common proper motion companions to *Hipparcos* stars with good (>  $3\sigma$ ) parallaxes, thereby effectively providing new parallaxes for these companions. Compared to stars in the *Hipparcos* catalog, these stars are substantially dimmer. The catalog includes 20 WDs and an additional 29 stars with  $M_V > 14$ , the great majority of the latter being M dwarfs.

Subject headings: astrometry – catalogs – stars: fundamental parameters – stars: late-type – white dwarfs

## 1. Introduction

Hipparcos (ESA 1997) revolutionized astrometry in three ways. First, its mass-production mode increased the sheer number of stars with mas parallaxes by several orders of magnitude to  $10^5$ . Second, it directly obtained absolute parallaxes rather than having to convert from relative to absolute parallax as is necessary from the ground. Third, it measured homogeneous parallaxes over the whole sky.

However, the Hipparcos catalog is notably deficient in dim stars, containing only 3 stars with  $M_V \geq 14$  and only 14 with  $M_V > 13$ . While the project attempted to observe all known dim stars down to its operational limit  $V \sim 12$ , at  $M_V = 14$  this limit corresponds to a distance of 4 pc. Hence, dim-star parallaxes generally continue to require the painstaking one-at-a-time methodology that had characterized this subject for the two centuries prior to Hipparcos. The classic work of Monet et al. (1992) remains crucial for tracing out the bottom of the main sequence (MS) and the cool subdwarfs (SDs). Bergeron, Leggett & Ruiz (2001) assembled a catalog of about 150 white dwarfs (WDs) with parallaxes, the vast majority from the ground. Jao et al. (2003) have targeted almost 200 nearby stars for parallaxes. Gizis (1997) assembled almost 100 ground-based parallaxes for SDs. Brown-dwarf parallaxes are at present obtained exclusively from the ground (Dahn et al. 2002).

While nothing can replace this crucial ground-based work, at least until a new generation of astrometry satellites is launched, it is in fact possible to obtain many additional parallaxes of dim stars using *Hipparcos* data. Common proper-motion (CPM) companions of *Hipparcos* stars should have the same parallaxes as their primaries up to a fractional error equal to their separation in radians, which is generally small compared to the measurement error and, in any event, always less than 1%.

The idea of obtaining parallaxes in this way is not new. In describing the technique he had used to discover what was then (and remained for four decades) the dimmest known star, vB 10, van Biesbroeck (1944) recounted: "In 1940 the author started at the prime focus of the 82-inch reflector of the McDonald Observatory a systematic search for faint companions to known PM stars in order to extend our knowledge toward the lowest luminosity stars." The Yale Parallax Catalog (van Altena, Lee, & Hoffleit 1995) lists the proper motions of CPM pairs together with the parallax of the primary, clearly intending the same distance estimate to be applied to the fainter star as well. Oppenheimer et al. (2001) applied this technique to CPM companions of Hipparcos stars to establish distances to some stars in their 8 pc sample. However, no one has attempted to systematically search for CPM companions of Hipparcos stars. Indeed the Villanova White Dwarf web site<sup>1</sup> continues to list Yale parallaxes for CPM companions of stars that now have Hipparcos parallaxes.

The primary problem in applying this technique is establishing a physical association between the two components of the binary. This is not difficult for very nearby stars, which usually have very large proper motions: the chance that two unrelated stars lying within a few arcmin of each other would have roughly similar proper motions of order  $\mu \sim 1000\,\mathrm{mas\,yr^{-1}}$  is vanishingly small. Hence, precise proper motions are not generally required to establish a physical connection between these stars. However, the projected density of stars as a function of proper motion grows extremely rapidly toward lower proper motions, so much higher precision is required to effectively reject spurious unrelated pairs. Until recently, such high-precision proper-motion catalogs were not available.

Several developments over the past 18 months have radically altered this situation thereby permitting a much more aggressive search for CPM companions of Hipparcos stars. First, Gould & Salim (2003) and Salim & Gould (2003) have published the revised New Luyten Two-Tenths (rNLTT) catalog, which identifies virtually all *Hipparcos* counterparts of NLTT (Luyten 1979, 1980; Luyten & Hughes 1980) stars, and which gives new more accurate ( $\sigma_{\mu} \sim 5.5 \,\mathrm{mas}\,\mathrm{yr}^{-1}$ ) proper motions for the vast majority of NLTT stars in the 44% of the sky covered by the intersection of the first Palomar Observatory Sky Survey (POSS I)

<sup>&</sup>lt;sup>1</sup>http://www.astronomy.villanova.edu/WDCatalog/index.html

and the Second Incremental Release of 2MASS (Skrutskie et al. 1997). Very importantly in the present context, each rNLTT entry indicates whether the given star has a NLTT CPM companion according to the NLTT Notes compiled by Luyten and, if so, whether that companion is identified in rNLTT itself. Moreover, for entries with CPM companions, it gives the positional offset of the companion, both as given by the NLTT Notes and as measured by rNLTT itself.

Second, 2MASS has now been released for the whole sky. Hence, by searching 2MASS at the CPM offsets as indicated by rNLTT, one can usually find the CPM companions of *Hipparcos* stars, at least those that satisfy the NLTT proper motion threshold,  $\mu \geq 180\,\mathrm{mas}\,\mathrm{yr}^{-1}$ .

Third, USNO-B1.0 (Monet et al. 2003), with its roughly 10<sup>9</sup> proper motion measurements, each having typical errors of a few mas yr<sup>-1</sup>, has now been released. Because more than 99% of its high proper-motion entries are spurious (Gould 2003), USNO-B cannot be used to search blindly for high proper motion stars. However, if one knows the approximate position and proper motion of a star (as one does for CPM companions of *Hipparcos* stars) then the false background stars become much more manageable.

Fourth, Chanamé & Gould (2003) have compiled a catalog of NLTT binaries for the 44% of the sky covered by the intersection of POSS I and the Second 2MASS Incremental Release. For these areas, they have determined which NLTT pairs that are specified in the NLTT Notes as CPM binaries are in fact physical pairs and also which pairs of NLTT stars are CPM binaries despite the fact that they are not so designated in the Notes. Hence, for these areas, virtually all the work required to assemble a catalog of *Hipparcos* companions is already done.

In this paper, we make use of these various new data sources to compile a catalog of CPM companions to *Hipparcos* stars. The catalog is restricted to *Hipparcos* stars that have accurate parallaxes and are in the NLTT. However, we also discuss how it might be extended to companions of other *Hipparcos* stars in future work.

### 2. Catalog Construction

### 2.1. Philosophy

Our aim is to construct a catalog with as many genuine CPM companions as possible, while minimizing the number of false entries. The quality of our underlying sources varies dramatically over the sky, and so we do not aim to construct a catalog that is "complete"

or "homogeneous" in any sense. When we have excellent proper-motion information, we can be very confident that we have identified physical pairs. As we will describe, the available data are often far from adequate to achieve this high level of confidence. Hence, for many stars we must make a judgment call, making use of photometric as well as astrometric data. As this photometry is generally photographic, it is sometimes not as reliable as one would like. Hence, any individual CPM pair in the catalog may be spurious. We expect that the primary use of the catalog will be as a source of candidate dim stars with parallaxes, and that few will be found to be spurious when checked by obtaining radial velocities and/or CCD photometry.

We exclude most pairs that are so close that they are not resolved in 2MASS. These are so close that the proximity of their components is usually a more reliable guide to their physical association than is the similarity of their proper motions. Hence, our proper-motion based approach brings nothing new to the table. Moreover, the great majority of these companions are luminous stars (e.g.,  $M_V \sim 5$ ) that are already well represented in the *Hipparcos* catalog and hence are not of much interest in the present context. We make an exception to this rule only when the companion is very dim and hence is a rare object from a parallax perspective.

We count stars as being "Hipparcos" only if their parallax measurements satisfy  $\pi/\sigma_{\pi} > 3$ . Otherwise, they do not have  $3\sigma$  parallax detections and therefore cannot provide significant parallax information about their companion. On the other hand, if such stars are companions to other Hipparcos stars with good parallaxes, they can gain parallax information. In our search, we therefore treat Hipparcos stars with less precise parallaxes as "non-Hipparcos".

For completeness, we include CPM pairs of *Hipparcos* stars that both have good parallaxes. However, we consider these to be of less (or at any rate, different) intrinsic interest and report them in a separate table. Of the 508 CPM pairs presented here, 84 are in this category.

#### 2.2. Search Breakdown

We divide our search for *Hipparcos* CPM companions into four subcategories. First, pairs from the Chanamé & Gould (2003) catalog, which covers 44% of the sky (248). Second, other pairs from rNLTT for which the primary (i.e., *Hipparcos* star) has a 2MASS identification in rNLTT (16). Third, other pairs from rNLTT for which the primary lacks a 2MASS identification (88). Fourth, pairs designated as CPM binaries by Luyten, for which one star

is in *Hipparcos* (and is in rNLTT) but for which the companion is not in rNLTT (156). As we will describe, these subcategories require progressively more work.

## 2.3. Catalog of Chanamé & Gould (2003)

Chanamé & Gould (2003, hereafter CG) have already done most of the work to determine which Luyten CPM binaries are genuine and have also found additional genuine CPM binaries among NLTT stars that were not recognized as such by Luyten. However, since the objectives of CG and the present work are slightly different, some additional work is required. First, CG were interested primarily in obtaining clean samples of disk and halo stars and so excluded pairs that could not be reliably classified as one or the other. For example, those that straddled the disk/halo boundary on a reduced proper motion (RPM) diagram were excluded. They also excluded all pairs for which one component was a WD. Note that MS pairs and SD pairs can be vetted by checking to see whether they lie parallel to these respective sequences on an RPM diagram (see their figs. 2 and 3), but that MS/WD pairs cannot be subjected to this test. Thus, CG were not in a position to supply as severe vetting of pairs with WD components as they were for other pairs and did not attempt to do so.

Hence, our approach is to accept all pairs (having an *Hipparcos* component) regarded as genuine by CG and then to review all candidates that they did not accept to see if they should be regarded as genuine pairs. This review makes use not only of the information used by CG, but also of the *Hipparcos* parallax. For example, if a MS/WD pair is genuine, then when the WD is placed at its companion's distance, it should lie close to the WD sequence.

Finally, CG excluded all triples, including NLTT pairs for which one NLTT component was resolved by the Tycho Double Star Catalog (TDSC, Fabricius et al. 2002). These do not appear in their catalog, even as candidates. We make no special effort to recover these as part of this subsearch: they are recovered automatically as part of the other subsearches.

The proper-motion selection criterion of CG was to accept all pairs with vector propermotion differences satisfying  $\Delta \mu \equiv |\Delta \vec{\mu}| < 20 \,\mathrm{mas}\,\mathrm{yr}^{-1}$ , to further accept those with  $\Delta \mu < (112 - 51 \log(\Delta \theta / '')) \,\mathrm{mas}\,\mathrm{yr}^{-1}$ , and to accept all pairs with separations  $\Delta \theta < 10''$ . In constructing these relatively severe criteria, CG took advantage of the fact that rNLTT contains independent proper-motion measurements for the great majority of its stars and that the relative proper-motion errors for these are quite small (see Salim & Gould 2003).

In the present context, however, it is important to note that for some of CG's pairs, one component lacked an independent proper-motion measurement. For these,  $\Delta\mu$  was

determined by taking the difference in the vector separations of the components as recorded in the NLTT Notes and as measured by 2MASS and then dividing this difference by the approximately 45-year difference in epochs. The errors from this method are about twice as large as from direct measurements and, moreover, have significant outliers, which probably originate from transcription errors in the NLTT Notes. This is a relatively minor problem for this CG subsample: of the 248 binaries that we eventually accept from this subcategory, only 39 lacked independent proper-motion measurements. All of these satisfied the original  $\Delta\mu$  criterion of CG. However, in other subcategories, notably the fourth, a much larger fraction of the binaries lack independent proper-motion measurements. We describe our procedure for dealing with these below.

## 2.4. Non-CG pairs in rNLTT with 2MASS data

A total of 16 such pairs are selected from two ultimate sources, Second Incremental 2MASS areas south of POSS I (5), and pairs rejected by CG because they were part of triples (11). The first group is quite small because this area covers only 3% of sky. All information is already available to classify these pairs.

## 2.5. Non-CG pairs in rNLTT without 2MASS data

By construction, these pairs must have both components in one of three position-and-proper-motion catalogs of bright stars, *Hipparcos*, Tycho-2 (Høg et al. 2000), or Starnet (Röser 1996), which were the only avenues into rNLTT for stars in non-2MASS areas. These stars almost all have very accurate proper motions, and it is a simple matter at this point to search for them in the 2MASS all-sky release. Hence, their classification as genuine or not is straightforward. There are a total of 88 physical pairs, of which 45 have both components in *Hipparcos*.

Very few of these are of interest as dim stars with new parallaxes simply because they are so bright that if they had been of special interest, they would have been observable directly with *Hipparcos*.

### 2.6. Non-rNLTT CPM Companions of Hipparcos Stars

This subcategory presents the greatest difficulties. Given that, even in cases for which the companion is not recovered by rNLTT, the *Hipparcos* stars with NLTT companions are

already marked in rNLTT, it is straightforward to find the vast majority of these in 2MASS simply by searching at the vector separation given in the NLTT Notes (and recorded in rNLTT). We then use the 2MASS position to search for the companion in USNO-B. When the companion is in USNO-B, we obtain a second estimate of the proper motion in addition to the one based on the difference between vector separations in 2MASS and the NLTT Notes (see § 2.3). Whenever the smaller of these two estimates violates the CG proper-motion criterion (see § 2.3), we flag the binary. We inspect the RPM diagram of each binary and if the binary is not well aligned with either the MS or SD tracks, we also flag it. In particular, all WDs are thus flagged. Finally, we flag all binaries for which one component is lacking 2MASS data. Of the 184 initial candidates, a total of 82 are flagged, some several times. We then review each of the flagged cases individually, inspecting the binary on both the RPM diagram, and the color-magnitude diagram (CMD), the latter under the provisional assumption that the *Hipparcos* parallax applies to both components. Whenever the binary is under question primarily because its proper-motion difference is too high, we attempt to locate the companion in USNO-A (Monet 1996, 1998). Comparison of this position with the 2MASS position gives another estimate of the proper motion, which is generally more reliable than either USNO-B or the 2MASS/NLTT-Notes difference method (Salim & Gould 2003; Gould 2003). Unfortunately, this method usually fails for stars south of POSS I because they are usually not in USNO-A. See Salim & Gould (2003). In the end, we take all the available evidence and make our best judgment as to whether the binary is physical. Of the 82 flagged binaries, 28 are rejected and another 10 are regarded as plausible but not fully convincing cases. They (along with one other pair from the CG subsample) are flagged as "somewhat uncertain" in the catalog.

## 3. Catalog Description

The catalog is divided into two tables. Table 1 lists the non-Hipparcos CPM companions of Hipparcos stars, i.e., the stars with new Hipparcos-based parallaxes found by this work. Columns 1 through 9 describe the companion: column 1 gives the NLTT number, columns 2 and 3 give the R.A. and Dec (2000 epoch and equinox), columns 4 and 5 give the east and north proper-motion components in mas  $yr^{-1}$ . Columns 6 and 7 give the V magnitude and V-J color, and column 8 gives the absolute magnitude. Column 9 is a 3 digit source code, which is described below. Columns 10–18 are similar to columns 1–9, but for the Hipparcos primary. Columns 19 and 20 give the separation (in arcsec) and position angle (north through east in deg) of the companion with respect to the Hipparcos star. Columns 21–23 gives the Hipparcos number, parallax and parallax error (in mas). Column 24 is the adopted proper-motion difference and column 25 is a flag, "1" if the binary is "somewhat

uncertain" and "0" otherwise.

The three-digit source code is an expanded version of the source code used in rNLTT. There, the three digits refer to the sources of the position, proper motion, and V photometry. 1 = Hipparcos, 2 = Tycho-2, 3 = TDSC, 4 = Starnet, 5 = USNO/2MASS, 6 = NLTT, 7 = USNO-A (for position) or common proper motion companion (for proper motion). More specifically, "555" means 2MASS based position, USNO-A based V photometry, and USNO/2MASS based proper motion. In addition, we add 8 = USNO-B (for position and proper motion). A "9" in the position column means that there is no actual detection in any catalog and the position is inferred from the separation vector given in the NLTT Notes. USNO-A photometry and NLTT photometry are transformed to V photometry using the prescriptions of Salim & Gould (2003).

Table 2 lists the CPM pairs of Hipparcos stars, i.e., CPM pairs composed of two Hipparcos stars, each with a parallax better than  $3\sigma$ . Column 1–6 give information on the brighter component. Column 1 and 2 give the Hipparcos and rNLTT numbers. Column 3 and 4 give the parallax and parallax error (in mas). Columns 5 and 6 give the V magnitude and V-J color. Columns 7–12 give the same information for the fainter companion. Columns 13 and 14 give the separation and position angle of the fainter component.

These two tables are available at http://www.astronomy.ohio-state.edu/ $\sim$ gould/rNLTT\_binaries/new\_parallaxes.dat and http://www.astronomy.ohio-state.edu/ $\sim$ gould/rNLTT\_binaries/hip\_doubles.dat respectively. The Fortran format statements for the table records are respectively, (2(i5,2f10.5,2i6,3f7.2,i4,1x),2f6.1,i7,2f7.2,i5,i2) and (2(i6,i6,2f7.2,2f7.2,1x),2f6.1).

### 4. Dim Stars

Figure 1 is a CMD of the CPM companions of *Hipparcos* stars from Table 1. The error bars reflect the parallax errors only. That is, they show the limits of precision for the absolute magnitudes provided that one obtained good CCD photometry for the stars. The vast majority of the optical photometry is at present photographic. Stars without J photometry are shown as V - J = -1. These could be either M dwarfs or WDs. Note the high concentration of dim stars in the sample. They are, for example, much dimmer than their primaries, which are shown in Figure 2. Figure 1 also shows the WDs from the *Hipparcos* catalog. While *Hipparcos* contains roughly the same number of WDs as our catalog, ours tend to be dimmer.

A striking feature of the CMD is its breadth, roughly 3 mag. This is much broader than

the CMD of Monet et al. (1992), which is likely to be due in significant part to errors in the photographic photometry. However, the underlying proper-motion limited sample is likely to have a broader range of metallicities than typical samples, and this may also contribute to the breadth of the CMD. It would be straightforward to obtain CCD photometry for the entire sample and more than half could be done in a single few-night run on a 1m telescope. Perfect photometric conditions would not be required to achieve dramatic improvements over the present optical photometry.

In Figure 3, we compare histograms of the stars in Table 1 and of the 400 dimmest stars in the *Hipparcos* catalog (restricted to entries with  $\pi/\sigma_{\pi} > 3$ ). While both catalogs contain many moderately dim stars, our catalog contains substantially more extremely dim stars than *Hipparcos*.

What was previously known about these dim CPM companions of *Hipparcos* stars? We address this question in Tables 3 and 4. Table 3 lists the 20 CPM WD companions shown in Figure 1 and indicates whether each was previously identified as being a WD and, if so, whether it had a measured parallax. The classifications are taken from the Villanova White Dwarf web site except for NLTT 49859, which comes from SIMBAD<sup>2</sup>. The previous parallaxes come most directly from Villanova, but most come ultimately from van Altena, Lee, & Hoffleit (1995), with the one exception (NLTT 15768) coming from Bergeron, Leggett & Ruiz (2001). The identifiers come mostly from SIMBAD, except for NLTT 26462 and 29967, which come from Villanova. Column 1 gives the NLTT number. Columns 2 and 3 give the RA and Dec (2000 epoch and equinox). Columns 4-6 give the V, V-J, and  $M_V$ magnitude, color, and absolute magnitude. Columns 7 and 8 give the separation and position angle relative to the *Hipparcos* star, whose *Hipparcos* number, parallax, and parallax error are given in columns 9-11. Columns 12 and 13 give the previously tabulated parallax and error for the WD if any were found. Column 14 gives one of the identifiers (if any were found) and Column 15 gives the spectral type. Of the 20 WDs, five (NLTT 12412, 38926, 42785, 47097, 49859) are not listed at Villanova as WDs. Two of these are listed but not classified by SIMBAD, while NLTT 12412 and 38926 are not listed and NLTT 49859 is classified as an M dwarf. Of the remaining 15 WDs, seven have previous parallaxes of which only two are of comparable quality to the *Hipparcos* parallaxes assigned in the present work.

With one exception, the stars in Table 3 are classified as WDs based either on their position on the CMD or because they were so classified in the Villanova White Dwarf web site. The exception is NLTT 38926. We classify this as a WD in spite of the fact that we have no J data and the only available reference in the literature (Luyten) classified it as an

<sup>&</sup>lt;sup>2</sup>http://simbad.u-strasbg.fr/Simbad

"m" star. We do so because of its dim absolute magnitude,  $M_V = 13.5$ , and the fact that it is significantly bluer than 4 random neighboring field stars in B - R and R - I, based on Digitized Sky Survey<sup>3</sup> images. However, we record Luyten's contrary view in the spectral classification column with the denotation "L:m".

Table 4 lists the 29 stars with  $M_V \geq 14$  from Table 1, except that the WDs listed in Table 3 are excluded. The columns in Table 4 are the same as those in Table 3. Five of these 29 stars have previously recorded parallaxes, of which three (NLTT 923, 26247, 47621) were determined by Oppenheimer et al. (2001) based on their being CPM companions to Hipparcos stars, and the other two (NLTT 18218, 42494) were determined by van Altena, Lee, & Hoffleit (1995) based on their being CPM companions to brighter stars with parallaxes. Only 12 of the 29 have spectral types listed in SIMBAD, and many are not listed at all (and hence have no identifier in Table 4). Three of the stars (NLTT 28864, 40719, 41096) have no J data, and moreover have no classification in the literature. For these, we list the Luyten classification based on photographic colors (e.g., "L:f", meaning "F star"). If these classifications are correct, then two of these stars (NLTT 28864, 41096) are WDs. We confirm the existence of NLTT 41096 based on Digitized Sky Survey images, but the other two stars are too close to their companions to be seen. The identifier and spectral class of NLTT 8870 are assigned question marks because the catalogued star with this identifier has roughly the right characteristics but the wrong position in SIMBAD.

## 5. Hipparcos Doubles

The pairs in Table 2 are of modest interest. For a few, one can obtain a significantly better parallax for one star by using the parallax of the other. However, one important application of this table is to test the accuracy of the error bars listed in the *Hipparcos* catalog. Figure 4 is a histogram of the differences (brighter minus fainter) in the parallaxes of the binary components divided by the root sum square of their reported errors. The curve shows the distribution expected for Gaussian errors.

Figure 4 has three notable features. First, the curve is in overall rough agreement with the histogram, indicating that, on average, the errors are properly estimated. Second, there is a spike at zero difference. Third, there are two severe outliers at  $+3.6 \sigma$  and  $+5.3 \sigma$ , and a third mild outlier at  $-2.8 \sigma$  (respectively Hip 82817/09, 90355/65, 64444/3).

The spike is principally due to four binaries with exactly zero difference in parallax

<sup>&</sup>lt;sup>3</sup>http://archive.eso.org/dss/dss

(Hip 13714/6, 17749/50, 86961/3, 97099/6). Indeed the two components of these four pairs have exactly the same reported errors as well. While the *Hipparcos* errors are known to be correlated at close separations (e.g., Narayanan & Gould 1999), the degree of correlation observed in these four pairs would appear a bit extreme. It is not clear what singles out these four. While they all have separations  $\Delta\theta \sim 20''$ , and so are closer than typical binaries in Table 2, there are 14 other binaries in this table with  $\Delta\theta < 20''$ , and these appear to have normally distributed errors.

Each of the three outliers has components with almost identical proper motions and so can be taken to be a genuine physical pair with virtual certainty. None of the six components of these pairs is resolved in TDSC as two distinct stars.

#### 6. Future Possibilities

The technique developed here could be applied to other data sets to obtain parallaxes for additional dim stars. Here we briefly outline the potential possibilities and pitfalls of this approach.

The key characteristic of the *Hipparcos*-CPM binaries reported here that made them relatively easy to find is that someone (mostly Luyten, but in a few cases CG) had already tabulated them as probable binaries and had recorded separations and position angles. It was then relatively straightforward to find the CPM companions in various catalogs and to use the data so obtained to judge the genuineness of the candidates.

Hence, one would be well advised to apply the same approach to another catalog of wide binaries, the most obvious choice being the Luyten Double Star (LDS) Catalog (Luyten 1940-87). This is comprised primarily of candidate wide binaries found by Luyten in his search for high proper-motion stars ( $\mu \geq 180 \,\mathrm{mas}\,\mathrm{yr}^{-1}$ ), but including even those that did not meet this threshold. The only additional required step (relative to the work reported in this paper) would be to match LDS primaries to *Hipparcos* stars. This was unnecessary for the NLTT binaries because their *Hipparcos* counterparts had already been identified by rNLTT. The drawback of LDS is that it will most likely yield a lower fraction of dim stars: for dim stars to be recovered in NLTT or LDS, they must be relatively close to satisfy the  $V \sim 19$  magnitude limit of these catalogs. They then typically have high proper motions, and so would tend to be in NLTT as well as LDS.

Another option would be to search in USNO-B for CPM companions of *Hipparcos* stars. While USNO-B cannot be used for a blind search because most of its high proper-motion entries are spurious (Gould 2003), if the search is restricted to the narrow range of proper

motions compatible with the known proper motion of an *Hipparcos* star, the number of spurious entries can be drastically reduced. Nevertheless, CG found a similar search for NLTT CPM companions in USNO-B to be quite laborious because of the still large number of spurious entries. Moreover, the return of real binaries (not already in NLTT) was relatively low. See their figure 13. On the plus side, however, most of those found were fainter than the NLTT companions, and so (at fixed distance) would also be dimmer. The false detections from such a USNO-B search could be drastically reduced by demanding detection of the candidates in 2MASS. Unfortunately, this would have the effect of eliminating most WDs, although it would preserve most red dwarfs.

We thank Samir Salim for making several very valuable comments on the manuscript. We thank D. Monet and the USNO-B team for providing us with a copy of the USNO-B1.0 catalog. This publication has made use of catalogs from the Astronomical Data Center at NASA Goddard Space Flight Center, VizieR and SIMBAD databases operated at CDS, Strasbourg, France, and data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the IPAC/Caltech, funded by NASA and the NSF. The ESO Online Digitized Sky Survey is based on photographic data obtained using The UK Schmidt This work was supported by grant AST 02-01266 from the NSF and by JPL contract 1226901.

### REFERENCES

- Bergeron, P., Leggett, S. K., & Ruiz, M.T. 2001, ApJS, 133, 413.
- Chanamé, J. & Gould, A. 2003, ApJ, submitted (astroph/0307284)
- Dahn, C.C., et al. 2002, AJ, 124, 1170
- European Space Agency (ESA). 1997, The Hipparcos and Tycho Catalogues (SP-1200; Noordwijk: ESA)
- Fabricius, C., Høg, E., Makarov, V. V., Mason, B. D., Wycoff, G. L., & Urban, S. E. 2002, A&A, 384, 180.
- Gizis, J.E. 1997, AJ, 113, 806
- Gould, A. 2003, AJ, 126, 472
- Gould, A. & Salim, S. 2003, ApJ, 582, 1001
- Høg, E. et al. 2000, A&A, 355, L27.
- Jao, W.C, Henry, T.J., Subasavage, J.P., Bean, J.L., Costa, E., Ianna, P.A., & Méndez, R.A. 2003, ApJ125, 332
- Luyten, W.J. 1979, 1980, New Luyten Catalogue of Stars with Proper Motions Larger than Two Tenths of an Arcsecond (Minneapolis: University of Minnesota Press)
- Luyten, W.J. 1940-87, The LDS Catalogue: Double Stars with Common Proper Motion (Minneapolis: University of Minnesota Press)
- Luyten, W.J. & Hughes, H.S. 1980, Proper Motion Survey with the Forty-Eight Inch Schmidt Telescope. LV. First Supplement to the NLTT Catalogue (Minneapolis: University of Minnesota Press)
- Monet, D.G. 1996, American Astronomical Society Meeting, 188, 5404.
- Monet, D. G. 1998, American Astronomical Society Meeting, 193, 112003
- Monet, D.G., Dahn, C.C., Vrba, F.J., Harris, H.C., Pier, J.R., Luginbuhl, C.B., & Ables, H.D. 1992, AJ, 103, 638
- Monet, D.G. 2003, AJ, 125, 984
- Narayanan, V.K. & Gould, A. 1999, ApJ, 523, 328

- Oppenheimer, B. R., Golimowski, D.A., Kulkarni, S.R., Matthews, K., Nakajima, T., Creech-Eakman, M., & Durrance, S.T. 2001, AJ, 121, 2189
- Röser, S. 1996, IAU Symp. 172: Dynamics, Ephemerides, and Astrometry of the Solar System, 172, 481.
- Salim, S. & Gould, A. 2003, ApJ, 582, 1011
- Skrutskie, M. F. et al. 1997, in The Impact of Large-Scale Near-IR Sky Survey, ed. F. Garzon et al (Kluwer: Dordrecht), p. 187
- van Altena, W.F., Lee, J.T., & Hoffleit, E.D. 1995, General Catalog of Trigonometric Parallaxes, Fourth Edition (Yale: New Haven)
- van Biesbroeck, G. 1944, AJ, 51, 61

This preprint was prepared with the AAS LATEX macros v5.0.

 ${\it Table 1.} \quad {\it White Dwarf Companions to Hipparcos Stars}$ 

NLTT	RA	Dec	V	V- $J$	$M_V$	$\Delta\theta$	p.a.	Hip. #	$\pi$	$\sigma(\pi)$	$\pi_{\mathrm{old}}$	$\sigma(\pi)$	Name	Type
	(2000)	(2000)				"	$\deg$		mas	mas	mas	mas		
1762	8.26164	44.73697	16.33	-0.33	11.22	28.8	30.7	2600	9.52	1.63			EGGR 306	DA4
12412	60.49868	-34.46993	17.82	0.99	14.23	64.1	313.0	18824	19.15	1.13		2.0	~ ~ ~ ~ ~	D. 1
13599	69.18708	27.16430	15.99	1.50	14.73	123.9	338.9	21482	56.02	1.21	60.2	$\frac{2.9}{2.9}$	G 39-27	DA
15768	88.47462	12.40402	15.60	0.20	11.95	90.0	232.0	27878	18.64	1.05	15.1	5.0	EGGR 44	DC5
16355	94.05943	-59.20764	13.65	-0.65	10.85	40.4	301.1	29788	27.50	0.50	42.0	15.0	LTT 2511	DB4
18414	116.41036	-33.93111	16.70	1.92	15.79	870.9	2.8	37853	65.79	0.56	57.5	3.1	GJ 288B	$\overline{DC9}$
26462	167.37547	-26.01849	16.79	0.41	13.93	100.2	180.0	54530	26.76	1.09			LP 849-059	
29967	183.15007	-6.37170	17.26	1.19	14.06	202.8	102.4	59519	22.94	1.63			LP 674-029 <del>□</del>	<del>i</del> DC
38926	224.46685	29.88101	17.66	-9.00	13.51	27.5	314.0	73224	14.79	1.24			ı	L:m
41169	236.87544	-37.91897	13.46	-9.00	12.54	15.0	130.0	77358	65.60	0.77	74.6	10.1	GJ 599B	DA7
42785	246.57123	2.18123	17.42	1.83	13.68	9.3	319.5	80522	17.86	1.91			LHS 3195	
47097	282.17372	68.87733	17.18	0.65	12.80	33.9	273.4	92306	13.33	0.85			LDS 2421B	
49859	311.52328	33.97068	14.29	0.67	12.57	87.9	271.2	102488	45.26	0.53			GJ 9707C	MV4
50189	314.19911	-4.84440	16.67	1.67	15.43	14.5	309.8	103393	56.56	4.03	64.6	5.1	vB 11	DC9
51482	323.06767	0.25407	15.27	0.37	11.80	133.0	29.0	106335	20.26	2.00			PHL 28	$\overline{DB}$
53526	334.86906	21.37246	17.69	1.25	14.21	83.3	282.3	110218	20.12	1.71			LP 460-3	DС
55288	343.95613	-7.83403	16.99	1.36	14.28	41.8	190.4	113231	28.72	1.30	36.7	5.3	G 156-64	DA
55300	343.98225	5.75618	15.71	0.53	13.78	17.1	285.2	113244	41.15	2.65			GJ 4305B	DB
55701	345.65880	76.50108	16.35	0.55	12.26	13.2	137.6	113786	15.22	0.81			EGGR 454	DC6
57958	356.79796	-26.39293	16.83	0.54	13.24	13.2	139.2	117308	19.11	2.93			G275-102	DB

Table 2.  $M_V > 14$  Companions to Hipparcos Stars

NLTT	RA	Dec	V	V- $J$	$M_V$	$\Delta\theta$	p.a.	Hip. #	$\pi$	$\sigma(\pi)$	$\pi_{\mathrm{old}}$	$\sigma(\pi)$	Name	Type
	(2000)	(2000)				"	$\overline{\deg}$		mas	mas	mas	mas		v <b>-</b>
638	3.42971	80.66376	16.38	5.45	14.92	12.8	126.1	1092	51.12	3.31			G 242-48B	M5
923	4.60755	44.02723	12.75	5.96	14.99	35.1	64.0	1475	280.27	1.05	280.27	1.05	HD 1326B	M3.5
2859	12.89642	-22.90851	16.35	5.45	15.44	16.8	71.4	4022	65.81	1.44			LDS 1082B	
8870	41.42178	44.95083	16.19	5.03	14.34	18.1	65.6	12886	42.59	2.21			GJ 3179B ?	M5?
8960	42.42210	75.02788	17.24	5.55	14.31	28.5	57.6	13179	25.98	2.00			LDS 1559B	
10761	50.63969	-30.19058	17.20	5.15	14.79	24.9	253.1	15725	32.99	1.29				
15867	89.57157	-4.63374	16.49	5.36	14.66	89.3	313.1	28267	43.10	0.93			BD-04 1310	
18218	114.90280	33.46461	18.83	6.01	16.10	13.0	57.4	37312	28.48	3.59	26.40	2.50	GJ 3458B	
19472	126.21852	-3.68378	17.16	5.57	14.99	356.4	47.1	41211	36.75	0.87				
22015	143.20107	26.99552	16.03	5.69	14.78	65.2	67.3	46843	56.35	0.89			GJ 354.1B	M5.5
24550	157.87822	57.08834	15.31	5.57	14.10	141.9	225.7	51547	57.18	1.31			LDS 2314B	M4.5
26247	166.37876	43.52164	14.62	5.89	16.20	31.4	126.1	54211	206.94	1.19	206.94	1.19	GJ 412B_	M6
28453	176.39744	-20.35138	18.20	6.47	16.68	15.4	60.8	57361	49.63	3.55			LP 793-34	late M
28513	176.63621	-40.49657	15.43	6.68	15.60	22.9	53.5	57443	108.23	0.70			vB 5	M4
28864	178.31940	-7.37521	18.67	-9.00	17.10	9.0	110.0	57959	48.63	3.34			GJ 452B	L:f
29580	181.44429	-18.82557	16.20	4.97	14.44	187.7	342.3	59000	44.41	1.51				
29661	181.84996	13.03719	18.06	5.06	14.97	66.9	283.3	59126	24.08	1.38				
29896	182.85170	43.55919	18.61	4.50	14.54	10.5	228.7	59428	15.35	1.64				
32366	194.12921	21.00776	19.04	4.96	14.56	35.2	244.0	63168	12.74	2.24			LDS 2891C	
34218	202.08720	30.05528	19.14	5.82	15.72	52.6	51.4	65706	20.74	1.97			LDS 1390B	
35919	209.84001	25.23418	17.26	4.86	14.18	20.3	177.8	68342	24.23	2.09			LHS 2838	$\mathbf{M}$
40719	234.21327	37.58035	19.28	-9.00	15.58	10.0	91.0	76452	18.21	1.65				L:+m
41096	235.54552	72.79309	18.06	-9.00	14.01	15.7	358.0	76902	15.48	4.29			LDS 1828B	L:a-f
42209	242.42474	65.82235	18.96	5.00	14.50	29.0	147.2	79180	12.82	0.77			LDS 2369B	
42494	245.01351	-37.53030	14.99	6.65	15.35	$_{5.3}$	227.2	80018	118.03	2.28	130.00	9.70	GJ 618B	M5.5
47621	289.24002	5.15044	17.05	7.15	18.21	74.0	150.0	94761	170.26	1.37	170.26	1.37	vB 10	M8V
49961	312.30737	32.28100	17.45	5.68	15.58	34.4	246.0	102766	42.23	0.98			LDS 2931B	
53379	334.01075	54.66652	16.72	7.00	15.06	76.8	107.4	109926	46.62	0.67			GJ 4269B	M4
57088	352.71868	27.77598	17.65	5.26	14.06	26.7	250.5	116052	19.13	1.21				




Fig. 1.— Color-magnitude diagram of 424 common proper motion (CPM) companions of Hipparcos stars with good (>  $3\sigma$ ) parallaxes. Error bars reflect only the parallax errors (and not the photometry errors) and so reflect the precision possible if the present, mostly photographic, photometry is replaced by CCD photometry. Stars without J photometry are displayed at V - J = -1. In addition all Hipparcos WDs with good parallaxes are shown as crosses. Note that these tend to be significantly more luminous than the CPM WDs from the catalog.




Fig. 2.— Color-magnitude diagram of the Hipparcos primaries to the stars shown in Fig. 1. Note that the CPM companions from Fig. 1 tend to fill in the region at the lower right, which is devoid of Hipparcos primaries.




Fig. 3.— Distribution of absolute magnitudes of the 424 stars in the catalog of CPM companions to *Hipparcos* stars (*bold histogram*), compared to that of the 400 dimmest stars in *Hipparcos* itself (*solid histogram*). Note that the CPM catalog contains many more extremely dim stars.




Fig. 4.— Differences between Hipparcos parallaxes divided by the root-sum square of the errors of the two components for 84 binaries with both components in Hipparcos. Histogram shows the actual distribution while the curve shows the expectation based on Gaussian statistics. There is rough agreement, but the actual distribution contains 4 pairs with exactly zero difference and three outliers at 2.8, 3.6, and 5.3  $\sigma$ , neither of which can be explained on the basis of Gaussian statistics.