Tycho Brahe's supernova: light from centuries past

Pilar Ruiz–Lapuente ^{1,2}

Received

accepted.

Running title: Tycho Brahe supernova

¹Department of Astronomy, University of Barcelona, Martí i Franqués 1, E–08028 Barcelona, Spain. E–mail: pilar@am.ub.es

²Max–Planck–Institut für Astrophysik, Karl–Schwarzschild–Strasse 1, D–85740 Garching, Federal Republic of Germany. E–mail: pilar@MPA–Garching.MPG.DE

ABSTRACT

The light curve of SN 1572 is described in the terms used nowadays to characterize SNeIa. By assembling the records of the observations done in 1572–74 and evaluating their uncertainties, it is possible to recover the light curve and the color evolution of this supernova. It is found that, within the SNe Ia family, the event should have been a SNIa with a normal rate of decline, its stretch factor being $s \sim 0.9$. Visual light curve near maximum, late–time decline and the color evolution sustain this conclusion. After correcting for extinction, the luminosity of this supernova is found to be $M_V = -19.58 - 5 \log (D/3.5 \text{ kpc}) \pm 0.42$.

Subject headings: cosmology: —stars: supernovae: general

1. Introduction

SN 1572 was well observed in Europe (as well as in the Far–East) for almost two years. It added a new aspect to the debate at the time over the Aristotelian cosmological views, as it forced to reconsider the immutability of the heavens and the solid nature of the celestial spheres: the "star" gained brightness and lost it during a period of two years, but it showed no detectable parallax. According to prevalent views about the heavens, mutability would only happen in the sublunar region. This was even the place where comets were assumed to originate. The appearence of SN 1572 challenged the order of the celestial spheres. The observers who followed it up³ took sides with respect to established Aristotelian views. Today we can still use their observations to see whether that supernova would be of use for cosmology in a different way, as a distance indicator if seen by observers billions of years away from us. It is indeed possible to have a clear idea of where SN 1572 stands among its class.

After several centuries of questioning the nature of SN 1572, the identification as a Type I supernova came through the revision of the light curve done by Baade (1945) and based on data taken or quoted by Tycho Brahe (1603a). Before that time there were still speculations on whether it was a variable star of some kind, a nova (Morgan 1945) and it was even still considered the suggestion of its cometary nature (Lynn 1883). The cometary idea expressed in 1573 by Jerónimo Muñoz was based on the fact that the event is aligned with the Milky Way, so its decay in luminosity could be explained if it was a comet born

³The observers who mostly contributed to measure the position and luminosity, Tycho Brahe, Thomas Digges, Thaddeus Hagecius, Michael Mästlin, Jerónimo Muñoz, Caspar Peucer & Johannes Prätorius held very different views on the meaning of SN 1572. A comparison of their measurements and an account of their views is given elsewhere (Ruiz–Lapuente 2003).

among the stars that would first approach and then move away from us just along the line of sight. Tycho's *stella nova* lies indeed only 49–98 pc above the Galactic plane.

In modern times, its comparison with other SNe allowed its classification as a Type I supernova (Baade 1945). Later on, this class was shown to contain events of very different nature: those identified with the explosion of white dwarfs (Type Ia) and those corresponding to the collapse of massive stars whose envelope had lost its hydrogen content in the interaction with a binary companion (Type Ib or c). SN 1572 was of the Type Ia class as discussed by de Vaucouleurs (1985) and by van den Bergh (1993). No further doubts that it is a Type Ia SN can now be held in view of the X-ray spectrum, which clearly differentiates SNIa events from SNe coming from the collapse of massive stars (Hughes et al. 1995). A first comparison of its luminosity at maximum with the bulk of SNeIa would have led to think that it was fainter than normal SNeIa. van den Bergh (1993) considers whether it could be a peculiar, subluminous SNIa, like SN 1991bg. However, SN 1572 was heavily obscured. It was reddened by $E(B-V)=0.6\pm 0.04$ as it corresponds to the reddening of the stars near its position (Ruiz-Lapuente et al. 2003a,b). After taking into account the extinction undergone across the Milky Way, as well as the measurement of its decline rate, it is concluded that SN1572 was not a 91bg-like event. Neither was it an overluminous SNIa like SN 1991T or similars, but rather an event in the middle of the SNIa class.

2. New evidence on visual, color evolution and late decline

2.1. Visual light curve till 60 days

The date of maximum of SN 1572 is, from the available accounts, the most uncertain record. The first observation was done on Nov 6 though there is no record of the magnitude at that date (see for a full discussion Stephenson & Green 2002). On Nov 2 it was not

yet noticeable as reported by Muñoz (1573). In Nov 11, date of the first observation done by Tycho Brahe (1603a) and the first reported observation by Muñoz (1573), it seems to have achieved a magnitude between those of Jupiter and Venus. The Nov 5 detection given in the report of the *Progymnasmata* by Tycho Brahe (1603b) is an erroneous date, as we have checked in the original. On Nov 16 and 17, when Caspar Peucer and Johannes Prätorius saw it, it seems to have a magnitude close to Venus (Table 1 and notes). Later on, in Jan 7, it was already fainter than Jupiter (Muñoz 1573). The reconstruction of the observations suggests an apparent visual magnitude at maximum of -4.0 ± 0.3 (Baade 1945; de Vaucouleurs 1985). The overall light curve fitting done in the present work by trying different decline rates (see below) indicates that the maximum was broad and that it should have taken place around Nov 21. On Nov 11, and even on Nov 16, when Peucer and Prätorius observed it, it would be still on the rise. This is consistent with the mean risetime to maximum of SNeIa of $17.8^{+1.9}_{-1.0}$ days (Goldhaber et al. 2001). While most authors have taken arbitrarily Nov 15 as the date of visual maximum, it seems obvious that the maximum could not have happened before Nov 20.

The SN 1572 data are compared in this section to templates using the stretch factor s as characterization of the rate of decline (Perlmutter et al. 1999; Goldhaber et al 2001; Nobili et al. 2003). The stretch factor s method is used by the Supernova Cosmology Project to quantify the decline rate of the supernova from data extending up to 60 days after maximum. Even in absence of a measurement of the brightness at maximum, the method allows to produce a fine description of the supernova within the family of decline rates. The magnitudes and limits on magnitudes for SN 1572, when compared to templates, lead us to conclude that this supernova was not fast. The best agreement found for SN 1572 corresponds to $s \sim 0.9$. In this regard, SN 1572 is very similar to SN 1996X which had a s = 0.889. A comparison can be seen in Figure 1, where the data of SN 1572 are shown overplotted to a s = 0.9 SNIa template in V. We also show for comparison the visual light

curves of SN 1996X (normal SNIa of s = 0.889) and SN 1991bg (a fast subluminous event of s = 0.62). Therefore, these first 60 days show that there is no similarity with the fast subluminous SN 1991bg. A broad light curve as the one of SN 1991T can also be excluded from the constraints about its decline (see Figure 1).

2.2. Late-time decline

An additional proof that SN 1572 is a standard SNIa comes from its slow late decline between 100 and 450 days after maximum, similar to that of the bulk of SNeIa. SN 1572 had a late decline of 1.4 mag in 100 days during that epoch, in concordance with normal SNeIa and far from the fast decline of SN 1991bg (Filippenko et al. 1992; Leibundgut et al. 1993; Ruiz–Lapuente et al. 1993; Turatto et al. 1996). SN 1990N and other normal SNeIa had decline rates in such period of 1.38–1.5 mag in 100 days. In consistency with the findings near maximum, late-time and overall similarity to the light curves of normal SNIa is found. In order to test the adequacy of the fit using the whole light curve, we construct new templates for a normal event of s = 0.9 (well exemplified by SN 1996X), a subluminous 91bg-type event and an overluminous 91T-type event matching the available early and late-time observations. Before 60 days those templates are equal to the Hamuy et al. 1996b templates and the SCP templates for the corresponding stretches of those SNeIa. At later times they follow the available late-time photometry (Schmidt et al. 1993; Salvo et al. 2001) to their latest available epoch. The fit of the SN 1572 data to the SNIa template corresponding to s = 0.9 has a χ^2 of 14.44 for 10 degrees of freedom, which is acceptable. In contrast, the fit to the template of a fast declining, underluminous SNIa like SN 1991bg (s = 0.62) has an exceedingly high χ^2 of 53.55 for 4 degrees of freedom (the two premaximum points as well as the last 4 points have been excluded because of the lack of data for those subluminous SNeIa at such stage). In the other extreme, the fit to the

template of a slow-declining, overluminous SNIa like SN 1991T (s = 1.2) is almost as bad, with a χ^2 of 82.99 for 10 degrees of freedom.

In Figure 2, we show the overall visual light curve till almost 500 days compared with that of SN 1996X and the templates for SN 1991bg and SN 1991T. The last upper limit given by Tycho Brahe indicates that the supernova had not yet leveled off at 480 days past maximum due to a light echo produced in an intervening dust cloud. Not all the SNeIa in dusty regions have shown light echoes associated with the scattering of the supernova light in neighboring dust clouds, though. The production of light echos depends on the spatial distribution of dust relative to both observer and supernova. In the case of SN 1986G, a very heavily reddened supernova, such light echo was not observed (Schmidt et al. 1993). However, for SN 1572 one can not exclude a leveling off occuring later, at a fainter magnitude level. SN 1991T and SN 1998bu have indeed shown a slowing down of the V magnitude decline at around 400 days as result of the light echo (Schmidt et al. 1993, Cappellaro et al. 2001), and the leveling off occurs at 500 days about 10 magnitudes below the maximum brightness. For SN 1572 an equivalent step in the level of brightness would correspond to V=6, which is about the naked eye limit. SN 1998bu is a SNIa very similar to SN 1996X and SN 1572 but with a reddening of $E(B-V) = 0.32 \pm 0.04$ (Hernandez et al. 2000; Cappellaro et al. 2001). In fact, we can think of Tycho Brahe's supernova as SN 1998bu with twice the reddening.

Moreover, the late-time decline of SN 1572 informs us of deposition of the energy from the decay ${}^{56}\text{Co} \rightarrow {}^{56}\text{Fe}$ in the ejecta of this supernova. Its slow decline would fit the declines predicted for SNeIa coming from the explosion of Chandrasekhar-mass WDs. The late-time decline in V indicates a significant deposition of the energy of the positrons which arise in the decay ${}^{56}\text{Co} \rightarrow {}^{56}\text{Fe}$. After 200 days since explosion, positrons become the dominant source of luminosity. Departures from full trapping of ${}^{56}\text{Co}$ decay are seen in all SNeIa and they are expected. They arise from lack of confinement of positrons to the ejecta of the supernova and/or incomplete thermalization of their energies. Various effects can lead to the diversity in the late-time SN bolometric light curves: nucleosynthesis, kinematic differences, ejected mass, degree of mixing of the ejecta, and configuration and intensity of the magnetic field. In particular, as suggested in Ruiz–Lapuente & Spruit (1998), those late-time declines can serve to investigate the magnetic field configuration of the SN and trace it back to the original configuration of the WD right before explosion. It is found that departures from the full trapping of 56 Co decay of the order of 10–15% at about 400 days can be explained by the distribution of radioactive material. Larger departures, of 30-40% or larger, have to be interpreted in terms of lack of magnetic field confinement of positrons, or even enhancement of the escape favored by a radially combed configuration of the magnetic field. In the case of SN 1991bg the very fast drop of the late light curve suggests not only a total ejected mass well below the Chandrasekhar mass, but an absence of a significant tangled magnetic field. An exploration of the late-time decline predicted for different explosion models compared with a sizeable sample of bolometric light curves was done by Milne, The and Leising (1999). For SN 1572, the availability of data is limited to the visual band. Though, analogies can be drawn with similar SNeIa for which the bolometric data are available.

The late decline is, as seen above, a major proof which confirms that this historical supernova is not part of the estimated 16% of intrinsically subluminous SNeIa (Li et al. 2001), likely arising from a different class of explosion ejecting a smaller amount of mass (Ruiz–Lapuente et al. 1993). If it were part of the class of peculiar subluminous SNeIa, caution should be given when used for cosmology, because it is not clear whether those subluminous events are in the linear relation brightness–rate of decline.

2.3. Reddening and color

The most direct estimate of the reddening comes from our measurement of the reddening and extinction of the stars around the geometrical center of SN 1572 at the distance of the supernova (Ruiz–Lapuente et al. 2003a,b). The stars which are compatible with a distance in the range of 3 ± 1 kpc (see de Vaucouleurs 1985 for a review of the distances towards SN 1572) have reddenings which average to $E(B-V)=0.6 \pm 0.04$. Those values were derived in the programme of search of the companion star of Galactic SNeIa (Ruiz–Lapuente et al. 2003a,b). Star candidates within a radius compatible with the degree of precision on the center and distance gained during the 431 years since explosion are modeled to determine T_{eff} , log g, metallicity, distance and E(B-V). The radial velocities are measured. The individual reddening values found in that programme for the stars within 35% of the radius of the remnant go from E(B-V) = 0.50 to E(B-V)=0.8, the values increasing with distance.

The above mean reddening to SN 1572 corresponds to an extinction $A_V = 1.86 \pm 0.12$ mag if we take $R_V = 3.1$ (Sneden et al. 1978; Riecke & Lebofsky 1985). The Galactic extinction data based on COBE/DIRBE observations (Drimmel & Spergel 2001) give a value of $A_V = 1.77$ mag in that direction (the maximum Galactic extinction being $A_V = 1.90$ mag there). Thus the extinction measured at the distance of SN 1572 is in agreement with the COBE results. Correcting the apparent brightness from the dimming by dust, we find again that Tycho's SN was not subluminous but a "normal" SN Ia.

We can correct as well the color of the supernova from the records of the epoch and derive the intrinsic color evolution. SNeIa are found to show a similar color evolution two months after maximum, with a well–established law of low intrinsic dispersion valid from 30 to 90 days, explored in Phillips et al. 1999 (Ph99) based on the work of Lira (1995):

$$(B - V)_0 = 0.725(\pm 0.05) - 0.0118(t_V - 60) \tag{1}$$

where t_V is the time since visual maximum.

The color of SN 1572 two months after discovery was reported to be similar to Mars and Aldebaran (thus B–V in the range 1.36–1.54). Previous and subsequent color estimates are also shown in Table 2. After correcting the observed color by our measurement of reddening $E(B - V) = 0.6 \pm 0.04$, the intrinsic color at 55 ±10 days is $(B - V)_0 = 0.76$ \pm 0.24, very much in agreement with the expected $(B - V)_0 = 0.78 \pm 0.15$ for that epoch. Such color fits very well in the Ph99 law for the time evolution of the intrinsic color of the SNeIa. In agreement with that law, Nobili et al. (2002) have shown that the color evolution of the bulk of SNeIa has a low dispersion of 0.1 mag in the tail. Before maximum the corrected color of SN 1572 would be $(B - V)_0 = 0.22 \pm 0.29$, thus consistent with the fact that normal SNeIa have $(B - V)_0 \sim 0$. SN 1991bg and other subluminous SNeIa clearly deviate from the standard color evolution at early epochs, being intrinsically red at maximum $(B - V)_0 = 0.6$ (Leibundgut et al. 1993; Ph99). Moreover, the color evolution derived from the B and V templates for a s = 0.9 SNIa gives a good agreement with the supernova. In Figure 3 we show the color light curve of SN 1572, corrected by its E(B-V)=0.6, compared with that of "normal" SNeIa. Both the peculiar SN 1991bg and the normal SN 1996X are also shown in their reddening–corrected colors.

The data displayed in this figure come from different observers (see Table 2 and notes). Color estimates by eye are difficulted by the lack of an established system which encompases the whole range of colors. The conversion of eye–estimated colors to our filter scale is addressed in Pskovskii (1977) and Schaefer (1996). To add to the inherent eye uncertainty in estimating color, it has to be noted that the first comparisons done by the observers of the time using planets like Jupiter, Venus or Saturn as references, were difficulted by the fact that the planets were often very far from the position of SN 1572 (away in the sky by 59^{0}). De Vaucouleurs (1985) discussed the effect of that on the early visual estimates (later estimates since they are done through comparison with particular stars are more reliable).

In the nebular phase, at 175 days according to Tycho Brahe's records (Table 2) the supernova went back to a white color, which is consistent, after correcting for extinction, with what is found in normal SNeIa as well. Uncertainties in the color estimates have been assigned and new records included.

2.4. The luminosity of Tycho's supernova

Given the apparent visual magnitude, the luminosity of Tycho Brahe's supernova would have corresponded to a visual absolute magnitude of $M_V = -17.72 - 5 \log(d/3.5 \ kpc) - A_V$. If corrected for $A_V = 1.86 \pm 0.12$, this means $M_V = -19.58 - 5 \log(d/3.5 \ kpc) \pm 0.42$.

De Vaucouleurs (1985) gives as the most likely estimate to the distance to SN 1572 3.2 \pm 0.3 kpc. Taking 3.2 kpc for the distance, the luminosity would be -19.38 ± 0.42 . That compares well with the value $M_V = -19.12 \pm 0.26$, the mean magnitude from the Calán/Tololo sample (Hamuy et al. 1996a).

To get a more refined distance to SN 1572, one would need to follow a similar procedure as in Winkler, Gupta & Knox (2003) for SN 1006. These authors obtained a distance to SN 1006 of 2.18 ± 0.08 kpc using the SNR expanding parallax method (Kirshner, Winkler & Chevalier 1987) with precise measurements of the proper motion of the filaments in SN 1006 and narrower estimates of the shock velocity of the expanding remnant.

The proper motion of the filaments in SN 1572 was measured by Kamper & van den Bergh (1978), and it is about 0.22 ± 0.02 " yr⁻¹. There is, however, a larger uncertainty in the estimate of the shock velocity, which is obtained through modeling of the broad $H\alpha$ emission of the material swept up by the SN. The estimate is model dependent and improves with a better knowledge of the composition of the fully ionized plasma behind the shock. Kirshner, Winkler & Chevalier (1987) had suggested from the observations available at that time a distance to SN 1006 between 1.4 and 2.1 kpc and a distance to SN 1572 of 2.0–2.8 kpc. The estimates of the shock velocity, though, could soon be revised from the wealth of spectral observations on SN 1572's filaments taken during the last years.

A rough estimate of the distance to Tycho's SN can be made by comparison of the angular diameters of the two historical SNeIa. The angular diameter of G327.6+14.6 (the remnant of SN 1006) is $\theta(1006) = 30.60 \pm 0.06$ arcmin whereas that of Tycho is $\theta(Tycho) = 8.65 \pm 0.05$ arcmin. We can place $\theta(Tycho) / \theta(1006) = d(1006) / d(Tycho) t(Tycho) / t(1006) \eta$, where t(Tycho) and t(1006) are the respective ages of the remnants and $\eta = v(Tycho)/v(1006)$ is the ratio of the average expansion velocities of the two SN ejecta. Such approach gives, if we use the distance obtained by Winkler, Gupta & Long (2003) to SN 1006, a distance to SN 1572 of $d(Tycho) \simeq 3.3$ kpc, well in the middle of other estimates.

All of the above shows that it is at present hard to escape from the uncertain range in the distance to SN 1572, i.e. from 2 to 4 kpc with the analyses done so far. Such limitation does not prevent to infer the right place of SN 1572 within the luminosity sample: the conclusion on the brightness-rate of decline for SN 1572 is based on the gathered empirical evidence and the well-established use of the peak luminosity-decline correlation.

Incidentally, if a good distance would be available for SN 1572, one could obtain the distance to SN 1996X, which is not well determined with the present available methods. The $D_n-\sigma$ relation gives a distance modulus $\mu = 31.32 \pm 0.4$ (Faber et al. 1989), whereas the SNeIa calibration using $H_0 = 65$ km s⁻¹ Mpc⁻¹ would place it a significantly larger distance (Salvo et al. 2001). The very good agreement between the late time light curves of SN 1572 and SN 1996X allows us to infer a $\Delta \mu$ (SN1572–SN1996X) = 18.9 ± 0.12 between both SNeIa.

3. Conclusions

We have found that SN 1572 was a supernova very close to the template with a *stretch* factor $s \sim 0.9$ (see Table 3 for a summary). The light curve grows in precision towards the late times, being highly uncertain around maximum brightness. An overall agreement between early, late decline and color with the expected evolution of normal SNeIa supports our conclusion.

Type Ia supernovae with stretch factors between 0.9 and 1.1 make the vast majority of the observed population. They are not only those most frequently found in nearby searches, but also the bulk of discoveries in cosmological searches at high–z, as can be seen in the sample of SNe at z > 0.3 found by the Supernova Cosmology Project (Perlmutter et al. 1999). Among SNeIa of s ~ 0.9 in nearby galaxies for which very late–time data are available, we have found a close resemblance to SN 1996X in rate of decline. SN 1572 likely has a slightly slower rate. However, whereas SN 1996X was not heavily reddened, the reddening in SN 1572 is $E(B-V)=0.60 \pm 0.05$. While at present heavily reddened SNe Ia involve a larger difficulty than unreddened ones for their proper use in cosmology (Riess et al. 2000; Knop et al. 2003), a better use of multicolor light curves should allow to overcome the problem in their calibration for cosmology.

Tycho Brahe's supernova belongs to the SNIa type that are used for the determination of the nature of dark energy. Its light, traveling already light years away from us, carries the information on our acceleration epoch to an even more dark–energy dominated future. If used for cosmology by observers in galaxies billions of light years distant, it would require, certainly, a careful handling of its intrinsic brightness.

Acknowledgements This paper amongst others to come is dedicated to my father. Past, present and future merged for him just a few months ago, as sadly as unexpectedly. His life was devoted to help the souls of many and to carry research in brain and mind studies. His inner light lives forever with us. I would like to express my gratitude to M. A. Granada, from the Department of History of Philosophy of the U. Barcelona, for helping to locate the XVI century historical sources.

REFERENCES

- Baade, W. B. 1945, ApJ, 102, 309
- Brahe, T. 1603a, Astronomiae Instauratae Progymnasmata, in Opera omnia 2, 307, ed. I.L.E. Dreyer (Amsterdam: Swets & Zeitlinger 1972)
- Brahe, T. 1603b, Astronomiae Instauratae Progymnasmata, in Opera omnia 3, 81, ed. I.L.E. Dreyer (Amsterdam: Swets & Zeitlinger 1972)
- Cappellaro, E., Patat, F., Mazzali, P. A., Benetti, S., Danziger, I.J., Pastorello, A., Rizzi, L., Salvo, M., & Turatto, M. 2001, ApJ, 549, L215
- de Vaucouleurs, G. 1985, ApJ, 289, 5
- Drimmel, R., & Spergel, D.N. 2001, ApJ, 556, 181
- Faber, S.M., et al. 1989, ApJS, 69, 763
- Filippenko, A.V., et al. 1992, AJ, 104, 1543
- Goldhaber, G., et al. 2001, ApJ, 558, 359
- Hamuy, M., Phillips, M.M., Schommer, R.A., Suntzeff, N.B., Maza, J., & Avilés, R. 1996a, AJ, 112, 2391
- Hamuy, M., et al. 1996b, AJ, 112, 2438
- Hernandez, M., et al. 2000, MNRAS, 319, 223
- Hughes, J.P., Hayashi, I., Helfand, D., Hwang, U., Itoh, M., Kirshner, R., Koyama, K., Markert, T., Tsunemi, H., & Woo, J. 1995, ApJ, 444, L81
- Kamper, K.W., & van den Bergh, S. 1978, ApJ, 224, 851

- Kirshner, R.P., Winkler, P.F., & Chevalier, R.A 1987, ApJ, 315, L135
- Knop, R., et al. (SCP2003) 2003, ApJ, submitted
- Leibundgut, B., et al. 1993, AJ, 105, 301
- Li, W., Filippenko, A.V., Treffers, R.R., Riess, A.G., Hu, J., & Qiu, Y. 2001, ApJ, 546, 734
- Lira, P. 1995, Master's Thesis, Univ. of Chile
- Lynn, W.T. 1883, The Observatory, 6, 151
- Milne, P.A., The, L.-S. The, & Leising, M.D. 1999, ApJS, 124, 503
- Morgan, F.P. 1945, JRAS, 39, 370
- Muñoz, J. 1573, Libro del nuevo cometa (Valencia: Pedro de Huete), re-ed. V. Navarro-Brotons (Valencia: Hispaniae Scientia 1981)
- Nobili, S, Goobar, A., Knop, R., & Nugent, P. 2003, A&A, 404, 901
- Nugent, P., Kim, A., & Perlmutter, S. 2002, PASP, 114, 803
- Perlmutter, S., et al. (SCP99) 1999, ApJ, 517, 565
- Phillips, M.M., et al. 1987, PASP, 99, 592
- Phillips, M.M., Lira, P., Suntzeff, N.B., Schommer, R.A., Hamuy, M., & Maza, J. 1999, AJ, 118, 1766 (P99)
- Pskovskii, Y.P. 1977, Soviet Astron., 22, 420
- Rieke, G.H., & Lebofsky, M.J. 1985, ApJ, 288, 618
- Riess, A., et al. 2000, ApJ, 536, 62

- Ruiz–Lapuente, P. 2003, in Astronomy as a Model for the Sciences in Early Modern Times, ed. B. Fritscher, in press
- Ruiz-Lapuente, P., Jeffery, D.J., Challis, P.M., Filippenko, A.V., Kirshner, R.P., Ho, L.C., Schmidt, B.P., Sanchez, F., & Canal, R. 1993, Nature, 365, 728
- Ruiz-Lapuente, P., & Spruit, H.C. 1998, ApJ, 500, 360
- Ruiz–Lapuente, P., Comeron, F., Smartt, S., Kurucz, R., Méndez, J., Canal, R., Filippenko,
 A.V., & Chornock, R. 2003a, in From Twilight to Highlight: the Physics of
 Supernovae, ed. W. Hillebrandt & B. Leibundgut (Berlin: Springer), 140

- Salvo, M., et al. 2001, MNRAS, 321, 254
- Schaefer, B.E. 1996, ApJ, 459, 438
- Schmidt, B. P., Kirshner, R.P., Leibundgut, B., Wells, L.A., Porter, A.C., Ruiz–Lapuente, P., Challis, P., & Filippenko, A.V. 1994, ApJ, 434, 19
- Sneden, C., Gehrz, R.D., Hackwell, J.A., York, D.G., & Snow, T.P. 1978, ApJ, 223, 1689
- Stephenson, F. R., & Green, D.A. 2002, Historical Supernovae and their Remnants (Oxford: Oxford University Press)
- Turatto, M., Benetti, S., Cappellaro, E., Danziger, I.J, Della Valle, M., Gouiffes, C., Mazzali, P.A., & Patat, F. 1996, MNRAS, 283, 1
- van den Bergh, S. 1970, Nature, 225, 503
- van den Bergh, S. 1993, ApJ, 413, 67
- Winkler, P.F., Gupta, G., & Knox, S.L 2003, ApJ, 585, 324

This manuscript was prepared with the AAS ${\rm IAT}_{\rm E}{\rm X}$ macros v4.0.

NOTES TO TABLE 1

¹ Jerónimo Muñoz writes that he was certain that on November 2, 1572 the "comet" was not in the sky, as he was *teaching to know the stars to a numerous group of people that* evening (Muñoz 1573).⁴

² Nov 11 is the first observation quoted by Muñoz in his book. The apparent magnitude appeared somewhat greater than that of Jupiter, which was 59 degrees distant from the comet, and almost equal to that of the Morning Star. Jupiter was at that time V=-2.61 and Venus V=-4.35 (Baade 1945). We assign -3.0 ± 0.3 .

³ On Nov 11 Tycho Brahe also saw it, and his preliminar report in *De nova stella* said that *equaled Venus when this planet was at maximum brightness*. He was, however, more conservative in the account he gives in *Progymnasmata*. We give -3.5 mag to the estimate of Tycho ± 0.3 . In the following we must keep in mind that Tycho Brahe observes from a

⁴Jerónimo Muñoz (?-1592) was professor of Hebrew and Mathematics at the University of Valencia and later on of *Astrology* at the University of Salamanca. The records assembled here are from his Book of the New Comet (*Libro del nuevo cometa*) written as King Philip II consulted him on SN 1572. He determines the absence of parallax of the star and gives estimates of color and magnitudes, though his account is mostly addressed to criticize the Aristotelian cosmology. He wrote that the heavens and stars are not made of quintessence, but related to the elements, and advocated to reconsider the presocratic cosmological views of Democritus and Anaxagoras. Tycho Brahe (1603b) compiled parts of Muñoz's observations and discussed those views in the third volume of the *Progymnasmata*. He left out, however, some useful information that we have added here. We corrected as well the premaximum quote. latitude of ~ N55⁰, as he stays in the area of Copenhague for two years since the discovery of SN 1572. From that location the supernova can be observed well above the horizon through the whole year. De Vaucouleurs notes that when comparing SN 1572 with Venus one has to account from the fact that the earlier is not affected by extinction while the second was significantly affected. This places V=-3.8 or fainter if one considers the apparent magnitude of Venus affected by extinction.

⁴ Tycho Brahe reports in the *Progymnasmata* on the observations done by Caspar Peucer and Johannes Prätorious on Nov 16. According to the first, *it surpassed in brightness all stars and planets with the exception of Venus*, for the second, it was *brighter than Jupiter but easily fainter than Venus*. Michael Mästlin who saw it before Nov 17 states that *almost equalled Venus*. It has been given to these estimates a value of V=-4.0 \pm 0.3.

⁵ Tycho Brahe does not give the exact date of the observations. He quotes the month or an interval encompassing two months. It is reasonable to think that he refers to dates in the middle of the period. Therefore, we follow here Baade (1945) and assign the middle of the period as date of observation. However, we add \pm 7 days of uncertainty in the case a month is quoted, and \pm 14 days if the period encompasses two months.

⁶ This is an upper limit given by Muñoz. He writes, when it began to be visible, it looked larger than Jupiter, and now, on January 7, 1573, it already looks smaller than Jupiter; this could have happened because it had risen higher than where it was when it first appeared. This upper limit to the rate of decline as well as the data around it by Tycho Brahe constrain the light curve stretch factor below s = 1.1.

⁷ This is a difficult estimate. The interval quoted encompasses 2.46 mag (see Baade 1945). It is uncertain the interpolation done, from Jupiter with V=-2.18 to the brightest stars of the first class which are +0.28. We assign to the Baade estimate of V=-1.4 an uncertainty

of ± 0.4 .

⁸ This a bit better estimate, as encompasses an smaller range in magnitudes ± 0.2 if one selects the brighter range of first magnitude stars (0.2–0.3).

⁹ It is a very ambiguous quote, one should give it ± 0.5 . The stars of second magnitude had a range encompassing 1 mag, $m_V = 4.0$ and $\sigma_V = 0.5$.

This is the point that lies far from the expected curve. It has been checked whether a problem of differential extinction could have affected SN 1572 with respect to the second magnitude stars, but there were second magnitude stars at similar altitude as SN 1572. The offset of 0.6 mag can not be attributed to extinction problems, but to ambiguous description of the record.

¹⁰ Tycho Brahe's estimates in magnitude are more precise now as he refers to particular stars not far from the supernova with a low scatter in magnitude. The mean of those stars is $m_V = 2.48$ and $\sigma_V = 0.23$. We assign ± 0.2 to the error in the mags of those stars.

¹¹ It becomes a very precise estimate by comparing to the fourth mag stars in Cassiopeia, when knowing that it was brighter than the nearby eleventh star of Cassiopeia. An error of 0.2 mag is assigned.

¹² This seems to be the most precise description as it says it is equal to κ Cas.

 13 Errors of 0.2 and 0.4 mag respectively seem reasonable for these last estimates.

¹⁴ It can be noted that the quote that it was not observable later than March 1574 (V > 6.0) allows to exclude that the luminosity levelled off, due to a light echo, at ~ 10 mag below the supernova maximum brightness.

Date		Phase	Description		
	(days)	Adopted	1	m_V	Ref
1572 Nov 2	Nov 2	-	No detection	> 5	1
1572 Nov	Nov 11	-10	Somewhat brighter than	$-3.\pm0.2$	2
			Jupiter		
			and almost equal to the Morning Sta	ır	
1572 Nov	Nov 11	-10	Equalled Venus when this	$-3.\pm0.2$	
			planet was at maximum brightness		3
1572 Nov 16,17	Nov 16 ± 1	$1-5\pm 1$	Almost as bright as Venus	-4.0 ± 0.3	4
			and Jupiter		
1572 Dec	Dec 15 ± 7	24 ± 7	About as bright as Jupiter	-2.4 ± 0.3	6
1573 Jan 7	Jan 7	47	Already fainter than Jupiter	>-2.	7
1573 Jan	Jan 15 \pm 7	55 ± 7	A little fainter than Jupiter	-1.4 ± 0.4	7
			and brighter than the brighter		
			stars of first mag		
1573 Feb–Mar	Mar 2 ± 14	4101 ± 14	Equal to brighter stars of first mag	$+0.3 \pm 0.1$	2^{8}
1573 Apr–May	May 1 ± 14	161 ± 14	Equal to stars of second mag.	$+1.6 \pm 0.$	5^{9}
1573 July–Aug	Aug 1 ± 14	253 ± 14	Equal to $\alpha, \beta, \gamma, \delta$ Cas	$+2.5 \pm 0.1$	2^{10}
1573 Oct–Nov	Nov 1 ± 14	345 ± 14	Equal to stars of fourth mag.	$+4.0 \pm 0.1$	2^{11}
			(among Cassiopea stars)		
1573 Nov	Nov 15 ± 7	7359 ± 7	Equal to κ Cas	$+4.2 \pm 0.2$	12
1573 Dec–1574 Jε	$n Jan 1 \pm 14$	406± 14	Hardly exceeding stars of fifth mag.	$+4.7 \pm 0.1$	2^{13}

Table 1: Visual estimates of Tycho's Supernova

1574 Feb Feb 15 \pm 7 451 \pm 7 Equal to stars of sixth mag. $+5.3 \pm 0.4^{13}$

Table 2: Colors of Tycho's Supernova

Date		Phase	Description	$\mathrm{B}-\mathrm{V}$	Observer
		Adopted		Adopted	Reference
1572 Nov	Nov 21 ± 7	-6 ± 7	Like Venus $(B - V = 0.83)$	$0.82 {\pm} 0.25$	1
			and Jupiter $(B - V = 0.82)$		
1572 Dec	Dec 1 ± 5	10 ± 5	Yellowish	1.0 ± 0.25	2
1572 Nov-Dec	Dec 2^{7}_{-22}	11^{7}_{-22}	Between Saturn $(B - V = 1.04)$	$1.2{\pm}~0.25$	3
			and Mars $(B-V)=1.36$,		
			closer to Mars		
1572 Jan	Jan 1 \pm 7	41 ± 7	Like Aldebaran (B–V=1.52)	1.52 ± 0.25	4
1573 Jan	Jan 15 ±7	55 ± 7	Similar to Mars $(B - V = 1.36)$	1.36 ± 0.2	5
1573 Feb	Feb 28 ± 7	$99{\pm}7$	Return to original	$0.82 {\pm} 0.25$	6
1573 May	May 15 ± 7	175 ± 7	Like the original $(B - V = 0.83)$	0.82 ± 0.25	7

NOTES TO TABLE 2

¹Various reports (see Baade 1945) give a color similar to the colors of Venus and Jupiter.

² Tycho Brahe (1603a) states that after maximum it became yellowish in color. By comparison to the description as whitish (like Jupiter) and red like Aldebaran, we assign $B-V = 1.0 \pm 0.25$.

³ Muñoz (1573) states that the color was between those of Saturn and Mars, closer to Mars. One could identify then the color as 1.2 ± 0.1 . Unfortunately, there is ambiguity in the date he is referring to: he mentions that he first saw it on Dec 2, and checked with observers that it was already there on Nov 11. He uses the expression "the color then was", so it is not clear whether he refers to Nov 11 or Dec 2. It seems that the description, because of its detail, is based on his own appreciation of color on Dec 2 (we place an errorbar in the dates towards Nov 11, and a 7 days error after Dec 2).

⁴Tycho Brahe notes that shortly after the beginning of the year it was red like Aldebaran (Brahe 1603a). We assign 0.25 mag of uncertainty to those estimates.

⁵ According to Prätorius it was like Mars two months after its appearance (around Jan 15).

⁶Mästlin refers that at the end of February it is going back to its whitish color. Tycho states that it had a red color along February and March, becoming whitish after May. However, this seems to be a worse account of the color than the one given by Mästlin. In April it is described as silverish by some observers in Spain, and as similar to Saturn (van den Bergh 1970).

 7 The last estimate is given 0.82 ± 0.25 in account for Tycho Brahe on the color in May.

Table	3:	SN	1572
10010	.	\sim \cdot	T O I D

Table 5. 51 157	_				
Parameter	8	$\Delta m(V)_{20}$	$\Delta m(V)_{60}$	$\gamma^{late}_{(V)}$	$B - V_0$
SN 1572	0.9 ± 0.05	1.1 ± 0.4	2.6 ± 0.6	1.6 ± 0.04	0.22 ± 0.25
SN 1991bg–like	0.62	1.68	3.3	2.7	0.6
SN 1991T–like	1.2	0.72	2.46	1.68	-0.1
$Normal^1$	0.9	0.9	2.8	1.7	0.0

Normal SNe Ia of s = 0.9. Whereas the $\Delta m(V)_{20}$ is not well defined for SN 1572 due to a lack of accurate observations in the week around the maximum, the rate of decline in terms of stretch can be well determined by fitting premaximum and postmaximum up to 60 days.




Fig. 1.— The visual light curve of SN 1572 till 60 days. The solid curve is the V light curve of a s = 0.9 SNIa, which gives the best account for the decline. Such *stretch factor* is typical of normal SNe Ia. We show for comparison the V light curve of the normal SNIa SN 1996X, whose stretch factor is s = 0.889 and of the fast-declining SN 1991bg. SN 1572 was significantly slower than SN 1991bg. The light curves plotted in dashed lines are the templates of 91bg-type events and 91T-like events, wich depart significantly from the data.




Fig. 2.— The visual light curve of SN 1572 till 500 days. Its late rate of decline is the one of normal SNeIa. It is very similar to the decline of the s = 0.889 SN 1996X. The visual data of SN1991bg and the template light curves of this SNIa and SN1991T are shown for comparison.




Fig. 3.— Color evolution of SN 1572 corrected from extinction as compared with normal SNe Ia and SN 1991bg. SN 1572 is consistent with the color evolution of a s = 0.9 SNIa (template for this stretch is plotted as dot-dashed curve). SN 1996X has been corrected for its very small reddening $E(B-V)=0.01 \pm 0.02$ as well as SN 1991bg ($E(B-V)=0.03 \pm 0.05$) (Ph99).