# Star Formation in Satellite Galaxies

José G. Funes, S.J.

Vatican Observatory, Univ. of Arizona, Tucson, AZ 85721, USA

Carlos M. Gutiérrez

Instituto de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Spain

Francisco Prada

Instituto de Astrofísica de Canarias & Isaac Newton Group of Telescopes, Spain

Marco Azzaro

Isaac Newton Group of Telescopes, Santa Cruz de La Palma, Spain

Marcelo B. Ribeiro

Instituto de Física, Universidade Federal do Rio de Janeiro, Brazil

Abstract. The study of satellite galaxies can provide information on the merging and aggregation processes which, according to the hierarchical clustering models, form the larger spiral galaxies we observe. With the aim of testing hierarchical models of galaxy formation, we have conducted an observational program which comprises H $\alpha$  imaging for both the parent and the satellite galaxies, taken from the compilation by Zaritsky et al. (1997) that contains 115 galaxies orbiting 69 primary isolated spiral galaxies. We have observed a subsample of 37 spiral and irregular galaxies taken from the compilation mentioned above. The aim of this study is to determine star formation properties of the sample galaxies. In this work we present the preliminary results of this program that we have carried out with the 1.8-m Vatican Telescope (VATT).

#### 1. Introduction

The hierarchical galaxy formation scenario implies that galaxies are formed by merging and accretion of previously collapsed systems. This framework has been successful in predicting the clustering of galaxies as a function of redshift (Silk 2000); however semianalytic models (Kauffmann, White, & Guiderdoni 1993) and numerical simulation (Klypin et al.1999; Moore et al.1999) of hierarchical galaxy formation predict satellite number counts an order of magnitude larger than the observed counts in the Local Group. The problem of the missing 2

number of observed satellites may be related to processes that limit the star formation. For example, the gas accretion by low-mass halos can be squelched by a strong photoionizing background (Somerville 2002).

The formation and evolution of satellite galaxies poses questions that are still open:

• What are the relationships between the morphologies of the parent galaxies and their satellite populations?

• How do the kinematic, dynamical, and chemical properties of the satellites compare to those of the parent galaxies?

• How do the properties of the satellites vary with projected distance from the primary galaxy?

• What are the star formation histories of the satellite galaxies? Are they related to the star formation history of the primary galaxies?

• What is the role of interactions among satellites or between satellites and their parent galaxies in the evolution of the system?

In this work we present a project aimed at addressing those questions related to the star formation in satellites of isolated spiral galaxies.

### 2. Observational Program

We have conducted an observational program which comprises broadband photometry in the optical and in the H $\alpha$  narrow-band for satellites of external giant spiral galaxies (Gutiérrez, Azzaro, & Prada 2002). In this work we present the preliminary results of an H $\alpha$  imaging program aimed at studying the star formation properties of satellites.

## 2.1. The Sample

The galaxies have been selected from the catalog by Zaritsky et al (1997). This catalog is the most complete compilation of satellite galaxies in the literature and contains 115 galaxies orbiting 69 primary isolated spiral galaxies. The limiting magnitude for this catalog is  $M_B \sim 15.5$ . The catalog includes satellites:

- that are 2.2 magnitude fainter than the parent galaxy
- with a projected distances  $\Delta D = 500$  kpc from their primaries

 $\bullet$  with a difference in recessional velocity between the parents and the satellites of  $\Delta v=500\ kms^{-1}$ 

Gutiérrez et al (2002) have observed a subsample of 60 satellites taken from the catalog mentioned above. This sample includes most of the objects situated at decl.  $\geq$  - 22. The satellites span a large range of morphologies and surface brightness profiles. Hubble types vary from elliptical to irregular with 35 objects were classified as spirals. For each galaxy the effective radius, the scale radius and the B/D ratio are tabulated. The broadband photometry will be published in a forthcoming paper (Gutiérrez et al., in preparation).

# **2.2.** $H\alpha$ Imaging

We have obtained  $H\alpha + [N II]$  emission-line images for a subsample of 37 spiral and irregular satellite galaxies. The narrow-band images of the sample were acquired during three runs in December 2001, May 2002, and December 2002



Figure 1. Left panel: R-band image. NGC 2718a is on left-lower corner and NGC 2718b is on the opposite corner. Right panel: continuum subtracted  $H\alpha + [N II]$  image. In both panels the field of view is 2.0 x 1.6.North is up and east left.

with the 1.8-m Vatican Advanced Technology Telescope (VATT) at the Mt. Graham International Observatory. A back illuminated 2048 x 2048 Loral CCD was used as the detector at the aplanatic Gregorian focus, f/9. It yielded a field of view was 6.4 x 6.4 with an image scale of 0.4 pixel<sup>-1</sup> after 2 x 2 pixel binning. The seeing varied between 0.9 - 1.8 with a typical value of 1.2.

For each galaxy we have obtained 3 x 1800 second narrow-band images using a set of interference filters with 70 Å widths that isolate the spectral region characterized by the redshifted H $\alpha$  and [N II]  $\lambda$ 6548,6583 Å emission lines. To subtract the stellar continuum we subtracted a 1200-second R-band image, suitably scaled, from the narrow-band image. The mean scale factor for the continuum image was estimated by comparing the intensity of the field stars in the two bandpasses. The data from the two first runs have been reduced. The images will be flux-calibrated using the spectrophometric standard stars observed during the different runs.

#### 3. Interaction Among Satellites

Gutierrez et al. (2002) pointed out two cases of interaction in pairs of satellites. In Figures 1 and 2 we show these cases.

NGC 2718a and NGC 2718b. In the *R*-band image (Figure 1, right panel), a bridge connecting the satellites is evident. In the H $\alpha$ +[N II] image (Figure 1, left panel), line emission is only observed in NGC 2718a, with no evidence for emission along the bridge.

NGC 4541b and NGC 4541e. Two tidal tails emerging from NGC 4541b are seen in the *R*-band image (Figure 2, right panel). One tail is pointing to NGC 4541e and the other one is in the opposite direction. In the H $\alpha$ +[N II] image (Figure 2, left panel) there is no line emission seen in NGC 4541b.

In addition, our preliminary results indicate that interacting satellite galaxies exhibit a higher level of star formation than satellites with no signs of interaction. Our work extends to satellite galaxies the idea - established previously for



Figure 2. Left panel: R-band image. NGC 4541a is the spiral on the right-upper corner. NGC 4541e is in the middle and NGC 4541b is on the left-lower corner. Right panel: continuum subtracted  $H\alpha + [N \ II]$  image. In both panels the field of view is 4.0 x 3.2. North is up and east left.

luminous galaxies (Kennicutt et al. 1987)- that interactions trigger star formation. On the other side it would be interesting to search for starburst activity in the parent galaxies produced by minor mergers as suggested by numerical simulations (Mihos & Hernquist 1994).

## References

Gutierrez, C.M., Azzaro, M., & Prada, F., 2002, ApJS, 141, 61

- Kauffmann, G., White, S.D.M., & Guiderdoni, B., 1993, MNRAS, 264, 201
- Kennicutt, R.C., Roettiger, K. A., Keel, W.C., van der Hulst, J.M., & Hummel, E., 1987, AJ, 93, 1011

Klypin, A., Kravtsov, A.V., Valenzuela, O., & Prada, F., 1999, ApJ, 522, 82

Mihos, J.C.& Hernquist, L. 1994, ApJ, 425, L13

- Moore, B., Ghigna, S., Governato, F., Lake, G., Quinn, T., Stadel, J., & Tozzi, P., 1999, ApJ, 524, L19
- Silk, J., 2000, PASP, 112, 1003
- Somerville, R.S., 2002, ApJ, 572, L23

Zaritsky, D., Smith, R., Frenk, C., & White, S. D. M. 1997, ApJ, 478, 39