AA DOR — AN ECLIPSING SUBDWARF – BROWN DWARF BINARY

 $T. Rauch^{1,2}$

AA Dor is an eclipsing, close, post commonenvelope binary (PCEB) consisting of a sdOB primary star and an unseen secondary with an extraordinary small mass ($M_2 \approx 0.066 \rm M_{\odot}$) – formally a brown dwarf (see Rauch 2004 for details). In a spectral analysis of AA Dor, Rauch (2000) determined $T_{\rm eff} = 42 \, \rm kK$ and $\log g = 5.2$ (cgs). The determination of the components' masses by comparison of these results with evolutionary models of Driebe et al. (1998) shows a discrepancy to masses derived from radial-velocity and the eclipse curves (Hilditch et al. 2003) – $\log g = 5.5$ would be necessary in order to achieve an intersection at $M_1 \approx 0.330 \rm M_{\odot}$.

Possible reasons for this discrepancy may be too optimistic error ranges in Rauch (2000) or in the analysis of light curve and radial-velocity curve, or that the evolutionary models of Driebe et al. (1998) are not appropriate in the case of AA Dor since these are post-RGB models for non-CE stars.

Since the decrement of the hydrogen Balmer series is a sensitive indicator for $\log g$, 107 high-resolution échelle spectra with short exposure times (180 sec) have been taken in Jan 2001 with UVES (UV-visual échelle spectrograph) attached to the ESO VLT. Additional medium-resolution longslit spectra have been taken at the 2.3m telescope at SSO in Sept 2003 with the DBS (double beam spectrograph). However, the analysis of both, the UVES spectra (Rauch & Werner 2003) and the DBS spectra, shows that a $\log g$ higher than 5.2 results in a worse fit to the observation.

Since the secondary is heated by irradiation of the primary up to $\approx 20\,\mathrm{kK}$, one can expect a weak H β emission in the UVES spectra. However, we do not find any signature of the secondary. The emission in the line core of H β (Fig. 1) comes clearly from the primary, its phase dependence is likely due to an irradiation effect of the heated secondary on the primary which increases its $T_{\rm eff}$ by $\approx 7\,\mathrm{kK}$, i.e. the primary – taken as an isolated star – would have only $T_{\rm eff} \approx 35\,\mathrm{kK}$, resulting in a $\approx 10\%$ smaller mass. A phase-dependent spectral analysis is presently performed in order to investigate on this effect. It ap-




Fig. 1. Section of the UVES spectra around H β compared with synthetic spectra. A weak H β emission is used to represent the secondary's radiation.

pears possible that this is one of the main reasons for the disagreement in the mass-radius relation described above.

REFERENCES

Driebe, T., Schönberner, D., Blöcker, T., & Herwig, F. 1998, A&A, 339, 129

Hilditch, R.W., Kilkenny, D., & Lynas-Gray, A.E., Hill, G. 2003, MNRAS, 344, 644

Rauch, T. 2000, A&A, 356, 665

Rauch, T. 2004, Ap&SS, in press

Rauch, T., & Werner, K. 2003, A&A, 400, 271

¹Dr.-Remeis-Sternwarte, Sternwartstraße 7, D-96049 Bamberg, Germany (Thomas.Rauch@sternwarte.uni-erlangen.de).

²Institut für Astronomie und Astrophysik, Sand 1, D-72076 Tübingen, Germany.