To appear in the

Proceedings of the 4th Cologne-Bonn-Zermatt-Symposium

"The Dense Interstellar Medium in Galaxies"

eds. S. Pfalzner, C. Kramer, C. Straubmeier, & A. Heithausen (Springer: Berlin)

Perspective from a Younger Generation – The Astro-Spectroscopy of Gisbert Winnewisser

Karl M. Menten

Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany kmenten@mpifr-bonn.mpg.de

Gisbert Winnewisser's astronomical career was practically coextensive with the whole development of molecular radio astronomy. Here I would like to pick out a few of his many contributions, which I, personally, find particularly interesting and put them in the context of newer results.

1 Beginnings – When Molecular Radio Astronomy was still Radio Astronomy

1.1 Introduction

The late 1960s and early 70s were an era of great excitement in all branches of radio astronomy. In spectral line research, 12 years after the detection of the first line, the 21 cm HI hyperfine structure (hfs) transition (Ewen & Purcell 1951), finally the first molecule was found, the hydroxyl (OH) radical (Weinreb et al. 1963).

The OH Λ -doublet transitions, as well as many other lines at m-, cm-and(!) mm-wavelengths from a variety of molecules, had been identified as possible astronomical targets in a visionary article by Charles Townes $(1957)^1$.

Shortly after the detection of OH, Hoglund & Mezger (1965) discovered the first radio recombination line, H109 α at 5009 MHz. *Molecular* radio astronomy started in earnest in 1968/9, with the discoveries of ammonia (NH₃; Cheung et al. 1968) and water vapor (H₂O; Cheung et al. 1969) by a group around Townes², as well as the first complex organic species, formaldehyde (H₂CO; Snyder et al. 1969), breaking loose a real gold rush that has yet to stop. Even today, approximately 2 – 4 "new" interstellar or circumstellar³ molecules are found each year, totaling 126 species (not counting isotopomers)⁴.

¹ see also Shklovsky 1952.

² Both lines were on Townes' (1957) list!

³ Circumstellar molecules exist in the envelopes of evolved giant stars. Many species are detected there that are also found in interstellar clouds, whereas some others, have so far *only* been detected in a circumstellar environment.

 $^{^4}$ See http://www.ph1.uni-koeln.de/vorhersagen/molecules/main_molecules.html

2

The early history of "molecule-hunting", interesting us here, is well documented by the review by Snyder (1972; "Molecules in Space"), which gives a very useful introduction to molecular radio astronomy, that is still very much readable today. In the same volume, the review following Snyder's is by Winnewisser, Winnewisser, & Winnewisser and presents an equally excellent overview of the laboratory side of "Millimetre Wave Spectroscopy".

Name Species Transition Frequency (MHz) Reference ОН $^{2}\Pi_{\frac{3}{6}}, J = \frac{3}{2}$ Hydroxyl 1665.4018(2)^a Weinreb et al. 1963 NH_3 Cheung et al. 1968 Ammonia $(1, \bar{1})$ 23694.4955(1)^b ${\rm H_2CO}$ 4829.660(1)^b Formaldehyde $1_{11} - 1_{10}$ Snyder et al. 1969 Water H_2O $6_{16}-5_{23}$ 22235.08(2)b Cheung et al. 1969 Methyl alcohol CH_3OH $1^{+} - 1^{-}$ 834.267(2) Ball et al. 1970 $1_{11} - 1_{10}$ Formic acid **HCOOH** 1638.804(1) Zuckerman et al. 1971 $2_{11} - 2_{12}$ $1_{11} - 1_{11}$ 4916.319(3) GWi & Churchwell 1975 Formamide NH_2CHO 1539.58(2) Palmer et al. 1971 $2_{11} - 2_{12}$ 4618.55(2) Rubin et al. 1971 HC_3N Cyanoacetylene 9098.6537(3)° Turner 1971 $\mathrm{H}^{13}\mathrm{C}_{3}\mathrm{N}$ 1 - 0 $\sim 9060^{\circ}$ Gardner & GWi 1975a $H^{12+13}C_3N$ $\sim 9000^{\rm d}$ 1 - 0Churchwell et al. 1977 CH₃CHO Acetaldehyde $1_{10} - 1_{11}$ 1065.075(5)Ball et al. 1971 ${}^{2}\Pi_{\frac{1}{2}}, J = \frac{1}{2}$ Methylidine CH3263.788(10)e Rydbeck et al. 1973 $2_{11}^{2} - 2_{12}$ $1_{10} - 1_{11}$ Thioformaldehyde 3139.38(3) Sinclair et al. 1973 H_2CS $\overline{\mathrm{CH}_{2}\mathrm{NH}}$ 5290.31(4) Methanimine Godfrev et al. 1973 $\mathrm{CH_2NH_2}$ Fourikis et al. 1974 Methylamine $2_{02}-1_{10}$ $8776.2(1)^{1}$ Vinyl cyanide H_2CCHCN $2_{11}-2_{12}$ 1371.8262(1) Gardner & GWi 1975b $\mathrm{HCOOCH_3}\text{-}A\ 1_{10}-1_{11}$ Methyl formate 1610.2445(7) Brown et al. 1975 ${
m HCOOCH_3}\text{-}E \ 1_{10} - 1_{11}$ 1610.91(10) Churchwell & GWi 1975 $2_{02}-1_{11}$ Dimethyl ether $(CH_3)_2O$ 9119.668(15) GWi~&~Gardner~1976Cyanodiacetylene $\dot{H}C_5\dot{N}$ 4 - 310650.650(4) Avery et al. 1976 2 - 15325.328(2) Gardner & GWi 1978

Table 1. The First Interstellar "Radio" Molecules

Only molecules first detected at cm-wavelengths are listed. For early detections in the mm-range, see Snyder 1972. All molecules were first detected in the transitions listed, except for dimethyl either, which was discovered in its $6_{06} - 5_{15}$ line at 90938.10(3) MHz by Snyder et al. 1974. No detections for isotopomers are listed, except for HC₃N.

Contributions coauthored by GWi are in *italics*. In most cases these contain essential laboratory spectroscopy work by GWi.

^a Apart from the F=1-1 line listed, also the 2-2 line at 1667.3590(2) MHz was reported in the detection paper. ^b Line with hfs, intensity-weighted frequency is given ^c Line with hfs, average of lowest and highest hfs line frequency is given ^d Various lines with differently placed ¹³C between 8815 and 9100 MHz. ^e Apart from the F=0-1 line listed, also the F=1-1 and F=1-0 lines at 3335.475(10) and 3349.185(10) MHz, resp., were reported in the detection paper (frequencies from Rydbeck et al. 1974). ^f Frequency from Fourikis et al. 1974.

As shown in Table 1^5 , many of the early molecular detections were made at low radio frequencies (and toward Sagittarius B2). A reason for the propensity to find *such* lines *there* is given in $\S 1.2$

Almost concurrently with cm-wavelength molecular radio astronomy, shorter wavelength exploration started. Driven by new high-frequency receiver technology and, generally, higher line intensities and a much higher "line density" (i.e. lines per GHz) the "millimeter explosion" started with the discoveries of CO by R. Wilson et al. and HCO⁺ ("X-ogen") by Buhl & Snyder, both in 1970 (the latter published in 1971), and continues through today, having reached the submillimeter range in the 1980s.

Once the millimeter window was opened, (most) people spontaneously lost interest in the cm-range because:

- Most molecules have the bulk or the entirety of their rotational spectrum at (sub)millimeter wavelengths
- Almost all "simple" (di- or triatomic), light molecules have (sub)mm or far-IR (FIR) rotational spectra
- Many of the complex molecules were (for a long time) only found in Sgr B2

Interest remained high in special cm-wavelength lines, including the NH₃ inversion transitions near 24 GHz, a number of prominent maser lines from OH (hfs, 1.7 GHz and higher frequencies), $\rm H_2O$ (22.2 GHz), $\rm CH_3OH$ (25, 12.2, and 6.7 GHz), as well as the CH hfs lines (3.3 GHz). All of these, except for CH₃OH, have all or the bulk of their *rotational* spectrum at submillimeter and far-infrared wavelengths.

1.2 Sagittarius B2

Considering the mm-wavelength data coming in, soon following the radio observations, it became clear that the emission from many low frequency, low-J lines of complex molecules toward Sgr B2 must be (weakly) inverted. This can be understood (at least for linear or near-linear species) from the fact that "when selection rules for collisional transitions in linear molecules are such that dipole collisions do not dominate, the J=1-0 transitions may have population inversion over a wide range of physical conditions (see, for example Goldsmith 1972)" (quoted from Morris et al. 1976). This is borne out by their statistical equilibrium calculations.

However, direct proof for this assertion came only very recently, from cm-wave interferometry of the CH₃CHO $1_{11}-1_{10}~(1065~\mathrm{MHz})$ and HC₃N 1-0

⁵ The frequencies taken from the $_{
m JPL}$ catalog (Pickare 1998; $_{
m ett}$ $_{
m et}$ al. see http://spec.jpl.nasa.gov/) or the Cologne Database for Molecular Spectroscopy (Müller et al.http://www.ph1.uni-koeln.de/vorhersagen/molecules/main_molecules.html). Numbers in parentheses are uncertainties in the last significant digit(s).

4

(9100 MHz) lines with the Giant Metre Wave Radio Telescope (GMRT; Chengalur & Kanekar 2003) and the Australia Telescope Compact Array (ATCA; Hunt et al. 1999), respectively. The molecular emission is extended over a $2' \times 4'$ region and shows a one-to-one correspondence with the continuum emission, proving that it is inverted. For the CH₃CHO line a typical optical depth of 0.035 is derived.

Apart from the mentioned lines, Gardner & Winnewisser (1975b) concluded that the vinyl cyanide line listed in Table 1 is inverted, as did Winnewisser & Gardner (1976) for the dimethyl ether line and Sinclair et al. (1973) for the thioformaldehyde line. Broten et al. (1976) inferred the same for the 2.663 GHz J=1-0 transition of HC₅N.

Once the first complex molecules were found in Sgr B2, this source turned into a Bonanza! The described inversion mechanism together with a relatively strong, extended continuum background "boosted" many otherwise undetectable lines into "observability". As a consequence, the vast majority of interstellar molecules was and still is found in Sgr B2, many of the most complex ones *only* there. All of the molecules listed in Table 1 are found in this source.

To put GWi's Sgr B2 effort into perspective: Several of the lines measured by Gardner and him confirmed "shaky" detections, based on noisy single line observations. The vinyl cyanide work presents the first detection of this molecule in space. The study on different ¹³C isotopes of HC₃N by Churchwell et al. (1977) indicated chemical fractionation and put a cautionary remark on attempts at determining the carbon isotope ratio in the Galactic center.

1.3 Extended and Compact Organic Molecular Emission in Sgr B2

Using the (low) rotation temperatures and column densities determined for many of the species in Table 1 by Cummins et al. (1986) based on their 2-3 mm wavelength line survey it is completely clear that *all* of the lines in that table must be inverted, which greatly helped their discovery at the time.

Why are these "old" Sgr B2 results still highly relevant today? In large beam observations, such as those of Cummins of al. (angular resolution 90"–180") one observes very low rotation temperatures for many species. However, interferometer observations by Snyder and collaborators with resolutions < 5" indicate $T_{\rm rot} \sim 200$ K for the same species in a very compact region, dubbed the "Large Molecule Heimat" (LMH) by Snyder et al. (1994). This source, which is in the immediate vicinity of the Sgr B2(N) continuum source, is the most prolific source of complex molecules known.

Perhaps most interestingly, in recent years arcsecond resolution interferometry with the BIMA Array has resulted in the detection and imaging of increasingly complex organic species, such as CH₂CHCN, HCOOCH₃, and CH₃CH₂CN; all Miao et al. 1995, 1997), NH₂CHO, HNCO, and HCOOCH₃ (Kuan & Snyder 1996), CH₃COOH (acetic acid; Mehringer et al. 1997; Remijan et al. 2002), HCOOH (formic acid; Liu et al. 2001), and (CH₃)₂CO (ace-

tone; Snyder et al. 2002); recent single dish detections include CH₂OHCHO (glycolaldehyde, the first sugar; Hollis et al. 2000; 2001); HOCH₂CH₂OH (ethylene glycol: interstellar "antifreeze"; Hollis et al. 2002), and CH₂CHOH (vinyl alcohol: Turner & Apponi 2001).

The elevated complex molecule abundances in a confined, hot, region, such as the LMH, may be explainable by the evaporation of icy grain mantles, during which molecules that have grown there to complexity by continual hydrogenation over a long time are suddenly released into the gas phase by the ignition of a high-mass embedded star⁶. However, such an event cannot possibly explain (for luminosity reasons alone) the unique large-scale distribution of complex molecules in (and maybe around) Sgr B2, which is the most prolific source of organic molecules anywhere and, so far, has only been probed by the low frequency lines described above. If these molecules come off grain mantles, other processes, such as shocks created, e.g., by cloud-cloud collisions must be invoked. There actually is evidence for the latter in Sgr B2 (Mehringer & Menten 1997; Sato et al. 2000).

1.4 A Giant Organic Cloud around the Galactic Center?

There is actually evidence for a giant repository of organic molecules beyond and including Sgr B2, coextensive with the Central Molecular Zone (CMZ, see e.g. Morris & Serabyn 1996), i.e., stretching from l = +1.6 to -1.1 in a $\sim \pm 0.3$ wide band around the Galactic center.

The first evidence for extended organic material in the CMZ came from widespread 4.8 GHz $\rm H_2CO$ absorption (Scoville et al. 1974). Given the ubiquity of formaldehyde in molecular clouds, one might dismiss this "as nothing special". Methanol (CH₃OH) on the other hand has usually quite low abundance and is difficult to detect outside hot, dense cloud cores. Nevertheless, Gottlieb et al. (1979) find the emission in the 834 MHz (1⁺ – 1⁻) line (see Table 1) in emission and extended relative to their 40'(!) beam, concluding it is inverted.

Furthermore, mapping of the HNCO $5_{05} - 4_{04}$ transition (made serendipitously simultaneously with a C¹⁸O survey), shows that the emission in this line is extending continuously from $l = -0^{\circ}2$ to $+1^{\circ}7$ (Dahmen et al. 1997). The possible existence of such a huge mass of organic material in our Galactic center is extremely exciting and its extent, chemistry, and excitation should be studied with suitable tracers.

1.5 Randbemerkung: TMC-1

After Sgr B2, GWi and his collaborators turned their attention to complex molecules in cold dark clouds. In the abstract of a paper by Churchwell,

 $^{^6}$ A review of complex molecule formation involving grain mantle processes has recently been presented by Ehrenfreund & Charnley (2000).

Winnewisser, & Walmsley (1978) titled fetchingly "Molecular observations of a possible proto-solar nebula in a dark cloud in Taurus", these authors reported "a small molecular condensation near the south eastern edge of Heiles' cloud 2" in which they find "strong emission from the J=9-8 transition of $\mathrm{HC}_5\mathrm{N}$ and the J=1-0 transition of $\mathrm{HC}_3\mathrm{N}$ ", and go on to "refer to this small cloud as the Taurus Molecular Cloud 1, or TMC 1". It turned out that they started a vogue: As of this writing (2003 October 25) the Astronomical Data System⁷ lists a total of 267 abstracts containing TMC 1 in their text.

Churchwell et al. were by no means the first to notice this spectacular source, which lies in the heart of the Taurus molecular cloud complex, and is well known for its high abundance of linear carbon chains, in particular the cyanopolyynes, the longest of which, $HC_{11}N$ has *only* been found there (Bell et al. 1997). Citing them: "Morris ⁸ et al. (1976) first drew attention to this region by their detection of the J = 5 - 4 transition of HC_3N ; this was followed by the detection of the J = 4 - 3 transition of HC_5N by MacLeod et al. (1978)[sicl] and the J = 9 - 8 transition of HC_7N by Kroto et al. (1977)".

A great amount of work on TMC 1 was done by the group at the Herzberg Institute of Astrophysics, GWi's onetime home, and at the Nobeyama Radio Observatory (NRO), whose 45 m telescope is ideally suited for the 22 – 50 GHz range in which many heavy species at TMC 1's cold (10 K) temperature have their strongest emission. In a single paper, Kaifu et al. (1987) reported seven(!), strong(!) unidentified lines, which, aided by laboratory work of Saito et al. (1987) and Yamamoto et al. (1987), led to the first identifications of CCS and CCCS.

The NRO's TMC 1 work culminated with a complete 8800 – 50000 MHz line survey, described by Ohishi & Kaifu (1998), but unfortunately not (yet) published in its entirety, which contains a total of 404 lines from 38 molecules (not counting isotopomers). Eleven of the molecules were newly identified in the course of the survey. It is a great testimony to the laboratory spectroscopists involved that only a single line remains unidentified!

2 Light Intermezzo – Interstellar Hydrides

When I was a student, an article by G. Winnewisser, Aliev & Yamada (1982) drew my attention. To quote from its abstract: "The great progress molecular astronomy has made in the past decade has focused attention to one special class of molecules: the hydrides. The presently known spectra of the hydrides are reviewed, both interstellar and laboratory. No interstellar metal hydrides

⁷ http://adsabs.harvard.edu/

⁸ Mark Morris tells me that the first person to point out TMC 1 to him was Nick Scoville.

⁹ That paper actually appeared in 1979.

are presently known. A large number of important hydrides exhibit spectra only in the submillimetre and far infrared region. Further technical advances in this area are expected to yield new interstellar detections."

2.1 Metal-bearing Molecules in the Interstellar Medium

Observations of species containing metals, as well as other refractory species such as SiO, deliver important information on depletion into dust grains. Moreover, when released either from sputtering of the grains or their mantles by violent dynamics, such as protostellar outflows, they can be used as probes of these phenomena (see, e.g. Bachiller 1996).

Today the third sentence quoted above, "No interstellar metal hydrides ..." is, unfortunately, still true, if we use the strict (chemistry) meaning of "metal" (e.g. Na, K, Mg, Al, Fe,...). Four diatomic, non H-bearing metal compounds (NaCl, KCl, AlCl, and AlF and three triatomic ones (MgNC, MGCN, NaCN, and AlNC) have been detected in the dense envelope of the carbon star IRC+10216 (Cernicharo & Guélin 1987; Guélin et al. 1993; Kawaguchi et al. 1993; Ziurys et al. 1995; Turner et al. 1994; Guélin et al. 1996; Ziurys et al. 2002). MgNC, NaCN, AlF, and NaCl were also found, recently, toward the proto-planetary nebula (PN) CRL 2688 (Highberger et al. 2001, 2003; Ziurys et al. 2002; see Petrie 1999 for a discussion of other candidate species for detection). Still, only one metal-containing molecule, FeO, has been detected in the interstellar medium, toward Sgr B2 (Walmsley et al. 2002).

As to metal hydrides: Radio and (sub)millimeter wavelength searches for interstellar NaH (Plambeck & Erickson 1982) and LiH¹⁰ have so far been unsuccessful, as was a search for MgH in IRC+10216's envelope (Avery et al. 1994). Optical wavelength absorption from CaH, AlH, FeH, and MgH has been found in cool stars and the Sun, MgH even in the integrated light of external galaxies (see, e.g., Spinrad & Wood 1965)

2.2 Light Hydrides – Present Status

The history of interstellar hydrides begins with the detection by Dunham (1937) of three optical absorption lines at 3957.7, 4232.6, and 4300.3 Å from diffuse interstellar clouds, which subsequently were found to arise from CH (4300.3 Å; McKellar 1940; suggested by Swings & Rosenfeld 1937) and CH⁺ (the former two; Douglas & Herzberg 1941).

It has been thought for a long time that hydrides might play an important rôle in cooling the highest density regions of interstellar clouds. Hydrides have, firstly, large level spacings, allowing them to emit a relatively large

The ground state J=1-0 transition of LiH is near 444 GHz, an atmospherically very unfavorable frequency. A tentative detection of this line was made in the redshifted (z=0.68) dense absorbing cloud toward the B0218+357 gravitational lens system (Combes & Wiklind 1998).

amount of energy per quantum. Secondly, their large dipole moments require high critical densities for their excitation. Thus, their rotational levels are only excited in high density regions with $n > 10^6$ cm⁻³, making them effective coolants in this regime, where even rare CO isotopes are so optically thick that CO cooling becomes negligible compared to that of (the hydride) H₂O.

This regime was excluded in the classical study by Goldsmith and Langer (1978) on "Molecular Cooling and Thermal Balance in Dense Molecular Clouds", but was re-addressed by Neufeld et al. (1995), who found that for elevated temperatures, $T=100~\rm K,\,H_2O$ cooling dominates and the hydrides' contribution is comparable to that of all other molecules combined, i.e. ≈ 30 % of the former.

Recent detections of hydride rotational transitions were made with the Caltech Submillimeter Observatory (CSO), namely $\mathrm{H_3O^+}$ [$J_K=1_1^--2_1^+,3_2^+-2_2^-,3_0^+-2_0^-/307,$ 365, 396 GHz, Wootten et al. 1991; Phillips et al. 1992], SiH ($^2H_{\frac{1}{2}},J=\frac{3}{2}-\frac{1}{2}/625,$ 628 GHz; Schilke et al. 2001)¹¹, HCl (J=1-0/626 GHz); Blake et al. 1985), $\mathrm{H_2O}$, $\mathrm{H_2^{18}O}$ ($\mathrm{I_{10}-I_{01}/557},$ 548 GHz; Melnick et al. 2000), HDO (the $\mathrm{I_{01}-0_{00}/465}$ GHz and $\mathrm{I_{11}-0_{00}/894}$ GHz ground-state transitions; Schulz et al. 1991, Pardo et al. 2001)¹², and NH₂ ($J=\frac{3}{2}-\frac{3}{2},\frac{1}{2}-\frac{1}{2},\frac{3}{2}-\frac{1}{2}/462,$ 459, 461 GHz); van Dishoeck et al. 1993). $\mathrm{H_2D^+}$ ($\mathrm{I_{10}-I_{11}/372}$ GHz), the deuterated isotope of the key $\mathrm{H_3^+}$ molecule was discovered with the James-Clerck-Maxwell telescope (Stark et al. 1999)¹³. Observations with the Long Wavelength Spectrometer aboard the Infrared Space Observatory led to the discoveries of HF (J=2-1/2.47 THz; Neufeld et al. 1997), HD (J=1-0/2.68 THz and 2-1/5.36 THz; Polehampton et al. 2002), plus various $\mathrm{H_2O}$, OH, and CH rotational and some $\mathrm{H_3O^+}$ inversion transitions (many toward Sgr B2; Goicoechea & Cernicharo 2001 and these proceedings).

A preliminary conclusion is that (for the only two sources observed, Orion-KL and Sgr B2) the measured HCl and HF abundances indicate that only a few percent of all chlorine is in these molecules, suggesting significant depletion. If this should hold true for other hydrides as well, their cooling contribution could be much smaller than discussed above. More sensitive observations of more transitions toward more lines of sight are needed. These will be possible with the Herschel Space Observatory¹⁴ and the Stratospheric Observatory For Infrared Astronomy (SOFIA¹⁵).

¹¹ Tentative detection

 $^{^{12}}$ A number of higher excitation cm and mm-wavelength $\mathrm{H}_2\mathrm{O},\,\mathrm{H}_2^{18}\mathrm{O},\,\mathrm{and}$ HDO transitions have also been found.

¹³ H₃⁺ itself was finally detected, after many arduous attempts, in absorption at near-infrared wavelengths by Geballe & Oka (1996), benefitting from a monumental body of laboratory work, mainly gathered by Oka and collaborators (see these proceedings).

 $^{^{14}\ \}mathrm{http://astro.estec.esa.nl/SA-general/Projects/First/first.html}$

¹⁵ http://sofia.arc.nasa.gov/

Under GWi's direction the KOSMA Spectroscopy Group has produced definitive submillimeter spectroscopy of NH₂, C, ¹³C, NH, SH, SD, and CF.

3 The quest for high spectroscopic precision – "Deutsche Gründlichkeit" gone loose or urgent necessity?

Over the years, the KOSMA lab, with GWi as the driving force, has produced "definitive" spectroscopy (i.e. frequencies to a few times 10 kHz rms up to 1 THz or higher) for a large number of astrophysically important molecules, including the first useful data on many rare isotopomers and various vibrationally excited states.

These are 16 : HCS⁺, HC¹⁴⁺¹⁵N, $^{12+13}$ CO, C¹⁷⁺¹⁸O, H₂¹²⁺¹³C¹⁶⁺¹⁸O, ground-state + vibrationally excited H¹²⁺¹³C₃¹⁴⁺¹⁵N, C₂H₅OCH3, CCS, CCC, ground-state + vibrationally excited HNC, CH2NH, CH₃CCH, CCH, $^{32+33}$ SO₂, S¹⁶⁺¹⁷O₂, C¹⁴⁺¹⁵N, CH₂CCHCN, CCCO, CH₃C₄H, NH₂D (hfs), HC₅N, HNCO, and the seven simple species with submillimeter spectra listed above¹⁷.

The first astronomical detections of $\rm C^{13}CH$ and $\rm HC^{15}N$ were made by the Cologne group with the KOSMA telescope (Saleck et al. 1992, 1993) and of $\rm ^{13}C^{17}O$ with the Swedish-ESO-Submillimeter Telescope (Bensch et al. 2001).

Among others, the Cologne spectroscopy effort is characterized by, first, comprehensive presentations of the *complete* spectrum of a given species, in general up to higher (several THz) frequencies than many other listings, and, second, significantly higher accuracy data for many "common" molecules, such as CO and its isotopomers, than available before.

Here we argue, using two examples, that both high frequency accuracy and comprehensiveness are necessary for progress in various different areas.

3.1 Accuracy

Ambipolar diffusion – ion/neutral slip – is an important mechanism in the star-formation process by which magnetic fields initially supporting clouds against collapse slip relative to the neutral gas, leading to the formation of dense cloud cores (see, e.g. Shu et al. 1987). Theory predicts velocity differences < 1 km s⁻¹, corresponding to 1 MHz at 300 GHz, between ionic and neutral species (Roberge et al. 1995). In particular, at submillimeter wavelengths present frequency uncertainties for many potentially suitable lines are much too high to draw any reliable conclusions.

For references, see http://www.ph1.uni-koeln.de/vorhersagen/catalog/catdir.html
Note the tenacity with which cyanoacetylene was pursued!; cf. §1

3.2 Comprehensiveness and Line Confusion

In a recent paper, Kuan et al. (2003) claim to have detected the simplest amino acid, glycine (H₂NCH₂COOH) in three warm, dense interstellar clouds, including Sgr B2. If true, this would be the conclusion of a long quest (see, e.g., Snyder et al. 1983; Berulis, Winnewisser, Krasnov, & Sorochenko 1985; Combes et al. 1996; Ceccarelli et al. 2000).

Any such "identification" must rely on a large number of lines. Even so (or maybe because of this) the confusion problem is a formidable one. Inspecting any figure presented by Kuan et al. we find comparably many glycine as unidentified ("U") lines. For a wider view, see the 218 – 263 GHz line survey by Nummelin et al. (1998, 2000), which, for Sgr B2(N), is at or near the confusion limit throughout that band, even at their relatively moderate sensitivity (compared to the interferometric data by Snyder and his collaborators).

In the high density ($n \sim 10^7~{\rm cm}^{-3}$) Sgr B2 "Large Molecule Heimat" (see §1.2) level populations are in Local Thermodynamic Equilibrium characterized by rotation temperatures, $T_{\rm rot}$, around 200 K (Liu et al. 2001). At this temperature, the lower vibrationally excited states of a large number of complex molecules are populated, giving rise to a plethora of emission lines, probably with comparable intensities to the putative glycine lines. The width of the LMH's lines, typically $\sim 6-10~{\rm km~s}^{-1}$, adds additional complication. Given the gigantic partition functions, many of the lines are weak.

In my opinion, it is completely impossible to conclusively identify *very* complex molecules in a "line jungle", such the one found in the Sgr B2 Large Molecule Heimat¹⁸, if one has not a much better knowledge of, a.o., vibrationally excited lines of many more species than is available today. This means that much more (tedious and little-appreciated) lab work is needed to make progress here: A big job for the Cologne group, which has already measured the spectra (although not necessarily those of the vibrationally excited states) of a number of extant molecules.

Apology

This contribution concerned itself mostly with (some aspects of) GWi's early astronomical career. Among many others, I didn't discuss such interesting and important topics as

 NH₃ hfs anomalies (Stutzki et al. 1982, 1984; Stutzki & Winnewisser 1985a,b,c)

¹⁸ High spatial resolution interferometry as employed by L. Snyder and collaborators helps significantly in situations as complicated as the LMH: Individual molecules or groups of molecules in general have slightly different spatial distributions, so that imaging adds an additional dimension to a confident identification.

- his pushing the Cologne Acousto-Optical Spectrometers and receivers to become world-class, frontline instrumentation (see, e.g., Schieder et al. 1989; Klumb et al. 1995; Graf et al. 1998, 2003; see also Gary Melnick's SWAS contribution in these proceedings)
- any of all the nice KOSMA/Gornergrat results
- any of his non-astronomical spectroscopy; little pre-Cologne spectroscopy.

Acknowledgement: I would like to thank Malcolm Walmsley and Lew Snyder for their encouragement.

References

- Avery, L. W., Bell, M. B., Cunningham, C. T., Feldman, P. A., Hayward, R. H., McLeod, J. M., Matthews, H. E., Wade, J. D.: ApJ, 426, 737 (1994)
- Avery, L. W., Broten, N. W., MacLeod, J. M., Oka, T., Kroto, H. W.: ApJ, 205, L173 (1976)
- 3. Bachiller, R.: ARA& 34, 111 (1996)
- 4. Ball, J. A., Gottlieb, C. A., Lilley, A. E., Radford, H. E.: ApJ, 162, L203 (1970)
- Ball, J. A., Gottlieb, C. A., Lilley, A. E., Radford, H. E.: IAU Circ. No. 2350 (1971)
- Bell, M. B., Feldman, P. A., Travers, M. J., McCarthy, M. C., Gottlieb, C. A., Thaddeus, P.: ApJ, 483, L61 (1997)
- Bensch, F., Pak, I., Wouterloot, J. G. A., Klapper, G., Winnewisser, G.: ApJ, 562, L185 (2001)
- 8. Berulis, I. I., Winnewisser, G., Krasnov, V. V., Sorochenko, R. L.: Soviet Astronomy Letters, 11, 251 (1985)
- 9. Blake, G. A., Keene, J., Phillips, T. G.: ApJ, 295, 501 (1985)
- Broten, N. W., MacLeod, J. M., Oka, T., Avery, L. W., Brooks, J. W., McGee, R. X., Newton, L. M.: ApJ, 209, L143 (1976)
- Brown, R. D., Crofts, J. G., Godfrey, P. D., Gardner, F. F., Robinson, B. J., Whiteoak, J. B.: ApJ, 197, L29 (1975)
- 12. Buhl, D., Snyder, L. E.: Nature, 232, 161 (1971)
- Ceccarelli, C., Loinard, L., Castets, A., Faure, A., Lefloch, B.: A&A, 362, 1122 (2000)
- 14. Cernicharo, J., Guélin, M.: A&A, 183, L10 (1987)
- 15. Chengalur, J. N., Kanekar, N.: A&A, 403, L43 (2003)
- 16. Cheung, A. C., Rank, D. M., Townes, C. H., Thornton, D. D., Welch, W. J.: Phys. Rev. Letters, 21, 1701 (1968)
- Cheung, A. C., Rank, D. M., Townes, C. H., Thornton, D. D., Welch, W. J.: Nature, 221, 626 (1969)
- 18. Churchwell, E., Walmsley, C. M., Winnewisser, G.: A&A, 54, 925 (1977)
- 19. Churchwell, E., Winnewisser, G.: A&A, 45, 229 (1975)
- 20. Churchwell, E., Winnewisser, G., Walmsley, C. M.: A&A, 67, 139 (1978)
- 21. Combes, F., Wiklind, T.: A&A, 334, L81 (1998)
- 22. Combes, F., Q-Rieu, N., Wlodarczak, G.: A&A, 308, 618 (1996)
- 23. Cummins, S. E., Thaddeus, P., Linke, R. A.: ApJS, 60, 819 (1986)
- 24. Dahmen, G. et al. 1997, A&AS, 126, 197 (1997)

- 25. Douglas, A. E., Herzberg, G.L.: ApJ, 94, 381 (1941)
- 26. Dunham, T. PASP, 49, 26 (1937)
- 27. Ehrenfreund, P., Charnley, S. B.:ARA&A, 38, 427 (2000)
- 28. Ewen, H. I., Purcell, E. M.: Nature, 168, 356 (1951)
- 29. Fourikis, N., Takagi, K., Morimoto, M.: ApJ, 191, L139 (1974)
- 30. Gardner, F. F. Winnewisser, G.: ApJ, 195, L127 (1975a)
- 31. Gardner, F. F. Winnewisser, G.: ApJ, 197, L73 (1975b)
- 32. Gardner, F. F. Winnewisser, G.: MNRAS, 185, 57P (1978)
- 33. Geballe, T. R., Oka, T.: Nature, 384, 334 (1996)
- Godfrey, P. D., Brown, R. D., Robinson, B. J., & Sinclair, M. W.: Ap. Lett., 13, 119 (1973)
- 35. Goicoechea, J. R., Cernicharo, J.: ApJ, 554, L213 (2001)
- 36. Goldsmith, P. F.: ApJ, 176, 597 (1972)
- 37. Goldsmith, P. F. Langer, W. D.: ApJ, 222, 881 (1978) eds. M. A. Gordon L. E. Snyder (New York, NY: John Wiley and Sons 1973) p 181
- Gottlieb, C. A., Ball, J. A., Gottlieb, E. W., Dickinson, D. F.: ApJ, 227, 422 (1979)
- Graf, U. U., Haas, S., Honingh, C. E., Jacobs, K., Schieder, R., Stutzki, J.: Proc. SPIE, 3357, 159 (1998)
- 40. Graf, U. U. et al.: Proc. SPIE, 4855, 322 (2003)
- 41. Guélin, M., Lucas, R., Cernicharo, J.: A&A, 280, L19 (1993)
- Highberger, J. L., Savage, C., Bieging, J. H., Ziurys, L. M.: ApJ, 562, 790 (2001)
- Highberger, J. L., Thomson, K. J., Young, P. A., Arnett, D., Ziurys, L. M.: ApJ, 593, 393 (2003)
- 44. Hoglund, B., Mezger, P. G.: Science, 150, 339 (1965)
- 45. Hollis, J. M., Lovas, F. J., Jewell, P. R.: ApJ, 540, L107 (2000)
- Hollis, J. M., Vogel, S. N., Snyder, L. E., Jewell, P. R., Lovas, F. J.: ApJ, 554, L81 (2001)
- 47. Hollis, J. M., Lovas, F. J., Jewell, P. R., Coudert, L. H.: ApJ, 571, L59 (2002)
- Hunt, M. R., Whiteoak, J. B., Cragg, D. M., White, G. L., Jones, P. A.: MNRAS, 302, 1 (1999)
- Kaifu, N., Suzuki, H., Ohishi, M., Miyaji, T., Ishikawa, S., Kasuga, T., Morimoto, M., Saito, S.: ApJ, 317, L111 (1987)
- Kawaguchi, K., Kagi, E., Hirano, T., Takano, S., Saito, S.: ApJ, 406, L39 (1993)
- Klumb, M., Frerick, J., Horn, J., Schieder, R., Winnewisser, G. F.: Proc. SPIE, 2583, 420 (1995)
- 52. Kuan, Y. Snyder, L. E.: ApJ, 470, 981 (1996)
- Kuan, Y., Charnley, S. B., Huang, H., Tseng, W., Kisiel, Z.: ApJ, 593, 848 (2003)
- 54. Liu, S., Mehringer, D. M., Snyder, L. E.: ApJ, 552, 654 (2001)
- 55. MacLeod, J. M., Avery, L. W., Broten, N. W.: ApJ., 233, 584 (1979)
- 56. McKellar, A.: PASP, 52, 187 (1940)
- 57. Mehringer, D. M., Menten, K. M.: ApJ, 474, 346 (1997)
- 58. Mehringer, D. M., Snyder, L. E., Miao, Y., Lovas, F. J.: ApJ, 480, L71 (1997)
- 59. Melnick, G. J. et al.: ApJ, 539, L87 (2000)
- 60. Miao, Y., Mehringer, D. M., Kuan, Y., Snyder, L. E.: ApJ, 445, L59 (1995)

- 61. Miao, Y. Snyder, L. E.: ApJ, 480, L67 (1997)
- 62. Morris, M., Serabyn, E.: ARA&A, 34, 645 (1996)
- 63. Morris, M., Turner, B. E., Palmer, P., Zuckerman, B.: ApJ, 205, 82 (1976)
- Müller, H. S. P., Thorwirth, S., Roth, D. A., Winnewisser, G.: 2001, A&A, 370, L49 (2001)
- 65. Neufeld, D. A., Lepp, S., Melnick, G. J.: ApJS, 100, 132 (1995)
- Neufeld, D. A., Zmuidzinas, J., Schilke, P., Phillips, T. G.: ApJ, 488, L141 (1997)
- Nummelin, A., Bergman, P., Hjalmarson, Å., Friberg, P., Irvine, W. M., Millar,
 T. J., Ohishi, M., Saito, S.: ApJS, 117, 427 (1998)
- Nummelin, A., Bergman, P., Hjalmarson, Å., Friberg, P., Irvine, W. M., Millar,
 T. J., Ohishi, M., Saito, S.: ApJS, 128, 213 (2000)
- 69. Ohishi, M. & Kaifu, N.: Faraday Discussions No. 109 (1998)
- 70. Palmer, P., Gottlieb, C. A., Rickard, L. J., Zuckerman, B.: BAAS, 3, 499 (1971)
- Pardo, J. R., Cernicharo, J., Herpin, F., Kawamura, J., Kooi, J., Phillips, T. G.: ApJ, 562, 799 (2001)
- 72. Petrie, S.: MNRAS, 302, 482 (1999)
- 73. Phillips, T. G., van Dishoeck, E. F., Keene, J.: ApJ, 399, 533 (1992)
- Pickett, H. M., Poynter, R. L., Cohen, E. A., Delitsky, M. L., Pearson, J. C., Müller, H. S. P.: J. Quant. Spectrosc. & Rad. Transfer, 60, 883 (1998).
- 75. Plambeck, R. L., Erickson, N. R.: ApJ, 262, 606 (1982)
- Polehampton, E. T., Baluteau, J.-P., Ceccarelli, C., Swinyard, B. M., Caux, E.: A&A, 388, L44 (2002)
- Remijan, A., Snyder, L. E., Liu, S.-Y., Mehringer, D., Kuan, Y.-J.: ApJ, 576, 264 (2002)
- 78. Roberge, W. G., Hanany, S., Messinger, D. W.: ApJ, 453, 238 (1995)
- Rubin, R. H., Swenson, G. W., Benson, R. C., Tigelaar, H. L., Flygare, W. H.: ApJ, 169, L39 (1971)
- 80. Rydbeck, O. E. H., Elldér, J., Irvine, W. M.: Nature, 246, 466 (1973)
- Rydbeck, O. E. H., Elldér, J., Irvine, W. M., Sume, A., Hjalmarson, Å.: A&A, 34, 479 (1974)
- Saito, S., Kawaguchi, K., Yamamoto, S., Ohishi, M., Suzuki, H., Kaifu, N.: ApJL, 317, L115 (1987)
- 83. Saleck, A. H., Eigler, K., Simon, R., Vowinkel, B., Winnewisser, G.: Astronomische Gesellschaft Abstract Series, 7, 83 (1992)
- Saleck, A. H., Simon, R., Schneider, N., Winnewisser, G.: ApJ, 414, L133 (1993)
- 85. Sato, F., Hasegawa, T., Whiteoak, J. B., Miyawaki, R.: ApJ, 535, 857 (2000)
- 86. Schieder, R., Tolls, V., Winnewisser, G.: Experimental Astronomy, 1, 101 (1989)
- 87. Schilke, P., Benford, D. J., Hunter, T. R., Lis, D. C., Phillips, T. G.: ApJS, 132, 281 (2001)
- Schulz, A., Güsten, R., Walmsley, C. M., Serabyn, E. 1991, A&A, 246, L55 (1991)
- 89. Scoville, N. Z., Solomon, P. M., Jefferts, K. B.: 1974, ApJ, 187, L63 (1974)
- 90. Shklovsky, I. S.: Astr. Zhur., 29, 144 (1952)
- 91. Shu, F. H., Adams, F. C., Lizano, S.: 1987, ARA&A, 25, 23 (1987)
- 92. Sinclair, M. W., Fourikis, N., Ribes, J. C., Robinson, B. J., Brown, R. D., & Godfrey, P. D.: Australian J. Phys., 26, 85 (1973)

- Snyder, L. E.: Molecules in Space. In: MTP International Review of Science vol 3, Spectroscopy, ed by D. A. Ramsay (Butterworths, London 1972) pp 193–240
- Snyder, L. E., Buhl, D., Zuckerman, B. Palmer, P.: Phys. Rev. Lett., 22, 679 (1969)
- Snyder, L. E., Buhl, D., Schwartz, P. R., Clark, F. O., Johnson, D. R., Lovas, F. J., Giguere, P. T.: ApJ, 191, L79 (1974)
- Snyder, L. E., Hollis, J. M., Brown, L. W., Buhl, D., Suenram, R. D., Lovas,
 F. J.: ApJ, 268, 123 (1983)
- 97. Snyder, L. E., Kuan, Y.-J., Miao, Y.: LNP Vol. 439: The Structure and Content of Molecular Clouds, 187 (1994)
- Snyder, L. E., Lovas, F. J., Mehringer, D. M., Miao, N. Y., Kuan, Y., Hollis, J. M., Jewell, P. R.: ApJ, 578, 245 (2002)
- 99. Spinrad, H., Wood, D. B. 1965: ApJ, 141, 109 (1965)
- 100. Stark, R., van der Tak, F. F. S., van Dishoeck, E. F.: ApJ, 521, L67 245 (1999)
- 101. Stutzki, J., Ungerechts, H., Winnewisser, G.: A&A, 111, 201 (1982)
- Stutzki, J., Olberg, M., Winnewisser, G., Jackson, J. M., Barrett, A. H.: A&A, 139, 258 (1984)
- 103. Stutzki, J., Winnewisser, G.: A&A, 144, 1 (1985a)
- 104. Stutzki, J., Winnewisser, G.: A&A, 144, 13 (1985b)
- 105. Stutzki, J., Winnewisser, G.: A&A, 148, 254 (1985c)
- 106. Swings, P., Rosenfeld, L.: ApJ, 86, 483 (1937)
- 107. Townes, C. H.: Microwave and radio-frequency resonance lines of interest to radio astronomy. In: IAU Symp. 4: Radio astronomy, ed. H. C. van de Hulst (Cambridge University Press 1957) pp 92–103
- 108. Turner, B. E.: ApJ, 163, L35 (1971)
- 109. Turner, B. E., Apponi, A. J.: ApJ, 561, L207 (2001)
- 110. Turner, B. E., Steimle, T. C., Meerts, L.: ApJ, 426, L97 (1994)
- van Dishoeck, E. F., Jansen, D. J., Schilke, P., Phillips, T. G.: ApJ, 416, L83 (1993)
- Walmsley, C. M., Bachiller, R., Forêts, G. P. d., Schilke, P.: ApJ, 566, L109 (2002)
- 113. Weinreb, S., Barrett, A. H., Meeks, M. L.: Nature, 200, 829 (1963)
- 114. Wilson, R. W., Jefferts, K. B., Penzias, A. A.: ApJ, 161, L43 (1970)
- 115. Winnewisser, G., Aliev, M. R., Yamada, K.: Interstellar Hydrides. In: Scientific Importance of Submillimetre Observations (ESA: SP-189) pp 23 28 (1982)
- 116. Winnewisser, G. Churchwell, E.: ApJ, 200, L33 (1975)
- 117. Winnewisser, G. Gardner, F. F.: A&A, 48, 159 (1976)
- Winnewisser, G., Winnewisser, M., Winnewisser, B. P.: Millimetre wave spectroscopy.
 In: MTP International Review of Science vol 3, Spectroscopy, ed by D. A. Ramsay (Butterworths, London 1972) pp 241–296
- Wootten, A., Turner, B. E., Mangum, J. G., Bogey, M., Boulanger, F., Combes, F., Encrenaz, P. J., Gerin, M.: ApJ, 380, L79 (1991)
- Yamamoto, S., Saito, S., Kawaguchi, K., Kaifu, N., Suzuki, H.: ApJ, 317, L119 (1987)
- 121. Ziurys, L. M., Apponi, A. J., Guélin, M., Cernicharo, J.: ApJ, 445, L47 (1995)
- Ziurys, L. M., Savage, C., Highberger, J. L., Apponi, A. J., Guélin, M., Cernicharo, J.: ApJ, 564, L45 (2002)
- 123. Zuckerman, B., Ball, J. A., Gottlieb, C. A.: ApJ, 163, L41 (1971)