N on-extensive statistics and the stellar polytrope index

R. Silva

Departamento de F sica, Universidade do Estado do Rio Grande do Norte, 59610-210, Mossoro - RN, Brasil

J. S. A lcaniz^y

Departamento de F sica, Universidade Federal do Rio Grande do Norte, C.P. 1641, Natal – RN, 59072–970, Brasil (Dated: February 7, 2020)

W e use physical constrains in posed from the H-Theorem and from the negative nature of the heat capacity of self-gravitating therm odynam ically isolated systems to investigate some possible lim its on the stellar polytrope index n within the dom ain of a classical non-extensive kinetic theory.

PACS num bers: 98.10.+ z, 04.40.-b, 05.70.Ln

Polytropic distribution functions provide the sim plest m odels for self-gravitating stellar systems [1]. These m odels are characterized by an equation of the state of the form p = K, where K and are constants. The polytropic index n is de ned by the following expression: = $1 + \frac{1}{n}$ which in the limit n ! 1 or, equivalently, = 1 results in an equation of state identical to the one of an isotherm albody of gases, i.e., p = K. The distri-

bution function associated to these systems is given by [1]

$$f() = \frac{1}{(2^{2})^{3=2}} e^{-2}$$
(1)

where $_1$ and = $(k_B\ T$ =m $)^{1=2}$ are, respectively, the density and the velocity dispersion. The quantity is equal to a constant m inus the total energy per unit m ass of an individual starm oving in the global galactic gravitational potential. In fact, is the relative energy of a star de ned by = v^2 =2, where is the relative potential, given by = $_0$, is the star potential energy and $_0$ is a arbitrary constant. In particular, the integrability condition $\ fd^3v < 1$ results in the constraint n > 1=2 (for a complete study on the gaseous polytropes and isotherm algas spheres, see [2]).

On the other hand, a non-extensive statistical form alism has been proposed as a possible extension of the classical one, which intends to study more properly system s that possess long-range interactions. Such a form alism is based on the entropy form ula β]

$$S_{q} = k_{B} \sum_{i}^{X} p_{i}^{q} \ln_{q} p_{i}; \qquad (2)$$

where p_i is the probability of the ith m icrostate and q is the non-extensive parameter. The Botzm ann-G ibbs extensive form ula, $S_1 = k_B \quad p_i \ln p_i$, is readily recovered in the extensive lim it q = 1 and the q-logarithm ic function above is de ned as [4]

$$\ln_q f = \frac{f^{1-q} - 1}{1-q}; f > 0;$$
 (3)

Som e attempts to build a kinetic counterpart for this non-extensive statistical form alism has been recently proposed in Refs. [5, 6]. The main result of these works consists in showing that this extended form ulation leads to a new velocity distribution for free particles given by

$$f_0(v) = B_q - 1 - (1 - q) \frac{m v^2}{2k_B T} ;$$
 (4)

where the quantity B_q is a q-dependent normalization constant whose expressions for q 1 and q 1 are respectively given by

$$B_{q-1} = G \frac{(\frac{1}{2} + \frac{1}{1-q})}{(\frac{1}{1-q})} \frac{m}{2 k_{B} T}$$
(5a)

and

$$B_{q-1} = H \frac{\left(\frac{1}{q-1}\right)}{\left(\frac{1}{q-1} - \frac{3}{2}\right)} \frac{m}{2 k_{\rm B} T}^{3=2}$$
(5b)

where the coe cients G and H are given by $G = n_p (1 q)^{1=2} (\frac{5}{2} \frac{3q}{2}) (\frac{3}{2} q)$ and $H = n_p (q 1)^{3=2}$, T is the tem – perature and n_p is the particle number density. This q-distribution can be derived at least from two di erent m ethods, namely, a generalization of the M axwell ansatz, f (v) \notin f (v_x) f (v_y) f (v_z), which follows from the introduction of statistical correlations between the com – ponents of the velocities [5] and a new form ulation for the Boltzm ann H -theorem, which requires q > 0 [6].

In the astrophysical dom ain, both the non-extensive statistical form alism and its kinetic counterpart have been applied in a wide range of problem s. The very rst application of this non-extensive statistics was done in connection with stellar polytropes [11], with several authors suggesting di erent expressions for the polytropic index n as a function of the non-extensive parameter q [8, 9, 10, 12]. Recently, a num erical study of stellar dynam ical evolution for self-gravitating system s was perform ed by Taruya and Sakagam i [13] while Chavanis [7] considered the T sallis' entropy as a particular H -function corresponding to isotherm al stellar system s and stellar polytropes. In this latter approach, the m axim ization of the H -function at xed m ass and energy reveals a

E lectronic address: rsilva@uem.br

^yE lectronic address: alcaniz@ dfte.ufm.br

therm odynam ical analogy with the study of the dynam ical stability in collisionless stellar systems. The nonextensive kinetic approach has been used to study the Jeans' gravitational instability [14, 15].

A nother recent application of this non-extensive kinetic theory was to investigate the negative nature of the heat capacity for isolated self-gravitating system [16]. This study was performed by considering the follow-ing steps: rst, consider a simple analogy between self-gravitating system and ideal gas (IG) or, in other words, the equivalence between IG kinetical and internal energies, given by $\frac{1}{2}$ m < v² > = $\frac{3}{2}$ k_BT, where m is the mass of a particle (e.g., a star). If such a system is composed by N particles, its total kinetic energy is K = $\frac{3}{2}$ N k_BT which, according to the viral theorem, is equal to m inus total energy, i.e., E = K. Therefore, the heat capacity of the system is given by (see [1] for details)

$$C_{V} = \frac{dE}{dT} = -\frac{3}{2}N k_{B} : \qquad (6)$$

Second, in order to investigate the q-dependence of the heat capacity, we consider a cloud of ideal gas within the non-relativistic gravitational context, which is analog to a self-gravitating collisionless gas. The kinetic energy of this system is simply given by K = $\frac{1}{2}m < v^2 >$, with the statistical content included in the average square velocity of the particles $< v^2 >$. Indeed, the non-extensivity m ay be introduced through a new derivation of expectation value [21]

$$< v^{2} >_{q} = \frac{R_{v_{m}}}{R_{v_{m}}} \frac{f^{q} v^{2} d^{3} v}{P_{v_{m}}};$$
 (7)

where $v_m = \frac{2k_B T}{m(1 q)}^{1-2}$ is a thermal cuto on the maximum value allowed for the velocity of the particles (q < 1), whereas for the power law without cuto (q > 1) $v_m ! 1$. This q-expectation value can be easily evaluated for $q \in 1$, resulting in

$$< v^2 >_q = \frac{6}{5 \quad 3q} \frac{k_B T}{m}$$
; for q < 5=3: (8)

Now, combining the q-expectation value for the square velocity of the particles with the de nition of the heat capacity $C_{\rm V}$ one nds

$$C_{\rm V} = \frac{3}{5 \ 3q} N k_{\rm B};$$
 (9)

which clearly places an upper lim it to the non-extensive parameter, i.e., q < 5=3. Following this reasoning, it is therefore natural that our kinetic approach uses the free particle distribution, with the long range nature of gravity being introduced via the virial theorem (see [16] for details). We note that, in the domain of ensemble theory, there are some analyses in the literature dealing with the classical IG within this nonextensive scenario [17, 18, 19, 20]. In particular, Ref. [18] showed that the



FIG.1: The quantity C_V =N $k_B\,$ for a self-gravitating collisionless gas obeying the non-extensive T sallis' q-statistic is shown as a function of the polytropic index n. The polytropic index and non-extensive parameter q are related by Eq. (12). A s indicated, the shadowed regions stand for the constraints from the second law of therm odynamics and the negative nature of C_V .

IG naturally exhibits negative heat capacity and that its pressure and particle num ber density obey a polytropetype relation. Here, how ever, we assume that all IG properties are accounted for by the q-M axwellian (see [5, 16] for details).

In this Letter, by considering the di erent expressions relating the polytropic index n and the non-extensive parameter q we discuss some possible physical constraints on n which arise from the second law of them odynam – ics (q > 0) and the negativeness of the heat capacity (q < 5=3). We show that the existent expressions between n and q (see [8, 9, 10]) can be naturally linked with the heat capacity in posing limits on the polytropic index. It is worth m entioning that the constraint q > 0, arising from the generalized H -theorem, does not provide a strong case for limiting the polytropic index n. This corresponds to a weak condition of the generalized H - theorem since it is possible to show that there are two arrow of time for q > 0 and q < 0.

In order to investigate such constraints we rst present the di erent expressions for the polytropic index n as a function of the non-extensive parameter q discussed in



FIG. 2: The same as in Fig. 1 for the case (ii). Here the polytropic index and non-extensive parameter q are related by Eq. (13).

the literature, i.e.,

(i)
$$n = \frac{1}{2} + \frac{1}{1 q}$$
 [8]; (10)

and

(ii)
$$n = \frac{5 \quad 3q}{2(1 \quad q)}$$
 [9;10]; (11)

The case (i) arises from an extension of Padmanabhan's classical analysis of the Antonov instability for the case of polytropic distribution function [22]. By using kinetic argum ents of the T sallis' non-extensive form alism, the case (ii) is obtained from an extension of the polytropic Lane-Em den spheres as well as from a study of maximum entropy solutions of the V lasov-Poisson equations describing self-gravitating system s. W e em phasize that the polytropic family of solutions to the collision less Boltzmann equation (parameterized by the index n) is always the same, which means that expressions (10) and (11) constitute only di erent ways of param eterizing (in term s of non-extensive param eter q) this polytropic fam ily. In both cases, Eqs. (10) and (11), the sam e choice of the statistical average, the so-called q-expectation value, has been used. As expected, for all the above expressions the M axwellian isotherm alspheres are obtained in the $\lim_{t \to 0} \frac{1}{2} = 1$ or, equivalently, n = 1.

By combining Eqs. (10)-(11) with Eq. (9), one obtains the following expressions relating the heat capacity and

TABLE I:Limitston

Thermodynamic constraints	case (i)	case (ii)
q > 0	$\frac{1}{2} < n < \frac{3}{2}$	$\frac{3}{2} < n < \frac{5}{2}$
q < 5=3	$0 < n < \frac{1}{2}$	$0 < n < \frac{3}{2}$

the polytropic index:

(i)

$$C_{V} = \frac{(3 \quad 6n)}{4n + 4} N k_{B};$$
 (12)

and

(ii)
$$C_V = \frac{(9 \text{ 6n})}{4n} N k_B$$
: (13)

From the above expressions, it is straightforward to see that by in posing the constraints on the non-extensive param eterq from the second law of therm odynam ics (q > 0) and the negative nature of $C_{\rm V}~(q < 5=3)$ it is possible to lim it regions in the $C_{\rm V}~$ n plane. To better visualize such constraints, in Figs. 1 and 2 we show, respectively for the cases (i) and (ii), the dimensionless quantity $C_{\rm V}$ =N $k_{\rm B}$ as a function of the polytropic index n.

In Figures 1 [case (i)] and 2 [case (ii)] we show that the constraint q > 0 results, respectively, in the forbidden intervals for the polytropic index $\frac{1}{2} < n < \frac{3}{2}$ and $\frac{3}{2} < n < \frac{5}{2}$ while the lim it resulting from Eq. (9), i.e., $q < \frac{5}{3}$ in plies $0 < n < \frac{1}{2}$ and $0 < n < \frac{3}{2}$. In all cases the bound on q from the second law of therm odynam ics (q > 0) restricts only negative regions in the plane $C_V = N k_B$ n while, as expected, the them odynam ical lim it $C_V < 0$ forbids any possible positive region in the plots. In particular, we also note that the constraints on n from Eq. (12) (case i) seem to be in agreem ent with the theorem proved by Antonov and Dremus [1] who showed that the equilibrium distribution is dynamically stable for values of the polytrope index n > 3=2 or, equivalently, q > 0 (see Fig. 1). The bounds on the polytopic index n obtained in this Letter are sum marized in Table I.

Finally, we emphasize that in a more quantitative case, i.e., in which a self-gravitating ideal gas is contained in a spherical container of radius R our constraints on the polytropic index n must be valid only in the negative region of the heat capacity. As widely known, when the potential energy is taking into account or, equivalently, the virial theorem is modiled to introduce a pressure term, 2K + W ' p(R), the heat capacity can assume both positive and negative values. We also note that a possible connection with stability of spherical systems can be introduced through the D orem us-Feix-B aum ann theorem which in plies, for the extensive case, that polytropes with $\frac{1}{2} < n < \frac{3}{2}$ are unrealistic (see, how ever, [23, 24]).

A cknow ledgm ents

The authors are very grateful to A. Taruya, J. R. da Silva and Prof. J. A. S. Lim a for helpful discussions and critical reading of the manuscript. We also

- J.B inney, S.Tram aine G alactic D ynam ics (Princeton U. Press, 1994).
- [2] S.Chandrasekhar An introduction to the Theory of Stellar Structure (University of Chicago Press, 1958).
- [3] C.Tsallis, J.Stat. Phys. 52 (1988) 479.
- [4] See also http://tsallis.cat.cbpfbr/biblio.htm for an up to date bibliography.
- [5] R. Silva, A. R. Plastino, J. A. S. Lima, Phys. Lett. A 249 (1998) 401.
- [6] J.A.S.Lima, R.Silva, A.R.Plastino, Phys. Rev. Lett. 86 (2001) 2938. cond-m at/0210072
- [7] P.H.Chavanis, A&A 386 (2002) 732; 401 (2003) 15.
- [8] A. Taruya, M. Sakagami, Physica A 322 (2003) 285.
- [9] A.R.P lastino, in Nonextensive Statistical Mechanics and Its Aplications, eds. S. Abe and Y.O kam oto (Springer, Berlin, 2001)
- [10] J. A. S. Lima, R. E. de Souza, On the Nonextensive Isotherm al Stellar Spheres. Submitted for publication.
- [11] A. Plastino, A. R. Plastino, Phys. Lett. A 174 (1993) 384.

- thank the anonymous referee for valuable comments. JSA is supported by the Conselho Nacional de Desenvolvimento Cient coe Tecnologico (CNPq-Brasil) and CNPq (62.0053/01-1-PADCT III/M ilenio).
- [12] A. Taruya, M. Sakagami, Physica A 318 (2003) 387
- [13] A. Taruya, M. Sakagam i, Phys. Rev. Lett. 90 (2003) 181101.
- [14] J.A.S.Lima, R.Silva, J.Santos, A&A 396 (2002) 309. astro-ph/0109474
- [15] J.L.Du, Phys. Lett. A 320 (2004) 347
- [16] R. Silva, and J. S. Alcaniz, Phys. Lett. A 313, (2003) 393. astro-ph/0306009
- [17] A.R.Plastino, A.Plastino and C.Tsallis, J.Phys.A 27, (1994) 5707.
- [18] S.Abe, Phys. Lett. A 263, (1999) 424.
- [19] S.Abe, Physica A 269, (1999) 403.
- [20] L.C.L.Botelho, M od. Phys. Lett. B 17, (2003) 733.
- [21] C.T sallis, R.S.M endes and A.Plastino, Physica A 261 (1998) 534.
- [22] T.Padm anabhan, Phys.Rep.188 (1990) 285.
- [23] M.Henon, A&A 24 (1973) 229.
- [24] J.Bames, P.Hut, Nature 324 (1986) 446.