

Particle-In-Cell simulations of circularly polarised Alfvén wave phase mixing: a new mechanism for electron acceleration in collisionless plasmas

David Tsiklauri¹, Jun-Ichi Sakai², and Shinji Saito²

¹ Institute for Materials Research, School of Computing, Science and Engineering, University of Salford, Salford, Greater Manchester, M5 4WT, United Kingdom.

² Laboratory for Plasma Astrophysics, Faculty of Engineering, Toyama University, 3190, Gofuku, Toyama, 930-8555, Japan

Received ??? 2004 / Accepted ??? 2004

Abstract. Particle-In-Cell simulations of a circularly polarised Alfvén wave interaction with one dimensional, across the uniform magnetic field, plasma density inhomogeneity (phase mixing) in collisionless plasmas were performed. In our preliminary work (Tsiklauri et al. 2004) we reported discovery of a new electron acceleration mechanism. In this mechanism, progressive distortion of Alfvén wave front, due to the differences in local Alfvén speed, generates oblique (nearly parallel to the magnetic field) electrostatic fields, which accelerate electrons through the Landau resonance. Here we report detailed study of this novel effect, including analysis of broadening of ion distribution function due to the presence of Alfvén waves and the generation of compressive perturbations due to weak non-linearity and plasma density inhomogeneity. The amplitude decay law in the inhomogeneous regions, in the kinetic regime, is shown to be the same as in the MHD approximation described by Heyvaerts and Priest (1983).

Key words. Sun: oscillations – Sun: Corona – (Sun:) solar wind

1. Introduction

The study of interaction of Alfvén waves (AWs) with plasma inhomogeneities is important for both astrophysical and laboratory plasmas. This is because both AWs and inhomogeneities often coexist in a number of these physical systems. AWs are believed to be good candidates for plasma heating, energy and momentum transport. On one hand, in many physical situations AWs are easily excitable (e.g. through convective motions of the solar interior) and they are present in a number of astrophysical systems. On the other hand these waves dissipate on shear viscosity as opposed to compressive fast and slow magnetosonic waves which dissipate on bulk viscosity. In astrophysical plasmas shear viscosity is extremely small as compared to bulk one. Hence, AWs are notoriously difficult to dissipate. One of the possibilities to improve AW dissipation is to introduce progressively decreasing spatial scales, $\delta l \rightarrow 0$, into the system (recall that the classical dissipation is $\propto \delta l^{-2}$). Heyvaerts and Priest have proposed (in astrophysical context) one such mechanism called AW phase mixing (Heyvaerts & Priest 1983). It occurs when

a linearly polarised AW propagates in the plasma with one dimensional, transverse to the uniform magnetic field density inhomogeneity. In such situation initially plane AW front is progressively distorted because of different Alfvén speeds across the field. This creates progressively strong gradients across the field (effectively in the inhomogeneous regions transverse scale collapses to zero), and thus in the case of finite resistivity, dissipation is greatly enhanced. Hence, it is believed that phase mixing can provide significant plasma heating. Phase mixing could be also important for laboratory plasmas: Hasegawa & Chen (1974) proposed the heating of collisionless plasma by utilising a spatial phase mixing by shear Alfvén wave resonance and discussed potential applications to toroidal plasma. A significant amount of work has been done in the context of heating open magnetic structures in the solar corona (Heyvaerts & Priest 1983, Nocera et al. 1986, Parker 1991, Nakariakov et al. 1997, DeMoortel et al. 2000, Botha et al. 2000, Tsiklauri et al. 2001, Hood et al. 2002, Tsiklauri & Nakariakov 2002, Tsiklauri et al. 2002, 2003). All phase mixing studies so far have been performed in the MHD approximation, however, since the transverse scales in the AW collapse progressively to zero, MHD approximation is inevitably violated, first, when

the transverse scale approaches ion gyro-radius r_i and then electron gyro-radius r_e . Thus, we proposed to study phase mixing effect in the kinetic regime, i.e. we go beyond MHD approximation. Preliminary results were reported in (Tsiklauri et al. 2004), where we discovered new mechanism for the acceleration of electrons due to wave-particle interactions. This has important implications for various space and laboratory plasmas, e.g. coronal heating problem and acceleration of solar wind. In this paper we present full analysis of the discovered effect and include analysis of broadening of ion distribution function due to the presence of Alfvén waves and the generation of compressive perturbations due to weak non-linearity and plasma density inhomogeneity.

2. the model

We used 2D3V, the fully relativistic, electromagnetic, particle-in-cell (PIC) code with MPI parallelisation, modified from 3D3V TRISTAN code (Buneman 1993, p.67). The system size is $L_x = 5000\Delta$ and $L_y = 200\Delta$, where $\Delta (= 1.0)$ is the grid size. The periodic boundary conditions for x - and y -directions are imposed on particles and fields. There are about total of 15 million electrons and ions in the simulation. The average number of particles per cell is 100 in low density regions (see below). Thermal velocity of electrons is $v_{th,e} = 0.1c$ and the same for ions is $v_{th,i} = 0.025c$. The ion to electron mass ratio is $m_i/m_e = 16$. The time step is $\omega_{pe}\Delta t = 0.05$. Here ω_{pe} is the electron plasma frequency. The Debye length is $v_{th,e}/\omega_{pe} = 1.0$. The electron skin depth is $c/\omega_{pe} = 10\Delta$, while the ion skin depth is $c/\omega_{pi} = 40\Delta$. Here ω_{pi} is the ion plasma frequency. The electron Larmor radius is $v_{th,e}/\omega_{ce} = 1.0\Delta$, while the same for ions is $v_{th,i}/\omega_{ci} = 4.0\Delta$. The external uniform magnetic field, $B_0 (= 1.25)$, is in the x -direction and the initial electric field is zero. The ratio of electron cyclotron frequency to electron plasma frequency is $\omega_{ce}/\omega_{pe} = 1.0$, while the same for ions is $\omega_{ci}/\omega_{pi} = 0.25$. The latter ratio is essentially V_A/c – the Alfvén speed normalised to the speed of light. Plasma $\beta = 2(\omega_{pe}/\omega_{ce})^2(v_{th,e}/c)^2 = 0.02$. Here all plasma parameters are quoted far away from the density inhomogeneity region. The dimensionless (normalised to some reference constant value of $n_0 = 100$ particles per cell) ion and electron density inhomogeneity is described by

$$n_i(y) = n_e(y) = 1 + 3 \exp \left[- \left(\frac{y - 100}{50\Delta} \right)^6 \right] \equiv F(y). \quad (1)$$

This means that in the central region (across y -direction) density is smoothly enhanced by a factor of 4, and there are strongest density gradients of width about 50Δ around the points $y = 51.5\Delta$ and $y = 148.5\Delta$. The background temperature of ions and electrons, and their thermal velocities are varied accordingly

$$T_i(y)/T_0 = T_e(y)/T_0 = F(y)^{-1}, \quad (2)$$

$$v_{th,i}/v_{i0} = v_{th,e}/v_{e0} = F(y)^{-1/2}, \quad (3)$$

such that the thermal pressure remains constant. Since the background magnetic field along x -coordinate is also constant, the total pressure remains constant too. Then we impose current of the following form

$$\partial_t E_y = -J_0 \sin(\omega_d t) (1 - \exp[-(t/t_0)^2]), \quad (4)$$

$$\partial_t E_z = -J_0 \cos(\omega_d t) (1 - \exp[-(t/t_0)^2]). \quad (5)$$

Here ω_d is the driving frequency which was fixed at $\omega_d = 0.3\omega_{ci}$, which insures that no significant ion-cyclotron damping is present. Also, ∂_t denotes time derivative. The t_0 is the onset time of the driver, which was fixed at $50/\omega_{pe}$ or $3.125/\omega_{ci}$. This means that the driver onset time is about 3 ion-cyclotron periods. Imposing such current on the system results in the generation of left circularly polarised AW, which is driven at the left boundary of simulation box and has spatial width of 1Δ . Initial amplitude of the current is such that relative AW amplitude is about 5 % of the background (in the low density homogeneous regions), thus the simulation is weakly non-linear.

3. main results

Because of no initial (perpendicular to the external magnetic field) velocity excitation was imposed in addition to the above specified currents (cf. Tsiklauri & Nakariakov (2002), DelZanna et al. (2001), Turkmani & Torkelsson (2003, 2004)), the excited (driven) at the left boundary circularly polarised AW is split into two circularly polarised AWs that travel in opposite directions. The dynamics of these waves as well as other physical quantities is shown in Fig. 1. (cf. with Fig. 1 from (Tsiklauri et al. 2004) where three time snapshots of B_z and E_y , circularly polarised Alfvén wave components were shown). Typical simulation, till $t = 875/\omega_{ce} = 54.69/\omega_{ci}$ takes about 8 days on the parallel 32 dual 2.4 GHz Xeon processors. It can be seen from the figure that because of the periodic boundary conditions circularly polarised AW that was travelling to the left has reappeared from the right side of the simulation box. The dynamics of the AW (B_z, E_y) progresses in a similar manner as in MHD, i.e. it phase mixes (Heyvaerts & Priest 1983). In other words middle region (in y -coordinate), i.e. $50\Delta \leq y \leq 150\Delta$, travels slower because of the density enhancement (note that $V_A(y) \propto 1/\sqrt{n_i(y)}$). This obviously causes distortion of initially plain wave front and the creation of strong gradients in the regions around $y \approx 50$ and 150 . In MHD approximation, in the case of finite resistivity η , in these regions AW is strongly dissipated. This effectively means that the outer and inner parts of the travelling AW are disattached from each other and propagate independently. This is way the effect is called phase mixing – after long time, in the case of developed phase mixing, phases in the wave front become effectively uncorrelated. Before study of (Tsiklauri et al. 2004), it was not clear what to expect from our PIC simulation. The code is collisionless and there are no sources of dissipation in it (apart

from possibility of wave-particle interactions). It is evident from Fig. 1 that in the developed stage of phase mixing ($t = 54.69/\omega_{ci}$) AW front is substantially damped in the strongest density gradient regions. Contrary to the AW (B_z and E_y) dynamics we do not see any phase mixing for B_y and E_z which behave similarly. It should be noted that E_z contains both driven (see Eq.(4) above) and non-linearly generated fast magnetosonic wave component, with the former being dominant over the latter. While since the B_x is not driven, and initially its perturbations were absent, we see only non-linearly generated slow magnetosonic perturbations confined to the regions of strongest density gradients (around $y \approx 50$ and 150). Note that these also have rapidly decaying amplitude. Also, we gather from Fig. 1 that the density perturbation ($\approx 10\%$) which is also generated through the weak non-linearity is present too. These are propagating density oscillations with variation both in overall magnitude (perpendicular to the figure plane) and across y -coordinate, and they are mainly confined to the strongest density gradients regions (around $y \approx 50$ and 150). Note that dynamics of B_x , B_y and B_z with appropriate geometrical switching (because in our geometry uniform magnetic field is along x -coordinate) is in qualitative agreement with (Botha et al. 2000) (cf their Fig. 9). Dynamics of remaining E_x component is treated separately in the next figure. It should be mentioned that it is the inhomogeneity of the medium $n_i(y)$, $V_A(y)$, i.e. $\partial/\partial y \neq 0$, is the cause of weakly non-linear coupling of the AWs to the compressive modes (see Nakariakov et al. (1997), Tsiklauri et al. (2001) for further details).

In Fig. 2 we try to address the question where did the AW energy go (as we saw strong decay of AWs in the regions of strong density gradients)? Thus in Fig. 2 we plot E_x longitudinal electrostatic field, and electron phase space (V_x/c vs. x and V_x/c vs. y) for the different times. We gather from this figure is that in the regions around $y \approx 50$ and 150 for later times significant electrostatic field is generated. This is the consequence of stretching of AW front in those regions because of difference in local Alfvén speed. In the middle column of this figure we see that exactly in those regions where E_x is generated electrons are accelerated in copious amounts along x -axis. We also gather from the right column that for later times ($t = 54.69/\omega_{ci}$) number of high velocity electrons is increased around strongest density gradient regions (around $y \approx 50$ and 150). Thus, the generated E_x field is somewhat oblique (not exactly parallel to the external magnetic field). Hence, we conclude that energy of the phase mixed AW goes into acceleration of electrons. In fact line plots of E_x show that this electrostatic field is strongly damped. i.e. energy is channelled to electrons via Landau damping.

In Fig. 3 we investigate ion phase space. In particular we plot ion phase space (V_z/c vs. x and V_z/c vs. y) for the different times. The reason for choice of V_z will become clear below. We gather from this plot that in V_z/c vs. x phase space clear propagating oscillations are present (left

column). These oscillations are incompressible, Alfvénic "kink" type, i.e. for those x 's where there is an increase of greater positive velocity ions, there is also corresponding decrease of lower negative velocity ions. In the V_z/c vs. y plot we also see no clear acceleration of ions.

As our next step, we look at the distribution functions of electrons and ions before and after the phase mixing took place. In Fig. 4 we plot distribution functions of electrons and ions at $t = 0$ and $t = 54.69/\omega_{ci}$. Note that at $t = 0$ the distribution functions do not look as purely Maxwellian because of the fact that temperature varies across y -coordinate (to keep total pressure constant) and the graphs are produced for the entire simulation domain. Also, note that for electrons in $f(V_x)$ there is a substantial difference at $t = 54.69/\omega_{ci}$ from its original form because of the aforementioned electron acceleration took place. In fact we see that number of electrons having velocities $V_x = \pm(0.1 - 0.3)c$ is increased. Note that the acceleration of electrons takes place mostly along the external magnetic field (along x -coordinate). Thus, very little electron acceleration occurs in V_y or V_z (solid and dotted curves practically overlap on each other). For the ions the situation is different: we see broadening of the ion velocity distribution functions in V_z and V_y (that is why we have chosen to present V_z component of ion phase space in Fig. 3).

The reason for this broadening of the ion distribution function becomes clear in Fig. 5 where we plot kinetic energies x, y, z components ($\propto V_{x,y,z}^2$) and total kinetic energies for electrons (top row) and the same for ions (bottom row). For ions we gather that y and z components of the kinetic energy (bottom left figure) oscillate in anti-phase and their oscillatory part perfectly cancels out in the total energy (bottom right figure). Thus, the broadening of the y and z components of the ion velocity distribution functions is due to presence of AWs (usual wave broadening, which is actually observed e.g. in the solar corona and solar wind, Belcher & Davis (1971), Smith et al. (1995), Bavassano et al. (2000)), and hence there is no ion acceleration present. Note that y and z components and hence total kinetic energy of ions is monotonously increasing due to continuous AW driving. Note that no significant motion of ions along the field is present. For electrons on the other hand we see significant increase of x component (along the magnetic field) of kinetic energy which is due to the new electron acceleration mechanism discovered by us (cf. Fig.10). Note that for ions y component attains lower values (than z component) because of lower AW velocity in the middle part of the simulation domain.

The next step is to check whether the increase in electron velocities really comes from the resonant wave particle interactions. For this purpose in Fig. 6, left panel, we plot two snapshots of Alfvén wave $B_z(x, y = 148)$ component at instances $t = 54.69/\omega_{ci}$ (solid line) and $t = 46.87/\omega_{ci}$ (dotted line). The distance between the two upper leftmost peaks (which is the distance travelled by the wave in time span between the snapshots) is about $\delta L = 150\Delta = 15(c/\omega_{pe})$. Time difference between the snapshots is $\delta t = 7.82/\omega_{ci}$. Thus, measured




Fig. 1. Contour (intensity) plots of electromagnetic field components and electron density at time $t = 54.69/\omega_{ci}$ (developed stage of phase mixing). Phase mixed Alfvén wave components are B_z and E_y . Excitation source is at the left boundary. Because of periodic boundary conditions, left-propagating AW re-appears from the right side of the simulation box. Note how (initially plain) AW is stretched because of differences in local Alfvén speed across y -coordinate. Significant ($\approx 10\%$) density fluctuations can be seen.

AW speed at the point of the strongest density gradient ($y = 148$) is $V_A^M = \delta L/\delta t = 0.12c$. Now we can also work out the Alfvén speed from the theory. In the homogeneous low density region the Alfvén speed was set to be $V_A(\infty) = 0.25c$. From Eq.(1) it follows that for $y = 148$ density is increased by a factor of 2.37 which means that the Alfvén at this position speed is $V_A(148) = 0.25/\sqrt{2.37}c = 0.16c$. The measured ($0.12c$) and calculated ($0.16c$) Alfvén speeds in the inhomogeneous region differ somewhat. This probably can be attributed to the fact that in the inhomogeneous regions (where electron acceleration takes place) because of momentum conservation AW front is decelerated as it passes on energy and momentum to the electrons. However, this may be not the case if wave-particle interactions play the same role as dissipation in MHD (Sagdeev & Galeev 1969): Then wave-particle interactions would result only in the decrease of the AW amplitude (dissipation) not in its deceleration. If we compare these values to Fig. 4 (top left panel for $f(V_x)$), we deduce that these are the veloci-

ties $> 0.12c$ above which electron numbers with higher velocities is greatly increased. This deviation peaks at about $0.25c$ which in fact corresponds to the Alfvén speed in the lower density regions. This can be explained by the fact the electron acceleration takes place in wide regions (cf. Fig. 2) along and around $y \approx 148$ (and $y \approx 51$) – thus the spread in the accelerated velocities. In Fig. 6 we also plot a visual fit curve (dashed line) in order to quantify the amplitude decay law for the AW (at $t = 54.69/\omega_{ci}$) in the strongest density inhomogeneity region. The fitted (dashed) curve is represented by $0.056 \exp[-(x/1250)^3]$. There is a surprising similarity of this fit with the MHD approximation results. Heyvaerts & Priest (1983) found that for large times (developed phase mixing), in the case of harmonic driver, the amplitude decay law is given by $\propto \exp[-(\frac{\eta\omega^2 V_A'^2}{6V_A^2})x^3]$ which is much faster decay law as compared to usual resistivity dissipation $\propto \exp(-\eta x)$. Here V_A' is the derivative of the Alfvén speed with respect to y -coordinate. The most interesting fact is that even




Fig. 2. Left column: contour plots of generated nearly parallel to the external magnetic field electrostatic field E_x at instances: $t = (0, 31.25, 54.69)/\omega_{ci}$. Central column: V_x/c versus x of electron phase space at the same times. Note, that in order to reduce figure size, only electrons with $V_x > 0.15c$ were plotted. Right column: V_x/c versus y of electron phase space at the same times. Note, that in order to reduce figure size, only electrons with $V_x > 0.12c$ were plotted (note the dip in the middle due to density inhomogeneity across y -coordinate).




Fig. 3. Left column: V_z/c versus x of ion phase space at instances: $t = (0, 31.25, 54.69)/\omega_{ci}$. Right column: V_z/c versus y of ion phase space at the same times. Note, that in order to reduce figure size, only ions with $V_z > 0.03c$ were plotted (note the dip in the middle due to density inhomogeneity across y -coordinate).

in the kinetic approximation the same $\propto \exp(-Ax^3)$ law holds as in MHD.

In MHD, finite resistivity and Alfvén speed non-uniformity are responsible for the enhanced dissipation via phase mixing mechanism. In our PIC simulations (ki-

netic phase mixing), however, we do not have dissipation and collisions (dissipation). Thus, in our case wave-particle interactions play the same role as resistivity η in the MHD phase mixing (Sagdeev & Galeev 1969). It should be noted that no significant AW dissipation was




Fig. 4. All three components of the distribution functions of electrons (top row) and ions (bottom row) at $t = 0$ (dotted curves) and $t = 54.69/\omega_{ci}$ (solid curves), i.e. for the developed stage of phase mixing.




Fig. 6. Left: two snapshots of Alfvén wave $B_z(x, y = 148)$ component at instances $t = 54.69/\omega_{ci}$ (solid line) and $t = 46.87/\omega_{ci}$ (dotted line). Dashed line represents fit $0.056 \exp[-(x/1250)^3]$. Center: $B_z(x, y = 10)$ (low density homogeneous region), $B_z(x, y = 100)$ (high density homogeneous region). Note the differences in amplitudes and propagation speeds, which are consistent with the equilibrium density and thus Alfvén speed dependence on y -coordinate.

found away from the density inhomogeneity regions (Fig. 6 middle and right panels, note also the differences in amplitudes and propagation speeds, which are consistent with the imposed density and hence Alfvén speed variation across y -coordinate). This has the same explanation as in the case of MHD – it is in the regions of density inhomogeneities ($V'_A \neq 0$) the dissipation is greatly enhanced,

while in the regions where $V'_A = 0$ there is no substantial dissipation (apart from classical $\propto \exp(-\eta x)$ one). In MHD approximation aforementioned amplitude decay law is derived from the diffusion equation, to which MHD equations reduce to for large times (developed phase mixing). It seems that kinetic description leads to the same type of diffusion equation. It is unclear although, at this




Fig. 5. Top row: kinetic energies (calculated by x (solid), y (dotted), z (dashed) velocity (squared) components) of electrons (left), and total kinetic energy for electrons (right) as a functions time. Bottom row: As above but for ions. The units on y -axis are arbitrary.

stage, what physical quantity would play role of resistivity η (from the MHD approximation) in the kinetic regime.

3.1. Homogeneous plasma case

In order to clarify issue with the broadening of the ion velocity distribution function and also for a consistency check we performed an additional simulation in the case of homogeneous plasma. Now the density was fixed at 100 ions/electrons per cell in the entire simulation domain and hence plasma temperature and thermal velocities were fixed as constants too. In such set up no phase mixing should take place as the AW speed is uniform.

In Fig. 7 we plot the only non-zero (above noise level) components at $t = 54.69/\omega_{ci}$, which are left circularly polarised AW fields: B_z, B_y, E_z, E_y . Note there are no B_x, E_x or density fluctuations present in this case (cf. Fig. 1) as it is the plasma inhomogeneity that facilitates the coupling between AW and the compressive modes.

In Fig. 8 we plot ion phase space (V_z/c vs. x and V_z/c vs. y) in the homogeneous plasma case for different times (cf. Fig. 3). We gather from the graph that still propagating incompressible, Alfvénic "kink" type oscillations are present (left column). While no significant ion acceleration takes place (right column). This is better witnessed from Fig. 9 where we plot electron and ion distribution functions for $t = 0$ and $t = 54.69/\omega_{ci}$ as in Fig. 4 for the homogeneous plasma case. There are three noteworthy points: (i) no electron acceleration is taking place because of the absence of phase mixing; (ii) there is again (as in the inhomogeneous case) broadening of the ion velocity distribution functions (in V_y and V_z) due to the present AW (wave broadening); (iii) The distribution looks now Maxwellian (cf. Fig. 4) because the distribution function is for the entire, homogeneous region.




Fig. 10. As is Fig. 5 but for the case of homogeneous plasma density (no phase mixing).

In Fig. 10 we plot kinetic energies x, y, z components ($\propto V_{x,y,z}^2$) and total kinetic energies for electrons (top row) and ions (bottom row). For ions we gather again (as in the inhomogeneous case) that y and z components of the kinetic energy (bottom left figure) oscillate in anti-phase and their oscillatory part perfectly cancels out in the total energy (bottom right figure). Thus, the broadening of the y and z components of the ion velocity distribution functions is due to the presence of AWs, and, in turn, there is no ion acceleration. Again y and z components and hence total kinetic energy of ions is monotonously increasing due to continuous AW driving. No significant motion of ions along the field is present. For electrons we do not observe any acceleration due to the absence of phase mixing (cf. Fig. 5). Note that y component attains now the same values as z component (bottom left figure) because of the same AW velocity in the entire part of the simulation domain.

4. discussion

In our preliminary work (Tsiklauri et al. 2004) we have outlined main results of the discovered new mechanism of electron acceleration. Here we presented more detailed analysis of the phenomenon. We have established the following:

- Progressive distortion of Alfvén wave front, due to the differences in local Alfvén speed, generates oblique (nearly parallel to the magnetic field) electrostatic fields, which accelerate electrons.
- The amplitude decay law in the inhomogeneous regions, in the kinetic regime, is shown to be the same as in the MHD approximation described by (Heyvaerts & Priest 1983).
- Due to the present weak non-linearity and plasma inhomogeneity, the density perturbations ($\approx 10\%$) are generated. These are propagating density oscillations with variation both in overall magnitude and across y -




Fig. 7. As is Fig. 1 but for the case of homogeneous plasma density (no phase mixing). Note only non-zero (above noise level) components are plotted. There are no B_x , E_x or density fluctuations present in this case.




Fig. 8. As is Fig. 3 but for the case of homogeneous plasma density (no phase mixing).

coordinate. They are mainly confined to the strongest density gradients regions (around $y \approx 50$ and 150). Longitudinal to the external magnetic field, B_x , perturbations are also generated in the same manner, but with smaller ($\approx 3\%$) amplitudes.

- Both in the homogeneous and inhomogeneous cases presence of AWs causes broadening the perpendicular (to the external magnetic field) ion velocities distribution functions, while no ion acceleration is observed.

It is worthwhile to mention that in MHD approximation authors of Ref.(Hood et al. 2002, Tsiklauri et al. 2003) showed that in the case of localised Alfvén pulses, Heyvaerts and Priest’s amplitude decay formula $\propto \exp(-Ax^3)$ (which is true for harmonic AWs) is replaced by the power law $B_z \propto x^{-3/2}$. A natural next step forward would be to check whether in the case of localised Alfvén pulses the same power law holds in the kinetic regime.




Fig. 9. As is Fig. 4 but for the case of homogeneous plasma density (no phase mixing).

Finally we would like to mention that after this study was complete we became aware of study by Vasquez & Hollweg (2004), who used hybrid code (electrons treated as neutralising fluid, while ion kinetics is retained) as opposed to our (fully kinetic) PIC code, to simulate resonant absorption. They found that a planar (body) Alfvén wave propagating at less than 90° to a background gradient has field lines which lose wave energy to another set of field lines by cross-field transport. Further, Vasquez (2004) found that when perpendicular scales of order 10 proton inertial lengths ($10c/\omega_{pi}$) develop from wave refraction in the vicinity of the resonant field lines, a non-propagating density fluctuation begins to grow to large amplitudes. This saturates by exciting highly oblique, compressive, and low-frequency waves which dissipate and heat protons. These processes lead to a faster development of small scales across the magnetic field, i.e. phase mixing mechanism, studied here.

Acknowledgements. Authors gratefully acknowledge support from CAMPUS (Campaign to Promote University of Salford) which funded J.-I.S.'s one month fellowship to the Salford University that made this project possible. DT acknowledges use of E. Copson Math cluster funded by PPARC and University of St. Andrews. DT kindly acknowledges support from Nuffield Foundation through an award to newly appointed lecturers in science, engineering and mathematics (NUF-NAL 04).

References

- Bavassano B., Pietropaolo E., Bruno R., 2000, *J. Geophys. Res.*, 105, 15959
- Belcher J.W., Davis L., 1971, *J. Geophys. Res.*, 79, 4174
- Botha G.J.J., Arber T.D., Nakariakov V.M., Keenan F.P., 2000, *Astron. Astrophys.*, 363, 1186
- Buneman O., 1993, p.67, in *Computer Space Plasma Physics: Simulation Techniques and Software*, Terra Scientific, New York
- DelZanna L., Velli M., Londrillo P., 2001, *Astron. Astrophys.*, 367, 705
- DeMoortel L., Hood A.W., Arber T.D., 2000, *Astron. Astrophys.*, 354, 334
- Hasegawa A., Chen L., 1974, *Phys. Rev. Lett.*, 32, 454
- Heyvaerts J., Priest E.R., 1983, *Astron. Astrophys.*, 117, 220
- Hood A.W., Brooks S.J., Wright A.N., 2002, *Proc. Roy. Soc. Lond. A*, 458, 2307
- Nakariakov V.M., Roberts B., Murawski K., 1997, *Sol. Phys.*, 175, 93
- Nocera L., Priest E.R., Hollweg J.V., 1986, *Geophys. Astrophys. Fl. Dyn.*, 35, 111
- Parker E.N., 1991, *Astrophys. J.*, 376, 355
- Sagdeev R.Z., Galeev A.A., 1969, *Non-Linear Plasma Theory*, W. A. Benjamin, Inc. Book Company, New York
- Smith E.J., Balogh A., Neugebauer M., McComas D., 1995, *Geophys. Res. Lett.*, 22, 3381
- Tsiklauri D., Nakariakov V.M., 2002, *Astron. Astrophys.*, 393, 321
- Tsiklauri D., Arber T.D., Nakariakov V.M., 2001, *Astron. Astrophys.*, 379, 1098
- Tsiklauri D., Nakariakov V.M., Arber T.D., 2002, *Astron. Astrophys.*, 395, 285
- Tsiklauri D., Nakariakov V.M., Rowlands G., 2003, *Astron. Astrophys.*, 400, 1051
- Tsiklauri D., Sakai J.I., Saito S., 2004, *Phys. Rev. Lett.* (submitted), astro-ph/0409537
- Turkmani R., Torkelsson U., 2003, *Astron. Astrophys.*, 409, 813

- Turkmani R., Torkelsson U., 2004, *Astron. Astrophys.*,
428, 227
- Vasquez B.J., 2004, *J. Geophys. Res.* (submitted)
- Vasquez B.J., Hollweg J.V., 2004, *Geophys. Res. Lett.*,
31, L14803