Relic Abundance of Mass-Varying Cold Dark Matter Particles

Rogerio Rosenfeld

Instituto de Física Teórica - UNESP, Rua Pamplona, 145, 01405-900, São Paulo, SP, Brazil

Abstract

In models of coupled dark energy and dark matter the mass of the dark matter particle depends on the cosmological evolution of the dark energy field. In this note we exemplify in a simple model the effects of this mass variation on the relic abundance of cold dark matter. We still do not know the origin and composition of the cold dark matter (CDM) in the universe. Recent precision measurements of the cosmological microwave background and of the large scale structure of the universe put a strict bound on the abundance of non-baryonic CDM [1]:

$$\Omega_{CDM}h^2 = 0.113 \pm 0.008,\tag{1}$$

resulting from a fit of several measurements combined in the framework of a Λ CDM model with a running spectral index.

Most probably CDM is made of particles (even though there are alternatives where dark matter is the manifestation of a fluid with a non-standard equation of state, such as the Chaplygin gas and quartessence models [2]) and the default candidate is the lightest supersymmetric particle (LSP) of supersymmetric (SUSY) extensions of the standard model of electroweak interactions, which is stable if R-parity is conserved.

The calculation of the LSP abundance in the universe has reached a very sophisticated level. Computer codes are now publicly available that take into account all the several LSP annihilation and co-annihilation processes that enter in the evaluation of its abundance today [3]. The comparison of the results of the computer codes with recent observations are placing strong constraints in the parameters of SUSY (actually, minimal supergravity models) [4]. In fact, accuracies of the order of 10% among differents codes are being sought [5].

We also know that the universe is accelerating today. There should exist a form of dark energy, comprising roughly 70% of the energy density of the universe, responsible for its acceleration. The simplest possibility, still consistent with cosmological data, is a cosmological constant. One tantalizing problem that arises in these models is the so-called coincidence problem: why dark energy starts to dominate the universe only at recent times? This has motivated the study of models in which dark energy is coupled to dark matter ¹.

In these models of coupled dark energy, the mass of the dark matter particle depends on the dark energy field and therefore it varies on a cosmological time scale. In this letter we point out in a general way what consequences this effect may have in the calculations of the cosmological abundance of cold dark matter, illustrating them with a particular simple model.

There are several different models of coupled dark energy, sometimes referred to as VAMPs (VAriable-Mass Particles) in the literature. The mass of the dark matter particles evolves according to some function of the dark energy field ϕ , as, for example, a linear function of the field [7, 8, 9, 10] with a inverse power law dark energy potential or an exponential function [11, 12, 13, 14, 15, 16] with an exponential dark energy potential. For instance, if the dark matter is a fermion one could have an interaction like $g(\phi)m_0\bar{\chi}\chi$, where the fermion mass $m_{\chi} = g(\phi)m_0$ is a function of the dark energy field. Since the dark energy field is dynamically evolving with time, one would have $m_{\chi} = m_{\chi}(\phi(a)) = m_{\chi}(a)$.

In order to have a rough idea of the possible magnitude of the mass-varying effect, we will follow a more phenomenological approach and assume that the dark matter particle mass

¹There are also recent models that couples dark energy to neutrinos so that the energy density of neutrinos tracks the dark energy density [6]

evolves with the cosmological scale factor a as [17]:

$$m_{\chi}(a) = m_{\chi}^{(0)} e^{\int_0^{\ln a} \delta(\alpha') d\alpha'}$$
(2)

where $m_{\chi}^{(0)}$ is the particle mass today and $\alpha = \ln a$. This results in the following equation for the evolution of the CDM energy density ρ_{CDM} :

$$\dot{\rho}_{CDM} + 3H\rho_{CDM} - \delta(a)H\rho_{CDM} = 0 \tag{3}$$

where $H = \dot{a}/a$ is the Hubble parameter. Conservation of the total stress-energy tensor them implies that the dark energy density should obey

$$\dot{\rho}_{DE} + 3H\rho_{DE}(1+w_{DE}) + \delta(a)H\rho_{CDM} = 0 \tag{4}$$

where w_{DE} is the equation of state of the dark energy fluid.

Many coupled dark energy models presents a so-called scaling solution where the dark matter and dark energy densities scale as simple functions of the scale factor. Following Majerotto et al. [17] we impose:

$$\frac{\rho_{DE}}{\rho_{CDM}} \propto a^{\xi} \tag{5}$$

This condition implies in the following expression for δ :

$$\delta = \frac{\delta_0}{\Omega_{DE}^0 + (1 - \Omega_{DE}^0)a^{-\xi}} \tag{6}$$

where

$$\delta_0 = -\Omega_{DE}^0(\xi + 3w_{DE}) \tag{7}$$

The free parameters of this model are taken to be Ω_{DE}^0 , w_{DE} and δ_0 . From a fit with the supernova Ia gold sample [18], Majerotto et al. obtain [17]:

$$\Omega_{DE}^{0} = 0.62; \quad w_{DE} = -1.9; \quad \delta_0 = -1.5.$$
(8)

which implies in $\xi = 8.1$.

With these values we can illustrate the effects of cold dark matter mass variation and we find that it saturates rather quickly for a < 0.1 at the value:

$$\frac{m_{\chi}(a)}{m_{\chi}^{(0)}} = 1.34\tag{9}$$

In order to estimate the consequences of the mass variation on the relic abundance of cold dark matter we have to solve the Boltzmann equation for the comoving number density Y = n/s, where n is the number density and s is the entropy density, as a function of the variable x = m(a)/T:

$$\frac{x}{Y_{EQ}}\frac{dY}{dx} = -\frac{\Gamma_A}{H}\left[\left(\frac{Y}{Y_{EQ}}\right)^2 - 1\right]$$
(10)

where Y_{EQ} is the equilibrium distribution and $\Gamma_A = n_{EQ} \langle \sigma_A v \rangle$ is the total annihilation width written in terms of the thermally averaged annihilation cross section times the relative velocity. The approximate solution of the Boltzmann equation gives the following relation for the CDM abundance today [19]:

$$\Omega_{CDM} \propto Y_{\infty} m_{\chi}^{(0)} \tag{11}$$

where Y_{∞} is the solution of the Boltzmann equation for large x, when the number density is frozen after decoupling. For an *s*-wave annihilation process, which we will consider below, $\langle \sigma_A v \rangle = \sigma_0$ and the solution can be well approximated by

$$Y_{\infty} \propto \frac{x_F}{m_{\chi}^{(F)} \sigma_0} \tag{12}$$

where x_F is the freeze-out value of the variable x and $m_{\chi}^{(F)}$ is the CDM mass at freeze-out. We have kept only terms which have dependence on the CDM mass. In fact, in the following we will neglect a logarithmic mass dependence in x_F to obtain

$$\Omega_{CDM} \propto \frac{m_{\chi}^{(0)}}{m_{\chi}^{(F)}} \frac{1}{\sigma_0}$$
(13)

At this point, the only particle physics assumption was that of CDM *s*-wave annihilation. Therefore, naively one would think that there is a simple linear difference due to the possibility of mass-variation.

However, the dependence can be potentially larger. As an example, we will consider a nonrelativistic annihilation process $\bar{\chi}\chi \to \bar{f}f$ with a vector interaction with a coupling constant g mediated by a vector boson of mass M_V . In this simple case, away from the resonance (for $M_V \gg m$) we have

$$\sigma_0(\bar{\chi}\chi \to \bar{f}f) = \frac{1}{2\sqrt{2\pi}} \frac{g^4}{M_V^4} (m_\chi^{(F)})^2 (1 + v^2 \beta_f^2/3)$$
(14)

where $\beta_f = \sqrt{1 - m_f^2 / (m_{\chi}^{(F)})^2}$. The important fact is the dependence on $(m_{\chi}^{(F)})^2$. Hence one would get:

$$\Omega_{CDM} = \left(\frac{m_{\chi}^{(0)}}{m_{\chi}^{(F)}}\right)^3 \Omega_{CDM}^{\text{c.m.}}$$
(15)

where $\Omega_{CDM}^{c. m.}$ is the usual result of the CDM abundance today with a constant mass particle.

Therefore, in this simple illustrative model a difference of a factor of $(1/1.34)^3 = 0.41$ in the cold relic energy density can be easily obtained. In these times of precision measurements and ruling out of parameter space for well defined models, the varying-mass phenomenom can be of importance. However, one should keep in mind that in realistic models there are several annihilation channels with different dependence on the CDM particle mass [20].

In summary, models that couple dark energy with dark matter predict mass-varying dark matter particles. Here we have exemplified the effect of mass variation in a simple model. Of course any realistic model can be solved to find the function m(a) and verify the effect on the abundance. It would be interesting to implement the possibility of dark matter mass variation in realistic computational codes of relic abundances.

Acknowledgments

I would like to thank Luca Amendola, Urbano França and Sasha Belyaev for useful comments. This work is partially supported by a CNPq grant.

References

- [1] D. N. Spergel et al. (WMAP Collaboration), Astrophys. J. Supp. 148, 175 (2003).
- [2] M. C. Bento, O. Bertolami and A. A. Sen, Phys. Rev. D66,043507 (2002); Phys. Rev. D70, 083519 (2004); J. C. Fabris, S. V. B. Goncalves and P. E. de Souza, Gen. Rel. Grav. 34, 53 (2002); M. Makler, S. Q. de Oliveira and I. Waga, Phys. Rev. D68, 123521 (2003); R. R. R. Reis, M. Makler and I. Waga, Phys. Rev. D69, 101301 (2004).
- P. Gondolo, J. Edsjo, P. Ullio, L. Bergstrom, M. Schelke and E. A. Baltz, JCAP 0407, 008 (2004); G. Belanger, F. Boudjema, A. Pukhov and A. Semenov, Comput. Phys. Commun. 149, 103 (2002) and hep-ph/0405253.
- [4] J. L. Feng, K. T. Matchev and F. Wilczek, Phys. Lett. B482, 388 (2000); H. Baer, C. Balazs and A. Belyaev, JHEP 0203, 042 (2002); J. R. Ellis, K A. Olive, Y. Santoso and V. C. Spanos, Phys. Lett. B565, 176 (2003); Phys. Rev. D70, 055005 (2004).
- [5] G. Belanger, S. Kraml and A. Pukhov, hep-ph/0502079.
- [6] R. Fardon, A. E. Nelson and N. Weiner, JCAP 0410, 005 (2004); X. Bi, B. Feng, H. Li and X. Zhang, hep-ph/0412002; R. D. Peccei, Phys. Rev. D71, 023527 (2005).
- [7] G. W. Anderson, S. M. Carrol, astro-ph/9711288 (unpublished).
- [8] J. A. Casas, J. García-Bellido, M. Quirós, Class. Quantum Grav. 9, 1371 (1992).
- [9] G. R. Farrar, P. J. E. Peebles, Astrophys. J. **604**, 1 (2004).
- [10] M. B. Hoffman, astro-ph/0307350 (unpublished).
- [11] L. Amendola, Phys. Rev. **D62**, 043511 (2000).
- [12] L. Amendola, D. Tocchini-Valentini, Phys. Rev. D64, 043509 (2001).
- [13] L. Amendola, Mon. Not. Roy. Astron. Soc. **342**, 221 (2003).

- [14] M. Pietroni, Phys. Rev. **D67**, 103523 (2003).
- [15] D. Comelli, M. Pietroni, A. Riotto, Phys. Lett. **B571**, 115 (2003).
- [16] U. França and R. Rosenfeld, Phys. Rev. **D69**, 063517 (2004).
- [17] E. Majerotto, D. Sapone and L. Amendola, astro-ph/0410543.
- [18] A. G. Riess *et al.*, Astrophys. J. **607**, 665 (2004).
- [19] E. W. Kolb and M. S. Turner, The Early Universe, Addison-Wesley (1994).
- [20] See, e.g., G. Jungman, M. Kamionkowski and K. Griest, Phys. Rep. 267, 195 (1996).