arXiv:astro-ph/0509538v2 15 May 2006

Neutron Star Equation of State and The Possibility of Complex Self-Energy in Landau Theory of Fermi Liquid in Presence of Strong Quantizing Magnetic Field

Soma Mandal^a), Roni Saha^a), Sutapa Ghosh^a) and Somenath Chakrabarty^b)

^{a)}Department of Physics, University of Kalyani, Kalyani 741 235, West Bengal, India

^{b)}Department of Physics, Visva-Bharati, Santiniketan 731 235,

West Bengal, India, E-mail:somenath@vbphysics.net.in

Using the relativistic version of Landau theory of Fermi liquid with $\sigma - \omega$ and ρ mesons exchange, we have obtained an equation state for dense neutron star matter in presence of strong quantizing magnetic field. It is found that in this scenario the self-energies of both neutron and proton components of dense neutron star matter become complex under certain physical conditions. To be more specific, it is observed that in the exchange diagrams of σ , ω and neutral ρ transfer processes and in the direct interaction diagram with ρ_{\pm} transfer reactions, the nucleon self-energies become complex in nature.

PACS numbers: 97.60.Jd, 97.60.-s, 75.25.+z

1. INTRODUCTION

With the recent discoveries of a few magnetars [1, 2, 3, 4], the study of dense neutron star matter in presence of strong quantizing magnetic field has gained a new life. These strange stellar objects are supposed to be strongly magnetized, relatively young (age ~ 10^4 yrs.) neutron stars and are also believed to be the possible sources of anomalous X-ray pulses and soft gamma ray emissions (AXP and SGR). As surface magnetic fields of these high magnetically-powered neutron stars are observed to be $\geq 10^{14} - 10^{15}$ G, then a theoretical estimation from virial theorem predicts that the field at the core region may be ~ 10^{18} G. Now, if the magnetic fields are really so strong, particularly at the central region, they must affect most of the physical properties of those strange stellar objects. Also, different physical processes taking place at the core region should get modified significantly.

The strong magnetic field at the core region affects the equation of state of the neutron star matter and as a consequence, some of the gross properties, e.g. mass-radius relation, moment of inertia, rotational frequency etc. of neutron stars should change to a great extent [5, 6, 7, 8].

In the case of compact neutron stars, the possibility of first order phase transition to quark matter at the core region is found to be completely forbidden if the magnetic field strength exceeds 10^{15} G. However, a second order phase transition is allowed, provided the field strength is $< 10^{20}$ G [9, 10].

We have noticed that the elementary processes, e.g. weak and electromagnetic reactions and decays are also strongly influenced by such intense magnetic field. As the cooling of neutron stars are mainly controlled by neutrino emissions through weak processes (URCA or modified URCA), the presence of strong quantizing magnetic field should significantly affect the thermal evolution of neutron stars. Further, the rates of the electromagnetic processes in presence of strong quantizing magnetic field should modify significantly the transport properties of dense stellar matter, in particular for electron gas present at the core region of the neutron stars [11, 12, 13, 14, 15]. We have noticed that the electron gas behaves like a neutral super fluid if the magnetic field strength exceeds 10^{16} G [16].

It has also been observed that chiral symmetry gets violated in presence of strong magnetic field. The strong quantizing magnetic field acts like a catalyst to generate the mass dynamically. A lot of investigations have already been done in this field [17, 18, 19, 20, 21, 22, 23, 24, 25].

In the general relativistic model some investigations have also been done to study the stability of neutron stars in presence of strong magnetic field. It is found that there are possibilities of geometrical deformations from the usual spherical shapes and in the extreme case of ultra strong magnetic field, they become either black disks or black strings [26, 27, 28].

In one of our recent studies, we have developed a relativistic formalism of Landau theory of Fermi liquid for dense neutron star matter in presence of strong quantizing magnetic field with $\sigma - \omega$ meson exchange [29]. In the present article we shall re-formulate our previous work as cited above, incorporating ρ -meson (both charged and neutral components) exchange interactions. To obtain the self-energies for both neutron and proton components from the Landau quasi-particle interaction functions considering nucleonic interactions via the exchange of ρ_{\pm} mesons, we have also included the effect of strong quantizing magnetic field on charged meson propagators.

To evaluate the self-energy of Landau quasi-particles, we have considered a typical many body fermionic system (in this case it is neutron star matter) in presence of strong quantizing magnetic field at zero temperature, interacting via scalar, vector and iso-vector meson fields (as mentioned above). This formalism is also useful to obtain the quasiparticle energy of other fermionic many body system, e.g., electron gas or even for exotic quark matter in presence

 $\mathbf{2}$

of strong magnetic field. The formalism is also applicable to study the magnetic properties and the non-equilibrium properties (e.g., the transport properties) of dense stellar plasma of fermions. Now similar to the conventional non-relativistic model of Landau theory of Fermi liquid, this modified version also deals with normal Fermi liquid and is applicable for the low-lying excited states of the system, which consists of a superposition of quasi-particle excitation close to the Fermi surface. The formalism we have developed in this article is basically a re-formulation of our previous work [29] and an extension of a very old paper by Baym and Chin [30]. Of course, in the present investigation, we have found a new and interesting result, the complex nature of nucleonic self energies in the mean field approximation.

The paper is arranged in the following manner. In the next section we are giving a general overview of the behavior of a fermionic system in a strong quantizing magnetic field. In Section 2, we shall consider a typical dense neutron star matter in presence of strong quantizing magnetic field interacting via $\sigma - \omega - \rho$ mesons exchange and develop a formalism to obtain the equation of state for dense neutron star matter in presence of strong quantizing magnetic fields. In the case of ρ_1 and ρ_2 (or ρ_{\pm} , the positive and negative components respectively) exchange interactions, we shall incorporate the effect of strong quantizing magnetic field on their propagators. In Section 3, we have concluded the results and discussed the future perspective of this work.

2. BASIC FORMALISM

1. Charged Fermion in Strong Magnetic Field

In the case of a typical many body fermionic system, let us consider the *i*th type fermion of charge q_i and mass m_i (of course, in the case of neutron component of neutron star matter, since $q_i = 0$, then one has to use the original work by Baym and Chin and the same formalism is also applicable if the magnetic field strength is less than the quantum critical value). We assume the gauge $A^{\mu} \equiv (0, 0, xB_m, 0)$, so that the constant magnetic field B_m is along z-axis. If the magnetic field strength B_m exceeds the quantum critical value $B_m^{(c)(i)} = m_i^2/q_i$, the Landau levels for the *i*th charged component will be populated and the matter is called the Landau diamagnetic system. In the relativistic region the charged matter shows Landau diamagnetic behavior, if the cyclotron quantum of the charged components exceeds the corresponding rest mass energy or equivalently the de Broglie wave length exceeds the Larmor radius of the charged particle. Under such circumstances the phase space integral will get modified and is given by

$$\frac{g_i}{(2\pi)^3} \int d^3 p F(p) = \frac{g_i}{(2\pi)^3} \int dp_z d^2 p_\perp F(p) \longrightarrow \frac{g_i q_i B_m}{(2\pi)^2} \sum_{\nu=0}^{\infty} (2 - \delta_{\nu_0}) \int_{-\infty}^{+\infty} dp_z F(p_z, \nu)$$
(1)

where g_i is the degeneracy, $\nu = 0, 1, 2, ...$ is the Landau quantum number and F is a function of particle momentum. At zero temperature, the above modified form reduces to

$$\longrightarrow \frac{g_i q_i B_m}{(2\pi)^2} \sum_{\nu=0}^{[\nu_{max}^{(i)}]} (2 - \delta_{\nu_0}) \int_{-p_{F_i}}^{+p_{F_i}} dp_z F(p_z, \nu)$$
(2)

where

$$[\nu_{max}^{(i)}] = \frac{(\mu_i^2 - m_i^2)}{2q_i B_m} \tag{3}$$

is an integer but less than the actual value of right hand side, μ_i is the chemical potential of the *i*th charged component and p_{F_i} is the corresponding Fermi momentum. Now it can very easily be shown from the solution of Dirac equation in presence of such strong magnetic field that the zeroth Landau level is singly degenerate, whereas all other levels are doubly degenerate (which is incorporated in eqn.(1) by introducing the factor $2 - \delta_{\nu 0}$). Further, the energy eigen value is given by

$$E_{\nu} = (p_z^2 + m_i^2 + 2\nu q_i B_m)^{1/2} = (p_z^2 + m_i^2 + p_{\perp,i}^2)^{1/2}$$
(4)

and therefore the chemical potential is simply given by

$$\mu_i = (p_{F_i}^2 + m_i^2 + 2\nu q_i B_m)^{1/2} \tag{5}$$

with $p_{\perp,i} = (2\nu q_i B_m)^{1/2}$ the orthogonal component of charged particle momentum (in the x - y plane), which varies in a discrete manner, hence the name, Landau quantization.

In presence of strong quantizing magnetic field as discussed above, it can very easily be shown that in standard notation, the modified form of Dirac spinors for the charged components are given by

$$\psi = \frac{1}{(L_y L_z)^{1/2}} \exp(-iE_\nu t + ip_y y + ip_z z) u^{\uparrow\downarrow}$$
(6)

where

$$u^{\uparrow} = \frac{1}{[E_{\nu}(E_{\nu}+m)]^{1/2}} \begin{pmatrix} (E_{\nu}+m)I_{\nu;p_{y}}(x) \\ 0 \\ p_{z}I_{\nu;p_{y}}(x) \\ -i(2\nu q_{i}B_{m})^{1/2}I_{\nu-1;p_{y}}(x) \end{pmatrix}$$
(7)

and

$$u^{\downarrow} = \frac{1}{[E_{\nu}(E_{\nu}+m)]^{1/2}} \begin{pmatrix} 0\\ (E_{\nu}+m)I_{\nu-1;p_{y}}(x)\\ i(2\nu q_{i}B_{m})^{1/2}I_{\nu;p_{y}}(x)\\ -p_{z}I_{\nu-1;p_{y}}(x) \end{pmatrix}$$
(8)

where the symbols \uparrow and \downarrow indicate the up and the down spin states respectively and

$$I_{\nu} = \left(\frac{qB_m}{\pi}\right)^{1/4} \frac{1}{(\nu!)^{1/2}} 2^{-\nu/2} \exp\left[-\frac{1}{2}qB_m\left(x - \frac{p_y}{qB_m}\right)^2\right] \\ H_{\nu}\left[(qB_m)^{1/2}\left(x - \frac{p_y}{qB_m}\right)\right]$$
(9)

with H_{ν} the well-known Hermite polynomial of order ν and L_y , L_z are length scales along y and z directions respectively.

Now for a dense relativistic many body system of fermions interacting via scalar, vector and iso-vector fields, represented by ϕ , V^{μ} and $\vec{\rho}^{\mu}$ with masses ζ , η and θ respectively, the basic Lagrangian density is given by

$$\mathcal{L} = \frac{i}{2}\bar{\psi}\gamma^{\mu}(\vec{D}_{\mu} - \overleftarrow{D}_{\mu})\psi - m\bar{\psi}\psi - g_{v}\bar{\psi}\gamma^{\mu}V_{\mu}\psi - g_{s}\bar{\psi}\phi\psi - g_{\rho}\bar{\psi}\gamma^{\mu}\rho_{\mu}.\frac{\vec{\tau}}{2}\psi + \mathcal{L}_{V} + \mathcal{L}_{S} + \mathcal{L}_{I-V} + \mathcal{L}_{EM}$$
(10)

where $D^{\mu} = \partial^{\mu} + iqA^{\mu}$, g_v , g_s and g_{ρ} are the coupling constants corresponding to vector, scalar and iso-vector interactions; \mathcal{L}_V , \mathcal{L}_S , \mathcal{L}_{I-V} are the free Lagrangian densities for vector, scalar and iso-vector field respectively and \mathcal{L}_{EM} is the Lagrangian density electromagnetic part corresponding to constant external magnetic field B_m along z-axis. Here the iso-vector fields behaves like a four vector in configuration space (represented by $\mu = 0, 1, 2, 3$) but like a three vector in iso-spin space (represented by the symbol \rightarrow).

Now in the Landau theory of Fermi liquid, to evaluate various physical quantities of the system, it is a common practise to switch on the interactions adiabatically. The normal Fermi liquid system will then evolve from noninteracting states to interacting states and there will be a one-to-one correspondence between the dressed or quasiparticle states of the interacting system to the bare non interacting states. To obtain f, the Landau Fermi liquid interaction function, also known as the quasi-particle interaction function, we have considered both direct and exchange type interactions. The exchanged bosons are either scalar or vector or iso-vector type. The quasi-particle interaction functions are related to the two-particle forward scattering via the exchange of any one of these bosons. We further assume that the external magnetic field acts as a constant background and is present everywhere. In the next subsection we consider the processes with scalar boson exchange.

2. Scalar Boson Exchange

To compute the Landau-Fermi liquid interaction function f from two particle forward scattering matrix with scalar boson (σ) exchange, we start with the definition of transition matrix element, given by

$$T_{fi} = i \int \rho_s(x)\phi(x)d^4x \tag{11}$$



FIG. 1: Basic Elementary Processes

where

$$\rho_s(x) = g_s \bar{\psi}(x) \psi(x) \tag{12}$$

is the scalar density and $\phi(x)$ is the scalar field satisfies the the well known Klein-Gordan equation

$$(\partial^2 + m^2)\phi(x) = \rho_s(x) \tag{13}$$

The solution of this equation can be obtained by Green's function technique and is given by

$$\phi(x) = -\int \frac{d^4q}{(2\pi)^4} \frac{\exp[-iq.(x-x')]}{q^2 - \zeta^2} \rho_s(x') d^4x'$$
(14)

where $q^{\mu} \equiv (q^0, \vec{q})$ is the exchanged four momentum.

Now to evaluate T_{fi} , we substitute the Green's function solution (eqn.14) into eqn.(12) to obtain $\rho_s(x)$ and finally putting it into eqn.(11). We consider the following strong interaction processes:

 $\begin{array}{ll} (i) & p+p \longrightarrow p+p, \\ (ii) & n+n \longrightarrow n+n \ \text{and} \\ (iii) & p+n \longrightarrow p+n. \end{array}$

The basic diagrams are as follows: We evaluate T_{fi} for both direct and exchange type interactions for all these processes. We first consider the scattering process represented by (i). The transition matrix element is given by

$$T_{fi} = -i \int \rho_s(x) \frac{d^q}{(2\pi)^4} \, \frac{\exp[-iq(x-x')]}{q^2 - \zeta^2} \rho_s(x') d^4x d^4x' \tag{15}$$

Now using the up and down spin positive energy spinors for proton, we have (since the temperature of the system \ll the typical chemical potential for nucleons ~ 1GeV, we have not considered the negative energy states).

$$\rho_s(x') = \frac{g_s}{L_y L_z} \exp[-i\{(E_{\nu_1} - E_{\nu_2})t' - (p_{1y} - k_{1y})y' - (p_{1z} - k_{1z})z'\}][\rho_s(x')]$$
(16)

Then substituting for $\phi(x)$, $\rho_s(x)$ and $\rho_s(x')$ and integrating over t', y' and z', we have the direct part for p - p (represented by (i)) scattering matrix element.

$$T_{fi,(pp\to pp)}^{(d,s)} = -i \int \frac{d^4q}{2\pi} \delta(q^0 - E_{\nu_1} + E_{\nu_2}) \delta(q_y - p_{1y} + k_{1y}) \delta(q_z - p_{1z} + k_{1z}) \frac{\exp(iq.x)}{q^2 - \zeta^2} \times [\rho(x')] \frac{g_s}{L_y L_z} \exp(iq_x x') dx' \frac{g_s}{L_y L_z} [\rho(x)] d^4x \\ \exp[i\{(E_{\nu'_1} - E_{\nu'_2})t - (p_{2y} - k_{2y})y - (p_{2z} - k_{2z})z\}]$$
(17)

Next integrating over t, y, z and q^0 , q_y , q_z we have

$$T_{fi,(pp\to pp)}^{(d,s)} = -i(2\pi)^{3}\delta(E)\delta(p_{y})\delta(p_{z})\frac{g_{s}^{2}}{(L_{y}L_{z})^{2}}\int \frac{dq_{x}}{2\pi}[\rho(x)][\rho(x')]\exp[-iq_{x}(x-x')] \times \frac{1}{q^{2}-\zeta^{2}}dxdx'$$
(18)

where the δ functions represent the abbreviated form of energy and momentum conservations along y and z directions. Finally, evaluating the contour integral over q_x and using the conditions for forward scattering given by $E_{\nu_1} = E_{\nu_2} = E_{\nu}$ (say); $E_{\nu'_1} = E_{\nu'_2} = E_{\nu'}$ (say); $p_{1y} = k_{1y} = p_y$ (say); $p_{2y} = k_{2y} = p'_y$ (say); $p_{1z} = k_{1z} = p_z$ (say) and $p_{2z} = k_{2z} = p'_z$ (say), we have the Landau-Fermi liquid interaction function corresponding to direct interaction

$$f_{pp \to pp}^{(d,s)} = \frac{g_s^2}{8\zeta} \int \exp[-\zeta |x - x'|] [\rho(x)] [\rho(x')] dx dx'$$
(19)

where

$$[\rho_s(x)] = \bar{u}(x, \nu_2, k_1)u(x, \nu_1, p_1)$$
(20)

Then summing over the final spin states and averaging over the initial spin states we have for direct diagram

$$[\rho_s(x)] = \frac{1}{2} \operatorname{Tr}[\Lambda(x,\nu_1,\nu_2,p_1,k_1)]$$
(21)

and similarly for x'. Whereas for the exchange diagram

$$[\rho_s(x)][\rho_s(x')] = \frac{1}{4} \operatorname{Tr}[\Lambda(x,\nu_1,\nu_2,p_1,k_1)\Lambda(x,\nu'_1,\nu_2i',p_2,k_2)]$$
(22)

where

$$\Lambda(x,\nu_1,\nu_2,p_1,k_1) = \sum_{\text{spin}} u(x,\nu_1,p_1)\bar{u}(x,\nu_2,k_2)$$
(23)

To obtain Λ 's for the process indicated by (i), we consider the direct diagram for forward scattering. Then substituting the up and down spin solutions of Dirac equation for proton (eqns.(7) and (8)) in the expression for projection operator (eqn.(23)), we get

$$\Lambda = \frac{1}{2E_{\nu}} [Ak_{\mu}\gamma^{\mu}(\mu = 0 \text{ and } z) + mA + Bk_{\mu}\gamma^{\mu}(\mu = y \text{ and } p_{y} = p_{\perp})]$$
(24)

The matrices A and B are given by

$$A = \begin{pmatrix} I_{\nu}I'_{\nu} & 0 & 0 & 0\\ 0 & I_{\nu-1}I'_{\nu-1} & 0 & 0\\ 0 & 0 & I_{\nu}I'_{\nu} & 0\\ 0 & 0 & 0 & I_{\nu-1}I'_{\nu-1} \end{pmatrix}$$
(25)

$$B = \begin{pmatrix} I_{\nu-1}I'_{\nu} & 0 & 0 & 0\\ 0 & I_{\nu}I'_{\nu-1} & 0 & 0\\ 0 & 0 & I_{\nu-1}I'_{\nu} & 0\\ 0 & 0 & 0 & I_{\nu}I'_{\nu-1} \end{pmatrix}$$
(26)

where the primes indicate the functions of x'. To the best of our knowledge, eqn.(24) is an entirely new result, first obtained in our previous publication [29] and has not been reported earlier. Further, these results play key roles in all kinds of calculation related to the electromagnetic and weak interactions in presence of strong quantizing magnetic field. In a recently completed work [25] we have used this results to study chiral symmetry violation in presence of strong magnetic field.

Since γ matrices are traceless and both A and B matrices are diagonal with identical blocks, it is very easy to evaluate the traces of the product of γ -matrices multiplied with any number of A and/or B, from any side with any order e.g.,

$$Tr(\gamma^{\mu}\gamma^{\nu}A_{1}A_{2}..B_{1}B_{2}..) = Tr(A_{1}A_{2}..B_{1}B_{2}..)g^{\mu\nu},$$
(27)

$$\operatorname{Tr}(\gamma^{\mu}\gamma^{\nu}\gamma^{\lambda}\gamma^{\sigma}A_{1}A_{2}..B_{1}B_{2}..) = \operatorname{Tr}(A_{1}A_{2}..B_{1}B_{2}..)(g^{\mu\nu}g^{\sigma\lambda} - g^{\mu\lambda}g^{\nu\sigma} + g^{\mu\lambda}g^{\nu\sigma}),$$
(28)

Tr(product of odd γ s with A and/or B) = 0 etc. The other interesting aspects of A and B matrices are: i) $k_{1\mu}k^{2\mu}\text{Tr}(A_1A_2) = (E_1E_2 - k_{1z}k_{2z})\text{Tr}(A_1A_2)$

- ii) $k_{1\mu}k^{2\mu}\text{Tr}(B_1B_2) = \vec{k}_{1\perp}.\vec{k}_{2\perp}\text{Tr}(B_1B_2)$
- iii) $k_{1\mu}k^{2\mu}\text{Tr}(A_1B_2) = k_{1\mu}k^{2\mu}\text{Tr}(B_1A_2) = 0$
- iv) $p_{1\mu}k^{1\mu}p_{2\nu}k^{2\nu}\operatorname{Tr}(A_1B_2) \neq 0 = (E_{\nu_1}E_{\nu'_2} p_{1z}k_{1z})\vec{p}_{2\perp}.\vec{k}_{2\perp}\operatorname{Tr}(A_1B_2)$

These set of relations are also totally new results and have been derived in our previous publication [29].

Then we have following Baym and Chin [30]

$$f_{pp\to pp}^{(d,s)} = -\frac{g_s^2 m^2}{16\zeta E_{\nu} E_{\nu'}} \int \exp[-\zeta |x - x'|] \text{Tr} A \text{Tr} A' dx dx'$$
(29)

where

$$\operatorname{Tr} A = 2(I_{\nu}^2 + I_{\nu-1}^2) \text{ and } \operatorname{Tr} A' = 2(I_{\nu'}^2 + I_{\nu'-1}^2)$$
 (30)

On the other hand if we consider the direct interaction corresponding to the process (ii), we have

$$f_{nn \to nn}^{(d,s)} = -\frac{g_s^2 m^2}{\zeta^2 E E'}$$
(31)

This result is same as that obtained by Baym and Chin long ago.

Next we consider a bit complicated case; scattering process represented by (iii). In this case only protons are affected by strong quantizing magnetic field, their Landau levels are populated. The transition matrix element is then given by

$$T_{fi} = i \int \rho_s^p(x)\phi(x)d^4x \tag{32}$$

where $\rho_s^{(p)}(x)$ is the scalar density corresponding to proton component and is given by eqn.(20). The scalar field

$$\phi(x) = -\int \frac{d^4q}{(2\pi)^4} \frac{exp[-iq.(x-x')]}{q^2 - \zeta^2} \rho_s^{(n)}(x') d^4x'$$
(33)

where the scalar density corresponding to neutron component is given by

$$\rho_s^{(n)}(x) = g_s \bar{\psi}_n(x) \psi_n(x) = \frac{g_s}{V} \exp[-i(p_2 - k_2) \cdot x] [\rho_s^{(n)}], \qquad (34)$$

where

$$[\rho_s^{(n)}] = \bar{u}(k_2)u(p_2),\tag{35}$$

here u and \bar{u} are the usual form of spinor and the adjoint for neutron, given by

$$u^{(s)} = \frac{1}{2E} \left(\begin{array}{c} \chi^{(s)} \\ \frac{\vec{\sigma} \cdot \vec{p}}{E+m} \chi^{(s)} \end{array} \right)$$
(36)

where s = 1 and 2 corresponding to the up and down spin cases respectively, and are given by

$$\chi^{(1)} = \chi^{\uparrow} = \begin{pmatrix} 1\\0 \end{pmatrix} \tag{37}$$

and

$$\chi^{(2)} = \chi^{\downarrow} = \begin{pmatrix} 0\\1 \end{pmatrix} \tag{38}$$

Using all these expressions and integrating over x, we have

$$T_{fi(p+n\to p+n)}^{(d,s)} = -i \int d^4q \frac{\exp[iq.x']}{q^2 - \zeta^2} \frac{g_s}{V} [\rho_s^{(n)}] \frac{g_s}{L_y L_z} \exp[-i\{(E_{\nu_1} - E_{\nu_2})t' - i(p_{1y} - k_{1y})y' - i(p_{1z} - k_{1z})z'\}] \\ [\rho_s^{(p)}] \delta^4(q - p_2 + k_2) d^4x'$$
(39)

Further integrating over t', y', z' and q_x , q_y , q_z , q_0 finally we have for the forward scattering case

$$f_{p+n \to p+n}^{(d,s)} = -\frac{g_s^2}{\zeta^2 E_\nu E} m^2$$
(40)

where we have used the same expression as before for $[\rho_s^{(p)}(x)]$ whereas

$$[\rho_s^{(n)}] = \frac{1}{2E} (p_\mu \gamma^\mu + m) \tag{41}$$

Here we have also used some obvious results as given below to obtain the above expression (eqn.(40)).

(a)
$$\operatorname{Tr}(\not p + m) = 4m$$
, (b) $\operatorname{Tr}(\Lambda) = m \operatorname{Tr} A'$ and (c) $\int_{-\infty}^{+\infty} \operatorname{Tr} A' dx' = 4$ (42)

Although the structural forms of both eqns.(40) and (31) are identical, the appearance of proton energy in the denominator of eqn.(40) makes it qualitatively different from eqn.(31). This is also obvious from the physics point of view. The interaction with σ -meson exchange can not distinguish neutron and proton. The difference comes from the change in proton parameters (spinor wave functions, energy eigen values etc.) in presence of strong quantizing magnetic field. As we will see later that in the evaluation of energy densities for proton and neutron separately the form of momentum integrals (phase space integrals) will differ significantly (which has also been discussed in the previous section).

We shall next evaluate the Landau interaction function for exchange part with σ -meson exchange. In this case the processes represented by (i) and (ii) give results in which there are nothing surprising. Whereas, as we shall show that the Landau interaction function obtained from the process (iii) is complex in nature. Which will give complex self energy of both proton and neutron. This is also true for exchange diagrams with vector boson and neutral iso-vector boson exchange, whereas in the case of charged iso-vector exchange, the direct term gives complex Landau interaction function and as a result the complex self-energies. The findings are to the best of our knowledge are new and not been reported earlier.

We start with the process (i). In this case the forward scattering condition gives $E_{\nu_1} = E_{\nu'_2} = E_{\nu}$ (say), $E_{\nu'_1} = E_{\nu_2} = E_{\nu'}$ (say), $p_{1y} = k_{2y} = p_y$ (say), $p_{1z} = k_{2z} = p_z$ (say), $p_{2y} = k_{1y} = p'_y$ (say), $p_{2z} = k_{1z} = p'_z$ (say). Then we have the Landau interaction function

$$f_{p+p\to p+p}^{(ex,s)} = \frac{g_s^2}{32E_{\nu}E_{\nu'}} \int dx dx' \frac{\exp[-k|x-x'|]}{K} [(E_{\nu}E_{\nu'} - p_z p_{z'} + m^2) \operatorname{Tr}(AA') - \vec{p_{\perp}}.\vec{p_{\perp}'}.\operatorname{Tr}(BB')]$$
(43)

where $K = (q_y^2 + q_z^2 + \zeta^2)^{1/2}$, $q_i = (p_i - p'_i)$ with i = y and z. The traces are obtained from eqns.(25) and (26). Similarly for the process (*ii*), the Landau quasi-particle interaction function is given by

$$f_{n+n\to n+n}^{(ex,s)} = \frac{g_s^2}{4EE'} \frac{p \cdot p' + m^2}{(p'-p)^2 - \zeta^2}$$
(44)

and is identical with zero field result of Baym and Chin.

Next we consider the exchange term corresponding to the process (iii) with scalar boson transfer. In this case the transition matrix element is given by

$$T_{fi} = i \int \rho_s^{(pn)}(x)\phi(x)d^4x \tag{45}$$

where

$$\rho_s^{(pn)}(x) = g_s \bar{\psi}_p(x) \psi_n(x)$$

= $\frac{1}{L_y L_z} \exp[-i\{(E_p - E_k)t - (p_y - k_y)y - (p_z - k_z)z\}] \exp(ip_x x) [\rho_s^{(pn)}(x)]$ (46)

and

$$\phi(x) = -\int \frac{d^4q}{(2\pi)^4} \, \frac{\exp[-iq(x-x')]}{q^2 - \zeta^2} \rho_s^{(np)}(x'd^4x' \tag{47}$$

Hence

$$T_{fi} = -i \int \frac{d^q}{(2\pi)^4} \frac{g_s}{L_y L_z} \exp[-i\{(E_{\nu_1} - E_{\nu'_2})t - (p_{1_y} - k_{2_y})y - (p_{1_z} - k_{2_z})z\}][\rho_s^{(pn)}(x)]$$

$$\frac{\exp[-iq(x - x')]}{q^2 - \zeta^2} \frac{g_s}{L_y L_z}[\rho_s^{(np)}(x')]$$

$$\exp[i\{(E_{\nu'_1} - E_{\nu_2})t' - (p_{2_y} - k_{1_y})y - (p_{2_z} - k_{1_z})z\}]\exp[-i(p_{1_x} - k_{1_x})x]d^4x d^4x'$$
(48)

Now integrating over $t, y, z, t', y', z', q_0, q_x, q_y$ and q_z , we have the Landau quasi-particle interaction function with the condition of forward scattering

$$f_{p+n\to p+n}^{(ex,s)} = \frac{g_s^2}{4} \int dx dx' \frac{\exp[i(q_x - p_x + p'_x)(x - x')]}{(p - p')^2 - \zeta^2} \operatorname{Tr}[\Lambda_{pn}(x, p)\Lambda_{np}(x', p')]$$
(49)

Now to evaluate the projection operators Λ_{np} and Λ_{pn} , we define

$$p_x + ip_y = p_\perp \exp(i\phi)$$
 and $p_x - ip_y = p_\perp \exp(-i\phi)$ (50)

Then from the ranges of p_x and p_y , both of which vary from $-\infty$ to $+\infty$ and the range of p_{\perp} , which is given by $(2\nu q B_m)^{1/2}$, with $0 \le \nu \le \nu_{\max}$ where ν_{\max} is a function of density of matter and the strength of magnetic field, we have $0 \le \phi \le \pi$. With these choices, we have

$$\Lambda_{np}(x,p) = u_n^{\uparrow} \bar{u}_p^{\uparrow} + u_n^{\downarrow} \bar{u}_p^{\downarrow} = \frac{1}{2E} \begin{pmatrix} I_{\nu}(E+m) & 0 & -p_z I_{\nu} & -ip_{\perp} I_{\nu-1} \\ 0 & I_{\nu-1}(E+m) & -ip_{\perp} I_{\nu} & p_z I_{\nu-1} \\ p_z I_{\nu} & p_{\perp} e^{-i\phi} I_{\nu-1} & \frac{(ip_{\perp}^2 e^{-i\phi} - p_z^2)}{E+m} I_{\nu} & \frac{p_z p_{\perp}(e^{-i\phi} - i)}{E+m} I_{\nu-1} \\ p_{\perp} e^{i\phi} I_{\nu} & -p_z I_{\nu-1} & -\frac{p_z p_{\perp}(e^{i\phi} + i)}{E+m} I_{\nu} & -\frac{(p_z^2 + ip_{\perp}^2 e^{i\phi}}{E+m} I_{\nu-1} \end{pmatrix}$$
(51)

Which after some simplification reduces to

$$\Lambda_{np} = \frac{1}{2E} \Big[(E\gamma_0 + m - p_z\gamma_z)\mathcal{A} + \frac{p_\perp^2}{E+m} \frac{\mathcal{A}}{2} \Sigma_z (1-\gamma_0) - \frac{p_z p_\perp}{2(E+m)} (1-\gamma_0) \Sigma_{yz} \mathcal{A} \\ + \frac{iP_\perp^2}{E+m} \frac{\mathcal{A}}{2} (1-\gamma_0) X + \frac{p_z p_\perp}{E+m} \Sigma_{zx} X^* \frac{\mathcal{A}}{2} (1-\gamma_0) \\ - ip_\perp \frac{(1+\gamma_0)}{2} \gamma_x \mathcal{A} - p_\perp \gamma_x \frac{\mathcal{A}}{2} (1+\gamma_0) X^* \Big]$$

$$(52)$$

where

$$X = \begin{pmatrix} e^{-i\phi} & 0 & 0 & 0\\ 0 & e^{i\phi} & 0 & 0\\ 0 & 0 & e^{-i\phi} & 0\\ 0 & 0 & 0 & e^{i\phi} \end{pmatrix},$$
(53)

9

$$\Sigma_{ij} = \begin{pmatrix} \sigma_i \sigma_j & 0\\ 0 & \sigma_i \sigma_j \end{pmatrix},\tag{54}$$

$$\mathcal{A} = \begin{pmatrix} I_{\nu} & 0 & 0 & 0\\ 0 & I_{\nu-1} & 0 & 0\\ 0 & 0 & I_{\nu} & 0\\ 0 & 0 & 0 & I_{\nu-1} \end{pmatrix},\tag{55}$$

and

$$\Sigma_i = \begin{pmatrix} \sigma_i & 0\\ 0 & \sigma_i \end{pmatrix} \tag{56}$$

whereas, γ 's are Dirac matrices in standard representation and σ_i are the usual Pauli spin matrices. Similarly

$$\Lambda_{pn}(x,p) = u_{p}^{\uparrow} \bar{u}_{n}^{\uparrow} + u_{p}^{\downarrow} \bar{u}_{n}^{\downarrow}
= \frac{1}{2E} \begin{pmatrix} I_{\nu}(E+m) & 0 & -p_{z}I_{\nu} & -p_{\perp}e^{-i\phi}I_{\nu} \\ 0 & I_{\nu-1}(E+m) & -p_{\perp}e^{i\phi}I_{\nu-1} & p_{z}I_{\nu-1} \\ p_{z}I_{\nu} & ip_{\perp}I_{\nu} & -\frac{[(E^{2}-m^{2})-p_{\perp}^{2}(ie^{i\phi}-1)]}{E+m}I_{\nu} & \frac{p_{z}p_{\perp}(i-e^{-i\phi})}{E+m}I_{\nu} \\ -ip_{\perp}I_{\nu-1} & -p_{z}I_{\nu-1} & \frac{p_{z}p_{\perp}(i+e^{i\phi})}{E+m}I_{\nu-1} & \frac{[p_{\perp}^{2}(ie^{-i\phi}+1)-(E^{2}-m^{2})]}{E+m}I_{\nu-1} \end{pmatrix}$$
(57)

After some simplification as before this can also be written as

$$\Lambda_{pn} = \frac{1}{2E} \Big[(E\gamma_0 + m - p_z\gamma_z)\mathcal{A} + \frac{p_\perp^2}{2(E+m)} (1-\gamma_0)\Sigma_z\mathcal{A} + \frac{ip_zp_\perp}{2(E+m)} (1-\gamma_0)\Sigma_x\mathcal{A} \\ + \frac{ip_\perp^2}{E+m} \frac{\mathcal{A}}{2} (1-\gamma_0)X^* + \frac{p_zp_\perp}{E+m}\Sigma_{yx}X\frac{\mathcal{A}}{2} (1-\gamma_0) \\ + p_\perp\mathcal{A}\Sigma_xX^* - \frac{(1-\gamma_0)}{2}\Sigma yz\mathcal{A} \Big]$$
(58)

Since diagonal elements for Σ_i , Σ_{ij} and γ_i with $i \neq j$ and i, j = x, y, z, are zero, the traces of all the quantities multiplied with these matrices are identically zero. Therefore in both Λ_{np} and Λ_{pn} only the first and the fourth terms will contribute in trace calculation. As a result the Landau quasi-particle interaction function will become complex in nature. The explicit form of the trace term is given by

$$Tr \left[\Lambda_{np}(p')\Lambda_{pn}(p)\right] = \frac{1}{E_{\nu}E_{\nu'}}\left\{\left(E_{\nu}E_{\nu'} + p_{z}p'_{z} + m^{2}\right)2\left(I_{\nu}I_{\nu'} + I_{\nu-1}I_{\nu'-1}\right) - \frac{p_{\perp}^{2}p'_{\perp}^{2}}{4(E_{\nu}+m)(E_{\nu'}+m)} - \frac{iE_{\nu}p'_{\perp}^{2}}{2(E_{\nu'}+m)}2\left(I_{\nu}I_{\nu'}\exp(i\phi) + I_{\nu-1}I_{\nu'-1}\exp(-i\phi)\right) + \frac{imp'_{\perp}^{2}}{2(E_{\nu'}+m)}2\left(I_{\nu}I_{\nu'}\exp(i\phi) + I_{\nu-1}I_{\nu'-1}\exp(-i\phi)\right) - \frac{iE_{\nu'}p_{\perp}^{2}}{2(E_{\nu}+m)}2\left(I_{\nu}I_{\nu'}\exp(-i\phi) + I_{\nu-1}I_{\nu'-1}\exp(i\phi)\right) + \frac{imp_{\perp}^{2}}{2(E_{\nu}+m)}2\left(I_{\nu}I_{\nu'}\exp(-i\phi) + I_{\nu-1}I_{\nu'-1}\exp(i\phi)\right) + \frac{imp_{\perp}^{2}}{2(E_{\nu}+m)}2\left(I_{\nu}I_{\nu'}\exp(-i\phi) + I_{\nu-1}I_{\nu'-1}\exp(i\phi)\right)\right]$$
(59)

Obviously the last four terms in the above expression are complex in nature.

Now the self-energies for proton corresponding to the direct and exchange diagrams of process (i) is given by

$$\varepsilon_{\nu}^{(d/ex)}(p_z) = E_{\nu}(p_z) + \frac{1}{(2\pi)^2} \sum_{\nu'=0}^{[\nu_{\max}]} (2 - \delta_{\nu'0}) \int_{p'_y = -\infty}^{\infty} \int_{p_{z'} = -p_f}^{p_f} dp_y dp_{y'} dp_{z'} f_{pp \to pp}^{(d/ex,s)}$$

= $E_{\nu}(p_z) + \Delta E_{\nu;\text{Dir./ex}}^{(s)}(p_z)$ (60)

For the process (ii) the neutron self-energy corresponding to direct or exchange digram is given by

$$\varepsilon^{(d/ex)}(p) = E(p) + \frac{g_n}{(2\pi)^3} \int f_{n+n \to n+n}^{(d/ex,s)} d^3 p' = E(p) + \Delta E_{\text{Dir./ex}}^{(s)}(p)$$
(61)

Next for the process (*iii*) with direct diagram, the proton self-energy is given by

$$\varepsilon_{\nu}^{(d)}(p_z) = E_{\nu}(p_z) + \frac{eB_m}{2\pi^2} \sum_{\nu'=0}^{|\nu_{\max}|} (2 - \delta_{\nu'0}) \int f_{p+n \to p+n}^{(d,s)} dp'_z$$

= $E_{\nu}(p_z) + \Delta E_{\nu;\text{Dir./ex}}^{(s)}(p_z)$ (62)

Similarly for neutron, it is given by

$$\varepsilon^{(d)}(p) = E(p) + \frac{g_n}{(2\pi)^3} \int f_{p+n \to p+n}^{(d,s)} d^3 p' = E(p) + \Delta E_{\text{Dir.}}^{(s)}(p)$$
(63)

Finally the exchange part of self-energy corresponding to process (iii) are same for both proton and neutron and it is given by

$$\varepsilon_{\nu}^{(ex)}(p_{z}) = E_{\nu}(p_{z}) + \sum_{\nu'=0}^{[\nu_{\max}]} (2 - \delta_{\nu'0}) \int_{0}^{\pi} d\phi d\phi' p_{\perp} p_{\perp}' \cos(\phi) \cos(\phi') \int_{-p_{F}}^{+p_{F}} dp_{z}' f_{p+n \to p+n}^{(ex,s)}$$

$$= E_{\nu}(p_{z}) + \Delta E_{\nu;ex}^{(s)}(p_{z})$$
(64)

This last expression for self energy for both proton and neutron is complex in nature. As we will see in the subsequent sections that similar results can also be obtained with vector and iso-vector exchange type interactions.

3. Vector Boson Exchange

We shall now consider the vector boson exchange type interaction. The transition matrix element is given by

$$T_{fi} = i \int j^{\mu}(x) V_{\mu}(x) d^4x$$
(65)

where

$$j^{\mu}(x) = g_v \bar{\psi}(x) \gamma^{\mu} \psi(x) \tag{66}$$

is the particle current and the vector $V^{\mu}(x)$ satisfies the Klein-Gordan equation given by

$$(\partial^2 + \eta^2) V^{\mu}(x) = j^{\mu}(x) \tag{67}$$

Hence the Green's function solution for $V^{\mu}(x)$ is given by

$$V^{\mu}(x) = -\int \frac{d^4q}{(2\pi)^4} \frac{\exp[-iq.(x-x')]}{q^2 - \eta^2} j^{\mu}(x') d^4x'$$
(68)

where η is the vector boson mass.

Let us first consider the direct term of the process (i). Using the spinor solutions for proton (eqns.(7) and (8)), we have

$$j^{\mu}(x) = \frac{g_s}{L_y L_z} \exp[-i\{(E_{\nu_1} - E_{\nu_2})t - (p_{1y} - k_{1y})y - (p_{1z} - k_{1z})z\}][j^{\mu}]$$
(69)

where in the direct term we define

$$[j^{\mu}] = \frac{1}{2} \sum_{\text{spin}} \bar{u}(x', \nu_2, k_1) \gamma_{\mu} u(x', \nu_1, p_1) = \frac{1}{2} \text{Tr}[\Lambda(x', \nu_1, \nu_2, p_1, k_1) \gamma^{\mu}]$$
(70)

whereas in the case of exchange term we use

$$[j^{\mu}(x)][j_{\mu}(x')] = \frac{1}{4} \operatorname{Tr}[\Lambda(x,\nu_{1},\nu_{2},p_{1},k_{1})\gamma^{\mu}\Lambda(x',\nu'_{1},\nu'_{2},p_{2},k_{2})\gamma_{\mu}]$$
(71)

Similar to scalar meson exchange case as discussed before, we first integrate over t', y', z'; then over q_0 , q_y , q_z and the contour integral is evaluated over q_x . The final integrations are over t, y and z. Thus the Landau interaction function for p - p (process (i)) forward scattering with vector boson exchange direct term is given by

$$f_{p+p\to p+p}^{(d,v)} = \frac{g_{v}^{2}}{32\eta E_{\nu}E_{\nu'}} \int \exp[-\eta |x - x'|] dx dx' [(E_{\nu}E_{\nu'} - p_{z}p_{z'}) \text{Tr}A\text{Tr}A' + p_{\perp}\vec{L}.p_{\perp}'] \text{Tr}B\text{Tr}B']$$
(72)

The above expression for process (ii) with vector boson exchange direct interaction is identical with the zero field case and is given by

$$f_{n+n\to n+n}^{(d,v)} = \frac{g_v^2}{\eta^2} \frac{1}{EE'} (EE' - \bar{p}.\bar{p'})$$
(73)

Next, for the process represented by (iii), the matrix element is given by

$$T_{fi} = i \int j^{\mu}_{(n)}(x) V_{\mu}(x) d^4x$$
(74)

In this case the vector field is given by

$$V^{\mu}(x) = -\int \frac{d^4q}{(2\pi)^4} \frac{\exp[-iq.(x-x')]}{q^2 - \eta^2} j^{(p)}_{\mu}(x') d^4x'$$
(75)

where

$$j_{(n)}^{\mu}(x) = g_{v}\bar{\psi}_{n}(x)\gamma^{\mu}\psi_{n}(x)$$

= $\frac{g_{v}}{V}\exp[-i(p_{2}-k_{2}).x][j_{(n)}^{\mu}]$ (76)

and

$$j_{\mu}^{(p)}(x') = \frac{g_v}{L_y L_z} \exp[-i\{(E_{\nu_1} - E_{\nu_2})t' - (p_{1y} - k_{1y})y' - (p_{1z} - k_{1z})z'\}][j_{\mu}^{(p)}]$$
(77)

Further for the direct term

$$[j_{(n)}^{\mu}] = \frac{1}{2} \operatorname{Tr}[\bar{u}(k_2)\gamma^{\mu}u(p_2)]$$
(78)

and

$$[j_{\mu}^{(p)}] = \frac{1}{2} \operatorname{Tr}[\bar{u}(k_1)\gamma_{\mu}u(p_1)]$$
(79)

Substituting all these terms in T_{fi} and subsequently integrating over x', t', y', z', x, t, y, z and d^4q , we have the direct type Landau-Fermi interaction function for ω meson exchange as

$$f_{p+n \to p+n}^{(d,v)} = -\frac{g_v^2}{4E_\nu E_{\nu'} \eta^2} p.p' \int_{-\infty}^{+\infty} dx' \text{Tr}(A+B)$$
(80)

where we have used

$$[j^{\mu}_{(n)}(x)] = \frac{1}{4E} \operatorname{Tr}[(\not p + m)\gamma^{\mu}] = \frac{p^{\mu}}{E}$$
(81)

and

$$[j^{(p)}_{\mu}(x')] = \frac{1}{4E} \operatorname{Tr}[(Ap^{\lambda}\gamma_{\lambda} + Am + Bp^{\lambda}\gamma_{\lambda})\gamma_{\mu}] = \frac{1}{4E} \operatorname{Tr}[A+B]p_{\mu}$$
(82)

It can now be shown easily that

$$\int_{-\infty}^{+\infty} dx' \operatorname{Tr} A(x') = 4, \quad \int_{-\infty}^{+\infty} dx' rm Tr B(x') = 0$$
(83)

Therefore

$$f_{p+n\to p+n}^{(d,v)} = -\frac{g_v^2}{E_\nu E' \eta^2} (E' E_\nu - p'_z p_z)$$
(84)

The structural form is same as that of non-magnetic case but qualitatively the result is entirely different as we have already discussed.

Now we shall consider exchange interaction terms corresponding to ω -meson transfer. For the process (i), i.e., $p + p \rightarrow p + p$, we have

$$f_{p+p\to p+p}^{(ex,v)} = -\frac{g_v^2}{16E_\nu E_{\nu'}} \int_{-\infty}^{+\infty} \int_{-\infty}^{+\infty} dx dx' \frac{\exp[-k|x-x'|]}{K} \times [(p_z p_{z'} - E_\nu E_{\nu'} + 2m^2) \operatorname{Tr}(AA') - 2\vec{p}_{\perp} . \vec{p}_{\perp}' \operatorname{Tr}(BB')]$$
(85)

In the case of $n + n \rightarrow n + n$, i.e., process (ii) the result is same as that of non-magnetic case and is given by

$$f_{n+n\to n+n}^{(ex,v)} = -\frac{g_v^2}{2} \frac{2m^2 - p \cdot p'}{EE'} \frac{1}{(p-p')^2 - \eta^2}$$
(86)

This is exactly the same as that of zero field result.

Let us now consider the exchange term corresponding to process (*iii*). In this case the calculation is almost identical with the scalar exchange model. Here unlike the scalar case we have to evaluate $\text{Tr}[\Lambda_{np}(p)\gamma^{\mu}\Lambda_{pn}(p')\gamma_{\mu}]$. where the projection operators are given by eqns.(52) and (58). We have noticed that from the non-zero contributions it is very easy to show

$$\operatorname{Tr}[\Lambda_{pn}(p)\gamma^{0}\Lambda_{np}(p')\gamma_{0}] = -\operatorname{Tr}[\Lambda_{pn}(p)\gamma^{x}\Lambda_{np}(p')\gamma_{x}] = -\operatorname{Tr}[\Lambda_{pn}(p)\gamma^{y}\Lambda_{np}(p')\gamma_{y}]$$

$$= -\operatorname{Tr}[\Lambda_{pn}(p)\gamma^{z}\Lambda_{np}(p')\gamma_{z}] =$$

$$= \frac{1}{4E_{\nu}E_{\nu'}}[(E_{\nu}E_{\nu'}+m^{2}+p_{z}p'_{z})+\frac{ip_{\perp}^{2}}{2(E_{\nu}+m)}(m-E_{\nu'})\operatorname{Tr}[\mathcal{A}X\mathcal{A}'] + \frac{ip_{\perp}^{2}'}{2(E_{\nu'}+m)}(m-E_{\nu})\operatorname{Tr}[\mathcal{A}\mathcal{A}'X']$$

$$- \frac{p_{\perp}^{2}p'_{\perp}}{2(E_{\nu}+m)(E_{\nu'}+m)}\operatorname{Tr}[\mathcal{A}\mathcal{A}']$$
(87)

where

$$Tr[\mathcal{A}\mathcal{A}'] = 2(I_{\nu}I_{\nu'} - I_{\nu-1}I_{\nu'-1})$$

$$Tr[\mathcal{A}\mathcal{A}'X] = 2(I_{\nu}I_{\nu'}\exp(-i\phi) - I_{\nu-1}I_{\nu'-1}\exp(i\phi))$$

$$Tr[\mathcal{A}\mathcal{A}'X] = 2(I_{\nu}I_{\nu'}\exp(i\phi') - I_{\nu-1}I_{\nu'-1}\exp(-i\phi'))$$
(88)

It is quite obvious that complex contribution is also coming to the Landau interaction function and as a consequence to the quasi-particle energy if we consider the exchange diagram with vector boson transfer in the process represented by (iii)

4. Iso-Vector Boson Exchange

Now, we consider the last part of our investigation; the iso-vector boson exchange in neutron star matter. Since this part was not included in the original work of Baym and Chin on conventional relativistic version of Landau theory of Fermi liquid, we shall first develop a formalism for the non-magnetic case. Here, as we have mentioned, the iso-vector field ρ^{μ} behaves like a four vector (indicated by symbol μ) in configuration space and a three vector (indicated by \rightarrow) in iso-spin space.

We start with the simplest case, the exchange of neutral iso-vector boson ρ_3^{μ} . The corresponding transition matrix element is given by

$$T_{fi} = i \int j_{3}^{\mu}(x) \rho_{3,\mu}(x) d^{4}x$$
(89)
where

$$\rho_{3,\mu}(x) = -\int \frac{d^4q}{(2\pi)^4} \frac{\exp[-iq.(x-x')]}{q^2 - \theta^2} j_{3,\mu}(x') d^4x'$$
(90)

where θ is the mass of iso-vector boson and

$$j_3^{\mu}(x) = g_{\rho}\bar{\psi}(x)\gamma^{\mu}\frac{\tau_3}{2}\psi(x) \tag{91}$$

is the third component of iso-vector fermion current. In the case of neutral ρ -meson exchange

$$j_{3}^{\mu}(x) = \frac{1}{2} [j_{3,(p)}^{\mu}(x) - j_{3,(n)}^{\mu}(x)]$$
(92)

where symbols p and n represent proton and neutron components respectively. Using the conventional spinor solutions for both proton and neutron we have,

$$j_3^{\mu}(x) = \frac{1}{2} [\bar{u_p}(k)\gamma^{\mu}u_p(p) - \bar{u_n}(k)\gamma^{\mu}u_n(p)] \exp[-i(p-k).x]$$
(93)

On substituting j_3^{μ} and ρ_3^{μ} in the transition matrix element and following the procedure by Baym and Chin for the case of vector meson exchange, we have the direct type Landau quasi-particle interaction function with neutral iso-vector boson exchange

$$f^{(d,iv)} = -\frac{g_{\rho}^2}{4\theta^2} \frac{p.p'}{4EE'}$$
(94)

Similarly, for exchange type interaction we have

$$f^{(ex,iv)} = -\frac{g_{\rho}^2}{8EE'} \frac{[2m^2 - p.p']}{[(p - p')^2 - \theta^2]}$$
(95)

Now in the non-magnetic case, it is very easy to show that the same expressions as above (eqns. (94) and (95)) will also be obtained for the charged iso-vector mesons exchange for the valid process represented by (iii).

Let us now consider the presence of strong quantizing magnetic field, the Landau interaction functions for neutral ρ -meson exchange are almost identical with the results of vector boson exchange (except a factor of 1/2). For the process (i), with direct type interaction, we have

$$f_{p+p\to p+p}^{(d,iv)} = \frac{g_v^2}{64\theta E_\nu E_{\nu'}} \int \exp[-\theta |x - x'|] dx dx' [(E_\nu E_{\nu'} - p_z p_{z'}) \text{Tr} A \text{Tr} A' + p_{\perp}^{-} . p_{\perp}^{'} \text{Tr} B \text{Tr} B']$$
(96)

For the process (ii), the direct interaction is again 1/2 times the non-magnetic result which we have already discussed and in the case of (iii) the Landau quasi-particle interaction function will again be 1/2 of that of vector boson case. In the last case, we again got complex contribution in Landau interaction function.

Next we consider the exchange of charged ρ -meson. For the sake of convenience we define

$$\rho_1^{\mu}(x) = \frac{1}{2} [\rho_+^{\mu}(x) + \rho_-^{\mu}(x)] \quad \text{and} \quad \rho_2^{\mu}(x) = \frac{1}{2} [\rho_+^{\mu}(x) - \rho_-^{\mu}(x)]$$
(97)

where $\rho_{\pm}^{\mu}(x)$ are related to the well known Ladder operators in iso-spin space (causes $p \leftrightarrow n$). Then in the lowest order diagram, the only valid scattering process is as represented by *(iii)*. The transition matrix element for non-magnetic case is given by

$$T_{fi} = -\frac{i}{16} \int \frac{d^4q}{(2\pi)^4} \quad (j_{\mu}^{(pn)}(x) + j_{\mu}^{(np)}(x)) \frac{\exp[-iq.(x-x')]}{q^2 - \theta^2} \\ (j_{(pn)}^{\mu}(x) + j_{(np)}^{\mu}(x))d^4xd^4x'$$
(98)

where

$$j^{\mu}_{(pn)}(x) = g_{\rho} \bar{\psi}_{n}(x) \gamma^{\mu} \psi_{p}(x)$$
(99)

and

$$j^{\mu}_{(np)}(x) = g_{\rho}\bar{\psi}_{p}(x)\gamma^{\mu}\psi_{n}(x)$$
(100)

After some simple algebraic steps we have

$$T_{fi} = -\frac{i}{16} \int \frac{d^4q}{(2\pi)^4} [j^{(pn)}_{\mu}(x) \frac{\exp[-iq.(x-x')]}{q^2 - \theta^2} j^{\mu}_{(np)}(x') + j^{(np)}_{\mu}(x) \frac{\exp[-iq.(x-x')]}{q^2 - \theta^2} j^{\mu}_{(pn)}(x')] d^4x d^4x'$$
(101)

Using the conventional form of spinor solutions for both proton and neutron, we have

$$j^{\mu}_{(pn)}(x) = \frac{g_{\rho}}{V} \exp[-i(p_p - p_n).x]\bar{u_n}(p_n)\gamma^{\mu}u_p(p_p)$$
(102)

and

$$j_{\mu}^{(np)}(x') = \frac{g_{\rho}}{V} \exp[-i(p'_n - p'_p).x'] \bar{u}_p(p'_p) \gamma_{\mu} u_n(p'_n)$$
(103)

Then integrating over d^4x , d^4x' and d^4q , the Landau-Fermi interaction function for direct type forward scattering

$$f^{(d,iv)} = -\frac{g_{\rho}^2 p \cdot p'}{8EE'\theta^2} \tag{104}$$

Similarly for the exchange case

$$f^{(ex,iv)} = \frac{g_{\rho}^2 (4m^2 + p.p')}{16EE'[(p-p')^2 - \theta^2]}$$
(105)

Let us now consider the effect of strong quantizing magnetic field on this process with charged ρ -meson exchange. In this case not only the Dirac spinor solutions of proton, but the charged meson propagators will also get modified. The meson propagator is given by

$$G_{\rho_{\pm}}(x) = \frac{eB_m}{4\pi^2} \sum_{\nu} \int dE_0 \Phi^*(x') \Phi(x) \exp[-i\{E_0(t-t') - q_y(y-y') - q_z(z-z')\}] \\ \frac{1}{E_0 - (\epsilon_q - i\delta)} \frac{1}{4} [j_{pn}(x') - j_{np}(x')] d^4x'$$
(106)

where

$$\Phi(x) = I_{\nu}(x, p) = \left(\frac{qB_m}{\pi}\right)^{1/4} \frac{1}{(\nu!)^{1/2}} 2^{-\nu/2} \exp\left[-\frac{1}{2}qB_m\left(x - \frac{p_y}{qB_m}\right)^2\right] \\ H_{\nu}\left[(qB_m)^{1/2}\left(x - \frac{p_y}{qB_m}\right)\right],$$
(107)

Then it is obvious that $\Phi^*(x) = \Phi(x)$. The quantity $\epsilon_q = (q_z^2 + \theta^2 + 2\nu eB_m)^{1/2}$ is the modified form of charged ρ -meson energy eigen value, obtained from the solution of Klein-Gordon equation in presence of strong quantizing magnetic field, with $2\nu = 2n + 1 + 2\alpha$, $\alpha = -1, 0$ or +1 and n = 0, 1, 2, ..., the Landau principal quantum number.

Now in the direct case, unlike the scalar and vector mesons exchange, we have to evaluate the product $\text{Tr}[\Lambda_{pn}\gamma^{\mu}]\text{Tr}[\Lambda_{np}\gamma_{\mu}]$. As before, we calculate this quantity term by term (i.e., for $\mu = 0, x, y$ and z). Using the expressions for the projection operators, we have

$$\operatorname{Tr}[\Lambda_{pn}\gamma^{0}]\operatorname{Tr}[\Lambda_{np}\gamma_{0}] = \left[\frac{1}{2}\operatorname{Tr}\mathcal{A} - \frac{ip_{\perp}^{2}}{4E_{\nu}(E_{\nu}+1)}\operatorname{Tr}(\mathcal{A}X)\right] \left[\frac{1}{2}\operatorname{Tr}\mathcal{A}' - \frac{ip_{\perp}'^{2}}{4E_{\nu}'(E_{\nu}'+1)}\operatorname{Tr}(\mathcal{A}'X^{*})\right]$$
(108)

$$\operatorname{Tr}[\Lambda_{pn}\gamma^{x}]\operatorname{Tr}[\Lambda_{np}\gamma_{x}] = \operatorname{Tr}[\Lambda_{pn}\gamma^{y}]\operatorname{Tr}[\Lambda_{np}\gamma_{y}] = 0$$
(109)

and

$$\operatorname{Tr}[\Lambda_{pn}\gamma^{z}]\operatorname{Tr}[\Lambda_{np}\gamma_{z}] = \left[-\frac{1}{2}p_{z}\operatorname{Tr}\mathcal{A}\right] \left[-\frac{1}{2}p_{z}'\operatorname{Tr}\mathcal{A}'\right]$$
(110)

Obviously, the term with $\mu = 0$ is complex in nature and it will make the Landau quasi-particle interaction function and as a consequence the quasi-particle energies for both neutron and proton complex in nature.

Fortunately, in the case of exchange type interaction with charged iso-vector meson exchange, we have to evaluate $\text{Tr}[\Lambda_{pp}\gamma^{\mu}\Lambda_{nn}\gamma_{\mu}]$ which gives real contribution to quasi-particle energy.

To obtain the variation of self energies (both real and imaginary parts) we have evaluated numerically the total self energy density for nucleons (both protons and neutrons separately) then plotted the average proton self energy as a function of proton density and are shown in the following two figures. To draw these figures, we have solved a few non-linear equations self-consistently. We assume that before the interaction is switched on, the matter is in β -equilibrium, given by

$$\mu_n = \mu_p + \mu_e$$

(Which is obtained from the weak processes $n \longrightarrow p + e^- + \bar{\nu}_e$ and $p + e^- \longrightarrow n + \nu_e$, known as the URCA and the modified URCA processes). We further assume the charge neutrality condition, given by

$$n_p = n_e$$

and assuming that the baryon number density $n_B = n_p + n_n$ remains constant. To solve for the chemical potentials μ_i , where i = n, p and e, we use

$$n_n = \frac{(\mu_n^2 - m_n^2)^{3/2}}{6\pi^2},$$

$$n_p = \frac{eB_m}{2\pi^2} \sum_{\nu=0}^{[\nu_{\max}]} (2 - \delta_{\nu 0}) (\mu_p^2 - m_p^2 - 2\nu eB_m)^{1/2}$$

and

$$n_e = \frac{eB_m}{2\pi^2} \sum_{\nu=0}^{[\nu_{\max}]} (2 - \delta_{\nu 0}) (\mu_e^2 - m_e^2 - 2\nu eB_m)^{1/2}$$

Hence, knowing the chemical potentials, we obtained numerically the total free energies (both real and imaginary parts) per particle for both protons and neutrons.



FIG. 2: Variation of the real part of average proton self energy (in MeV.) with proton density (expressed in terms of normal nuclear density $n_0 = 0.17 \text{fm}^{-3}$). The upper curve is for $B_m = 10^{14}$ G, middle one is for $B_m = 10^{16}$ G and the lower one is for $B_m = 10^{18}$ G.



FIG. 3: Variation of the imaginary part of average proton and neutron self energies (in MeV.) with proton density (expressed in terms of normal nuclear density $n_0 = 0.17 \text{fm}^{-3}$). The upper curve is for $B_m = 10^{18}$ G, middle one is for $B_m = 10^{16}$ G and the lower one is for $B_m = 10^{14}$ G.

As one can see from these two figures that the real part always increases with the increase of matter density but the matter becomes more stable if the magnetic field is stronger. Which is also true in all other cases, e.g., atoms, quark matter etc. [10, 31, 32] in strong quantizing magnetic fields. Whereas the imaginary part decreases with the increase of matter density and at very high density the imaginary part is almost negligibly small with respect to the corresponding real part. The reason is that at very high density, the effect of Landau levels become insignificant, in other ward, at high enough density the magnetic field no longer has quantum mechanical effect. On the other hand at low density, it is significant and the effect is more as the magnetic field becomes stronger.

For the sake of completeness we have also plotted the life time of nucleons (both neutron and proton) as a function of proton density, for three different magnetic field strengths. The life times are in the strong interaction time scale within a broad density range. Because of such small time scale we belive that the bulk neutron star matter remains stable against strong decay, just like a nucleus is a stable object although strong interaction processes are going on inside. The other extreme could be that the magnetic field strength at the core region of magnetars are much smaller



FIG. 4: Variation of proton and neutron life times (in sec. in unit of 10^{22}) with proton density (expressed in terms of normal nuclear density $n_0 = 0.17 \text{fm}^{-3}$). The upper curve is for $B_m = 10^{18}$ G, middle one is for $B_m = 10^{16}$ G and the lower one is for $B_m = 10^{14}$ G.

than the quantum critical value and the observed strong magnetic fields are concentrated at the crustal region.

3. CONCLUSIONS

We therefore conclude that in the relativistic version of Landau theory of Fermi liquid in presence of strong quantizing magnetic field, although the self-energies for both neutron and proton become complex in nature, the bulk neutron star matter still remain stable against strong decay. We have noticed from the complex nature of self-energies for both the dressed protons and neutrons, that the life time is within the strong interaction scale and we belive that just like a stable nucleus, the bulk neutron star matter will also remain stable in presence of strong quantizing magnetic field.

The other explanation, which is of course not so convincing is that, we can have a stable dense neutron matter at the central region of a magnetar with a very low magnetic field strength. The observed strong magnetic field in such objects are concentrated at the crustal region. However, in our opinion, the best explanation to have a magnetic field of strength $\sim 10^{14}$ G at the surface and several orders of magnitude large field at the core is that the conducting (and possibly not superconducting because the field strength is > the critical value) neutron star matter does not allow the magnetic flux lines to move from the core region to the crust. Since the density of neutron star matter at the core region is high enough, a large number of Landau levels will be populated, even if the field strength is extremely high. Which further means, that the quantum mechanical effect of strong magnetic field will be extremely weak.

Finally, we would like to add a few words to compare the results, in particular the complex nature of nucleon self energies we obtained in this work with some of the well known phenomena already discussed in the literature.

About sixty years ago L.D. Landau had shown the possibility of some kind of damping or dissipation in collisionless plasmas and this phenomenon is known as *Landau Damping* [33, 34]. Being independent of collisions, it is fundamentally different from dissipation in ordinary absorbing media; the collisionless dissipation does not involve an increase of entropy and is therefore a thermodynamically reversible process. In this phenomenon, there is a continuous exchange of energy between plasma waves (produced by some external perturbation) and plasma particles. As a consequence the self energy of plasma particles become complex and they loose energy to plasma wave reversibly and in the very next step again gain energy from the plasma waves and the process of loss and gain continues, but does not decay to other particles. This phenomenon makes the system mechanically stable.

The evaluation of nucleon self energy Σ in relativistic approaches has been restricted mainly to the real part of Σ in either the mean field theory or the relativistic Hartree approximation. In these approximations the self-energy is real, local and energy independent. It is completely determined once the constant effective mass and the mean constant value of vector potential is know [35].

However, the imaginary part of the self-energy, which in reality the imaginary part of optical potential has been studied to lowest order by Horowitz [36]. Here, only the on-shell properties of $Im(\Sigma(\varepsilon, \vec{p}))$ is important and is

restricted to a limited energy range. He obtained a quantity similar to the imaginary part of optical potential by the cancellations of large contributions coming from the exchange of scalar and vector mesons; between scalar, time-like and three-vector contributions which are also independently very large.

In ref. [37], real and imaginary parts of nucleon self-energy was obtained with relativistic Dirac-Bruckner approach assuming $\sigma - \omega$ meson exchange. In this paper fourth order meson-nucleon coupling was considered and studied the off-shell behaviour of imaginary part of the nucleon self energy.

In the reference [38] Dyson equation is used to obtain the nucleon self energy Σ^* . In this paper the irreducible self energy is calculated by summing over the Dyson expansion to all order. It is shown that in the lowest order diagrams the irreducible self-energy contains mean field part and the Hartree-Fock contribution. Both of them are real in nature. The dynamic part of the self energy is expressed in terms of the four point vertex function.

However, the complex nature of nucleon self-energy as we have obtained in this article, arises because of some kind of mixing in the iso-spin space (the modified form of *Projection Operators* e.g., Λ_{pn} and Λ_{np} are actually responsible for such strange phenomena. These modified form of projection operators are again coming because in presence strong quantizing magnetic field, the Landau levels of protons are only populated, whereas neutrons behave classically) and it is at present very difficult to predict what will happen to these nucleons with complex energy.

- R.C. Duncan and C. Thompson, Astrophys. J. Lett. **392**, L9 (1992); C. Thompson and R.C. Duncan, Astrophys. J. **408**, 194 (1993); C. Thompson and R.C. Duncan, MNRAS **275**, 255 (1995); C. Thompson and R.C. Duncan, Astrophys. J. **473**, 322 (1996).
- [2] P.M. Woods, C. Kouveliotou and J. van Paradijs, Astrophys. J. Lett. 519, L139 (1999); C. Kouveliotou, et. al., Nature 393, 235 (1999).
- [3] K. Hurley, et. al., Astrophys. Jour. **510**, L111 (1999).
- [4] S. Mereghetti and L. Stella, Astrophys. Jour. 442, L17 (1999); J. van Paradijs, R.E. Taam and E.P.J. van den Heuvel, Astron. Astrophys. 299, L41 (1995); S. Mereghetti, astro-ph/99111252; see also A. Reisenegger, astro-ph/01003010.
- [5] D. Bandopadhyaya, S. Chakrabarty, P. Dey and S. Pal, Phys. Rev. D58, 121301 (1998).
- [6] S. Chakrabarty, D. Bandopadhyay and S. Pal, Phys. Rev. Lett. 78, 2898 (1997); D. Bandopadhyay, S. Chakrabarty and S. Pal, Phys. Rev. Lett. 79, 2176 (1997).
- [7] C.Y. Cardall, M. Prakash and J.M. Lattimer, astro-ph/0011148 and references therein; E. Roulet, astro-ph/9711206; L.B. Leinson and A. Pérez, astro-ph/9711216; D.G. Yakovlev and A.D. Kaminkar, The Equation of States in Astrophysics, eds. G. Chabrier and E. Schatzman P.214, Cambridge Univ.
- [8] S. Chakrabarty and P.K. Sahu, Phys. Rev. **D53**, 4687 (1996).
- [9] S. Chakrabarty, Phys. Rev. D51, 4591 (1995); Chakrabarty, Phys. Rev. D54, 1306 (1996).
- [10] T. Ghosh and S. Chakrabarty, Phys. Rev. D63, 0403006 (2001); T. Ghosh and S. Chakrabarty, Int. Jour. Mod. Phys. D10, 89 (2001).
- [11] D.G. Yaklovlev and D.A. Shalybkov, Astrophys. Space Sci. 176 (1991) 171.
- [12] D.G. Yaklovlev and D.A. Shalybkov, Astrophys. Space Sci. 176 (1991) 191.
- [13] D.G. Yaklovlev and D.A. Shalybkov, Sov. Astron. Lett 16 (1990) 86.
- [14] V.G. Bezchastrov and P. Haensel, astro-ph/9608090.
- [15] S. Mandal and S. Chakrabarty (two papers on weak and electromagnetic processes are communicated for publication).
- [16] Sutapa Ghosh and Somenath Chakrabarty, Mod. Phys. Lett. A17, (2002) 2147.
- [17] V.P. Gusynin, V.A. Miransky and I.A. Shovkovy, Phys. Rev. D52, 4747 (1995).
- [18] D.M. Gitman, S.D. Odintsov and Yu.I. Shil'nov, Phys. Rev. D54, 2968 (1996).
- [19] D.S. Lee, C.N. Leung and Y.J. Ng, Phys. Rev. D55, 6504 (1997).
- [20] V.P. Gusynin and I.A. Shovkovy, Phys. Rev. D56, 5251 (1995).
- [21] D.S. Lee, C.N. Leung and Y.J. Ng, Phys. Rev. D57, 5224 (1998).
- [22] V.P. Gusynin, V.A. Miransky and I.A. Shovkovy, Phys. Rev. Lett. 83, 1291 (1999).
- [23] E.V. Gorbar, Phys. Lett. **B491**, 305 (2000).
- [24] T. Inagaki, T. Muta and S.D. Odintsov, Prog. Theo. Phys. **127**, 93 (1997), hep-th/9711084; T. Inagaki, S.D. Odintsov and Yu.I. Shil'nov, Int. J. Mod. Phys. **A14**, 481 (1999), hep-th/9709077; T. Inagaki, D. Kimura and T. Murata, Prog. Theo. Phys. **111**, 371 (2004), hep-ph/0312005.
- [25] Sutapa Ghosh and Somenath Chakrabarty, communicated.
- [26] A. Melatos, Astrophys. Jour. 519, L77 (1999); A. Melatos, MNRAS 313, 217 (2000).
- [27] S. Bonazzola and E. Gourgoulhon, Astron. & Atrophysics. 278, 421 (1993); M. Bocquet, S. Bonazzola, E. Gourgoulhon and J. Novak, Astron. & Atrophysics. 301, 757 (1995).
- [28] B. Bertotti and A.M. Anile, Astron. & Atrophysics. 28, 429 (1973); C. Cutler and D.I. Jones, gr-qc/0008021; K. Konno, T. Obata and Y. Kojima, gr-qc/9910038; A.P. Martinez, H. Perez Rojas and H.J. Mosquera Cuesta, hep-ph/0011399; M. Chaichian, S.S. Masood, C. Montonen, A. Perez Martinez and H. Perz Rojas, Phys. Rev. Lett. 84, 5261 (2000); Guangjun Mao, Akira Iwamoto and Zhuxia Li, astro-ph/0109221; A. Melatos, Astrophys. Jour. 519, L77 (1999); A. Melatos, MNRAS 313, 217 (2000); R. González Felipe, H.J. Mosquera Guesta, A. Perez Martinez and H. Perez Rojas, astro-ph/0207150;

Chin. J. Astron. Astrophys. 5, 399 (2005). and references therein.

- [29] S. Ghosh, S. Mandal and S. Chakrabarty, Ann. Phys. **312**, 398 (2004).
- [30] G. Baym and S.A. Chin, Nucl. Phys. A262, 527 (1976).
- [31] E.M. Lieb, J.P. Solovej and J. Yngvason, Phys. Rev. Lett. 69, 749 (1992).
- [32] S. Chakrabarty and A. Goyal, Mod. Phys. Lett. A9, 3611 (1994).
- [33] L.D. Landau, Jour. Phys. (U.S.S.R.) 10, 25 (1946).
- [34] Also see J. Neufeld and R.H. Ritchie, Phys. Rev. 98, 1632 (1955).
- [35] B.D. Serot and J.D. Walecka, Adv. Nucl. Phys. 16, 1 (1986).
- [36] C.J. Horowitz, Nucl. Phys. A412, 228 (1984).
- [37] A.E.L. Dieperink, J. Piekarewicz and K. Wihrberger, Phys. Rev. C41, 2479 (1990).
- [38] W.H. Dickhoff and C. Barbieri, Prog. in Part. and Nucl. Phys. 52, 377 (2004).