

Spitzer Discovery of Very Low Luminosity Objects

J. KAUFFMANN¹, F. BERTOLDI², N.J. EVANS II³, and THE C2D COLLABORATION

¹ Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

² Radioastronomisches Institut der Universität Bonn, Auf dem Hügel 71, D-53121 Bonn, Germany

³ Department of Astronomy, University of Texas at Austin, 1 University Station C1400, Austin, TX 78712-0259

Received; accepted; published online

Abstract. The Spitzer Space Telescope allows for the first time to search systematically for very low luminosity ($\lesssim 0.1 L_{\odot}$) objects (VeLLOs) associated with dense molecular cores. They may be the first candidate Class 0 sources with sub-stellar masses. We describe such a source in the dense molecular core L1148. VeLLO natal cores show properties that are unusual for star-forming cores. The low luminosity and in some cases the lack of prominent outflow could be the result of the small gas supply near the VeLLO.

Key words: stars: formation — ISM: globules — ISM: individual (L1148)

©0000 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim

1. Introduction

The “revealed” phase of early stellar evolution was discovered and classified in the late 1980’s (Lada 1987), but it took some years until objects in the “obscured” phase of star-formation were found and classified (André, Ward-Thompson & Barsony 1993). Such Class 0 protostars are so deeply embedded in their natal dense cores that they could not be detected with contemporary near- and mid-infrared detectors. They are believed to be in the main accretion phase of star formation (André, Ward-Thompson & Barsony 2000), drive collimated bipolar outflows (André et al. 2000) and can show maser emission (Furuya et al. 2001).

While the first Class 0 sources were studied, also the first brown dwarfs (BDs) and extrasolar planets were discovered (Nakajima et al. 1995). Many BDs are not orbiting a stellar mass primary (Oppenheimer, Kulkarni & Stauffer 2000). There is no proof that these BDs were ejected from forming protostellar systems. The large BD binary fraction of $\gtrsim 15\%$ (Martín et al. 2003, Pinfield et al. 2003) is furthermore hardly explained by ejection scenarios (Whitworth & Goodwin 2005). It thus appears plausible that at least some of the single brown dwarfs may have formed by processes similar to those by which normal stars form, i.e. the collapse of a dense core onto a compact object. Sub-stellar mass compact objects in the process of formation would appear as low-luminosity ($L_{\text{bol}} \ll 1 L_{\odot}$) Class 0 sources.

It is a challenge to search and study accreting very low luminosity objects (VeLLOs) that may turn into sub-stellar mass brown dwarfs. While a first such source (IRAM04191) was found serendipitously (André, Motte & Bacmann 1999), a systematic search for such objects has become possible only with the Spitzer Space Telescope (SST). Our SST Legacy Program “From Molecular Cores to Planet Forming Disks” (c2d; Evans et al. 2003), which images about 90 isolated dense cores and 5 large molecular clouds (Huard et al., this volume), is tailored to find such VeLLOs. The source L1014-IRS (Young et al. 2004) was the first low luminosity object discovered in our survey. We here present another candidate for an accreting sub-stellar object, L1148-IRS.

2. The VeLLO Candidate L1148-IRS

2.1. The compact IR Source

The dense core L1148 is part of the Cepheus flare at a distance of about 325 pc (Straizys et al. 1992). We have imaged the thermal dust emission of L1148 at 1.2 mm wavelength using the Max-Planck Millimeter Bolometer (MAMBO-2) array camera at the IRAM 30m-telescope on Pico Veleta (Spain). Such wide band continuum observations map the projected mass distribution, since the dust emission is nearly proportional to the H_2 column density and the dust temperature (e.g., Motte, André & Neri 1998). In our mm-continuum map we found two parallel, faint filaments in the south-east




Fig. 1. MAMBO dust emission (top) and $24\ \mu\text{m}$ Spitzer map of L1148 (bottom). The bright Spitzer source L1148-IRS is marked by a star in the MAMBO map. Offsets are in arcseconds.

and north-west of L1148 (Fig. 1). Assuming dust temperatures of 10 K we estimate their masses to 11 and $5 M_{\odot}$ and their peak visual extinctions to 7 and 5 mag, respectively.

The dust emission peak in the north-western filament is associated with a bright Spitzer source at $24\ \mu\text{m}$ and $70\ \mu\text{m}$ (Fig. 1), named L1148-IRS. From its appearance in the IRAS Faint Source Catalog Kun (1998) noted it as a potential low-luminosity protostar. A spatial, physical association of the dust emission peak and L1148-IRS is suggested by the $6''$ proximity. The probability of a chance alignment within $20''$ of the dust peak (about twice the separation of dust peak and L1148-IRS) and a background galaxy of comparable infrared brightness is very low (1:7700 at $24\ \mu\text{m}$, 1:3400 at $70\ \mu\text{m}$). Because L1148 is way above the galactic plane ($z \approx 80\ \text{pc}$ at $b = +15^{\circ}$) a chance alignment with a background protostar is also unlikely. Therefore L1148-IRS is most probably an infrared source associated with the L1148 dense core.

Using the IRAM 30m-telescope we have unsuccessfully searched for a CO outflow from L1148-IRS, which would have clearly identified L1148-IRS as an accreting protostar. This does not exclude the presence of a small-scale flow, which in the VELLO L1014-IRS could only be detected with an interferometer (Sec. 3).




Fig. 2. The spectral energy distribution of L1148-IRS. The dashed line shows our model SED for the compact emission from an embedded protostar (which does not apply to the MAMBO data point).

With the available archival 2MASS and IRAS data and our own OMEGA-Cass. (Calar Alto 3.5m-telescope) near-IR observations, in Fig. 2 we plot the observed spectral energy distribution (SED) of L1148-IRS. For $\lambda \lesssim 100\ \mu\text{m}$ the source L1148-IRS is compact (smaller than a few arcsec) and the fluxes listed hold for this compact source. For $\lambda \gtrsim 100\ \mu\text{m}$ the source is extended (several arcmin at $\lambda = 1\ 200\ \mu\text{m}$) and the fluxes hold for an aperture of $12''.9 = 4\ 200\ \text{AU}$ radius, to be consistent with previous work (André et al. (1999)).

To estimate the bolometric luminosity of the embedded star we integrated the flux density over the observed SED using a modified version (at $\lambda \gtrsim 200\ \mu\text{m}$ we assume the SED to follow a greybody of temperature 10 to 15 K and dust opacity $\kappa \propto \lambda^{-2}$ instead of a powerlaw) of the method of Chen et al. (1995). The integration yields $L_{\text{bol}} = 0.1$ to $0.2 L_{\odot}$ (out of which about 30 to 70% are due to heating by the interstellar radiation field [ISRF]), a submillimeter-to-bolometric luminosity ratio $L_{>350\ \mu\text{m}}/L_{\text{bol}} \geq 0.05$, and a bolometric temperature $T_{\text{bol}} \leq 140\ \text{K}$. As $L_{>350\ \mu\text{m}}/L_{\text{bol}} \gg 0.005$ L1148-IRS qualifies as a Class 0 protostar (André et al. 2000).

We tried to model the observed SED with an embedded young stellar object (Fig. 2). The continuum radiative transfer was calculated¹ for a star of luminosity L and temperature T , surrounded by a spherical core with a power-law density profile: $\rho \propto r^{-\alpha}$ with $\alpha = 3/2$ to 2 (Shu 1977) between radius r_{in} and r_{trans} , $\alpha = 0$ between r_{trans} and r_{out} , $\rho = 0$ for $r < r_{\text{in}}$ and $r > r_{\text{out}}$. While r_{in} and r_{trans} are free parameters, the outer radius, r_{out} , and density are constrained by the extended dust emission seen in the MAMBO maps. The core is exposed to the ISRF. We take the differential observing technique (e.g., a “background intensity” is subtracted from a Spitzer image before a flux measurement) into account when modelling the individual flux densities. This produces different model SEDs for the compact ($< 2\ \text{arcsec}$ at $\lambda \lesssim 100\ \mu\text{m}$) and the extended emission ($> 2\ \text{arcsec}$ at $\lambda \gtrsim 100\ \mu\text{m}$). The

¹ The code was kindly provided by E. Krügel.

main result from fitting our models is that independent of our choice of values for the free parameters, the stellar luminosity and temperature can be constrained to $L = 0.05$ to $0.15L_{\odot}$ and $T = 2000$ to 4000K .

In a detailed comparison we notice a discrepancy between the predicted and observed flux densities in the $8\ \mu\text{m}$ Spitzer band (Fig. 2), which may be dominated by silicate absorption and emission. Furthermore, our model fit implies an inner radius, r_{in} , that is much larger than the dust sublimation radius (the dust in our model has temperatures $\lesssim 100\text{K}$, where dust sublimates at $\approx 1500\text{K}$), possibly suggesting a more complicated geometry including the presence of a disk and cavities. The presence of significant amounts of hot ($\gg 100\text{K}$) dust, in which case we would observe a significant slope in the IRAC bands, can be excluded.

Following Young et al. (2004) we derive an upper limit to the protostar mass, M_* , using the constraint that the accretion luminosity, $L_{\text{accr}} = GM_*\dot{M}/R_*$ is smaller than the total luminosity, $L_{\text{accr}} < L$. Here G is the constant of gravity, R_* is the protostar radius, and \dot{M} the accretion rate. Using estimates for R_* and \dot{M} from Young et al. (2004), we find $M_* \ll 0.1 M_{\odot}$.

2.2. The Natal Dense Core

Our Effelsberg 100m telescope maps of L1148 in the 2_1-1_0 transition of CCS probe the dense core on large scales (Fig. 3). The velocity-integrated CCS intensity map reveals two filaments associated with the dust emission filaments. The CCS centroid velocity map shows a jump in velocity about $70''$ south-west of L1148-IRS. Since the velocity field is smooth elsewhere, the velocity-discontinuity suggests two neighbouring, possibly overlapping dense cores, separated in velocity by 0.1 km s^{-1} . The linewidth is $< 0.2\text{ km s}^{-1}$ in the north-western filament, except where the gradient of the centroid velocity is large, possibly due to overlapping velocity components.

Our molecular line observations with the IRAM 30m-telescope probe the dense core on smaller scales. The C^{18}O 2-1 emission, combined with the dust emission implies a CO depletion of $f_d = 8$ (Crapsi et al. 2005a); the abundance of CO in the gas phase appears to be reduced by a factor 8 relative to the “standard” abundance, probably through freeze-out onto dust grains during the evolution of the core. Some of the observed C^{18}O emission could come from velocity components unrelated to the dust peak, in which case f_d for the dense core would be underestimated.

Our map of the N_2H^+ 1-0 emission shows that the dense gas is closely associated with L1148-IRS (Fig. 4), further supporting the picture that the Spitzer source is physically associated with the L1148 dense core.

3. The confirmed Protostars IRAM04191 and L1014-IRS

Besides L1148-IRS, two other low-luminosity protostars have been studied in detail.




Fig. 3. CCS 2_1-1_0 integrated intensity, centroid velocity, and velocity dispersion. Overlaid contours are dust continuum intensity. The position of L1148-IRS is marked by a star. Offsets are arcseconds.




Fig. 4. Dense gas near L1148-IRS as probed in the N_2H^+ (1-0) transition. The position of L1148-IRS is marked by a star. Offsets are in seconds of arc.

Table 1. Properties of VeLLOs (top) and their natal dense cores (bottom) as observed with single dish telescopes. Dense cores are compared with criteria for “evolved” cores from Crapsi et al. (2005b); N is column density. A plus indicates that a criterion is met, a minus that it is not, a question mark that the data is insufficient to decide. Data are from Motte & André (2001), Belloche et al. (2002), Crapsi et al. (2005b), and this work.

Property	IRAM04191	L1014	L1148
<i>protostar properties:</i>			
prominent outflow	+	–	–
$M_{\text{env}}^{4200 \text{ AU}} / M_{\odot}$	0.45	0.27	0.11
<i>dense core properties:</i>			
$f_d > 10$?	–	–
$N(\text{N}_2\text{D}^+) / N(\text{N}_2\text{H}^+) \geq 0.1$?	+	?
$N(\text{N}_2\text{H}^+) \geq 8.5 \cdot 10^{12} \text{ cm}^{-2}$	+	–	–
contr. motions, infall	+	–	+

IRAM04191 was the first known protostar with a very low luminosity ($L \approx 0.1 L_{\odot}$) and therefore mass (André et al. 1999). It was found through its prominent CO outflow, that is associated with a VLA cm radio source and a dust emission peak. A near- and far-infrared counterpart was recently identified in our Spitzer data (Dunham et al. 2005).

The VeLLO L1014-IRS was discovered through the association of an unusually bright Spitzer source with a dust emission peak (Young et al. 2004). It has a low luminosity of $\sim 0.1 L_{\odot}$. The physical association of the Spitzer source with the L1014 dense core has been established by the detection of an infrared outflow cone (Huard et al. 2005; see also Huard et al., this volume) and the discovery of a small-scale CO outflow with the SMA interferometer (Bourke et al. 2005).

4. The Physics of Very Low Luminosity Objects

Compared with previously discovered Class 0 protostars, VeLLOs appear to be surrounded by small amounts of molecular gas. Motte & André (2001) suggest a comparison based on the mass within a projected radius of 4200 AU from the protostar, $M_{\text{env}}^{4200 \text{ AU}}$.

In the Class 0 source sample of Motte & André (2001), IRAM04191 has the smallest inner envelope mass (and luminosity), and the masses for L1014 and L1148 are even smaller than that (Tab. 1). It is suggestive that for VeLLOs the low luminosity and lack of outflow activity somehow correlate with the low envelope mass. However, such conclusion is yet based on too small a sample.

VeLLOs are found in cores that were thought not to be able to form stars. Crapsi et al. (2005a) suggested criteria for properties of starless cores just before the onset of star-formation. According to this, cores must be exceptionally dense and chemically evolved, e.g., they show high degrees of deuteration and depletion. Table 1 shows that the star-forming

cores L1014 and L1148 meet only some of Crapsi’s criteria. Since these were put forward on the basis of single-dish observations, they may be insufficient to probe the VeLLO natal dense cores on those scales relevant for the star-formation process.

We remark (as André et al. 1999 did for IRAM04191 and Young et al. 2004 did for L1014) that VeLLOs are embedded in dense cores with sufficiently large masses that they could in principle grow to stellar masses ($> 0.1 M_{\odot}$), and thus do not necessarily remain sub-stellar objects.

5. Summary

The Spitzer Space Telescope allows for the first time a systematic search for very low luminosity ($L \approx 0.1 L_{\odot}$) embedded, compact objects (VeLLOs), accreting Class 0 sources of apparently sub-stellar mass. With L1148-IRS we here present a further candidate for such an object.

The presently known VeLLOs have a very low inner envelope mass. This could relate to their low luminosities and – in the case of L1014-IRS and L1148-IRS – to their low outflow activity. Two of the three sufficiently studied VeLLO natal cores (L1014-IRS and L1148-IRS) have properties that do not meet the proposed criteria for starless cores before the onset of star-formation. They appear neither very dense nor chemically evolved when probed with single-dish telescopes. VeLLOs seem to form in clouds that were not expected to form stars.

Acknowledgements. We thank E. Krügel, T. Bourke, P. Myers, C.W. Lee, T. Huard, and P. André for insightful discussions, and the Instituto de Astrofísica de Canarias (IAC) for providing a very stimulating atmosphere at the ULMSF05 workshop.

References

- André, P., Ward-Thompson, D., Barsony, M.: 1993, ApJ 406, 122
 André, P., Ward-Thompson, D., Barsony, M.: 2000, in “Protostars and Planets IV”, eds. V. Mannings, A.P. Boss, S.S. Russell (Tucson: University of Arizona Press), 59
 André, P., Motte, F., Bacmann, A.: 1999, ApJ 513, L57
 Belloche, A., et al.: 2002, A&A 393, 927
 Bourke, T.L., et al.: 2005, accepted by ApJ Letters
 Chen, H., et al.: 1995, ApJ 445, 377
 Crapsi, A., et al.: 2005a, ApJ 619, 379
 Crapsi, A., et al.: 2005b, A&A 439, 1023
 Dunham, M., et al.: 2005, in preparation
 Evans, N.J., II, et al.: 2003, PASP 115, 965
 Furuya, R.S., et al.: 2001, ApJ 559, L143
 Huard, T.L., et al.: 2005, accepted by ApJ
 Kun, M.: 1998, ApJS 115, 59
 Lada, C.J.: 1987, in IAU Symp. 115, “Star Forming Regions”, eds. M. Peimbert & J. Jugaku (Dordrecht: Kluwer), 1
 Martín, E.F., et al.: 2003, ApJ 594, 525
 Motte, F., André, P., Neri, R.: 1998, A&A 336, 150
 Motte, F., André, P.: 2001, A&A 365, 440
 Nakajima T., et al.: 1995, Nature 378, 463
 Oppenheimer, B.R., Kulkarni, S.R., Stauffer, J.R.: 2000, in “Protostars and Planets IV”, eds. V. Mannings, A.P. Boss, S.S. Russell (Tucson: University of Arizona Press), 1313
 Pinfield, D.J., et al.: 2003, MNRAS 342, 1241

Shu, F.H.: 1977, ApJ 214, 488

Straizys, V., Cernis, K., Kazlauskas, A., Meistas, E.: 1992, BaltA 1, 149

Whitworth, A.P., Goodwin, S.P.: 2005, Memorie della Societa Astronomica Italiana, v. 76, p. 211

Young, C.H., et al.: 2004, ApJS 154, 396