

A search for spectral variations of planetary nebulae and related objects

L.N. KONDRATYEVA

Fessenkov Astrophysical Institute, 480020, Almaty, Kazakhstan

The results of long-term spectral observations were used to search for changes in planetary nebulae and emission-line stars. Significant increase of excitation degree is found in two objects: M1-6 and M1-11.

KEY WORDS planetary nebulae, – spectra: variability

1 INTRODUCTION

It is widely known, that physical parameters of planetary nebulae and their central stars change gradually during evolution, but these effects are rather difficult for detection in a short time interval. Sometimes much more exotic processes, such as throwing out of a secondary envelope, jets and so on, take place in planetary nebulae. Similar processes cause appreciable changes in spectra, but these events are rare and unpredicted. In other words everyone knows, that planetary nebulae are changing, but these effects are hard to be found out. The problem is more complicated, because results, obtained simultaneously, but by different authors may sometimes differ sufficiently. We decided to use the fact, that spectral observations of the large group of planetary nebulae and other emission objects were carried out in the Astrophysical Institute for a long time and with the same equipment. The very first spectrograms have been obtained in 1970 - 1973, and the last observations were conducted in 2004 - 2005. The aim of this paper is to search for any variations in spectra of chosen objects within 30 - 35 years. We expected, that even small fluctuations of T_{eff} , T_e or N_e in some young objects can affect essentially the intensities of forbidden lines. We also hoped, to find any traces of fast dynamic processes, such a jet, if they have taken place within the specified period.

2 The sample of objects

The objects of different types and ages were chosen for this study. There are three classical planetary nebulae of high and moderate excitation (K3-58, K3-3, M1-5), three low -excitation planetary nebulae (M1-6, M1-11, M1-12) and M1-77. The last object has the spectrum of very low-excitation with the [O II], [N II] and [S II] emission lines, weak emission and absorption lines of neutral metals. It was identified as a young planetary nebula by [1]. Later some irregular changes of brightness and radial velocity were found out by [2] and [3]. Our observations showed that there are TiO bands in the spectrum of M1-77, and it was considered to be the nebula around the symbiotic star [4]. In addition three emission objects were included in our program: MWC 137, M1-15 and He2-446.

3 Observations

All observations have been carried out by the author with the 70-cm telescope of the Astrophysical Institute (Almaty, Kazakhstan). We used a slit spectrograph, equipped with a three-cascade image tube until 1998, and with a CCD matrix ST8 since 1999. A sample of gratings and objective lenses

provided a spectral range from 3700 to 8200 Å, with a dispersion in the range of 20 – 200 Å/mm and with 0.3 – 1.1 Å/pixel after 1998. Wavelength calibration has been done using a laboratory source of He I, Ne and Ar emission lines.

The earlier data base has been obtained in 1970 - 1975, and the most of results (but not all) were published. Later, observations of objects were continued but quite irregularly - from time to time until 2005.

4 Observational data for the planetary nebulae

At the first stage of our study we compared the results of the earliest and the latest observations. One or two rows of data for the middle of specified period were included as well (Table 1). The ratios of $I(\text{O III}, 4959+5007) / I(\text{H}_\beta)$ and $I(\text{N II}, 6548+6583)/I(\text{H}_\alpha)$ were used, as an initial criterion. These lines were chosen because, as a rule, they are the strongest in a spectrum, and secondly, an influence of interstellar absorption on these ratios is minimal. The designation of the objects, observation date and ratios of line intensities are given in the 1 - 4 columns of Table 1. Corresponding references are presented in the last column. Our unpublished results are marked by "AFI".

One can see, that line ratios for the most of the objects in Table 1 do not show significant changes, i.e. ratios of the maximal values to minimal do not exceed 1.05. By the way this result confirms the reliability of our method of observations and measuring. But spectra of two objects (M1-6 and M1-11) have undergone essential changes: $I(\text{O III}, 4959+5007) / I(\text{H}_\beta)$ increased during the specified period by 40% and 110% , respectively. A little bit smaller increasing of this ratio (about 14%) is noticed in the spectrum of K3-58. Below we will consider the spectral changes of M1-6 and M1-11 in details. In the Table 2, 3 results of spectral observations for these objects are presented. The year of observation is given in the first line, The next line contains references. All values of emission line intensities are dereddened.

Six low lines contain available parameters of nebulae and central stars: EW – the equivalent widths, $\log(F_\beta)$ –logarithm of the absolute flux at H_β (in $\text{erg cm}^{-2} \text{sec}^{-1}$) N_e – the electron density determined from the $[\text{S II}], 6717/6731$ line ratio, T_e – the electron temperature calculated from $[\text{N II}]$ line ratio and/or $[\text{O III}]$ line ratio and T_{eff} determined by Zanstra method. Analysis of Table2 shows that in M1-6 the first essential increase of $[\text{O III}]$ intensities have taken place between 1975 and 1986, and at that time changes did not concern other zones of the nebula. The strengthening of He I lines began rather later, and decrease of $[\text{S II}]$ was registered only in 2004. All these facts give evidence of gradual increase of excitation degree of nebula. But we may say nothing about its reason. Both the temperature of central star T_{eff} , and electron density N_e in S^+ region look like rather constant. Some other parameters, such as $\text{EW}(\text{H}_\beta)$ and $F(\text{H}_\beta)$ show fluctuations, but without any tendency. It is appeared that an emission spectrum is more sensitive to changing of conditions in a nebula than usual methods of parameters estimation.

Spectral data for M1-11 are presented in Table 3. In spite of low precision there is good agreement between our the earliest values of $[\text{O III}]$ intensities, and data of [13]. Then some weakening of $[\text{O III}]$ intensities was registered, and errors were decreased significantly. Intensity of $[\text{O III}]$ lines began to increase gradually since 1988. Up to now other lines do not show essential change. Only the intensity of He I has a little raised. In other word the raising of excitation degree takes place only in the most inner region of nebula M1-11. Some physical parameters of M1-11 have undergone fluctuations. Increasing of $T_e(\text{O III})$ as high as twice after 1986 is very interesting fact. It may be the result of any dynamic process (shock wave?) in the inner zone of the nebula. From the other hand high value of T_e may be the reason of strengthening of forbidden lines. The value of $\text{EW}(\text{H}_\beta)$ is practically constant, but $\text{EW}(5007)$ increases in according to the intensity behavior.

5 Spectra of related objects

Three other objects from our list are identified as B[e] stars. Their spectra contain the strong continuum and Balmer emission lines of hydrogen. In addition there are [O I] 6300,6364Å lines in the spectrum of He2-446 and weak emission lines of Fe II in the spectrum of M1-15 and MWC 137. All three objects are embedded into large H II regions and have infrared excesses due to warm dust. Emission lines H_α and H_β are broad in all three objects.

Filter photometric and high resolution spectral observations have been carried out in order to study the stellar brightness and profiles of emission lines. The corresponding procedure is described in details in [17].

Selective data for these objects are presented in Table 4. This information, is not complete, but it allows to estimate the behavior of studied stars.

He2-446 is classified as a B[e] star [16]. It shows a photometric variability within 0^m2 . There is no correlation between change of brightness and EW values. One can see that between 1988 and 1995 $EW(H_\alpha)$ and $EW(H_\beta)$ have increased as high as twice. At the same time FWHM has remained practically unchanged. The profile of H_α consists of two components with the following ratio of their intensities: $I_{blue}/I_{red} = 0.70 \pm 0.05$. It may be explained as a result of stellar wind and rotating. Variations of EW is probably connected with a gradual increase of ionized mass of stellar envelope.

M1-15 is the faintest among the chosen stars, and thus it is poorly studied. All that we know are the values of $EW(H_\alpha)$, and their changes are within errors.

MWC 137 – the central star of H II region Sh2-266. According to [14] it is an evolved B[e] supergiant, probably of B0 spectral class. As it is seen from Table 4, $EW(H_\alpha)$ and $EW(H_\beta)$ show significant changes, as far as V-magnitude Spectral and photometric variability of the object may be caused by change of stellar wind power.

6 Conclusions

Long-term spectral observations of planetary nebulae and some emission-line stars were carried out within 30-35 year. The results were analyzed in order to find out any change in these objects. Fluctuations of some emission line intensities in the most of objects did not exceed 5%. Essential changes were revealed in two planetary nebulae: M1-6 and M1-11. Significant strengthening of [O III] and He I lines and weakening of [S II] (in M1- 6) testifies about change of physical conditions in the nebulae. It is reasonable to assume that all observable effects are caused by increase of T_{eff} , however available estimations of this stellar parameter don't confirm such conclusion. In case of M1-11 a sudden increase of $N_e(S II)$, $T_e(O III)$ and $T_e(N II)$ was registered in 1996. Probably these changes are connected with some dynamic events in the nebula Our study of the emission-line stars shows, that MWC137 has undergone as photometric and spectral variability. All changes are irregular and have rather low amplitude. M1-15 is appeared to be quite stable object. He2-446 showed significant increase of EW of hydrogen lines, connected probably with a growth of ionized mass of circumstellar envelope.

References

- [1] F. Sabbadin, et al., *Astron&Astrophys.***123** 147 (1983).
- [2] G. Handler. *Astron&Astrophys.***135** 493 (1999).
- [3] O. De Marco, et al., *Astrophys. J.* **602L** 93(2004).
- [4] L. N. Kondratyeva. *Izvestiya of AFIF.* **4** 100 (2005).

- [5] L. N. Kondratyeva. *Astron. Zh.* **55** 334 (1978).
- [6] A. Acker. *Catalogue of Galactic Planetary nebulae.* (1992).
- [7] T. Barker. *Astrophys. J.* **219** 914 (1978).
- [8] L. N. Kondratyeva. *Astron. Zh.* **56** 345 (1979).
- [9] J. Kaler, K. Kwitter. *Publ. Astron. Soc. Pacif.* **108** 980 (1996).
- [10] M. Dopita, C. Hua. *Astrophys. J. Supple Ser.* **108** 515 (1997).
- [11] M. Peimbert, S. Torres-Peimbert. *Rev. Mex. Astron Astrof.* **14** 540 (1987).
- [12] F. Cuisiner, A. Acker, and J. Koppen. *SAstron&Astrophys.* **307** 215 (1996).
- [13] R. Shaw, J. Kaler. *Astrophys. J. Supple Ser.* **69** 495 (1989).
- [14] C. Esteban, M. Fernandez. *Mon. Not. R. Astron. Soc.* **298** 185 (1998).
- [15] R. Oudmajjer, J. Drew. *Mon. Not. R. Astron. Soc.* **305** 166 (1999).
- [16] D. de Winter, M. Ancker, et al., *Astron&Astrophys.* **380** 609 (2001).
- [17] L. N. Kondratyeva. *Astron&Astrophys. Transactions.* **22** 181 (2003).

Table 1. Spectral data for the chosen planetary nebulae

Name	Date	$I[\text{O III}]/I(\text{H}_\beta)$	$I[\text{N II}]/I(\text{H}_\alpha)$	References
K3-58	1974-1975	14.5 ± 2.0	1.73 ± 0.24	AFI
	1988	15.2 ± 1.5	1.62 ± 0.16	[6]
	2004-2005	13.3 ± 1.0	1.59 ± 0.10	AFI
M1-5	1972-1973	4.82 ± 0.48	0.44 ± 0.05	[7]
	1971-1973	5.02 ± 0.50	0.43 ± 0.04	[5]
	2004-2005	5.08 ± 0.45	0.44 ± 0.04	AFI
M1-6	1971-1973	1.25 ± 0.12	0.60 ± 0.07	[8]
	1986	1.60 ± 0.16	0.68 ± 0.07	[6]
	2004-2005	1.72 ± 0.11	0.76 ± 0.04	AFI
M1-11	1986	0.10 ± 0.02	0.88 ± 0.10	[9]
	1996	0.14 ± 0.01	0.98 ± 0.10	[10]
	2004-2005	0.21 ± 0.04	0.95 ± 0.05	AFI
M1-12	1975	0.26 ± 0.06	0.74 ± 0.06	[8]
	1993	0.25 ± 0.04		[12]
	2004	0.25 ± 0.03	0.72 ± 0.04	AFI
M1-77	1972-1973		0.37 ± 0.05	AFI
	1983		0.33 ± 0.06	[1]
	1988		0.41 ± 0.05	[6]
	2004-2005		0.41 ± 0.04	AFI
M3-3	1972-1974	11.8 ± 1.2	4.45 ± 0.60	AFI
	1987	11.6 ± 0.9	4.36 ± 0.40	[11]
	2005	12.1 ± 0.8	4.40 ± 0.40	AFI

Table 2. Results of spectral observations of M1-6

Date of observations	1971-1975	1986	1989-1990	1995	1996	2004-2005
References	[8]	[6]	AFI	AFI	[10]	AFI
4101 H δ	23.6 \pm 3.0		25 \pm 4			25.2 \pm 2.5
4340 H γ	47 \pm 3.0		48 \pm 3	47.5 \pm 5		44.0 \pm 3.2
4471 He I	7.9 \pm 1.0					10.1 \pm 1.5
4861 H β	100 \pm 3	100	100 \pm 3	100 \pm 4	100 \pm 2	100 \pm 3
4959 [O III]	29.0 \pm 3.0		42 \pm 4	39.4 \pm 4		44.1 \pm 2.3
5007 [O III]	86.7 \pm 7.7	122	117 \pm 7	122 \pm 6		128 \pm 6.7
6548 [N II]	47.2 \pm 3.5		46.4 \pm 4.0		45.8 \pm 4.1	48.5 \pm 3.5
6563 H α	285 \pm 15	285	285 \pm 7		286 \pm 20	286 \pm 11
6583 [N II]	124 \pm 10	140	130 \pm 11		135.6 \pm 11.1	146 \pm 8
6678 He I	2.1 \pm 0.5		2.3 \pm 0.5			4.2 \pm 0.4
6717 [S II]	5.4 \pm 0.5	1.11	4.8 \pm 0.6		5.04 \pm 0.5	2.9 \pm 0.3
6731 [S II]	7.9 \pm 0.7	2.46	8.4 \pm 0.9		7.16 \pm 0.7	4.2 \pm 0.4
EW(H β)	260 \pm 60		187 \pm 23		257 \pm 20	230 \pm 20
EW(5007)			213 \pm 23			260 \pm 30
log F(H β)	-12.20 \pm 0.10				-12.269 \pm 0.005	
N $_e$ [S II]	8300 \pm 1000		8000 \pm 1000		7500 \pm 1000	8000 \pm 1000
T $_e$ [N II]	12000 \pm 1000					
T $_{eff}$	\geq 30000					\geq 29300

Table 3. Results of spectral observations of M1-11

Date of observ.	1971- 1973	1984	1986	1988- 1989	1996	1999	2004- 2005
Refer.	[8]	[13]	[9]	AFI	[10]	AFI	AFI
4101 H δ	24.2 \pm 3.2		23.1 \pm 0.5			25.8 \pm 2.7	26.4 \pm 1.7
4340 H γ	51 \pm 5.2		43 \pm 0.6			45.5 \pm 4.9	44.0 \pm 3.3
4861 H β	100 \pm 3	100	100 \pm 4	100 \pm 2	100 \pm 3	100 \pm 4	100 \pm 3
4959 [O III]	6.0 \pm 2	5.2 \pm 2	2.2 \pm 0.1	3.4 \pm 0.5	3.3 \pm 0.04	5.5 \pm 0.7	5.3 \pm 0.3
5007 [O III]	19 \pm 8		6.8 \pm 0.2	10.2 \pm 1.4	10.4 \pm 0.1	15.4 \pm 1.9	15.6 \pm 0.7
6548 [N II]	62.5 \pm 5.5		63.7 \pm 2.2	65 \pm 6	66 \pm 4	69.5 \pm 8	65.5 \pm 3.5
6563 H α	285 \pm 15	285 \pm 7	285 \pm 10	286 \pm 14	285 \pm 10	286 \pm 20	285 \pm 10
6583 [N II]	186 \pm 10	192 \pm 5	187 \pm 10	190 \pm 12	201 \pm 10	200 \pm 20	204 \pm 10
6678 HE I	0.8 \pm 0.4		0.85 \pm 0.04	1.05 \pm 0.10	1.19 \pm 0.05	1.1 \pm 0.2	1.3 \pm 0.1
6717 [S II]	0.42 \pm 0.08		0.63 \pm 0.04	0.70 \pm 0.09	0.46 \pm 0.03	0.46 \pm 0.03	0.51 \pm 0.05
6731 [S II]	0.96 \pm 0.15		1.1 \pm 0.04	175 \pm 30	0.96 \pm 0.05	0.95 \pm 0.11	1.06 \pm 0.08
EW(H β)				18 \pm 1.4	173 \pm 17	175 \pm 20	175 \pm 18
EW(5007)					19 \pm 2	40 \pm 10	30 \pm 7
logF(H β)	-11.76 \pm 0.10	-11.839 \pm 0.005	11.85 \pm 0.05		-11.781 \pm 0.003		
N $_e$ [S II]	10000 \pm 2000		4500 \pm 500	5100 \pm 900	9300 \pm 900	8300 \pm 900	8300 \pm 800
T $_e$ [N II]	10500 \pm 1000	9800 \pm 1000	10300 \pm 400		22800 \pm 800		
T $_e$ [O III]			11300 \pm 400		22000 \pm 800		
T $_{eff}$	\geq 26500	29000	27000				\geq 27000

Table 4. Spectral data for the chosen B[e] stars

Name	Date of observations	EW(H $_{\beta}$) in Å	EW(H $_{\alpha}$) in Å	FWHM(H $_{\alpha}$) in Å	Vmag	References
He2-446	1971-1973	36±5	280 ±30	6.6±0.4	14. ^m 7±0.03	AFI
	1988	35±4	330±30		14. ^m 6±0.2	AFI
	1995-1996	70±15	700±40			AFI
	2004-2005	68±7	730±60			AFI
M1-15	1971-1973		2453±30	7.8±0.4		AFI
	1991		230±17			AFI
	2004-2005		245±12			AFI
MWC137	1971-1973	47±10	300±40	5.5±0.4		AFI
	1977-1980	46±10			11. ^m 67-11. ^m 98	[16]
	1981		311±15			AFI
	1988	54.7±10	240±25	4.8±0.4		AFI
	1989-1994		254±27	4.6±0.4		[14]
	1993	60±10	295±10			AFI
	1996		550±10		11. ^m 2	[15]
	1998	39±10	200±20			AFI
2004-2005	37±5	196±10	5.0±0.3		AFI	