

**THE NEAR-INFRARED SPECTRUM OF THE BROWN
DWARF GLIESE 229B**

T.R. Geballe

Joint Astronomy Centre, 660 N. A'ohoku Pl., Hilo, HI 96720

S.R. Kulkarni

Palomar Observatory 105-24, California Institute of Technology, Pasadena, CA 91125

Charles E. Woodward^{1,2}

University of Wyoming, Department of Physics & Astronomy, Laramie, WY 82071-3905

G.C. Sloan¹

NASA Ames Research Center, MS 245-6, Moffett Field, CA 94035-1000

¹ Visiting Astronomer, United Kingdom Infrared Telescope.

² NSF Presidential Faculty Fellow.

Received _____; accepted _____

ABSTRACT

A medium resolution $1.0 - 2.5 \mu\text{m}$ spectrum of the brown dwarf, Gliese 229B has been obtained using CGS4 on UKIRT. In addition to the broad spectral structure seen in earlier low resolution observations, the new spectrum reveals a large number of absorption lines, many of which can be identified with water vapor. Water and methane are both shown to be strong absorbers in the near-infrared spectrum of the object. Several spectral features in Gl 229B that are attributable to methane match ones seen in reflection in the giant outer planets and, in particular, Titan.

Subject headings: infrared: general — infrared: stars — infrared: lines and bands — stars: individual (Gl 229B) — stars: low mass, brown dwarfs

1. INTRODUCTION

Recently, Nakajima et al. (1995) reported the first detection of a cool brown dwarf, Gliese 229B (hereafter Gl 229B), a proper motion companion of Gliese 229A. Oppenheimer et al. (1995) obtained a low resolution ($\lambda/\Delta\lambda \approx 150$) near-infrared ($1 - 2.5 \mu\text{m}$) spectrum of Gl 229B and found a number of strong absorption bands. Similar absorption bands are seen in the spectrum of Jupiter and are attributed to methane. Methane is not seen in stars. Tsuji, Ohnaka & Aoki (1995) conclude that methane will be seen only in objects cooler than about 1800 K, lower than the effective temperatures of even the least massive stars (Burrows & Liebert 1993). From the measured broad-band spectrum of Gl 229B and assuming a radius equal to that of Jupiter, Matthews et al. (1996) infer $T_{\text{eff}} = 900$ K and a bolometric luminosity of $6.4 \times 10^{-6} L_{\odot}$. The low luminosity and the presence of methane in the photosphere require that Gl 229B is a cool brown dwarf.

Here we present a new $1.0 - 2.5 \mu\text{m}$ spectrum of Gl 229B with significantly higher resolution and signal-to-noise ratio than the original spectrum presented by Oppenheimer et al. (1995). The new data provide a considerably more detailed view of Gl 229B. We compare our spectrum to spectra of the Jovian planets and Titan which are known to show strong absorption features due to methane in the near-infrared.

2. OBSERVATIONS AND DATA REDUCTION

We obtained spectra of Gl 229B in the $0.99 - 2.52 \mu\text{m}$ interval at the United Kingdom 3.8 m Infrared Telescope (UKIRT) on 1995 October 28-29 UT and December 12-13 UT, using the facility spectrometer CGS4 (Mountain et al. 1990). The 75 l/mm grating and the 150 mm focal length camera optics in CGS4 imaged a slit $90''$ long onto a 256×256 InSb array. Each pixel spanned $1''.23$, and the slit was one pixel wide. Table 1 provides details of

the observations. The resolution of CGS4 was approximately $390 \times \lambda (\mu\text{m})$ in first order and $780 \times \lambda (\mu\text{m})$ in second order. To calibrate the spectra, we observed HR 1849 (A0V, $V=5.55$) each night just before observing Gl 229B.

Gl 229B is about $8''$ from Gl 229A. In the J, H, and K bands Gl 229A outshines Gl 229B by approximately 10 magnitudes. At UKIRT the halo of diffracted and scattered light from Gl 229A contributed a large amount of radiation to the array rows containing the Gl 229B spectrum. In order to minimize the difficulties of removing this background, we observed with the slit of CGS4 perpendicular to the line joining the two sources on the sky and obtained alternate spectra of Gl 229B by nodding the telescope 10 rows along the slit. Subtracting these alternate spectral images removed much of the halo, allowing Gl 229B to be detected clearly at nearly all wavelengths, typically at greater strength than the residual halo from the primary. We extracted the spectrum of Gl 229B using standard infrared reduction procedures to correct for curvature of the spectra and to remove residual background. Spectra of HR 1849 were utilized to correct for atmospheric and instrumental absorption, after editing them to remove atomic hydrogen absorption lines (with the exception of Pa α at $1.875 \mu\text{m}$).

Figure 1 presents the complete spectrum of Gl 229B from 0.99 to $2.52 \mu\text{m}$ (a numerical file of this spectrum is available from the authors upon request). A more expanded version of the spectrum is provided in Fig. 2. To produce the spectrum, we slightly smoothed and rebinned the observed spectra in steps of $0.0005 \mu\text{m}$ for the second order segments and $0.0010 \mu\text{m}$ for the first order segments. The smoothing lowers the spectral resolving power by $\approx 10\%$ from the actual instrumental resolution cited above. In the region of overlap between the two orders, we chose to use the first order spectrum beyond $1.585 \mu\text{m}$. We scaled the individual segments (by factors ranging from 0.7 to 1.2) to match the spectra in overlapping sections and at $1.585 \mu\text{m}$. The average scaling factor was unity, but we

scaled the final spectrum by a factor of 0.76 to more accurately match (to within about 0.1 mag) the recent Palomar photometry of Gl 229B (J=14.2, H=14.3, Ks=14.3, K=14.4) as reported by Matthews et al. (1996a).

The noise in the CGS4 spectrum varies considerably with wavelength, as illustrated in Fig. 2, due to large variations in atmospheric transmission and in the background (caused by both thermal and OH emissions). In particular, in the spectral regions dominated by strong telluric absorption bands (1.12 – 1.14 μm , 1.36 – 1.42 μm , 1.81 – 1.93 μm , and 2.48 – 2.52 μm), apparent spectral features may not be real. In these spectral regions the data probably are only useful for estimating the average flux level.

3. DESCRIPTION OF THE INFRARED SPECTRUM

The new spectrum of Gl 229B is consistent with the lower resolution spectrum of Oppenheimer et al. (1995). However, the latter spectrum was incorrectly normalized, as noted by Matthews et al. (1996a). Matthews et al. (1996b) present a revised spectrum with the proper normalization, with which the CGS4 spectrum agrees well.

Two new results are immediately apparent from the new spectrum. First, emission is seen across the entire IR band from 1.0 to 2.5 μm . Specifically, emission is detected through the wavelength intervals of strong telluric H₂O absorption (see discussion above) and in the range 2.2 – 2.5 μm including a notable rise in flux at wavelengths longward of 2.4 μm . The data of Oppenheimer et al. (1995) were restricted to the usual near-infrared windows (Z, J, H, K) and furthermore were noisy in the range 2.2 – 2.5 μm . We emphasize that, with the possible exception of the wavelength interval 1.36 – 1.40 μm , where the detection is marginal, the emission detected is from Gl 229B and is not contamination by Gl 229A; crosscuts through the differenced spectral images in these wavelength intervals

show local maxima in the rows of the array corresponding to the location of Gl 229B. Second, numerous narrow lines are seen (see Fig. 2). These appear in regions of high emission and on all of the absorption edges with the exception of the very steep edge at $1.6 \mu\text{m}$. Broader features occur in the $1.6 - 1.9 \mu\text{m}$ absorption trough at about 1.63 , 1.67 , 1.71 , and $1.80 \mu\text{m}$, and at 2.17 , and $2.20 \mu\text{m}$. Most of the broad features were seen in the spectrum of Oppenheimer et al. (1995), albeit at lower resolution.

4. DISCUSSION

4.1. Methane: comparison with Solar System objects

We have compared the new spectrum of Gl 229B to the solar reflectance spectra of three of the giant gaseous planets (Jupiter, Saturn, and Uranus) and Titan, using the data of Fink & Larson (1979), and recent CGS4 spectra of Titan (Owen, unpublished) and Saturn (Geballe, unpublished). Methane dominates the $1.0 - 2.5 \mu\text{m}$ spectra of all of the above outer planets and Titan, but the degree of domination varies, increasing in the sequence Jupiter, Saturn, Titan, and Uranus.

The spectrum of Titan provides the best overall spectral match to that of Gl 229B and the spectra of the two exhibit several similar details. Most remarkably, their very sharp cut-offs at $1.61 \mu\text{m}$ are essentially identical (see Fig. 3). The matches with the spectra of the other Solar System objects do not appear as precise; for example, the cut-off in Saturn is shifted by $0.01 \mu\text{m}$ to longer wavelength. Between 1.62 and $1.72 \mu\text{m}$ the wavelengths of some of the absorption maxima in Titan and Gl 229B agree. Both Titan and Gl 229B show absorption bands at $\sim 2.17 \mu\text{m}$ and $2.20 \mu\text{m}$ (as do Jupiter and Saturn). In Titan each of these features arises from CH_4 (Fink & Larson 1979; C. Griffith, pers. comm.). Thus, despite the considerably different physical conditions of the CH_4 in Gl 229B compared

to those in the atmospheres of the giant gaseous planets and Titan, detailed comparison provides clear confirmation of the strong influence of methane on the spectrum of Gl 229B.

4.2. H₂O in Gl 229B

Theoretical model spectra of brown dwarfs with effective temperatures similar to that of Gl 229B show considerable absorption due to steam (Marley et al. 1996; Allard et al. 1996). We note, however, that these model spectra only include low resolution methane opacities shortward of 1.6 μm (R.S. Freedman, personal communication). Jones et al. (1994) have compiled low resolution spectra of the coolest stars and attribute a broad absorption centered at about 1.4 μm and its sharp absorption edge, whose wavelength they report as 1.34 μm , to H₂O. The 1.4 μm water band also appears strongly in the above model brown dwarf spectra. Following Jones et al., Oppenheimer et al. (1995) identified the absorption edge seen in Gl 229B at about this wavelength as due to H₂O.

In the present spectrum, the absorption edge in Gl 229B is seen to be centered at about 1.31 μm . The H₂O absorption edge in late M stars actually occurs at 1.33 μm (Walker et al. 1996) and is much steeper than the absorption edge in Gl 229B; indeed it is expected to strengthen and steepen with decreasing temperature. The methane absorption edges of Titan and Saturn are centered at 1.30 μm and 1.31 μm , respectively. The work of Jones et al. is restricted to stars which are cool, but certainly not cool enough to have strong photospheric methane bands. Thus, confusion between CH₄ and H₂O does not arise in the interpretation of stellar spectra. We conclude on the basis of the low resolution spectra alone that the absorption edge and trough longward of 1.3 μm in Gl 229B cannot be attributed unambiguously to water alone and that methane could contribute significantly. However, when compared to the spectra of all of the giant outer planets and Titan, the emission bump at 1.5 – 1.6 μm is badly eaten away on its short wavelength side (Fig. 3).

Absorption by H₂O in Gl 229B can explain this systematic difference between Gl 229B and the giant planets and Titan. The sharpest portion of the 1.3 μm absorption edge of Gl 229B, at 1.33 μm , would then be due to H₂O.

Although the low resolution spectra are ambiguous as to the presence of water vapor, comparison of the new higher resolution spectrum of Gl 229B with plots of H₂O opacity at similar resolution (kindly supplied by D. Saumon, R.S. Freedman, and D. Schwenke) demonstrate its presence clearly. An illustration of this is provided in Fig. 4. Almost every absorption line in the spectrum of Gl 229B in the intervals 1.30 – 1.34 μm , 1.53 – 1.58 μm , and 1.95 – 2.07 μm has a counterpart in the water opacity spectrum. Moreover, there is good overall correlation between observed and modelled line strengths. The opacities of the lines in the above wavelength intervals are relatively low, implying that the depressions in Gl 229B at about 1.4 μm and 1.9 μm are indeed due to absorption by H₂O, which has much higher opacity at those wavelengths.

Thus, the new spectrum allows a firm identification of water vapor in the photosphere of Gl 229B. It is clear that, as predicted in the models, both H₂O and CH₄ produce strong absorptions in the near-infrared spectrum of Gl 229B. However, until the line parameters of methane are better known, it will be difficult both to disentangle the relative contributions of methane and water and to search for other absorbers at near-infrared wavelengths.

5. Acknowledgements

We thank the many individuals responsible for the excellent construction, maintenance, and operational support of CGS4 and the staff of UKIRT for its support of these observations. We are grateful to R.S. Freedman, D. Schwenke, and D. Saumon for providing us with detailed opacity plots of H₂O and CH₄ and to T. Owen for permitting us to publish

a portion of the spectrum of Titan. We thank the referee, F. Allard, for several helpful comments. We also thank D. Cruikshank, M. Marley, K. Matthews, T. Nakajima, and B. Oppenheimer for their assistance. The United Kingdom Infrared Telescope is operated by the Joint Astronomy Centre on behalf of the U.K. Particle Physics and Astronomy Research Council. SRK’s research is supported by Caltech, the US NSF, NASA, and the Packard Foundation. CEW acknowledges support from the US National Science Foundation.

6. REFERENCES

- Allard, F., Hauschildt, P.H, Baraffe, I. & Chabrier, G. 1996, ApJ (Letters), in press.
- Burrows, A. & Liebert J. 1993, Rev. Mod. Phys., 65, 301
- Fink, U., & Larson, H. P. 1979, ApJ 233, 1021
- Jones, H. R. A., Longmore, A. J., Jameson, R. F., & Mountain, C. M. 1994, MNRAS 267, 413
- Marley, M.S., Saumon, D., Guillot, T., Freedman, R.S., Hubbard, W.B., Burrows, A. & Lunine, J.I. 1996, Science, submitted
- Matthews, K., Nakajima, T., Kulkarni, S., & Oppenheimer, B. 1996a, I.A.U. Circ. No. 6280
- Matthews, K., Nakajima, T., Kulkarni, S.R. & Oppenheimer, B. 1996b, ApJ, submitted
- Mayor, M., & Queloz, D. 1995, I.A.U. Circ. No. 6251
- Mountain, C.M., Robertson, D.J., Lee, T.J. & Wade, R. 1990, in Instrumentation in Astronomy VII, D.L. Crawford Ed., (Proc. SPIE, 1235), 25
- Nakajima, T., Oppenheimer, B.R., Kulkarni, S.R., Golimowski, D.A., Matthews, K. & Durrance, S.T. 1995, Nature, 378, 463
- Tsuji, T., Ohnaka, K., Aoki & W. 1995, in The Bottom of the Main Sequence - and Beyond, C.G. Tinney, Ed. (Springer, Berlin), p. 45
- Walker, H.J., Tsikoudi, V., Clayton, C.A., Geballe, T.R., Wooden, D.H. & Butner, H.M. 1996, A&A, submitted

7. FIGURE CAPTIONS

Fig. 1.— Combined 1.0 – 2.5 μm spectrum of Gl 229B. The intensity scale is logarithmic; negative values near 1.4 μm were removed prior to plotting

Fig. 2.— Spectral segments for Gl 229B, covering 0.99 – 2.52 μm . Representative noise levels are shown as vertical lines along the bottom of each panel. a. 0.99 – 1.41 μm . b. 1.39 – 1.91 μm . c. 1.89 – 2.52 μm .

Fig. 3.— Spectra of Gl 229B (darker line) and Titan (scaled by a factor of ~ 0.003) near the 1.6 μm methane absorption edge.

Fig. 4.— Spectrum of Gl 229B and the opacity of water vapor in the 1.95 – 2.10 μm region. The opacity is for $P = 1$ bar and $T = 700$ K, convolved to a resolving power of 1000, and is courtesy of R.S. Freedman, D. Schwenke, and D. Saumon.

Table 1. Observation Log

Date (UT)	Grating order	$\lambda(\mu\text{m})$	Int. Time	Conditions
1995 Dec. 12	second	0.99-1.33	18 m	photometric
1995 Dec. 12	second	1.27-1.61	24 m	photometric
1995 Oct. 29	first	1.48-2.15	24 m	photometric
1995 Oct. 28	first	1.85-2.52	30 m	partly cloudy
1995 Dec. 13	first	1.85-2.52	36 m	photometric