## Andreev scattering in the asymmetric ladder with preformed bosonic pairs

Karyn Le Hur

Département de Physique Théorique, Université de Genève, CH-1211, Genève 4, Switzerland

(October 26, 2018)

We discuss the phase coherence which emanates from the ladder-like proximity effect between a "weak superconductor" with preformed bosonic pairs (here, a single-chain Luther-Emery liquid with superconducting correlations that decay approximately as  $x^{-1}$ ) and a Fermi gas with unpaired fermions. Carefully studying tunneling mechanism(s), we show that the boson-mediated Cooper pairing between remaining unpaired electrons results in a quasi long-range superconductivity: Superconducting correlations decay very slowly as  $x^{-\eta}$  with  $\eta \approx 1/2$ . This process is reminiscent of the coupling of fermions to preformed bosonic pairs introduced in the context of high- $T_c$  cuprates.

PACS numbers: 71.10.Pm, 74.50.+r

Surprisingly, the excitation spectrum in underdoped high- $T_c$  cuprates exhibits a pseudogap at a temperature  $T^*$  far above the superconducting temperature  $T_c$ . Experimental results suggest that the gap formation is due to the pairing of electrons in the corners of the Fermi surface into preformed bosonic pairs (with a small spectral weight), whereas electrons in the "diagonals" would still remain unpaired (See Fig. 1) [1]. Theoretically, the truncation of the Fermi surface has been predicted by Rice et al. from Renormalization Group arguments in two dimensions [2] and from the behavior of lightly doped ladders [3]. This has been also emphasized by Lee and Wen based on gauge theory calculation [4]. The superconductivity which emanates from the Bose condensation of these preformed pairs coexisting with unpaired fermions has been examined phenomenologically by Geshkenbein, Ioffe and Larkin [5]. Below, we introduce a simple asym*metric* two-leg ladder system which allows to rigorously exemplify the (almost) Bose condensation of preformed bosonic pairs in the vicinity of unpaired fermions.

We discuss the quasi long-range superconductivity emerging at  $T_c$  from the ladder-like proximity effect between a one-dimensional (1D) weak superconductor which already displays a spin gap and preformed Cooper pairs below  $T^*$  — and a 1D Fermi gas.



FIG. 1. A ladder system which allows to investigate the Bose condensation of preformed pairs coexisting with unpaired fermions (discussed for high- $T_c$  cuprates): A single-chain Luther-Emery liquid weakly-coupled to a Fermi gas.

To keep the discussion as clear as possible, here we assume that the weak superconductivity emerges from the single-chain Hubbard model with small attractive interactions, producing the usual Luther-Emery liquid [6]. Above  $T_c$  (but below  $T^*$ ), the superconducting correlations are weak in the sense that they still decay approximately as the charge density wave correlations, i.e., roughly as  $x^{-1}$ . Such a prototype system has some similarities to the one introduced by Emery, Kivelson and Zachar in another context [a metallic stripe coupled to an active spin-gapped doped Mott insulator] [7]. Following the methodology developed in Ref. [8], we build the relevant tunneling process(es) between the two chains. Here, this corresponds to an Andreev scattering mechanism [9]. Then, we show how the boson-induced Cooper pairing between unpaired fermions results below  $T_c \propto 1/\{T^*\}^2$ both in a spinon-pairing (spin gap) in the Fermi gas [7] and in a quasi long-range phase coherence. In contrast with Ref. [7] (page 6131), we report that the superconducting correlations now decay as  $x^{-\eta}$  with  $\eta \approx 1/2$ , whereas the charge density wave correlations at the wave vector  $q = 2k_F$  arise only at short distances. The system behaves as a conventional two-leg ladder [10,11,8].

For temperatures  $T \gg T^*$ , the kinetic energy for fermions takes the standard form  $\mathcal{H}_{kin} = \mathcal{H}_o + \mathcal{H}_{\perp}$  where

$$\mathcal{H}_{o} = -t \sum_{j,\alpha} \psi_{j\alpha}^{\dagger}(x+1)\psi_{j\alpha}(x) + \text{H.c.},$$
$$\mathcal{H}_{\perp} = -t_{\perp} \sum_{\alpha} \psi_{2\alpha}^{\dagger}(x)\psi_{1\alpha}(x) + \text{H.c.}$$
(1)

Here j = (1,2) denote the chains and  $\alpha = (\uparrow, \downarrow)$  the spin projections of an electron. The bare short-distance cutoff is equal to a = 1. The longitudinal and transverse hopping amplitudes are respectively t and  $t_{\perp}$ .

We consider the case of a weak-proximity effect between the two chains, i.e., we start with  $t_{\perp}/T^* \ll 1$ . For weak interactions in chain 1, our temperature  $T^*$  will be already very small compared to the Fermi energy  $E_F \sim t$ .

Focusing on electronic states near the Fermi points,

we expand  $\psi_{j\alpha}(x) = \psi_{+j\alpha}(x)e^{ik_Fx} + \psi_{-j\alpha}(x)e^{-ik_Fx}$ ; Below  $p = \pm$  denote respectively right and left excitations. Unlike in Ref. [7], the two chains are here equally doped.

Weak superconductor at  $\mathbf{T}^*$ . Let us first consider that  $t_{\perp} = 0$ . As emphasized in the introduction, we start with an asymmetric situation where in chain 2 electrons are completely "free" whereas electrons of chain 1 are subjected to a weak attractive Hubbard interaction of the form:  $\mathcal{H}_1 = g_c J_{+11} J_{-11} - g_s \mathbf{J}_{+11} \mathbf{J}_{-11}$ . We have rewritten the Hubbard interaction in chain 1 as a function of the usual charge and spin currents [For the definitions: See Ref. [8], Eq. (5)]. Note that  $g_c$  and  $g_s$  describe chargeand spin backscatterings respectively.

The bare interactions are of the order of the *negative* Hubbard coupling, U < 0.

Bosonizing the 1D Fermi fields, we can write [11]

$$\psi_{pj\alpha} \approx \exp\left(i\sqrt{\frac{\pi}{2}}\left[p(\Phi_{jc} + \alpha\Phi_{js}) - (\theta_{jc} + \alpha\theta_{js})\right]\right), \quad (2)$$

 $\alpha = \pm$  for spin up and spin down respectively. The electron spectrum exhibits spin-charge separation. The charge part of the Hamiltonian results in the Luttinger Gaussian model [12]:

$$H_{oj}^{c} = \frac{v_F}{2\pi} \int dx \, \frac{1}{K_{jc}} (\rho_{jc} - \rho_o)^2 + K_{jc} (\nabla \theta_{jc})^2.$$
(3)

Here  $v_F = 2t \sin k_F$  denotes the Fermi velocity,  $\partial_x \Phi_{jc} = (\rho_{jc} - \rho_o)$  measures fluctuations of charge density in each chain, and  $\nabla \theta_{jc}$  is the conjugate momentum to  $\Phi_{jc}$ . The free spin Hamiltonians have the same forms as the charge Hamiltonian of Eq.(3) replacing  $\Phi_{jc}$  by  $\Phi_{js}$ . The Luttinger exponents are,

$$K_{1c} = 1 - g_c / \pi v_F > 1 \qquad K_{2c} = 1$$
(4)  
$$K_{1s} = 1 + g_s / \pi v_F < 1 \qquad K_{2s} = 1.$$

For attractive interactions in chain 1  $[K_{1s} < 1]$ , the spin backscattering term  $g_s$  provides another (relevant) contribution to the low-energy physics. Indeed, this produces the Sine-Gordon model [6,11]

$$H_1^s = H_{o1}^s - g_s \int dx \, \cos[\sqrt{8\pi}\Phi_{1s}](x). \tag{5}$$

The coupling  $g_s$  is known to be strongly renormalized at the temperature scale  $T^* = E_F \exp(-\pi v_F/g_s)$ . The spin field gets pinned at the classical value  $\Phi_{1s} \approx 0$ . The resulting spin gap is of the order of  $T^*$  (See Ref. [11] page 76). This produces the growth of the superconducting (SC) and charge density wave (CDW) correlations in chain 1. These are given by the operators

$$\mathcal{O}_{SC}^{1} = \psi_{+1\uparrow}\psi_{-1\downarrow} + \psi_{-1\uparrow}\psi_{+1\downarrow} \qquad (6)$$
$$\propto \exp(-i\sqrt{2\pi}\theta_{1c})\cos(\sqrt{2\pi}\Phi_{1s})$$

and

$$\mathcal{O}_{CDW}^{1} = \psi_{\pm 1\uparrow}^{\dagger}\psi_{-1\uparrow} + \psi_{\pm 1\downarrow}^{\dagger}\psi_{-1\downarrow} \qquad (7)$$
$$\propto \exp(-i\sqrt{2\pi}\Phi_{1c})\cos(\sqrt{2\pi}\Phi_{1s}).$$

The operator  $\mathcal{O}_{SC}^{1\dagger}$  describes the *preformed bosonic pairs* with charge Q = 2e and spin S = 0. Using the fact that there is no fluctuation of the spin field  $\Phi_{1s} \approx 0$  for temperatures smaller than  $T^*$ , one finds for the charge-density correlation function

$$<\mathcal{O}_{CDW}^{1\dagger}(x)\mathcal{O}_{CDW}^{1}(0)>\propto x^{-K_{1c}}$$
(8)

and for the SC pairing correlation function

$$<\mathcal{O}_{SC}^{1\dagger}(x)\mathcal{O}_{SC}^{1}(0)>\propto x^{-1/K_{1c}}.$$
(9)

As expected, since  $K_{1c} > 1$  the superconducting correlations in chain 1 are (slightly) more important than the CDW correlations for  $T < T^*$  (See Fig. 2). Here, pairing of spinons is somewhat equivalent to pairing of electrons inducing prevalent bosonic pairs in chain 1.

As long as the chain 2 remains weakly "coupled" to chain 1 (i.e., for  $T > T_c$ , see discussion below), all its density-density correlation functions decay as  $1/x^2$  as for any noninteracting 1D electron gas.

Fate of transverse electron motion. Now, we examine the influence of the finite (bare) transverse hopping amplitude. Again, we assume that the spin backscattering term  $g_s$  is flowing *first* to strong couplings (at the temperature  $T^*$ ;  $g_s(T^*) \approx t$ ) which corresponds to very small bare values of  $t_{\perp}$  ( $t_{\perp} \ll T^*$ ). The bosonic form of the term  $t_{\perp}$  (at q = 0) is given, e.g., in Ref. [8].

We find that  $t_{\perp}$  evolves according to

$$\frac{d\ln t_{\perp}}{dl} = \frac{7}{4} - \frac{1}{4} \left( K_c^- + \frac{1}{K_c^-} \right) - \frac{1}{8} \left( K_{1s} + \frac{1}{K_{1s}} \right)$$
(10)

The *l* describes the renormalization of the short-distance cutoff  $a(l) = \exp l$ . The renormalization procedure is stopped at lengths a(l) comparable to the thermal length  $v_F/T$ , which means  $l = \ln(E_F/T)$ . We have combined the charge boson fields in the two chains into a symmetric "+" and antisymmetric "-" part:  $\Phi_c^{\pm} = [\Phi_{1c} \pm \Phi_{2c}]/\sqrt{2}$ and similarly for the conjugate momenta. The resulting Luttinger exponents obey

$$K_c^{\pm} = \sqrt{K_{1c}K_{2c}} \approx 1 - \frac{g_c}{2\pi v_F} > 1.$$
 (11)

Furthermore, we obtain

$$\frac{dK_{1s}}{d\ln(\frac{E_F}{T})} = -\frac{1}{2}(g_s K_{1s})^2.$$
 (12)

As long as  $T > T^*$ , integrating Eq.(10) produces a linear growth of  $t_{\perp}(T)$  [i.e.,  $g_s(T) \ll t$  and  $K_{1s} \approx 1$ ]

$$t_{\perp}(T) \approx t_{\perp} E_F / T \ll t. \tag{13}$$

Second, from Eq.(12) we deduce that the explicit divergence of the spin backstattering  $g_s$  at the opening of the spin gap (i.e., for  $T = T^*$ ) results formally in  $K_{1s}\{T < T^*\} = 0$ . Using Eq.(10), this also implies

$$t_{\perp} \{ T < T^* \} = 0. \tag{14}$$

This produces a jump in the electron hopping amplitude. This can be interpreted as a consequence of the disappearance of the electron states on chain 1. At  $T = T^*$ , in contrast chain 1 exhibits preformed bosonic pairs due to the occurrence of the spin gap. In consequence, one expects the vanishing of the single-particle transverse susceptibility  $\chi_{\perp,s}$  [13] (See Ref. [11] page 224). This reflects the suppression of the electron motion in the transverse direction. Such a phenomenon can also emerge due to the presence of strong Umklapp scattering (leading to a charge gap) [8].

On the other hand, the Hamiltonian must be supplemented by extra terms (i.e., tunneling processes) which are generated in the course of renormalization by expanding the partition function as a function of  $t_{\perp}$  [11]. For a two-chain model, those are precisely known and have been classified by Khveshchenko and Rice using the bosonic language [14]. Following the scheme introduced in Ref. [8], we now examine which tunneling processes are relevant below  $T^*$ . We will show that because  $K_c^- > 1$ , these correspond to an incoming electron from chain 2 being reflected back as a hole, thereby injecting an additional Cooper pair in chain 1. This is an Andreev scattering mechanism [9].

Andreev scattering amplitude. Terms containing the spin operator  $\exp[i\beta\theta_{1s}]$  with  $\beta = \sqrt{2\pi}$  may be already dropped. Indeed, below  $T^*$  these terms naturally acquire a scaling dimension greater than 2 and thus become irrelevant in the sense of the renormalization procedure. Using the classification scheme of Ref. [14] away from half-filling, one must then keep the following interchain "pair-hopping" terms

$$\Delta H = \cos\sqrt{4\pi}\theta_c^- \left\{ g_2 \cos\sqrt{4\pi}\Phi_s^- + g_3 \cos\sqrt{4\pi}\Phi_s^+ \right\}$$
(15)  
+  $g_5 \cos\sqrt{4\pi}\Phi_c^- \cos\sqrt{4\pi}\Phi_s^-.$ 

To fix the values of the coupling constants for  $T \approx T^*$ , one must proceed as follows [8].

Above  $T^*$ , the couplings  $g_i$  with i=(2,3) evolve in a similar manner as  $(z(l) = t_{\perp}(l)/E_F)$ 

$$\frac{dg_i}{dl} = \lambda g_i - \lambda z^2, \tag{16}$$

with

$$\lambda = 2 - [K_s^{\pm} + \frac{1}{K_c^{-}}] \approx -(g_c + g_s)/2\pi v_F > 0 \qquad (17)$$

being slightly renormalized. As long as the spin backscattering is small, one can use the exponents  $(K_s^{\pm} < 1)$ :

$$K_s^{\pm} = \sqrt{K_{1s}K_{2s}} \approx 1 + \frac{g_s}{2\pi v_F} = \frac{1}{2}(K_{1s} + K_{2s}).$$
 (18)

The bare values are given precisely by  $z(0) = t_{\perp}/E_F$ and  $g_i(0) = 0$  [11]. Under renormalization, these pairhopping terms then acquire a small but *non-zero* value

$$g_{2,3}(l) = -\frac{\lambda}{2-\lambda} z(0)^2 \Big[ \exp(2l) - \exp(\lambda l) \Big]$$
(19)  
$$\approx -\frac{\lambda}{2-\lambda} z(l)^2,$$

with  $z(l) = z(0) \exp l$ . Using Eqs.(13) and (19) for  $T \to T^*$ , one finally finds  $(|g_{2,3}(T^*)| \ll 1)$ 

$$g_{2,3}(T^*) \propto -\frac{\{t_{\perp}(T^*)\}^2}{E_F^2} = -\frac{t_{\perp}^2}{\{T^*\}^2}$$
 (20)

The exact prefactor is not of interest here.

The evolution of the coupling  $g_5$  can be discussed in an identical way. Starting with *free (or almost free)* electrons on chain 2, we obtain the same equation as for  $g_{2,3}$ with  $\lambda$  replaced by [15]

$$\gamma = 2 - [K_s^- + K_c^-] \approx (g_c - g_s)/2\pi v_F = 0.$$
 (21)

Therefore, one reaches the important conclusion that in our case  $g_5(T) = 0$  whatever the temperature. Note that in principle, this does not remain true if (quite strong) repulsive interactions are added between unpaired remaining electrons. This provides both  $K_c^- < 1$  and  $K_s^- < 1$ [16]. In that case, the coupling  $g_5$  becomes strongly relevant leading to another spin-gapped (but à priori not superconducting) fixed point [8,17].

When T approaches  $T^*$ , the spin field  $\Phi_{1s}$  gets locked, resulting in  $\langle \cos \sqrt{2\pi} \Phi_{1s} \rangle \sim \{T^*a\}^{1/2}$  (we put a = 1) and then in the unique pair-tunneling mechanism

$$\Delta H = V \cos \sqrt{4\pi} \theta_c^- \cos \sqrt{2\pi} \Phi_{2s}.$$
 (22)

This term is the driving force for the quasi long-range Bose condensation of the preformed pairs. We stress that in 1D, the (bare) interchain hopping is sufficient to generate such (particle-particle) pair-hopping term, i.e.,  $V = -t_{\perp}^2/\{T^*\}^{3/2}$ . It is appropriate to write

$$\Delta H = V \Big( \mathcal{O}_{SC}^{1\dagger} \big( \psi_{+2\uparrow} \psi_{-2\downarrow} + \psi_{-2\uparrow} \psi_{+2\downarrow} \big) + \text{H.c.} \Big).$$
(23)

We immediately recognize the boson-mediated Cooper pairing between remaining unpaired electrons, introduced in the context of high- $T_c$  cuprates [5]. This also corresponds to an Andreev reflection [9].

It is worth to note that the coupling V does not affect the quantum coherence along the chains. The charge current of each chain  $J_j \propto v_F \int dx \nabla \theta_{jc}(x)$  commutes with the process V, i.e.,  $[J_j, \Delta H] = 0$ . For clean and infinitely long chains, the conductivity  $\sigma_{\parallel}$  remains infinite. To prompt for the question of the (in)coherence in the transverse direction below  $T^*$  one can examine the transverse susceptibility  $\chi_{\perp,p}$  induced by the Andreev process (transfer of pairs). As long as V is small, we can expand

the partition function up to terms of fourth order in V. Using Ref. [11] (page 223), we find  $\chi_{\perp,p} \propto T^{2(d_{\perp,p}-1)}$  and  $d_{\perp,p} = (K_{2s}/2 + 1/K_c^-)$  [13]. Near  $T^*$ , we immediately get  $d_{\perp,p} > 1$ . Thus,  $\chi_{\perp,p}$  is first reduced by decreasing the temperature, still reflecting a certain incoherence between the bosonic pairs and the unpaired electrons. At lower temperatures the system converges to a superconducting and completely coherent fixed point. Below  $T^*$ , the coupling V evolves according to  $dV/d\ln\left(\frac{T^*}{T}\right) =$  $(2 - d_{\perp,p})V$  [18]. This will be strongly renormalized at the temperature,  $T_c \approx T^* V^2 = a t_{\perp}^4 / \{T^*\}^2 \ll T^*$ . Close to  $T_c$ , the exponents  $1/K_c^-$  and  $K_{2s}$  drastically decrease [similarly as in Eq.(12)] resulting in  $d_{\perp,p}\{T_c\} \ll 1$ . A strong interchain coherence thus arises below  $T_c$ ;  $\chi_{\perp,p}$ diverges at T = 0. The proximity effect results in a fluid of coherent bosons with (almost) long-range pairing.

As in Ref. [7], the pre-existing spin gap is conveyed to the Fermi gas.

Superconductivity at  $\mathbf{T} = \mathbf{0}$ . The spin field  $\Phi_{2s}$  gets massive [See Eq.(22)], so the spin fluctuations now contribute a multiplicative constant to all the correlation functions. The SC phase below  $T_c$  thus becomes only a property of the charge degrees of freedom. Here, the And reev reflection imposes that the field  $\theta_c^-$  gets locked (for all x), i.e.,  $\theta_{1c}(x) = \theta_{2c}(x)$ . Only the superfluid phase  $\theta_c^+$  remains massless, thereby producing strong superconductivity. In each chain, one must write (j = 1, 2)

$$\sqrt{2}\theta_{jc} = \frac{1}{\sqrt{2}} \left\{ \theta_{1c} + \theta_{2c} \right\} \approx \theta_c^+.$$
(24)

Using Eq.(6), this gives

$$\mathcal{O}_{SC}^j \propto \exp(-i\sqrt{\pi}\theta_c^+).$$
 (25)

The two chains become obviously phase-coherent and the SC correlation functions decay very slowly with x,

$$<\mathcal{O}_{SC}^{j\dagger}(x)\mathcal{O}_{SC}^{j}(0)>\propto x^{-1/(2K_c^+)}.$$
(26)

(The exponent is in contrast with the one of Ref. [7] at Page 6131; Our  $T_c$  corresponds to their  $T_2^*$ ). This exemplifies the almost Bose condensation of preformed pairs due to the exchange of fermions even though a phase order is not strictly possible in 1D, i.e.,  $\langle \mathcal{O}_{SC}^j(x) \rangle = 0$ .

This behavior is reminiscent of a conventional twoleg ladder material (our chains behave as bands of the symmetric two-leg ladder with repulsive interactions) [10,11,8]. We deduce that the pairing susceptibility diverges approximately as  $\chi_{SC} \propto T^{-3/2}$  (See Fig. 2). Each chain is now characterized by the same superflow. Since  $\theta_c^- \approx 0$ , this indeed produces  $J_1 \approx J_2$ . The density of Cooper pairs in each chain  $n_i$  fluctuates strongly, i.e.,  $(n_1 - n_2) \propto \partial_x \Phi_c^-$ . Finally, the CDW fluctuations can develop only at very short distances [19]. To conclude, we have introduced an asymmetric two-leg ladder system which allows to rigorously investigate the (almost) Bose condensation of preformed bosonic pairs in the vicinity of unpaired fermions discussed in the context of high- $T_c$ cuprates. The Andreev scattering mechanism has been derived properly from the fermionic model (above  $T^*$ ).

We already like to push forward the fact that this asymmetric model can be generalized to preformed bosonic pairs having an approximate d-wave pairing and a very small spectral weight (and coexisting with chain(s) of unpaired fermions), e.g., by investigating asymmetric three-leg ladder systems [20–22].

Acknowledgment. We thank Lev Ioffe for inspiring us this work, and T.Maurice Rice for discussions on superconductivity with preformed pairs.

- [1] For a review, See T. Timusk and B. Statt, Rep. Prog. Phys. 62, 61 (1999).
- C. Honerkamp, M. Salmhofer, N. Furukawa, and T. M. |2|Rice, Phys. Rev. B 63, 035109 (2001), and references therein.
- T. M. Rice, S. Haas, M. Sigrist, and F. C. Zhang, Phys. [3] Rev. B 56, 14655 (1997); U. Ledermann, K. Le Hur, and T. M. Rice, *ibid.* **62**, 16383 (2000).
- P. A. Lee and X. G. Wen, Phys. Rev. Lett 78, 411 (1997).
- [5]V. G. Geshkenbein, L. B. Ioffe, and A. I. Larkin, Phys. Rev. B 55, 3173 (1997).
- A. Luther and V. J. Emery, Phys. Rev. Lett 33, 589 [6](1974).
- V. J. Emery, S. A. Kivelson, and O. Zachar, Phys. Rev. [7]B 56, 6120 (1997).
- [8] K. Le Hur, Phys. Rev. B 63, 165110 (2001).
- [9] P.-G. de Gennes, Superconductivity of Metals and Alloys, Benjamin, New York, 1966.
- [10] M. Fisher, in Topological Aspects of low-dimensional system, Les Houches LXIX, edited by A. Comtet et al. (EDP Sciences, Les Ulis & Springer, Paris, 1998).
- [11] A. O. Gogolin, A. A. Nersesyan, and A. M. Tsvelik, in Bosonization and Strongly Correlated Systems (Cambridge University Press, 1998)
- [12] We neglect corrections to the Fermi velocity for chain 1. [13] We have  $\chi_{\perp,s} \propto T \frac{\partial^2}{\partial t_{\perp}^2} \ln \mathcal{Z}_{\perp}$  and  $\chi_{\perp,p} \propto T \frac{\partial^2}{\partial V^2} \ln \mathcal{Z}_{\perp}$ where  $\mathcal{Z}_{\perp}$  is the  $\perp$ -part of the grand partition function.
- [14] D. V. Khveshchenko and T. M. Rice, Phys. Rev. B 50, 252 (1994).
- [15] Taking into account second order corrections to  $K_c^-$  and  $K_s^-$  would not change the further conclusions.
- [16] For repulsive interactions in chain 2, the rotation invariance implies that for  $T \ll E_F$ :  $K_{2s} = 1, K_s^{\pm} < 1$ . The term  $g_5$  produces a coupling  $\mathcal{O}_{CDW}^{2\dagger} \psi_{\pm 1\alpha}^{\dagger} \psi_{\pm 1\alpha}$ . This
- [17]would lead to a spin-gapped CDW ground state ( $\lambda \approx 0$ ).
- [18] For  $T \leq T^*$ , in Eqs.(16) and (18) one must formally replace  $g_i \to V$  and equate z(l) = 0,  $K_{1s}(l) = 0$ .
- [19] Similarly, we obtain:

$$\mathcal{O}_{CDW}^j \propto \exp(-i\sqrt{2\pi}\Phi_{jc})$$
 (27)

$$= \exp(-i\sqrt{\pi}\Phi_c^+)\exp(\mp i\sqrt{\pi}\Phi_c^-).$$

The two-point correlation functions of the disordered field  $\Phi_c^-$  decay exponentially (its dual field is pinned), therefore  $\langle \mathcal{O}_{CDW}^{j\dagger}(x)\mathcal{O}_{CDW}^{j}(0) \rangle \propto \exp(-xT_c)$  (in contrast with Ref. [7]).

- [20] K. Le Hur, unpublished.
- [21] But the stability of the "C2S1" phase in the symmetric 3-leg Hubbard ladder (or 3-band Hubbard model) of Ref.
  [3] is clear. Starting from the half-filled case, the unpaired fermions of band 2 have been found to be completely decoupled from bands 1 and 3 (which contain the bosonic pairs). Adding (by hand) a scattering term of the form

$$V\Big(\big(\mathcal{O}_{SC}^{1\dagger} + \mathcal{O}_{SC}^{3\dagger}\big)\Big(\psi_{+2\uparrow}\psi_{-2\downarrow} + \psi_{-2\uparrow}\psi_{+2\downarrow}\Big) + \text{H.c.}\Big) \quad (28)$$

would not affect the fixed point. The bosonic pairs in bands 1 and 3 indeed obey  $\mathcal{O}_{SC}^1 \approx -\mathcal{O}_{SC}^3$  and the unpaired fermions are located exactly on the diagonals of the 2D Fermi surface, i.e.,  $\Delta H \propto (V - V) = 0$ .

[22] V. J. Emery, S. A. Kivelson, and O. Zachar, Phys. Rev. B 59, 15641-15653 (1999).



FIG. 2. Schematic vue of the CDW and SC susceptibilities as a function of the temperature. In general, the corresponding susceptibilities vary as  $T^{\nu-2}$  (The associated correlation functions decrease as  $x^{-\nu}$ ). Below  $T_c$ , the CDW correlation in contrast exhibits an exponential decay resulting in the complete vanishing of the  $(2k_F)$  CDW susceptibility at T = 0. We have approximated  $(K_{1c}; K_c^+) \approx 1$ .