

**Cugliandolo and Iguain reply:** In [1] we stated precisely that the aim of our study was to rule out non-resonant hole-burning (NSHB) experiments as proofs for the existence of *spatially* heterogeneous regions in glasses. The results of NSHB experiments have been often cited as evidence for the existence of *spatially distinguishable, dynamical, regions* in the samples. The list of references is long, see *e.g.* [2]. Agreement about the limitation of NSHB to determine the existence of *spatial* heterogeneities is now reached [3,4].




FIG. 1. Auto - correlation of the SK mean-field spin-glass model in the absence of an external field for  $t_w = 1, 100, 1000, 25000, 50000$  MC steps from bottom to top, the last three curves collapse.




FIG. 2. The distortion  $\Delta C$  after a single pump oscillation.  $h_F = 1, t_w = 25000$  MC steps. The symbols  $\star$  (blue),  $\times$  (green) and  $+$  (red) correspond to  $\omega = 1.5, 0.1, 0.005$ , respectively.

Diezemann and Böhmer point out, as we have also done in our article, that our measurements were done out of equilibrium. At least two of the NSHB experiments have been done below  $T_g$ , where equilibration is impossible [5,6] (see [7] where it is proven that even a field-cooled spin-glass may show nonequilibrium relaxation). Aging effects and the relaxation due to an applied ac pumping field have not been explicitly disentangled in these references.

The question as to whether the NSHB protocol applied to an equilibrated fully connected model (without notion of space), can reproduce the results of these experiments is very interesting indeed and deserves careful inspection. We take this opportunity to present results from a Mon-

tecarlo simulation of the Sherrington-Kirkpatrick (SK) mean-field spin-glass model with  $N = 3000$  spins at a temperature  $T = T_g - \epsilon = 0.99$ . Figure 1 shows one of the standard checks of equilibration in monitoring the autocorrelation function  $C(t+t_w, t_w)$  for different waiting-times  $t_w$  and observing when it starts being stationary. This happens for  $t_{eq} \approx 25000$  MC steps. Other, more subtle methods can also be used [8] and confirm equilibration at this waiting time. In Fig. 2 we apply the NSHB protocol after  $t_w = 25000$  MC steps with one oscillation of an ac-field with  $h_F = 1$  and  $\omega = 1.5, 0.1, 0.005$ ;  $t_r = 0$ . This figure can be compared, for instance, to Fig. 2 in Ref. [6] and, as far as we can judge, all qualitative features, including the shift in the base of the spectral hole mentioned in [4], are reproduced.

The effect of several cycles has already been discussed in [1]. A thorough discussion of where in parameter space aging effects disappear under a permanent ac-field in this model appeared in [9]. Once the stationary regime (in stroboscopic time) is reached, the spectral modification will obviously saturate.

In the  $p$ -spin and SK models it is not possible to identify localized domains and, however, the response to NSHB looks pretty much the same as in real systems.

Leticia F. Cugliandolo<sup>1,2</sup> and José Luis Iguain<sup>2</sup>

<sup>1</sup>Laboratoire de Physique Théorique, École Normale Supérieure, 24 rue Lhomond, 75231 Paris Cedex 05, France

<sup>2</sup>Laboratoire de Physique Théorique et Hautes Energies, Jussieu, 5ème étage, Tour 24, 4 Place Jussieu, 75005 Paris France

Received 2001

PACS numbers: 64.70.Pf, 75.10.Nr

- 
- [1] L. F. Cugliandolo and J. L. Iguain, Phys. Rev. Lett. **85**, 3448 (2000).
  - [2] R. Böhmer, Current Opinion in Solid State and Mat. Sc. **3**, 278 (1998). H. Sillescu, J. Non-Cryst. Solids **243**, 81 (1999). M. D. Ediger, Annu. Rev. Phys. Chem. **51**, 99 (2000).
  - [3] G. Diezemann and R. Böhmer, unpublished.
  - [4] R. V. Chamberlin and R. Richert, unpublished.
  - [5] R. Richert and R. Böhmer, Phys. Rev. Lett. **83**, 4337 (1999).
  - [6] R. V. Chamberlin, Phys. Rev. Lett. **24**, 5134 (1999).
  - [7] T. Jonsson, K. Jonason and P. Nordblad, Phys. Rev. **B59**, 9402 (1999).
  - [8] See, *e.g.*, A. Billoire and E. Marinari, cond-mat/0101177 and refs. therein.
  - [9] L. Berthier, L. F. Cugliandolo and J. L. Iguain, Phys. Rev. **E** 2001.