

Softening the Extensive Postulates

L. Velázquez^{a,*} and F. Guzmán^{b,†}

^a*Departamento de Física, Universidad de Pinar del Río
Martí 270, esq. 27 de Noviembre, Pinar del Río, Cuba.*

^b*Departamento de Física Nuclear
Instituto Superior de Ciencias y Tecnologías Nucleares
Quinta de los Molinos. Ave Carlos III y Luaces, Plaza
Ciudad de La Habana, Cuba.
(December 2, 2024)*

We address the problem of softening the extensive postulates of the standard thermodynamics in order to extend it to the study of nonextensive systems. We do this in analogy with the traditional analysis, starting from the microcanonical ensemble, but this time, considering the equivalence of it with some generalized canonical ensemble in the thermodynamic limit by mean of the scaling properties of the fundamental physical observables.

PACS: 05.20.Gg;05.20.-y

I. INTRODUCTION

Traditionally, the Statistical Mechanics and the standard Thermodynamics have been formulated to be applied to the study of extensive systems, in which the consideration of the additivity and homogeneity conditions, and the realization of the thermodynamic limit are indispensable ingredients for their good performance. In spite of its great success, this theory seems to be inadequate to deal with many physical contexts. Today it is known that the above conditions can be considered as reasonable approximations for those systems containing a huge number of particles interacting by means of short range forces [1–3] when these systems are not suffering a phase transition [4,5].

In the last years considerable efforts have devoted to the study of nonextensive systems. The available and increasingly experimental evidences on anomalies presented in the dynamical and macroscopic behavior in plasma and turbulent fluids [6–9], astrophysical systems [10–14], nuclear and atomic clusters [15–17], granular matter [18] glasses [19,20] and complex systems [21] constitute a real motivation for the generalization of the Thermodynamics.

One of the most important tendencies in the resolution of this problem is the generalization of Boltzmann-Gibbs's formalism through the consideration of more general entropy form than the Shannon-Boltzmann-Gibbs extensive entropy:

$$S_{SBG} = - \sum_k p_k \ln p_k. \quad (1)$$

This is the case of the Renyi [22] and Sharma-Mittal [23] entropies and the so popular Tsallis non-extensive entropy [24], to mention some of them. It has been said that in the second half of last century close to 30 different entropic forms have been proposed! This number is increased continuously with the years.

The main advantage of all these Probabilistic Thermodynamic Formalisms (PThF), those based on the consideration of the concept of information entropy, is the versatility of applications, be already well inside the physical scenarios, in the study of the behavior of systems during the equilibrium or dynamical processes, as well as others interdisciplinary branches of the knowledge. However, in our opinion, this kind of theory suffers two important defects. The first, the no uniqueness, which is evident. The second, it is not possible to determine the application domain *a priori* for each specific entropic form. For example, in the case of the Tsallis nonextensive statistics, the determination of the entropic index, q . That is why we consider that the statistical description of nonextensive systems has to start from the microscopic characteristics of them.

Ordinarily, the macroscopic description of the system in thermodynamic equilibrium through its microscopic properties has the starting point in the consideration of the microcanonical ensemble. Although there is no way to justify

*luisberis@geo.upr.edu.cu

†guzman@info.isctn.edu.cu

this selection, it is generally accepted its great significance, since from this ensemble the Thermostatistics could be deduced without to invoke anything outside the mechanics.

Boltzmann [25], Gibbs [5], Einstein [26,27] and Ehrenfests [28] recognized the hierarchical primacy of the microcanonical ensemble with regard to the canonical and Gran-canonical ensembles. The last ones can be derived starting from the first by the consideration of certain particular conditions: *the extensivity postulates*. These are the essential ideas to generalize the traditional results for the justification of any generalized Boltzmann-Gibbs Formalism, to put these on the level of the microscopic description, in the ground of the Mechanics.

The present paper will be devoted to the analysis of the conditions to take into account in order to perform this derivation. The same will be done in analogy with the traditional case, starting from the microcanonical ensemble but this time softening the traditional postulates in order to cover systems with long-range interactions.

II. MICROCANONICAL ENSEMBLE: ITS GEOMETRIC ASPECTS AND SCALING PROPERTIES.

Let us now to show that a reasonable way to generalize the traditional Thermodynamic is assuming the **scaling postulates**: The **equivalence** of the microcanonical description with some generalized canonical one during the realization of the **Thermodynamic Limit** by mean of the **self-similarity scaling properties** of the fundamental physical observables.

A. The thermodynamic limit.

Of the three postulates that sustain the validity of the standard thermodynamic, it is the realization of the thermodynamic limit the one that possesses the most universal character. In fact, the additivity and homogeneity conditions are associated more indissolubly to the short-range nature of the interactions in comparison with the characteristic linear dimension of a sufficient large system.

However, in the real world we find examples of systems where the thermodynamic limit does not take place, since they are not composed by a huge number of particles. We find examples of these in the molecular and atomic clusters. A very interesting problem is the extension of the macroscopic description for this kind of systems as well as the question about when it could be considered that these systems are found in the thermodynamic limit.

An important step toward the resolution of these problems was accomplished by D.H.E. Gross with the development of the *Microcanonical Thermostatistics* [15,29,30], formalism based on the consideration of the microcanonical ensemble. In this approach is possible to accomplish the description of some finite systems, extending this way the study of phase transitions in them. On the other hand, it is attractive the possibility that the same one has, exploring the behavior of the macroscopic observables with the increase of the number of particles, analyzing the convergence in some cases in an ordinary extensive system, allowing this way to give a valuation about when the thermodynamic limit is established in the system.

A remarkable aspect of this approach is that the justification of the canonical ensemble is not supported by the consideration of the Gibbs's argument, a closed system composed by a subsystem with a weak interaction with a thermal bath, but by means of equivalence between the microcanonical and canonical ensemble in the thermodynamic limit for this kind of system. The Gibbs's argument is sustained by the possibility of the separation of this subsystem from the whole system, that is said, in the independency or weak correlation of this subsystem with the remaining part. It is easy to understand that this argument can not be applied for many nonextensive systems, since in them there are long-range correlations due to the presence of long-range interactions.

Thus, *the equivalence of the microcanonical ensemble with the canonical ensemble during the thermodynamic limit could be easily extended for the nonextensive systems because this argument has an universal character stronger than the Gibbs's ones*. We have to point out that this equivalence also supports the thermodynamic formalism based on the Legendre Transformation between the thermodynamic potentials, which is usually valid without restriction for the probabilistic interpretation of the thermodynamics [29].

The thermodynamic formalism of the Gross's theory have been defined in order to be equivalent in the thermodynamic limit with the ordinary form of the Thermodynamic for the case of the extensive systems. That is why this theory is appropriate to study systems that in the thermodynamic limit become in ordinary extensive system (see for example in ref. [31,32]). However it is not trivial its application to systems with long-range interactions because we do not know what are the asymptotical properties of them and what are the generalized canonical ensemble that is equivalent to the microcanonical ensemble in the thermodynamic limit for these systems.

B. Geometrical aspects of the PDF.

A feature distinguishing characteristic of the PThF is that the description of the macroscopic state of the systems is performed by means of the intensive parameters of the generalized canonical distributions, β , that is to say, they are supported by the validity of some *generalized zero principle of the Thermodynamics*. The next step is the analysis of the prescriptions for this kind of description.

Let us pay special attention to the geometrical aspects that posses the Probabilistic Distribution Functions (PDF) of the ensembles. For the case of the microcanonical ensemble its PDF is given by:

$$\omega_M(X; I, N, a) = \frac{1}{\Omega(I, N, a)} \delta[I - I_N(X; a)]. \quad (2)$$

Any function of the integrals of movement is itself an integral of movement. If we have a complete independent set of functions of the integrals of movement of the distribution, then this set of functions is equivalent to the other set of integrals of movement, that is, both represent the same macroscopic state. That is why it is more exactly to speak about an abstract space of integrals of movement for the microcanonical distribution, \mathfrak{S}_N . Therefore, every physical quantity or behavior has to be equally reproduced by any representation of the space of macroscopic state, $(\mathfrak{S}_N; a)$.

It is easy to show that the microcanonical distribution is invariant under the transformation group of *local reparametrizations or Diffeomorfism* of the space of the integrals of movement, $Diff(\mathfrak{S}_N)$. Let \mathfrak{R}_I and \mathfrak{R}_ϕ be two representations of \mathfrak{S}_N , where:

$$I \equiv \{I^1, I^2 \dots I^n\} \text{ and } \phi \equiv \{\phi^1(I), \phi^2(I) \dots \phi^n(I)\}. \quad (3)$$

By the property of the δ -function we have:

$$\delta[\phi(I) - \phi(I_N(X; a))] = \left| \frac{\partial \phi}{\partial I} \right|^{-1} \delta[I - I_N(X; a)], \quad (4)$$

hence:

$$\tilde{\Omega}(\phi, N, a) = \left| \frac{\partial \phi}{\partial I} \right|^{-1} \Omega(I, N, a), \quad (5)$$

and therefore:

$$\omega_M(X; \phi, N, a) = \omega_M(X; I, N, a) \equiv \omega_M(X; \mathfrak{S}_N, a). \quad (6)$$

From here is derived two interesting observations. *The first:* This local reparametrization invariance is the maximal symmetry that a geometrical theory could possess, and it is usually associated to the local properties of a curved general space. This property is in complete contradiction with the *ergodic chauvinism* of the traditional distributions. This is translated as the preference of the energy over all the integrals of movement corresponding to the system. In the microcanonical ensemble this preference is not justified since it does not depend on the representation of the space of the integrals of movement. *The second:* The movement integrals are defined by the commutativity relation with the Hamiltonian of the system. In the case of closed systems, the Hamiltonian is the total energy, a conserved quantity. In the microcanonical distribution it represents one of the coordinate of the point belonging to \mathfrak{S}_N , in a specific representation. When we change the representation the energy lose its identity. Since we can not fix the commutativity of the energy with the others integrals, we must impose that all movement integrals commute between them in order to preserve these conditions. As we see, still from the classical point of view, *the reparametrization invariance suggests that the set of movement integrals have to be simultaneously measurable*. In the quantum case, this is an indispensable request for the correct definition of the statistical distribution. This property is landed to the classical distribution by the correspondence between the Quantum and the Classic Physic.

On other hand, all the PDF derived from the PThF must depend on the integrals of movement in a lineal combination with the canonical intensive parameters:

$$p(X; \beta, N) = F(\beta, N; \beta \cdot I_N(X)), \quad (7)$$

therefore, the most general symmetry group that preserve this functional form is the *local unitary lineal group*, $SL_c(\mathbf{R}^n)$ (n is the dimension of \mathfrak{S}_N), which is related with the euclidean vectorial spaces. It is supposed that an *spontaneous symmetry breaking* takes place during the thermodynamic limit, from $Diff(\mathfrak{S}_N)$ to $SL_c(\mathbf{B}^n)$, where \mathbf{B}^n

is the space of the generalized canonical parameters. The way in which this hypothetical symmetry breaking happens will determinate the specific form of the PDF in the thermodynamic limit.

What do we understand by an spontaneous symmetry breaking? Ordinarily, the microcanonical ensemble could be described equivalently through any representation of the abstract space of the integrals of movement of the distribution. However, with the increasing of the degree of freedom of the system, some specific representation will be more addecuate to describe the macroscopic state because it reflects in better manner the general properties of the system: i.e. in the case of the traditional systems, *the extensivity*.

C. The scaling symmetry of the fundamental physical observables.

In complete analogy with the traditional analysis, we identify the cause of this spontaneous symmetry breaking with the *scaling properties of the fundamental macroscopic observables during the thermodynamic limit*: the behavior of the integrals of movement, external parameters and the accessible states density with the increasing of the system's degree of freedom.

As it was indicated previously, these symmetry properties will be alone expressed for certain set of representations, $\mathcal{M} = \{\mathfrak{R}_{I^*}\}$. Only in this set of admissible representations will be valid the equivalency the microcanonical description and the generalized canonical one. The remaining symmetry among the representations belonging to \mathcal{M} constitutes the local group $SL_c(\mathbf{R}^n)$ of the canonical description.

In order to access to the scaling behavior of the integrals of movement we have to obtain the asymptotical dependence of the states density for large values of the number of particles, $W_{asym}(I^*, N, a)$:

$$W(I^*, N, a) \xrightarrow[N \gg 1]{} W_{asym}(I^*, N, a), \quad (8)$$

where $\mathfrak{R}_{I^*} \in \mathcal{M}$. Let W_o be defined by:

$$W_o = W_{asym}(I_o^*, N_o, a_o), \quad (9)$$

where N_o could be considered to be in the asymptotical region $N \gg 1$. The scaling behavior of the macroscopic observables is dictated by the symmetry of the asymptotical states density under the **self-similar scaling transformations**:

$$\left. \begin{array}{l} N_o \rightarrow N = \alpha N_o \\ I_o^* \rightarrow I^* = \alpha^\varkappa I_o^* \\ a_o \rightarrow a = \alpha^{\pi_a} a_o \end{array} \right\} \Rightarrow W_{asym}(I^*, N, a) = \mathcal{F}[W_o, \alpha]. \quad (10)$$

In the above definition, α is the scaling parameter, \varkappa and π_a are real constants which characterizing the scaling transformations. As we see, all the integrals of movement are transformed by the same scaling law in order to satisfy the $SL(\mathbf{R}^n)$ invariance of the PDF in the thermodynamic limit. Finally, \mathcal{F} is a functional dependence of W_o and α satisfying the exigencies:

$$\begin{array}{l} 1.- \mathcal{F}[x, 0] = x \text{ (Identity),} \\ 2.- \mathcal{F}[\mathcal{F}[x, \alpha_1], \alpha_2] = \mathcal{F}[x, \alpha_1 \alpha_2] \text{ (Self-similarity).} \end{array} \quad (11)$$

The above conditions could be satisfied by the following generic form:

$$\mathcal{F}[x, \alpha] = f_\infty[\alpha^\sigma f_\infty^{-1}(x)], \quad (12)$$

where $f_\infty(x)$ is a increasing function of x with a monotonic first derivative in the asymptotical region $x \gg 1$, and σ , a real constant. The function $f_\infty(x)$ defines the specific **scaling law** of the system. Let us introduce the function $f(x)$, which is an increasing function of x with a monotonic first derivative for all the positive real values of x satisfying the following conditions:

$$\begin{array}{l} 1.- f(0) = 1 \text{ (Normalization),} \\ 2.- f(x) \xrightarrow{x \gg 1} f_\infty(x) \text{ (Convergency).} \end{array} \quad (13)$$

This function allows us to define the function $\gamma(I, N; a)$:

$$\gamma(I^*, N; a) = \frac{1}{N^\sigma} f^{-1}[W(I^*, N; a)], \quad (14)$$

which is scaling invariant during the realization of the thermodynamic limit. It is not difficult to understand that the objective of the canonical description is the knowledge of the set of admissible representations, \mathcal{M} , the scaling laws of the fundamental macroscopic observables (integrals of movement and external parameters) as well as the determination of the scaling invariant function $\gamma_\infty(I^*, N; a)$:

$$\gamma_\infty(I^*, N; a) = \lim_{\alpha \rightarrow \infty} \gamma(I^*, N; a) \equiv \frac{1}{N_o^\sigma} f_\infty^{-1}[W(I_o^*, N_o; a_o)]. \quad (15)$$

Although $\gamma_\infty(I^*, N; a)$ is not a physical observable, from this function could be derived others quantities which are experimentally mensurable, allowing this way to obtain information about the ordering of the system. This fact points to the following conclusion: *In the generalized canonical description, the function:*

$$S_f = f^{-1}[W(I^*, N; a)], \quad (16)$$

plays the role of the generalized entropy. From here and after we will refer the definition of the Eq.(16) as the **generalized Boltzmann's Principle**. The function $f^{-1}(x)$ defines the so called *entropy's counting rule*. Our proposition suggests the relative significance of the entropy concept. In the thermodynamic limit the generalized Boltzmann's entropy may possess a probabilistic interpretation with the same style of the Shannon-Boltzmann-Gibbs or the Tsallis nonextensive entropy. The specific form of the thermodynamic formalism have to be determined by mean of the conditions for the equivalence of the ensembles, maybe, through of some generalization of the ordinary Laplace's Transformation [29].

A final observation. Ordinarily, the validity of this equivalence between the microcanonical and the canonical ensemble have to satisfy an stability condition, which is ordinarily improve by the exigency of the concavity of the entropy function. This mathematical demand is intimately bound with the homogeneity condition, the non occurrence of phase transitions in the system. In the case of the finite system this exigency have to lead to the generalization of the microcanonical thermostatic of D.H.E. Gross.

III. CONCLUSIONS

We have analyzed a possible way to generalize the extensive postulates in order to extend the Thermodynamics to the study of some nonextensive systems. We showed the superiority of the argument of the equivalence of the microcanonical ensemble with some generalized canonical ensemble with regard to the Gibbs's one. The ergodic chauvinism of the classical thermodynamics is not justified in the microcanonical ensemble since this description is reparametrization invariant. This property has a relevant role in the passage from one description to the another, with the occurrence of an hypothetical spontaneous breaking of this symmetry in the thermodynamic limit by mean of the scaling properties of the fundamental physical observables. The generalized Boltzmann's Principle, the definition of the generalized Boltzmann's entropy [the Eq.(16)], have been defined in order to deal the study of systems with an arbitrary scaling laws [i.e., *exponentials, potentials (fractals)*, etc.].

The previous arguments can be applied for the study of some nonextensive systems, allowing this way to justify a thermodynamic probabilistic formalism analogue to the traditional one without the necessity of appealing to additional postulates.

-
- [1] R.K. Pathria, Statistical Mechanics , Butterworth Heinemann 1996.
 - [2] H.E. Stanley, Introduction to Phase Transitions and Critical Phenomena, Oxford University Press, New York 1971.
 - [3] P.T. Landsberg, Thermodynamics and Statistical Mechanics , Dover 1991.
 - [4] L. Tisza, Annal. Phys. B, 1, (1981); Generalized Thermodynamics, MIT Press, Cambridge, 166, page 123.
 - [5] J.W.Gibbs, Elementary Principles in Statistical Physics, Volume II of The Collected works of J. Williard Gibbs, Yale University Press, 1902.
 - [6] B.M. Boghosian, Phys. Rev. E 53 (1996) 4754.
 - [7] C. Beck, G.S. Lewis and H.L. Swinney , Phys. Rev. E 63 (2001) 035303(R).
 - [8] T.H.Solomon, E.R. Weeks and H.L. Swinney, Phys. Rev. Lett. 71 (1993) 3975.
 - [9] M.F. Shlesinger, G.M. Zaslavsky and U. Frisch Eds., Lévy flights and related topics, Springer-Verlag Berlin (1995).
 - [10] D. Lynden-Bell , Physica A 263 (1999) 293.

- [11] F. Sylos Labini, M. Montuori and L. Pietronero, Phys. Rep. 293 61 (1998) 61.
- [12] L. Milanovic, H.A.Posch and W. Thirring, Phys. Rev. E 57 (1998) 2763.
- [13] H. Koyama and T. Konishi, Phys. Lett. A 279 (2001) 226.
- [14] A. Torcini and M. Antoni, Phys. Rev. E 59 (1999) 2746.
- [15] D.H.E Gross, Microcanonical thermodynamics: Phase transitions in Small systems, 66 Lectures Notes in Physics, World scientific, Singapore (2001) and refs. therein.
- [16] M. D'Agostino et al., Phys. Lett. B 473 (2000) 219.
- [17] M. Schmidt, R. Kusche, T. Hippler, J. Donges and W. Kronmueller, Phys. Rev. Lett. 86 (2001) 1191.
- [18] A. Kudrolli and J. Henry, Phys. Rev. E 62 (2000) R1489.
- [19] G. Parisi, Physica A 280 (2000) 115.
- [20] P.G. Benedetti and F.H. Stillinger, Nature 410 (2001) 259.
- [21] G.M. Viswanathan , V. Afanasyev , S.V. Buldyrev, E.J. Murphy and H.E. Stanley, Nature 393 (1996) 413.
- [22] A. Renyi, in Proc . Fourth Berkeley Symposium ,1960, Vol. 1 (University of California Press, Berkeley ,Los Angeles, 1961), 547; Probability Theory (North-Holland, 1970) and references therein.
- [23] B.D. Sharma and D.P . Mittal, J. Math. Sci. 10 , 28(1975).
- [24] C.Tsallis, J. Stat. Phys. 52, 479 (1988).
- [25] L.Boltzmann, Über die Beziehung eines allgemeinen mechanischen Satzes Zum Hauptsatz der Wärmelehre. Sitzungsbericht der akademie der Wissenschaften, Wien, II:67-73, 1877.
- [26] A.Einstein, Eine Theorie der Crundlagen der Thermodynamik, Ann. Phys 11:170-187, 1903.
- [27] A.Einstein, Zur allgenmeinen molekularen. Theorie der Wärme, Ann. Phys 14:354-362, 1904.
- [28] P.Ehrenfest and T. Ehrenfest. The conceptual Foundation of the Statistical Approach in Mechanics, Cornell University Press, Ithaca NY, 1959.
- [29] D.H.E.Gross, Microcanonical Statistical Mechanics of some non extensive systems, Invited talk at the International Workshop on Classical and Quantum Complexity and non-extensive Thermodynamics, Denton-Texas, April 3-6, 2000, in press *cond-mat/0004268*.
- [30] D.H.E.Gross, Microcanonical Thermodynamics and statistical fragmentation of disipative systems- the topological structure of N-body phase space, Phys. Report, 257:133-221(1995).
- [31] D.H.E.Gross, A.Ecker, X.Z.Zhang, Microcanonical Thermodynamics of First Order Phase Transitions studied in the Potts Model, Annalen Phys. 5 (1996) 446-452.
- [32] D.H.E.Gross, M.E.Madjet, Fragmentation Phase Transition in Atomic Clusters IV – Liquid-gas transition in finite metal clusters and in the bulk, in press *cond-mat/9707100*.