

# Analytical expressions for the spin-spin local-field factor and the spin-antisymmetric exchange-correlation kernel of a two-dimensional electron gas

B. Davoudi<sup>1,2</sup>, M. Polini<sup>1</sup>, G. F. Giuliani<sup>3</sup> and M. P. Tosi<sup>1</sup>

<sup>1</sup>*NEST-INFM and Classe di Scienze, Scuola Normale Superiore, I-56126 Pisa, Italy*

<sup>2</sup>*Institute for Studies in Theoretical Physics and Mathematics, Tehran 19395-5531, Iran*

<sup>3</sup>*Physics Department, Purdue University, West Lafayette, Indiana*

We present an analytical expression for the static many-body local field factor  $G_-(q)$  of a homogeneous two-dimensional electron gas, which reproduces Diffusion Monte Carlo data and embodies the exact asymptotic behaviors at both small and large wave number  $q$ . This allows us to also provide a closed-form expression for the spin-antisymmetric exchange and correlation kernel  $K_{xc}^-(r)$  which represents a key input for spin-density functional studies of inhomogeneous electronic systems.

PACS number: 71.10.Ca, 71.15.Mb

The static spin-spin response function  $\chi_s(q)$  of a paramagnetic electron gas (EG) can be written in terms of the Lindhard function  $\chi_0(q)$  by means of the spin-antisymmetric many-body local field  $G_-(q)$  through the relationship

$$\chi_s(q) = -\mu_B^2 \frac{\chi_0(q)}{1 + v_q G_-(q) \chi_0(q)}, \quad (1)$$

where  $\mu_B$  is the Bohr magneton. Thus  $G_-(q)$  is a fundamental quantity for the determination of many properties of a general electron system. It is a key input, together with the charge-charge local-field factor  $G_+(q)$ , in the spin-density functional theory (SDFT) of the inhomogeneous electron gas<sup>1,2</sup> and in studies of quasiparticle properties (such as the effective mass and the effective Landè g-factor) in the electronic Fermi liquid<sup>3,4</sup>.

For what concerns SDFT calculations, a common approximation to the unknown exchange-correlation energy functional  $E_{xc}[n, \zeta]$  of electron density  $n$  and spin-polarization  $\zeta$  appeals to its second functional derivatives,

$$K_{xc}^{\sigma\sigma'}(\bar{n}, \bar{\zeta}; |\mathbf{r} - \mathbf{r}'|) \equiv \left. \frac{\delta^2 E_{xc}[n, \zeta]}{\delta n_\sigma(\mathbf{r}) \delta n_{\sigma'}(\mathbf{r}')} \right|_{n=\bar{n}, \zeta=\bar{\zeta}}, \quad (2)$$

where  $\bar{n} = \bar{n}_\uparrow + \bar{n}_\downarrow$  is the average local density of the EG and  $\bar{\zeta} = (\bar{n}_\uparrow - \bar{n}_\downarrow)/\bar{n}$  is the average local spin-polarization. The local field factors  $G_{\sigma\sigma'}(q)$  and the matrix of the exchange-correlation kernels are simply related in Fourier transform by

$$\tilde{K}_{xc}^{\sigma\sigma'}(q) \equiv \int d^d r e^{-i\mathbf{q}\cdot\mathbf{r}} K_{xc}^{\sigma\sigma'}(r) = -v_q G_{\sigma\sigma'}(q), \quad (3)$$

where  $d$  is the dimensionality of the system and  $v_q$  is the Fourier transform of the Coulomb potential  $e^2/r$ . In the paramagnetic state the following relation holds between the charge-charge and spin-spin local-field factors  $G_\pm(q)$  and  $G_{\sigma\sigma'}(q)$ :

$$G_\pm(q) = \frac{G_{\uparrow\uparrow}(q) \pm G_{\uparrow\downarrow}(q)}{2}. \quad (4)$$

Thus, from Eq. (3) and (4), one obtains that the spin-symmetric and the spin-antisymmetric exchange and correlation kernels  $K_{xc}^\pm(r)$  are related to  $G_\pm(q)$  by

$$\tilde{K}_{xc}^\pm(q) \equiv \frac{\tilde{K}_{xc}^{\uparrow\uparrow}(q) \pm \tilde{K}_{xc}^{\uparrow\downarrow}(q)}{2} \equiv \int d^d r e^{-i\mathbf{q}\cdot\mathbf{r}} K_{xc}^\pm(r) = -v_q G_\pm(q). \quad (5)$$

In what follows we shall only consider  $K_{xc}^-(r)$  in the case of two spatial dimensions, with  $d = 2$  and  $v_q = 2\pi e^2/q$ . An analytical expression for  $K_{xc}^+(r)$  has already been given in Ref. [5].

A number of exact asymptotic properties of  $G_-(q)$  in two dimensions are readily proven. In particular<sup>6</sup>,

$$\lim_{q \rightarrow 0} G_-(q) = A - \frac{q}{k_F} \quad (6)$$

with

$$A_- = \frac{1}{r_s \sqrt{2}} \left( 1 - \frac{\chi_0}{\chi} \right), \quad (7)$$

where  $k_F = \sqrt{2\pi n} = \sqrt{2}/r_s a_B$  is the Fermi wave number,  $r_s = \sqrt{\pi n a_B^2}$  is the usual EG density parameter with  $a_B$  the Bohr radius,  $\chi_0 = \mu_B^2 m/\pi \hbar^2$  is the (Pauli) spin susceptibility of the ideal Fermi gas while  $\chi$  is the spin susceptibility of the interacting system. By making use of the thermodynamic definition of  $\chi$  we can write

$$\frac{\chi_0}{\chi} = 1 - \frac{\sqrt{2}}{\pi} r_s + \frac{r_s^2}{2} \left. \frac{\partial^2 \epsilon_c}{\partial \zeta^2} \right|_{\zeta=0} \quad (8)$$

where  $\epsilon_c$  is the correlation energy per particle in the spin-polarized fluid. Unfortunately the correlation energy at finite spin polarization is poorly known. By extrapolating the Diffusion Monte Carlo (DMC) data<sup>7</sup> for  $G_-(q)$  to  $q = 0$  and by observing from Eq. (8) that  $A_- \rightarrow 1/\pi$  for  $r_s \rightarrow 0$ , we obtain the following parametrization of  $A_-$  in the range  $0 \leq r_s \leq 10$ :

$$A_- = \frac{1}{\pi + 1.4954 r_s + 0.3193 \sqrt{r_s}}. \quad (9)$$

The asymptotic behaviour of  $G_-(q)$  at large  $q$  is also known exactly<sup>6,8</sup>,

$$G_-(q) \sim C_- \frac{q}{k_F} + B_- \quad (10)$$

for  $q \rightarrow \infty$ , where  $C_-$  is proportional to the difference in kinetic energy between the interacting and the ideal gas,

$$C_- = \frac{t - t_0}{2\pi n e^2} k_F = -\frac{r_s}{2\sqrt{2}} \frac{d}{dr_s} \left( r_s \epsilon_c(r_s) \right). \quad (11)$$

The correlation energy per particle  $\epsilon_c(r_s)$  in the paramagnetic fluid is available from Monte Carlo data<sup>9</sup>. Finally  $B_- = g(0)$ ,  $g(0)$  being the value of the pair-correlation function at the origin<sup>10</sup>.

In this work we fit the values of  $G_-(q)$  originally obtained by DMC in Ref. [7] in such a way as to obtain analytical expressions for both  $\tilde{K}_{xc}^-(q)$  and  $K_{xc}^-(r)$ . The functional form previously employed<sup>5</sup> for the charge-charge local-field factor  $G_+(q)$  is also suitable for  $G_-(q)$ :

$$G_-(\bar{q}) = A_- \bar{q} \left[ \frac{\kappa_-(r_s)}{\sqrt{1 + (A_- \kappa_-(r_s) \bar{q}/B_-)^2}} + (1 - \kappa_-(r_s)) e^{-\bar{q}^2/4} \right] + C_- \bar{q} \left( 1 - e^{-\beta_- \bar{q}^2} \right) + P_-(\bar{q}) e^{-\alpha_- \bar{q}^2}, \quad (12)$$

where  $\bar{q} = q/k_F$ ,  $\kappa_-(r_s) = \sqrt{1 + 0.0082 r_s^2}$  and  $P_-(\bar{q})$  is the polynomial  $P_-(\bar{q}) = h_2 \bar{q}^2 + h_4 \bar{q}^4 + h_6 \bar{q}^6 + h_8 \bar{q}^8$ . The free parameters  $\alpha_-$ ,  $\beta_-$  and  $h_2, \dots, h_8$  are fitted so as to minimize the differences from the DMC numerical results. As for  $G_+(q)$ , it proves useful to have a continuous parametrization of these parameters at least in the range  $0 \leq r_s \leq 10$ . We propose the following:

$$\begin{aligned} \alpha_-(r_s) &= \exp \left( -\frac{0.2231 + 81.2115 (r_s/10)}{1 + 54.6665 (r_s/10) + 50.7534 (r_s/10)^2} \right), \\ \beta_-(r_s) &= 0.8089 - 0.4025 (r_s/10)^3 - 0.0941 (r_s/10)^{1/2}, \\ h_2(r_s) &= \exp \left( -\frac{12.6262 + 20.9673 (r_s/10)}{1 + 12.4002 (r_s/10)} \right), \\ h_4(r_s) &= 0.0531 \left( 1 - e^{-2.154 r_s^2} \right) - 0.4984 (r_s/10)^{3/2} + 0.4021 (r_s/10)^2, \\ h_6(r_s) &= -0.0076 \left( 1 - e^{-r_s} \right) + 0.0977 (r_s/10)^{3/2} - 0.0726 (r_s/10)^2, \\ h_8(r_s) &= -0.0027 (r_s/10). \end{aligned} \quad (13)$$

In Figure 1 we compare the fit given by Eq. (12) with the DMC data for  $r_s = 1, 2, 5$  and 10. In Figure 2 we show the local field factor  $G_-(q)$  as from Eq. (12) for various values of  $r_s$ .

We turn next to the evaluation of  $K_{xc}^-(r)$ . From Eq. (5) and (12), the expression of the spin-antisymmetric exchange-correlation kernel in real space (in Ryd) is readily obtained as

$$K_{xc}^-(r) = N_1 \frac{\delta^{(2)}(\mathbf{r})}{k_F^2} + N_2 \frac{\exp\left(-B_- k_F r / (A_- \kappa_-(r_s))\right)}{k_F r} + N_3 e^{-(k_F r)^2} + N_4 e^{-(k_F r)^2 / 4\beta_-} + \sum_{n=1}^4 N_{5,2n} F_{2n}(\alpha_-, k_F r), \quad (14)$$

where  $N_1 = -4\pi\sqrt{2}C_-/r_s$ ,  $N_2 = -2\sqrt{2}B_-/r_s$ ,  $N_3 = -4\sqrt{2}A_-(1-\kappa_-(r_s))/r_s$ ,  $N_4 = \sqrt{2}C_-/r_s\beta_-$  and  $N_{5,n} = -2^{\frac{3}{2}}h_n/r_s$ . The function  $F_n(\alpha, x)$  is given by

$$F_n(\alpha, x) = \int_0^\infty dy y^n J_0(xy) e^{-\alpha y^2} = \frac{1}{2} \alpha^{-(1+n)/2} \Gamma\left(\frac{1+n}{2}\right) {}_1F_1\left(\frac{1+n}{2}; 1; -\frac{x^2}{4\alpha}\right) \quad (15)$$

where  $\Gamma(z)$  is Euler's Gamma function and  ${}_1F_1(a; b; z)$  is Kummer's function. In practice the function  $F_n(\alpha, x)$  can be obtained via the recursive relation

$$F_{n+2}(\alpha, x) = -\frac{dF_n(\alpha, x)}{d\alpha}, \quad F_2(\alpha, x) = \frac{\sqrt{\pi}}{16\alpha^{5/2}} \left[ (4\alpha - x^2) I_0\left(\frac{x^2}{8\alpha}\right) + x^2 I_1\left(\frac{x^2}{8\alpha}\right) \right] e^{-x^2/8\alpha} \quad (16)$$

where  $I_n(z)$  is the modified Bessel function of order  $n$ . It is also useful to recall that  $dI_0(z)/dz = I_1(z)$  and that  $dI_1(z)/dz = I_0(z) - I_1(z)/z$ .

In Figure 3 we show the matrix elements  $K_{xc}^{\sigma\sigma'}(r)$  of the exchange-correlation kernel for various values of  $r_s$ . These kernels describe the local structure (Pauli-Coulomb hole) of the paramagnetic electron fluid around an electron of a given spin.  $K_{xc}^{\uparrow\uparrow}(r)$ , being the sum of  $K_{xc}^+(r)$  and  $K_{xc}^-(r)$ , has the same behavior of these two components, namely no structure at both small and high  $r_s$ . The situation is different for  $K_{xc}^{\uparrow\downarrow}(r)$ , which is instead given by the difference of  $K_{xc}^+(r)$  and  $K_{xc}^-(r)$ . Although these two components are structureless at small  $r_s$ , the difference between them around  $k_F r \simeq 1$  is large and this is reflected in the large structure shown in Figure 3.

In conclusion, we have presented an analytic parametrization of the local field factor entering the spin response of the two-dimensional electron gas in the paramagnetic state, incorporating the known asymptotic behaviors and giving an accurate representation of the available quantum Monte Carlo data. We have obtained from it and from our earlier results on the dielectric response<sup>5</sup> analytic expressions of the exchange-correlation kernels for spin-density-functional calculations on inhomogeneous two-dimensional electronic systems.

## ACKNOWLEDGMENTS

This work was partially supported by MURST through the PRIN 1999 program. We are grateful to Dr. S. Moroni for fruitful discussions and for providing us with the results of the Diffusion Monte Carlo study. Three of us (B. D., M. P. and M. P. T.) wish to thank the Condensed Matter Group of the Abdus Salam International Center for Theoretical Physics in Trieste for their hospitality during the final stages of this work.




FIG. 1. The local field factor  $G_-(\vec{q})$  for various values of  $r_s$  as computed according to Eq. (12), in comparison with the DMC data of Ref. 7




FIG. 2. The local field factor  $G_-(q)$  as from Eq. (12) for various values of  $r_s$ .




FIG. 3. The matrix elements  $K_{xc}^{\uparrow\uparrow}(r)$  and  $K_{xc}^{\uparrow\downarrow}(r)$  of the exchange-correlation kernel for various values of  $r_s$ .

- <sup>1</sup> U. von Barth and L. Hedin, *J. Phys. C* **5**, 1629 (1972); M. M. Pant and A. K. Rajagopal, *Solid State Commun.* **10**, 1157 (1972).
- <sup>2</sup> R. M. Dreizler and E. K. U. Gross, *Density Functional Theory, An Approach to the Quantum Many-Body Problem* (Springer, Berlin 1990).
- <sup>3</sup> S. Yarlagadda and G. F. Giuliani, *Phys. Rev. B* **40**, 5432 (1989); S. Yarlagadda and G. F. Giuliani, *Phys. Rev. B* **49**, 14188 (1994).
- <sup>4</sup> Direct measurements of the effective mass and the effective Landè  $g$ -factor in high-mobility Si MOSFETs over a wide range of carrier densities have been reported by V. M. Pudalov, M. Gershenson, H. Kojima, N. Butch, E. M. Dizhur, G. Brunthaler, A. Prinz, and G. Bauer, *cond-mat/0105081*.
- <sup>5</sup> B. Davoudi, M. Polini, G. F. Giuliani, and M. P. Tosi, *Phys. Rev. B*, in press.
- <sup>6</sup> G. E. Santoro and G. F. Giuliani, *Phys. Rev. B* **37**, 4813 (1988).
- <sup>7</sup> S. Moroni, D. M. Ceperley and G. Senatore, *Phys. Rev. Lett.* **69**, 1837 (1992); G. Senatore, S. Moroni and D. M. Ceperley in *Quantum Monte Carlo Methods in Physics and Chemistry*, eds. M. P. Nightingale and C. J. Umrigar (Kluwer, Dordrecht 1999); and private communication.
- <sup>8</sup> A. Holas, in *Strongly Coupled Plasma Physics*, eds. F. J. Rogers and H. E. DeWitt, (Plenum, New York 1986), p. 463; G. Vignale, *Phys. Rev. B* **38**, 6445 (1988).
- <sup>9</sup> F. Rapisarda and G. Senatore, *Aust. J. Phys.* **49**, 161 (1996).
- <sup>10</sup> M. Polini, G. Sica, B. Davoudi and M. P. Tosi, *J. Phys.: Condens. Matter.* **13** 3591 (2001).