

# Multi-particle excitations and spectral densities in quantum spin-systems

Christian Knetter, Kai P. Schmidt and Götz S. Uhrig

*Institut für Theoretische Physik, Universität zu Köln, Zùlpicher Str. 77, D-50937 Köln, Germany*

(November 20, 2018)

The excitation spectrum of the 2-leg  $S=1/2$  Heisenberg ladder is examined perturbatively. Using an optimally chosen continuous unitary transformation we expand the Hamiltonian and the Raman operator about the limit of isolated rungs leading to high order series expansions allowing to calculate spectral densities quantitatively. The 2-particle sector is examined for total momentum  $k = 0$ . We show that triplet-triplet interaction gives rise to a band splitting.

PACS numbers: 75.40.Gb, 75.50.Ee, 75.10.Jm

Antiferromagnetic quantum spin systems are investigated intensively both in theory and experiment. At  $T = 0$ , the so called spin liquids exhibit a non magnetic short ranged ground state and gapped excitations with unusual dynamics. The spin liquid phase is favoured by frustration, low dimensionality and low spin.

Perturbation theory directly allows to obtain information in the thermodynamic limit and is not afflicted by finite size effects. The perturbative Continuous Unitary Transformation Method (CUT) introduced previously [1] proved to be a versatile tool to calculate high order series expansions of low lying excitation energies in spin systems (see e.g. [2–4]). In this article, we present quantitative results for the spectral densities of multi-particle excitations.

The system considered is the 2-leg  $S=1/2$  Heisenberg ladder of which the Hamiltonian reads

$$H(J_1, J_2) = \sum_i [J_1 \mathbf{S}_{1,i} \mathbf{S}_{2,i} + J_2 (\mathbf{S}_{1,i} \mathbf{S}_{1,i+1} + \mathbf{S}_{2,i} \mathbf{S}_{2,i+1})], \quad (1)$$

where  $i$  denotes the rung (coupling  $J_1$ ) and 1,2 the leg (coupling  $J_2$ ). This system is realized in a number of substances [5] one of them being the superconducting compound  $\text{Sr}_{0.4}\text{Ca}_{13.6}\text{Cu}_{24}\text{O}_{41.84}$  [6].

For a given observable  $\mathcal{O}$  the  $T = 0$  spectral density is given by

$$S_{\mathcal{O}}(\omega) = -\frac{1}{\pi} \text{Im} \langle \psi_0 | \mathcal{O}^\dagger (\omega - H(J_1, J_2) + i0^+)^{-1} \mathcal{O} | \psi_0 \rangle, \quad (2)$$

$\psi_0$  is the system's ground state. Using the limit of isolated rungs  $x = J_2/J_1 = 0$ , we map the full Hamiltonian  $H(J_1, J_2)$  continuously and unitarily onto an effective Hamiltonian  $H_{\text{eff}}(x)$ ,  $x > 0$ , which conserves the number of triplets on the rungs [ $H_{\text{eff}}(x), H(J_1, 0) = 0$ ].

The effective Hamiltonian  $H_{\text{eff}}(x)$  is given as an exact operator series in  $x$  truncated after order 14 in the 0- and 1-triplet and after order 13 in the 2-triplet sector. For more details see Ref. [1]. The *same* transformation used to derive  $H_{\text{eff}}$  is applied to observables resulting in effective observables  $\mathcal{O}_{\text{eff}}$ , also given as an operator series truncated after 10<sup>th</sup> order acting on  $|0\rangle$ , the ground state of the effective model (details in Ref. [7]).

With the triplet vacuum  $|0\rangle$  the ground state energy  $E_0$  is a series in  $x$ ,  $E_0(x) = \langle 0 | H_{\text{eff}}(x) | 0 \rangle$ . Similarly the 1-(2-)triplet energies are obtained from the states  $|i\rangle$  ( $|i, j\rangle$ ) denoting the eigenstates of  $H(J_1, 0)$  with one (two) triplet(s) on rung  $i$  (and  $j$ ) [7]. Result for the energies have also been obtained by cluster techniques [8,9]. To calculate the Greens function in Eq. (2) for the effective operators we utilize the continued fraction technique [10] with proper termination [11]. In combination with standard series approximations our method allows an accuracy within 5% for  $x \approx 1$




FIG. 1. The 2-triplet spectral densities at total momentum  $k = 0$  and  $x = 1$  computed from Eq. (2) with  $\mathcal{O} = \mathcal{O}_R$  (black) and a test operator (gray) with triplet-triplet interaction switched on (solid) and off (dashed). All curves are scaled to give total weight 1. (Further details see text.) Inset: 1-triplet dispersion.

In Ref. [12] we show the  $k$ -resolved spectra for various observables at  $x = 1$  in great detail. Relevant for Raman scattering [13,14] is the  $S = 0$  operator  $\mathcal{O}_R = \mathbf{S}_{1,i} \mathbf{S}_{2,i}$ , locally exciting two triplets from the ground state  $|0\rangle$  on rung  $i$ , at total momentum  $k = 0$  depicted as solid black

line in Fig. 1 for  $x = 1$ . Particularly surprising is the appearance of a mid band singularity (MBS) at an energy of about  $3.4J_1$  splitting the 2-triplet continuum into two bands. The CUT method allows to identify the parts of the 2-triplet energies that solely arise from 1-triplet kinetics [7]. The remaining part is the pure triplet-triplet interaction energy. The dashed black and gray lines in Fig. 1 (denoted by “no I”) depict the spectral density of  $\mathcal{O}_R$  at  $k = 0$  with the interaction part set to zero. In both cases the MBS disappears, whereas it appears when the interaction is switched on (solid lines “with I”). Hence the band splitting is due to the 2-triplet interaction. To rule out a possible matrix element effect we calculate Eq. (2) for an artificial  $S = 0$  test operator at total momentum  $k = 0$  which creates the two triplets on adjacent sites. The gray (“naive”) lines in Fig. 1 show the results with / without interaction (solid / dashed). Comparing with the corresponding black lines (results for the “true” operator  $\mathcal{O}_R$ ) one finds that  $\mathcal{O}_R$  merely shifts weight towards the lower band edge, being clearly not responsible for the MBS. The inset of Fig. 1 shows the 1-triplet dispersion  $\omega(k)$  [12] which has a shallow dip at small  $k$ -values. This leads to a high density of states in the 2-triplet sector at an energy of  $2 \times \omega(0) = 3.6J_1$ . Here the dashed gray line displays a van-Hove singularity reduced to a finite value due to the remaining intrinsic hard-core interaction: two triplets cannot occupy the same rung. Due to this large DOS the system becomes extremely sensitive to an interaction induced orthogonality between  $\mathcal{O}|0\rangle$  and eigenstates  $|\omega\rangle$  with energy  $\approx 3.6 J_1$ . Indeed, the MBS remains detectable even if the interaction is lowered to 1% of its full strength. In summary we note that the CUT method allows to compute spectral densities of quantum multi-particle excitations in great detail. The nature of the structures revealed can be explained giving insight in the complex dynamics of the spin ladder system.

2999 (1987).

- [11] V. S. Viswanath and G. Müller, *The Recursion Method; Application to Many-Body Dynamics*, Vol. m23 of *Lecture Notes in Physics* (Springer-Verlag, Berlin, 1994).
- [12] C. Knetter, K. P. Schmidt, M. Grüninger, and G. S. Uhrig, cond-mat/0106077.
- [13] K. P. Schmidt, C. Knetter, and G. S. Uhrig, cond-mat/0107431.
- [14] S. Sugai and M. Suzuki, Phys. Stat. Sol. (b) **215**, 653 (1999).

- 
- [1] C. Knetter and G. S. Uhrig, Eur. Phys. J. B **13**, 209 (2000).
  - [2] C. Knetter and G. S. Uhrig, Phys. Rev. B **63**, 94401 (2001).
  - [3] C. Knetter, A. Bühler, E. Müller-Hartmann, and G. S. Uhrig, Phys. Rev. Lett. **85**, 3958 (2000).
  - [4] G. S. Uhrig and B. Normand, Phys. Rev. B **58**, R14705 (1998).
  - [5] D. Johnston *et al.*, cond-mat/0001147.
  - [6] M. Uehara *et al.*, J. Phys. Soc. Jpn. **65**, 2764 (1996).
  - [7] C. Knetter, K. P. Schmidt, and G. S. Uhrig, in prep..
  - [8] Z. Weihong, V. Kotov, and J. Oitmaa, Phys. Rev. B **57**, 11439 (1998).
  - [9] Z. Weihong *et al.*, cond-mat/0010354.
  - [10] E. R. Gagliano and C. A. Balseiro, Phys. Rev. Lett. **26**,