

Deconfinement transition and Luttinger to Fermi Liquid crossover in quasi one-dimensional systems

S. Biermann,^{1,2} A. Georges,^{2,1} A. Lichtenstein,³ and T. Giamarchi^{1,2}

¹Laboratoire de Physique des Solides, CNRS-UMR 8502, UPS Bât. 510, 91405 Orsay France

²LPTENS CNRS UMR 8549 24, Rue Lhomond 75231 Paris Cedex 05, France

³University of Nijmegen, NL-6525 ED Nijmegen, The Netherlands

(Dated: October 31, 2018)

We investigate a system of one dimensional Hubbard chains of interacting fermions coupled by inter-chain hopping. Using a generalization of the Dynamical Mean Field Theory we study the deconfinement transition from a Mott insulator to a metal and the crossover between Luttinger and Fermi liquid phases. One-particle properties, local spin response and inter-chain optical conductivity are calculated. Possible applications to organic conductors are discussed.

PACS numbers: 71.10.Pm, 71.10.Hf, 71.27+a, 71.30+h

The nature of the metallic phase of interacting electron systems depends strongly on dimensionality. In three dimensions, Fermi liquid (FL) theory applies, whereas in one dimension the quasi-particle concept breaks down, leading to a different kind of low-energy fixed point known as a Luttinger liquid (LL). For commensurate electron fillings, strong enough repulsive interactions destroy the metallic state altogether by opening a Mott gap. This phenomenon exists in all dimensions but the one-dimensional case is particularly favorable [1]. In quasi one-dimensional (Q1D) systems, interchain hopping can induce a (deconfinement) transition from the Mott insulating (MI) state to a metallic state and crossovers between different metallic behaviors (Fig. 1). Besides their intrinsic theoretical interest, understanding these phenomena is directly relevant for a number of compounds such as the organic (super)-conductors (Bechgaard salts), which are three dimensional stacks of quarter-filled chains[2]. Indeed, in these compounds some of the low temperature properties are well described by Fermi liquid theory, whereas optical [3] and transport properties [4] have shown that the high temperature phase is either a Luttinger liquid or a Mott insulator.

Describing these Q1D systems is not an easy task. The transverse hopping t_{\perp} is a relevant perturbation on the LL [5, 6, 7]. Hence, perturbative renormalization group calculations yield an estimate of the crossover scale [5, 8] but fail below that scale. In the MI state, electrons are confined on the chains by the Mott gap. A finite critical value of t_{\perp} is needed to induce an insulator to metal transition. Such a transition has been advocated [9, 10] to be at the root of the change from insulating to metallic behavior observed in Bechgaard salts when increasing pressure or when going from the TMTTF to the TMTSF family. Thus the effects of the inter-chain hopping that are the most important physically *cannot be handled reliably by perturbative methods*. Although some non-




FIG. 1: Schematic phase diagram of a quasi one-dimensional system, as a function of temperature T and inter-chain hopping t_{\perp} . The Mott-Insulator (MI), Luttinger Liquid (LL) and Fermi Liquid (FL) regimes are displayed. All lines are crossovers, except for the $T = 0$ transition from a MI to a FL. The MI phase is present only for commensurate fillings. Possible phases with long-range order have been omitted.

perturbative studies of deconfinement have been made for a finite number of chains [11] the case of an infinite system is still open. Some of the key questions yet to be answered are: (i) What is the crossover scale from the LL to MI, or from the LL to FL (Fig.1), and is there only one crossover scale for the different physical properties (transport, spin response, single-particle properties etc...)? (ii) What is the nature of the low-temperature FL state, and in particular is the shape of the Fermi surface (FS) affected by interactions? Do quasi-particle (QP) properties such as the quasi-particle residue $Z(k)$, vary significantly as the wavevector k moves along the FS?

In order to deal with these questions in a non-perturbative manner, a new method has been pro-

posed [12, 13]. This method, which generalizes the single site dynamical mean-field theory (DMFT) [14], replaces the quasi one-dimensional system by a single *effective chain* from which electrons can hop to a self-consistent bath. We thus nickname it *chain-DMFT* (ch-DMFT). It is exact in the limit of decoupled chains, while it reduces to usual DMFT in the opposite limit of zero-hopping along the chains, hence capturing the physics of a strongly correlated FL (or MI) [14]. Solving the mean field equations is a very challenging task however. In this paper, the first quantitative solution of the ch-DMFT equations is presented. We show that the above issues can be valuably addressed, and some of the above questions answered. In view of the difficulty of the problem only the Hubbard model is considered in this paper, but more realistic extensions are in sight, as discussed below.

Let us consider a system of coupled chains described by a sum of Hamiltonians of the isolated chains plus a hopping term between neighboring chains. The ch-DMFT approximates this system by the effective action:

$$S_{\text{eff}} = - \int \int_0^\beta d\tau d\tau' \sum_{ij,\sigma} c_{i\sigma}^+(\tau) \mathcal{G}_0^{-1}(i-j, \tau-\tau') c_{j\sigma}(\tau') + \int_0^\beta d\tau H_{1D}^{\text{int}}[\{c_{i\sigma}, c_{i\sigma}^+\}] \quad (1)$$

where H_{1D}^{int} is the interacting part of the on-chain Hamiltonian. \mathcal{G}_0 is an *effective propagator* determined from a self-consistency condition: the Green's function $G(i-j, \tau-\tau') \equiv -\langle c(i, \tau) c^+(j, \tau') \rangle_{\text{eff}}$ calculated from S_{eff} should coincide with the on-chain Green's function of the original problem, with the same self-energy $\Sigma = \mathcal{G}_0^{-1} - G^{-1}$. This reads:

$$G(k, i\omega_n) = \int d\epsilon_\perp \frac{D(\epsilon_\perp)}{i\omega_n + \mu - \epsilon_k - \Sigma(k, i\omega_n) - \epsilon_\perp} \quad (2)$$

Let $\epsilon_\perp(k_\perp)$ denote the Fourier transform of the inter-chain hopping $t_\perp^{mm'}$, $D(\epsilon_\perp) = \sum_{k_\perp} \delta[\epsilon_\perp - \epsilon_\perp(k_\perp)]$ the corresponding density of states, k the momentum in the chain direction and ω_n the Matsubara frequencies corresponding to the inverse temperature β . The ch-DMFT equations (1,2) fully determine the self-energy and Green's function of the coupled chains $G(k, k_\perp, i\omega_n)^{-1} = i\omega_n + \mu - \epsilon_k - \epsilon_\perp(k_\perp) - \Sigma(k, i\omega_n)$. The ch-DMFT approach can be rigorously justified in the formal limit where the lattice connectivity in the transverse direction is taken to infinity. In systems with finite transverse connectivity it can be viewed as an approximation neglecting the dependence of the self-energy on transverse momentum, keeping both frequency and in-chain momentum dependence: $\Sigma = \Sigma(k, i\omega_n)$. In the following, we specialize to the Hubbard model, i.e for each chain:

$$H_{1D} = -t \sum_i [c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i] + U \sum_i n_{i\uparrow} n_{i\downarrow} \quad (3)$$

and we consider a two-dimensional array of chains forming a square lattice with nearest neighbor hopping, so that the transverse dispersion is $\epsilon_\perp(k_\perp) = -2t_\perp \cos k_\perp$.

We solve the effective chain problem for chains of 16 or 32 sites with periodic boundary conditions using the QMC Hirsch-Fye algorithm [15]. Using 32 time-slices in imaginary time allows to reliably access temperatures down to $T/W \simeq 1/50 = 0.02$ with $U/W \simeq 1$ where $W = 4t$ is the bandwidth of the one-dimensional chain. Typically, about 5000 QMC sweeps and 10 to 15 ch-DMFT iterations are sufficient to reach convergence. The main quantities that we calculate and analyze are: a) On-chain (i.e summed over k_\perp) single particle Green's functions: $G(k, i\omega_n)$, and self-energies $\Sigma(k, i\omega_n)$ (obtained from $\mathcal{G}_0^{-1} - G^{-1}$ [21]). This allows to identify the location of the FS $k_\perp^F(k)$ by solving $\epsilon_\perp(k_\perp^F) = \mu - \text{Re}\Sigma(k, i\omega_{n=1}) - \epsilon_k$ and the QP residue $Z = Z(k_\perp^F)$ by fitting the slope ($= 1 - Z^{-1}$) of Σ vs. $i\omega_n$ at a specific FS point. b) On-chain spin and charge response functions and in particular the local spin response function: $\chi_s(\tau) = \langle S^z(j, 0) S^z(j, \tau) \rangle = \sum_{k, k_\perp} \chi_s(k, k_\perp, \tau)$. Indeed, in a LL (or with $K_\rho = 1$ in a FL) we have $\chi_s(\tau) = \chi_s(\beta/2) (\sin \pi\tau/\beta)^{-(1+K_\rho)}$ in the asymptotic regime where β, τ are larger than some coherence scale. By fitting our results to this form, we can define an effective K_ρ . Furthermore, the temperature dependence of $\chi_s(\beta/2)$ is closely related to the NMR relaxation rate $1/T_1 \equiv \lim_{\omega \rightarrow 0} \chi_s''(\omega)/\omega$. In a FL liquid at low-enough temperature [16]: $1/T_1 = \beta \chi_s(\beta/2)/2\pi^2$. In a LL $1/T_1$ and $\beta \chi_s(\beta/2)$ have the same T-dependence (albeit with a proportionality factor depending on K_ρ). c) Inter-chain optical conductivity. Within ch-DMFT vertex corrections drop out [13] and the conductivity is given by

$$\sigma_\perp(i\omega) \propto \frac{t_\perp^2}{\omega} \int \frac{dk_\perp}{2\pi} \sin^2 k_\perp \int \frac{dk}{2\pi} \frac{1}{\beta} \sum_m G(k, k_\perp, i\omega_m) \times G(k, k_\perp, i\omega_m + i\omega) \quad (4)$$

Note that we have taken into account the k_\perp -dependence of the current vertex in (4). In practice, we perform analytically the k_\perp -integration in (4) and numerically the k -integration and Matsubara summation. We then use the Maximum Entropy (MaxEnt) algorithm [17] to continue $\sigma_\perp(i\omega)$ to the real axis and obtain the inter-chain optical conductivity $\text{Re} \sigma_\perp(\omega, T)$.

We first discuss our results away from half-filling, for $U/W = 1$ and $\mu = 0.2$ (corresponding to a total density $n \simeq 0.8$). The numerical method was thoroughly tested for decoupled chains ($t_\perp = 0$), for which our fitting procedure of χ_s yields $K_\rho \simeq 0.7$, in agreement with the exact result [18]. We then study how inter-chain coherence develops as temperature is lowered for coupled chains with $t_\perp/W = 0.14$. Fig. 2 displays the effective K_ρ as a function of temperature: a clear crossover from a LL (with $K_\rho \simeq 0.7$) to a FL ($K_\rho = 1$) is seen as temperature is lowered below $T^*/W \simeq 1/44$. According to [5, 8], the




FIG. 2: Effective K_ρ vs. temperature in the doped case (filling $n \simeq 0.8$) for $U/W = 1.0$, $t_\perp/W = 0.14$.

TABLE I: QP weights $Z(k_\perp)$ for different points on the FS (doped case: $n \simeq 0.8$, $U/W = 1.0$, $t_\perp/W = 0.14$).

k_\perp/π	0.19	0.36	0.50	0.62	0.74	0.91
$Z(k_\perp)$	0.62	0.65	0.69	0.72	0.75	0.78

crossover scale is given by $T^* = \frac{t_\perp}{\pi} C(t_\perp/t)^{\theta/(1-\theta)}$, with $\theta = (K_\rho + 1/K_\rho)/4 - 1/2$ ($\simeq 0.03$ here). The interactions reduce its value compared to the naive one t_\perp/π . We cannot meaningfully test this renormalization effect because of the small values of θ in the Hubbard model. Our results are consistent with $T^* \simeq C t_\perp/\pi$ with a proportionality constant $C \simeq 0.5$. In the low temperature FL regime, we find that the location of the FS is essentially unaffected by interactions. The quasi-particle residue Z is not uniform along the FS. Its dependence on k_\perp^F is displayed in Table I. Z is larger for k_\perp close to π , while it is somewhat smaller for small k_\perp . This is not a very big effect however, and it would be inappropriate to speak of “hot spots” [19] in that region (even though it may be indicative that such a phenomenon appears for stronger interactions). Note that the single-chain approximation $\Sigma = \Sigma_{1D}$ used by many authors would lead to a much stronger variation of Z along the FS. Our results, in agreement with those of [12], show that Z is more uniform than this approximation would suggest.

We now turn to the commensurate (half-filled) case, and present results for $U = 0.65W$. By fitting the self-energy to an analytical form, we first checked that our numerical results are consistent with a Mott insulating state with a gap $\Delta_{1D}/W = 0.1$, in agreement with exact results. Turning on t_\perp , we estimate the t_\perp -dependence of the gap by performing simulations at one of the lowest accessible temperatures $T/W = 1/40$. The gap vanishes for $t_\perp/W \simeq 0.07$. Hence our method captures the insulator-to-metal transition induced by transverse hopping. The effective K_ρ , shown in Table II is also a clear indicator of the MI ($K_\rho = 0$) to FL ($K_\rho = 1$) transition, in agree-

TABLE II: Effective K_ρ at half-filling, as a function of t_\perp/W for $U/W = 0.65$ and $T/W = 1/40$.

t_\perp/W	0.00	0.04	0.07	0.11	0.14	0.16	0.18
K_ρ	0.00	0.02	1.01	1.09	1.07	1.06	1.04




FIG. 3: Inter-chain optical conductivity at half-filling for $U = 0.65W$, $\beta = 40/W$ and $t_\perp = 0.14$ and 0.07 (solid and dashed lines) and for $U = 1.0W$, $\beta = 40/W$ and $t_\perp = 0$. (dotted line).

ment with the qualitative expectations of Fig. 1. At this low temperature, the intermediate LL regime is too narrow to be visible. The location of the transition is in reasonable agreement with the naive criterion $\Delta_{1D} \sim t_\perp^*$ [9, 10, 11]. Fig. 3 displays the inter-chain optical conductivity for several values of t_\perp . In the MI phase, $\sigma_\perp(\omega)$ shows a gap, followed by an onset of absorption starting at approximately the gap and extending up to a scale of order U , where a broad second peak is apparent. A low-frequency Drude peak develops as the insulator to metal transition is crossed. Close to the transition, the weight in the Drude peak is small, while the Hubbard band feature is still clearly visible and carries a significant part of the spectral weight. The results for the NMR relaxation rate $1/T_1$ are displayed in Fig. 4 for a value of $t_\perp/W = 0.11$. At low temperatures the Korringa law $1/(T_1T) = \text{const}$ is recovered, while at higher temperatures $1/T_1T$ is temperature dependent. The observed T-dependence is consistent with $1/(T_1T) \simeq T^{K_\rho-1}$ [5] with a T-dependent exponent $K_\rho \leq 1$, indicative of a LL evolving gradually into a FL (cf. Fig. 1). As for the doped case, the FS is nearly indistinguishable from the non-interacting case (see Fig. 5). The QP residue Z_{k_\perp} depends only very weakly on the location on the Fermi surface as shown in Table III. The data may suggest very shallow minima, which in contrast to the doped case, are found at $k_\perp \sim \pm\pi/2$, corresponding to a vanishing inter-chain kinetic energy. We would like to contrast our findings with the conclusions drawn from the “single-chain” (RPA) approximation [6, 20]: $\Sigma = \Sigma_{1D}$. As recently




FIG. 4: $1/(T_1 T)$ versus T in the half-filled case with $t_{\perp}/W = 0.11$, $U/W = 0.65$. The crossover from a FL at low- T to a LL for $T/W \geq 0.025$ is clearly apparent.

TABLE III: QP weights $Z(k_{\perp})$ for different points on the FS (half-filled case, $t_{\perp} = 0.14W$, $U/W = 0.65$).

k_{\perp}/π	0.23	0.38	0.50	0.62	0.77
$Z(k_{\perp})$	0.79	0.77	0.76	0.77	0.79

pointed out [20], this approximation predicts that the FS close to the metal-insulator transition should consist of disconnected electron and hole “pockets”, as depicted schematically on Fig. 5. This is because the self-energy of a MI diverges at low-frequency for $k = \pi/2$, so that no FS point can correspond to this value of k . In contrast, our data find a conventional connected FS down to the transition point (up to our numerical accuracy). This shows that it is crucial, as done in ch-DMFT, to take into account the feedback effects of the interchain hopping in the self-energy.

In conclusion, we have shown in this paper that the ch-DMFT approach is a tool of choice for the study of the crossovers and insulator to metal transitions induced by transverse hopping in quasi one-dimensional systems. Several features found in our study are qualitatively reminiscent of experimental observations on organic conductors, such as: the deconfinement transition itself, the crossover from a LL at high- T to a FL at low- T (as revealed e.g in NMR) and most significantly the coexistence of a Drude feature with small spectral weight and of a Hubbard-band feature in optical conductivity. Extensions of the model including on-site and nearest neighbor interactions would allow to study the deconfinement transition at quarter-filling and thus to make a more realistic [3, 9] comparison with experimental results. Though this is computationally much more demanding, we hope to use ch-DMFT in this context in future work.

Acknowledgements: We acknowledge useful discussions with A. Tselvik, F. Essler, D. Jerome and C. Bourbonnais. This research has been supported by a Marie




FIG. 5: FS in the half-filled case with $t_{\perp}/W = 0.14$, $U/W = 0.65$ (circles), compared to the FS of the non-interacting case (dotted line) and of the purely 1d case ($t_{\perp} = 0$ -dashed-). The solid line depicts schematically the Fermi surface obtained when making the approximation $\Sigma = \Sigma_{1D}$ [20].

Curie Fellowship of the European Community Programme “Improving Human Potential” under contract number MCFI-2000804 and a grant of supercomputing time at NIC Jülich.

- [1] J. Voit, Rep. Prog. Phys. **58**, 977 (1995).
- [2] D. Jérôme and H. J. Schulz, Adv. Phys. **31**, 299 (1982).
- [3] A. Schwartz, M. Dressel, G. Grüner, V. Vescoli, L. Degiorgi, and T. Giamarchi, Phys. Rev. B **58**, 1261 (1998).
- [4] J. Moser, M. Gabay, P. Aubin-Senzier, D. Jerome, K. Bechgaard, and J. Fabre, Eur. Phys. J. B **1**, 39 (1998).
- [5] C. Bourbonnais, F. Creuzet, D. Jérôme, K. Bechgaard, and A. Moradpour, J. de Phys. (Paris) Lett. **45**, L755 (1984).
- [6] X. G. Wen, Phys. Rev. B **42**, 6623 (1990).
- [7] V. M. Yakovenko, JETP Lett. **56**, 510 (1992).
- [8] C. Bourbonnais and L. G. Caron, Int. J. Mod. Phys. B **5**, 1033 (1991).
- [9] T. Giamarchi, Physica B **230-232**, 975 (1997).
- [10] V. Vescoli, L. Degiorgi, W. Henderson, G. Grüner, K. P. Starkey, and L. K. Montgomery, Science **281**, 1191 (1998).
- [11] M. Tsuchiizu, P. Donohue, Y. Suzumura, and T. Giamarchi, Eur. Phys. J. B **19**, 185 (2001), and references therein.
- [12] E. Arrigoni, Phys. Rev. Lett. **83**, 128 (1999), and Phys. Rev. B **61**, 7909 (2000).
- [13] A. Georges, T. Giamarchi, and N. Sandlers, Phys. Rev. B **61**, 16393 (2000).
- [14] A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, Rev. Mod. Phys. **68**, 13 (1996).
- [15] J. E. Hirsch and R. M. Fye, Phys. Rev. Lett. **25**, 2521 (1986).

- [16] M. Randeria, N. Trivedi, A. Moreo, and R. T. Scalettar, Phys. Rev. Lett. **69**, 2001 (1992).
- [17] M. Jarrell and J. E. Gubernatis, Phys. Rep. **269**, 133 (1996).
- [18] H. J. Schulz, Phys. Rev. Lett. **64**, 2831 (1990).
- [19] A. T. Zheleznyak and V. M. Yakovenko, Synth. Met. **70**, 1005 (1995).
- [20] F. H. L. Essler and A. M. Tsvelik (2001), unpublished.
- [21] To get a reliable estimate for the self-energy we used an analytical Fourier transform of a spline interpolated Green's function with exact boundary conditions in the spirit of Joo et al., condmat-0009367.