

# Growing Surfaces with Anomalous Diffusion – Results for the Fractal Kardar-Parisi-Zhang Equation

Eytan Katzav

Raymond and Beverly Sackler Faculty of Exact Sciences

School of Physics and Astronomy, Tel Aviv University,

Ramat Aviv, Tel Aviv 69978, Israel

## Abstract

In this paper I study a model for a growing surface in the presence of anomalous diffusion, also known as the Fractal Kardar-Parisi-Zhang equation (FKPZ). This equation includes a fractional Laplacian that accounts for the possibility that surface transport is caused by a hopping mechanism of a Levy flight. I show that for a specific choice of parameters of the FKPZ equation, the equation can be solved exactly in one dimension, so that all the critical exponents, which describe the surface that grows under FKPZ, can be derived for that case. Afterwards, I use a Self-Consistent Expansion (SCE) to predict the critical exponents for the FKPZ model for any choice of the parameters and any spatial dimension. It is then verified that the results obtained using SCE recover the exact result in one dimension. At the end a simple picture for the behavior of the Fractal KPZ equation is suggested, and the upper critical dimension of this model is discussed.

The Kardar-Parisi-Zhang (KPZ) equation [2] for surface growth under ballistic deposition was introduced as an extension of the Edwards-Wilkinson theory [1]. The interest in the KPZ equation exceeds far beyond the interest in evolving surfaces because of the following reasons: (a) The KPZ system is known to be equivalent to a number of very different physical systems. Examples are the directed polymer in a random medium, the Schrodinger equation (in imaginary time) for a particle in the presence of a potential that is random in space and time and the important Burgers equation from hydrodynamics [1]. (b) The second reason, that is more important, to my mind, is that it serves as a relatively simple prototype of non-linear stochastic field equations that are so common in condensed matter physics.

The equation for the height of the surface at the point  $\vec{r}$  and time  $t$ ,  $h(\vec{r}, t)$ , is given by

$$\frac{\partial h}{\partial t} = v\nabla^2 h + \frac{\lambda}{2}(\nabla h)^2 + \eta(\vec{r}, t) \quad (1)$$

where automatically the constant deposition rate is removed and  $\eta(\vec{r}, t)$  is a noise term such that

$$\begin{aligned} \langle \eta(\vec{r}, t) \rangle &= 0 \\ \langle \eta(\vec{r}, t) \eta(\vec{r}', t') \rangle &= 2D_0 \delta(\vec{r} - \vec{r}') \delta(t - t') \end{aligned} \quad (2)$$

As can be seen in eq. (1) the basic relaxation mechanism in the KPZ equation is a Laplacian term that results from nearest neighbor hopping in the growing surface. During the last years, there has been a growing interest in other relaxational mechanisms [22-25], namely subdiffusive diffusion that seems to appear in the context of charge transport in amorphous semiconductors [26,27], NMR diffusometry in disordered materials [28], and the dynamics of a bead in polymer networks [29]. The theoretical effort to account for such phenomena led to the formulation of the

celebrated fractional Fokker-Planck equation (FFPE) [22]. This equation includes a fractional (Riemann-Liouville) operator instead of the standard derivative of the Fokker-Planck equation.

Recently, Mann & Woyczynski [3] have suggested that in order to account for experimental data, namely experiments in which impurities were present on the growing surface [4], a modification of the KPZ equation has to be considered. They used the observation that the presence of an impurity can act as a strong trap for an adatom migrating at room temperature, to conjecture that this process corresponds to Levy flights between trap sites. This conjecture then served as a justification for the introduction of a fractional Laplacian into the continuum equation of the growing surface as another relaxation mechanism. Actually, the fractional Laplacian dominates the standard Laplacian in the KPZ equation in the scaling regime (i.e. in the large scale limit), so that the standard Laplacian can be ignored from the beginning. To summarize this exposition and to be more specific, the equation they eventually suggest to describe the growing surface in the presence of self-similar hopping surface diffusion is the fractal KPZ (FKPZ) equation given by

$$\frac{\partial h(\vec{r}, t)}{\partial t} = v(\Delta_\alpha h) + \frac{\lambda}{2}(\nabla h)^2 + \eta(\vec{r}, t), \quad (3)$$

where  $\Delta_\alpha \equiv -(-\Delta)^{\alpha/2}$  (or in Fourier space  $\mathcal{F}(\Delta_\alpha f)(q) \equiv -(|q|)^{\alpha/2} \mathcal{F}f(q)$ ) is the fractional Laplacian, and in our context it is more convenient to choose  $\alpha = 2 - \rho$ , with  $0 \leq \rho < 2$  (where the special case  $\rho = 0$  corresponds to the standard KPZ equation). In addition,  $\eta(\vec{r}, t)$  is a noise term satisfying

$$\langle \eta(\vec{r}, t) \rangle = 0, \quad (4)$$

$$\langle \eta(\vec{r}, t) \eta(\vec{r}', t') \rangle = 2D_0 |\vec{r} - \vec{r}'|^{2\sigma-d} \delta(t - t'). \quad (5)$$

(actually Mann & Woyczynski [3] discussed the specific case of white noise that corresponds to  $\sigma = 0$  in the last equation).

The proposed FKPZ equation generalizes the FFPE equation mentioned above in that it is a field equation, rather than an equation for a single degree of freedom. Therefore, it is understood that such a generalization is essential in order to account for the dynamics of a whole medium experiencing anomalous diffusion, and not just an artificial problem. Obviously, the technical mathematical difficulties to be overcome in this nonlinear case are formidable in comparison to those for the linear fractal kinetic equations.

However, in their paper Mann & Woyczynski [3] were not able to predict the critical exponents that describe the surface that grows under FKPZ. But before I make any new statements about this model let me briefly summarize the various quantities of interest.

A very important quantity of interest is the roughness exponent  $\alpha$  that characterizes the surface in steady state. It is defined by the following relation

$$\langle [h(\vec{r}) - h(\vec{r}')]^2 \rangle \propto |\vec{r} - \vec{r}'|^{2\alpha} \quad (6)$$

The roughness exponent is sometimes also defined using  $w(L, t)$  the roughness of the surface (that is defined as the RMS of the height function  $h(\vec{r}, t)$ , in a system of linear size  $L$ ). Then, in terms of  $w(L, t)$ ,  $\alpha$  is given by

$$w(L, t) \propto L^\alpha \quad (7)$$

Another important quantity of interest is the growth exponent  $\beta$  that describes the short time behavior of the roughness  $w(L, t)$  (with flat initial conditions)

$$w(L, t) \propto t^\beta \quad (8)$$

Finally, I introduce the dynamic exponent  $z$  that describes the typical relaxation time scale of the system (i.e. the dependence of the equilibration time on the size of the system)

$$t_x \propto L^z \quad (9)$$

It is well known [1] that these three exponents are not independent and that under very general considerations one should expect the following scaling relation

$$z = \alpha/\beta \quad (10)$$

(this relation is a direct consequence of the Family-Vicsek scaling relation [5]).

In this paper I show that for a specific choice of parameters of the FKPZ equation (namely  $\rho = 2\sigma$ ), the equation can be solved exactly in one dimension, so that all the critical exponents can be derived easily for that case. This solution is accomplished using the Fokker-Planck equation associated with FKPZ. Afterwards, in order to give a more complete picture (i.e. for any dimension  $d$ , and any spatial correlation index  $\sigma$ ) I apply a method developed by Schwartz and Edwards [6-8] (also known as the Self-Consistent-Expansion (SCE) approach). This method has been previously applied successfully to the KPZ equation. The method gained much credit by being able to give a sensible prediction for the KPZ critical exponents in the strong coupling phase, where many Renormalization-Group (RG) approaches failed, as well as Dynamic Renormalization Group (DRG) [2, 9]. It is then verified that the results obtained using SCE recover the exact result in one dimension.

In order to derive the exact result, notice that it is naturally more convenient to work with the Fourier components of eq. (3)

$$\frac{\partial \mathbf{h}_q}{\partial t} = -vq^{2-\rho} \mathbf{h}_q - \frac{\lambda}{2\sqrt{\Omega}} \sum_{\ell, m} (\vec{\ell} \cdot \vec{m}) \delta_{q, \ell+m} \mathbf{h}_\ell \mathbf{h}_m + \eta_q, \quad (11)$$

(from now on I use  $v_q = vq^{2-\rho}$ , and  $M_{q\ell m} = \frac{\lambda}{2\sqrt{\Omega}} (\vec{\ell} \cdot \vec{m}) \delta_{q, \ell+m}$ )

where  $\eta_q$  is the noise term that satisfies:

$$\langle \eta_q(t) \rangle = 0 \quad (12)$$

$$\langle \eta_q(t) \eta_{q'}(t') \rangle = 2D_0 q^{-2\sigma} \delta_{q, -q'} \delta(t - t'). \quad (13)$$

I now turn from the FKPZ equation in Langevin form to a Fokker-Planck form

$$\frac{\partial \mathbf{P}}{\partial t} + \sum_q \frac{\partial}{\partial \mathbf{h}_q} \left[ D_0 q^{-2\sigma} \frac{\partial}{\partial \mathbf{h}_{-q}} + vq^{2-\rho} \mathbf{h}_q + \frac{\lambda}{2\sqrt{\Omega}} \sum_{\ell, m} (\vec{\ell} \cdot \vec{m}) \delta_{q, \ell+m} \mathbf{h}_\ell \mathbf{h}_m \right] \mathbf{P} = 0 \quad (14)$$

where  $\mathbf{P}(\{\mathbf{h}_q\}, t)$  is the probability of having a configuration  $\{\mathbf{h}_q\}$  at a specific time  $t$ .

By direct substitution, it can be verified that the following probability function (a Gaussian actually)

$$\mathbf{P}_0 = N \exp \left[ -\frac{1}{2} \sum_q \frac{\mathbf{h}_q \mathbf{h}_{-q}}{\phi_q} \right] \quad (15)$$

(where  $N$  is the normalization yielding  $\int \mathbf{P}_0 \{\mathbf{h}\} D\mathbf{h} = 1$ , and  $\phi_q$  is the two point function) is a solution of the steady-state Fokker-Planck equation (i.e. with  $\partial \mathbf{P} / \partial t = 0$ ) for the one-dimensional case ( $d = 1$ ), if  $\phi_q = Aq^{-\Gamma}$  with  $A = D_0/v$ ,  $\Gamma = 2$  and  $\rho = 2\sigma$ . This gives  $\alpha = \frac{1}{2}$  (the translation of  $\Gamma = 2$  into  $\alpha = 1/2$  is done using eq. (18) below) that coincides with the known exact solution of one-dimensional classical KPZ  $\alpha = 1/2$  - as it should be (since for classical KPZ  $\rho = \sigma = 0$ ).

Now, I apply the Self-Consistent Expansion (SCE) in order to learn about the behavior of this system in more general contexts (other dimensions and cases where

$\rho \neq 2\sigma$ ). SCE's starting point is the Fokker-Planck form of FKPZ (eq. (14) above), from which it constructs a self-consistent expansion of the distribution of the field concerned.

The expansion is formulated in terms of  $\phi_q$  and  $\omega_q$ , where  $\phi_q$  is the two-point function in momentum space, defined by  $\phi_q = \langle h_q h_{-q} \rangle_S$ , (the subscript S denotes steady state averaging), and  $\omega_q$  is the characteristic frequency associated with  $h_q$ . It is expected that for small enough  $q$ ,  $\phi_q$  and  $\omega_q$  are power laws in  $q$ ,

$$\phi_q = Aq^{-\Gamma} \quad (16)$$

and

$$\omega_q = Bq^\mu. \quad (17)$$

[Since dynamic surface growth is a remarkably multidisciplinary field, there are almost as many notations as there are workers in the field. Therefore I give a brief translation of our notations to those most frequently used:

$$\begin{aligned} \mu &= z, \\ \alpha &= \frac{\Gamma - d}{2}, \\ \beta &= \frac{\alpha}{z} = \frac{\Gamma - d}{2\mu}. \end{aligned} \quad (18) \quad ]$$

The main idea is to write the Fokker-Planck equation  $\partial P/\partial t = OP$  in the form  $\partial P/\partial t = [O_0 + O_1 + O_2]P$ , where  $O_0$  is to be considered zero order in some parameter  $\lambda$ ,  $O_1$  is first order and  $O_2$  is second order. The evolution operator  $O_0$  is chosen to

have a simple form  $O_0 = -\sum_q \frac{\partial}{\partial h_q} \left[ D_q \frac{\partial}{\partial h_{-q}} + \omega_q h_q \right]$ , where  $D_q/\omega_q = \phi_q$ . Note that

at present  $\phi_q$  and  $\omega_q$  are not known. I obtain next an equation for the two-point function. The expansion has the form  $\phi_q = \phi_q + c_q \{\phi_p, \omega_p\}$ , because the lowest order in the expansion already yields the unknown  $\phi_q$ . In the same way an expansion for  $\omega_q$  is also obtained in the form  $\omega_q = \omega_q + d_q \{\phi_p, \omega_p\}$ . Now, the two-point function and the characteristic frequency are thus determined by the two coupled equations

$$c_q \{\phi_p, \omega_p\} = 0 \quad \text{and} \quad d_q \{\phi_p, \omega_p\} = 0. \quad (19)$$

These equations can be solved exactly in the asymptotic limit to yield the required scaling exponents governing the steady state behavior and the time evolution. Working to second order in the expansion, one gets the two coupled integral equations

$$D_q - v_q \phi_q + 2 \sum_{\ell, m} \frac{M_{q\ell m} M_{q\ell m} \phi_\ell \phi_m}{\omega_q + \omega_\ell + \omega_m} - 2 \sum_{\ell, m} \frac{M_{q\ell m} M_{\ell m q} \phi_m \phi_q}{\omega_q + \omega_\ell + \omega_m} - 2 \sum_{\ell, m} \frac{M_{q\ell m} M_{m\ell q} \phi_\ell \phi_q}{\omega_q + \omega_\ell + \omega_m} = 0, \quad (20)$$

and

$$v_q - \omega_q - 2 \sum_{\ell, m} M_{q\ell m} \frac{M_{\ell m q} \phi_m + M_{m\ell q} \phi_\ell}{\omega_\ell + \omega_m + \omega_q} = 0, \quad (21)$$

where in deriving the last equation I have used the Herring consistency equation [10]. In fact Herring's definition of  $\omega_q$  is one of many possibilities, each leading to a different consistency equation. But it can be shown, as previously done in [7], that this does not affect the exponents (universality).

Detailed solution of equations (20) and (21) in the limit of small  $q$ 's (i.e. large scales) yields a rich family of solutions that I shall describe immediately.

First, there are two kinds of weak-coupling solutions – both with a dynamic exponent  $\mu = 2 - \rho$  (they are called weak-coupling because they are exactly the



solutions obtained in the case of the Fractal Edwards- Wilkinson equation, see [3]). Now, when the spatial correlations of the noise are relevant (i.e., when  $\sigma > 0$ ) I obtain the solution  $\Gamma = 2 - \rho + 2\sigma$  - provided that  $d > 2 - 3\rho + 2\sigma$ . But if the spatial correlations of the noise are not relevant (i.e., when  $\sigma \leq 0$ ) I obtain the simpler solution  $\Gamma = 2 - \rho$  - provided that  $d > 2 - 3\rho$ .

The second type of solutions is strong coupling solutions that obey the well known scaling relation  $\mu = \frac{d+4-\Gamma}{2}$  obtained from eq. (21). The first strong coupling solution is determined by the combination of the scaling relation and the transcendental equation  $F(\Gamma, \mu) = 0$ , where F is given by

$$F(\Gamma, \mu) = -\int d^d t \frac{\vec{t} \cdot (\hat{e} - \vec{t})}{t^\mu + |\hat{e} - \vec{t}|^\mu + 1} \left[ (\hat{e} \cdot \vec{t}) |\hat{e} - \vec{t}|^{-\rho} t^{-\Gamma} + \hat{e} \cdot (\hat{e} - \vec{t}) t^{-\rho} |\hat{e} - \vec{t}|^{-\Gamma} \right] + \int d^d t \frac{[\vec{t} \cdot (\hat{e} - \vec{t})]^2}{t^\mu + |\hat{e} - \vec{t}|^\mu + 1} t^{-\Gamma} |\hat{e} - \vec{t}|^{-\Gamma}, \quad (22)$$

and  $\hat{e}$  is a unit vector in an arbitrary direction.

This solution is valid as long as the solutions of the last equations satisfy the following condition  $\Gamma > \max \left\{ \frac{d+4+4\sigma}{3}, \frac{d+4}{3}, d+2\rho \right\}$ .

It turns out that for  $d = 1$  the equation  $F(\Gamma, \mu(\Gamma)) = 0$  is exactly solvable, and yields  $\Gamma = 2$  and  $\mu = \frac{3}{2}$  (it can be checked immediately by direct substitution). In this

case the validity condition reads  $2 > \max \left\{ \frac{5+4\sigma}{3}, 1+2\rho \right\}$  or equivalently  $0 < \rho < 1/2$

and  $\sigma < 1/4$ . By using eq. (18) I translate the results into  $\alpha = \frac{1}{2}$  and  $z = \frac{3}{2}$  that are precisely the exact classical KPZ results.

It should also be mentioned that for  $d \geq 2$  such an exact solution in closed form cannot be found, and one has to solve numerically the equation  $F(\Gamma, \mu(\Gamma)) = 0$ . For convenience I denote the numerical value of this solution by  $\Gamma_0(d)$ . For example, in two dimensions I obtain  $\Gamma_0(2) = 2.59$ .

The second strong coupling solution is obtained by power-counting, and it is relevant when  $\sigma > 0$  (i.e. when the spatial correlations of the noise are relevant),  $d < 2 + 2\sigma$  and  $F(\Gamma, \mu(\Gamma)) < 0$  (the last condition turns out to be equivalent to the condition  $\Gamma > \Gamma_0(d)$  because from this value of  $\Gamma$  and on the function  $F(\Gamma, \mu(\Gamma))$  is negative). This power counting solution can be written in closed form and is given by

$$\mu = \frac{d+4-2\sigma}{3} \text{ and } \Gamma = \frac{d+4+4\sigma}{3} = \mu + 2\sigma.$$

The third strong coupling solution (that is in some sense the only "genuine" FKPB solution, in the sense that it is the only solution that is dramatically influenced by the fractional Laplacian, and at the same time is not a solution of the Fractal Edwards-Wilkinson equation) is determined by the combination of the scaling relation

$$\mu = \frac{d+4-\Gamma}{2} \text{ and the extra relation } d+4-2\Gamma-\mu = 2-\rho-\Gamma. \text{ This solution can be}$$

written in closed form as  $\Gamma = d+2\rho$  and  $\mu = 2-\rho$ . It turns out that this phase is relevant when  $d > \max\{2-3\rho, 2-3\rho+2\sigma\}$ .

At this point, the full picture of possible solutions might seem too complex, so I want to suggest the following simple interpretation for the behavior of the Fractal KPZ equation. If you remember, the starting point of this discussion was the introduction of the fractional Laplacian into the KPZ equation. This immediately implies that faster relaxations are now possible (faster in the sense that a smaller

dynamical exponent is expected) when compared with the Edwards-Wilkinson equation. However, it is well known that already the KPZ nonlinearity introduces faster relaxations (at least for dimensions lower than the upper critical dimension). Therefore, in the FKPZ system, the dynamics is controlled by the fastest "component": if the dynamical exponent of the classical KPZ system is smaller than  $2 - \rho$  then it dominates, otherwise the new fractional dynamical exponent  $z = 2 - \rho$  controls the dynamics. However, at this point the picture gets a little bit more complicated (just like in the classical KPZ case), namely there are several possible phases with this new fractional dynamical exponent  $z = 2 - \rho$ , and the transition between these phases is controlled by the strength of dimensionless coupling constant.

These conclusions have an important implication for the upper critical dimension of the FKPZ model (i.e. the dimension above which the dynamical exponent is the same as that of the linear theory). Namely, it turns out that the FKPZ equation always has an upper critical dimension that is determined by the relation  $\Gamma_0(d_{uc}^{FKPZ}) = d_{uc}^{FKPZ} + 2\rho$  (or alternatively, using the roughness exponent  $\alpha$  - the upper critical dimension is the dimension where  $\alpha_{KPZ}(d_{uc}^{FKPZ}) = \rho$ ). Notice, that this result does not depend on the ongoing debate over the existence of the upper critical dimension for the KPZ system (see [11-20]). Actually, it merely requires that the roughness exponent of the classical KPZ system becomes arbitrarily small in higher dimensions – an assumption that is generally accepted (only Tu [21] had a different conjecture).

As one can see, the results obtained using the Self-Consistent Expansion (SCE) are quite general, and cover all possible values of the relevant parameters ( $\rho$

and  $\sigma$ ) as well as dimensions. It is also easily verified that these results recover the exact result obtained at the beginning of this paper for the case  $\rho = 2\sigma$ . This situation, suggest that the SCE method is generally appropriate when dealing with non-linear continuum equations.

Acknowledgement: I would like to thank Moshe Schwartz for useful discussions.

## REFERENCES

- [1] A.-L. Barabasi and H. E. Stanley, *Fractal Concepts in Surface Growth* (Cambridge Univ. Press, Cambridge, 1995).
- [2] M. Kardar, G. Parisi and Y.-C. Zhang, Phys. Rev. Lett. **56**, 889 (1986).
- [3] J.A. Mann Jr. and W.A. Woyczynski, Physica A **291**, 159 (2001).
- [4] G.L. Kellog, Phys. Rev. Lett. **72**, 1662 (1994).
- [5] F. Family and T. Vicsek, J. Phys. A **18**, L75 (1985).
- [6] M. Schwartz and S. F. Edwards, Europhys. Lett. **20**, 301 (1992).
- [7] M. Schwartz and S. F. Edwards, Phys. Rev. E **57**, 5730 (1998).
- [8] E. Katzav and M. Schwartz, Phys. Rev. E **60**, 5677 (1999).
- [9] Medina E., Hwa T. and Kardar M., Phys. Rev. A, **39** (1989) 3053.
- [10] J. R. Herring, Phys. Fluids **8**, 2219 (1965); **9**, 2106 (1966).
- [11] T. Ala-Nissila, Phys. Rev. Lett. **80**, 887 (1998).  
J. M. Kim, *ibid.*, **80**, 888 (1998).  
M. Lassig and H. Kinzelbach, *ibid.*, **80**, 889 (1998).
- [12] E. Perlsman and M. Schwartz, Physica A **234**, 523 (1996).
- [13] C. Castellano, M. Marsili and L. Pietronero, Phys. Rev. Lett. **80**, 3527 (1998).  
C. Castellano, A. Gabrielli, M. Marsili, M. A. Munoz and L. Pietronero, Phys. Rev. E **58**, 5209 (1998).
- [14] T. Halpin-Healy, Phys. Rev. A **42**, 711 (1990).
- [15] T. Blum and A. J. McKane, Phys. Rev. E **52**, 4741 (1995).
- [16] J-P. Bouchaud and M. E. Cates, Phys. Rev. E **47**, 1455 (1993);  
(erratum), Phys. Rev. E **48**, 653 (E) (1993).
- [17] E. Katzav and M. Schwartz, Physica A **309**, 69 (2002).
- [18] J. Cook and B. Derrida, J. Phys. A **23**, 1523 (1990).

- [19] F. Colaiori and M. A. Moore, Phys. Rev. Lett. **86**, 3946 (2001).
- [20] E. Marinari, A. Pagnani, G. Parisi, and Z. Rácz, Phys. Rev. E **65**, 026136 (2002)
- [21] Y. Tu, Phys. Rev. Lett. **73**, 3109 (1994).
- [22] R. Metzler, E. Barkai and J. Klafter, Phys. Rev. Lett. **82**, 3563 (1999).
- [23] R. Metzler and J. Klafter, Chem. Phys. Lett. **321**, 238 (2000).
- [24] R. Granek and J. Klafter, Europhys. Lett. **56**, 15 (2001).
- [25] M. Sokolov, J. Klafter and A. Blumen, Phys. Rev. E **64**, 21107 (2001).
- [26] H. Scher and E. Montroll, Phys. Rev. B **12**, 2455 (1975).
- [27] Q. Gu, E. A. Schiff, S. Grebner, and R. Schwartz, Phys. Rev. Lett. **76**, 3196 (1996).
- [28] A. Klemm, H.-P. Müller, and R. Kimmich, Phys. Rev. E **55**, 4413 (1997).
- [29] F. Amblard, A. C. Maggs, B. Yurke, A. N. Pargellis, and S. Leibler, Phys. Rev. Lett. **77**, 4470 (1996).