

ON THE ISING-HEISENBERG MODEL WITH THE DOUBLY DECORATED NETWORK STRUCTURE I

Jozef Strečka and Michal Jaščur

Department of Theoretical Physics and Astrophysics, Faculty of Science,
P. J. Šafárik University, Moyzesova 16, 041 54 Košice, Slovak Republic,
E-mail:jozkos@pobox.sk, jascur@kosice.upjs.sk.

Abstract

Using an exact mapping transformation method, magnetic properties of a spin-1/2 and spin-1 doubly decorated Ising-Heisenberg model are investigated in detail. By assuming that both interaction parameters are ferromagnetic, we found in addition to an usual ferromagnetic phase another two phases with a nontrivial long-range order.

Keywords: Ising-Heisenberg model; Mapping transformation

Low-dimensional quantum spin systems have become during the last few years the subject of immense interest, since their properties are strongly affected by quantum fluctuations. However, the role of quantum fluctuations can be understood from a quantum Heisenberg model (QHM) - basic theoretical model of magnetic insulators. Although much effort has been focused on this topic, a general exact solution has not been found for any 2D QHM, yet.

Owing to this fact, we will propose a simplified doubly decorated Ising-Heisenberg model, which can be exactly treated by means of a generalized mapping transformation technique [1, 2]. Moreover, the suggested model can also be viewed as useful model of magnetic insulators with the doubly decorated network structure. Experimentally, such an arrangement of metal ions has been recently reported in a novel $\text{Rh}^{2+}\text{-Co}^{3+}$ compound [3]. Hence, the considered model represents a feasible model for possible structural derivatives of the aforementioned bimetallic compound.

In this article, we will explore the Ising-Heisenberg model with the doubly decorated network structure (see Fig.1), in which the sites of original lattice are occupied by the spin-1/2 atoms (full circles) and the decorating sites by the spin-1 atoms (empty circles). We will further assume that the nearest-neighbour interaction $J(\Delta)$ between spin-1 atoms is an anisotropic Heisenberg interaction (Δ marks the XXZ exchange anisotropy), while the nearest-neighbour interaction J_1 between spin-1/2 and spin-1 atoms is purely an Ising-type interaction.

Then, it is very convenient to write the total Hamiltonian as a sum of bond Hamiltonians, namely $\hat{\mathcal{H}} = \sum_k \hat{\mathcal{H}}_k$, where each bond Hamiltonian $\hat{\mathcal{H}}_k$ involves all the interaction terms associated with k th couple of spin-1 atoms (Fig.1)

$$\begin{aligned} \hat{\mathcal{H}}_k = & - J[\Delta(\hat{S}_{k1}^x \hat{S}_{k2}^x + \hat{S}_{k1}^y \hat{S}_{k2}^y) + \hat{S}_{k1}^z \hat{S}_{k2}^z] \\ & - J_1 \hat{S}_{k1}^z \hat{\mu}_{k1}^z - J_1 \hat{S}_{k2}^z \hat{\mu}_{k2}^z. \end{aligned} \quad (1)$$

In above, $\hat{\mu}_k^z$ and \hat{S}_k^α ($\alpha = x, y, z$) denote well-known components of the standard spin-1/2 and spin-1 operators, respectively. Commutability between different

bond Hamiltonians $\hat{\mathcal{H}}_k$ ensures that the generalized decoration-iteration transformation [1, 2] maps the partition function \mathcal{Z}_{IHM} of Ising-Heisenberg model on the partition function \mathcal{Z}_{IM} of simple spin-1/2 Ising model

$$\mathcal{Z}_{IHM}(\beta J, J_1, \Delta) = A^{Nq/2} \mathcal{Z}_{IM}(\beta R), \quad (2)$$

with following mapping parameters A and βR

$$\begin{aligned} A^2 &= [V(J_1, 0) + W(0)][V(0, J_1) + W(J_1)], \\ \beta R &= 2 \ln \left[\frac{V(J_1, 0) + W(0)}{V(0, J_1) + W(J_1)} \right]. \end{aligned} \quad (3)$$

In above, N denotes a total number of spin-1/2 atoms, q their coordination number and the functions $V(x, y)$ and $W(x)$ are defined by

$$\begin{aligned} V(x, y) &= 2 \exp(\beta J) \cosh(\beta x) + \\ &\quad + 4 \cosh(\beta x/2) \cosh(\beta \sqrt{y^2 + 4(J\Delta)^2}/2), \\ W(x) &= \exp(-2\beta J/3) \sum_{n=1}^3 \exp[-\beta z_n(x)], \end{aligned} \quad (4)$$

where z_n label the roots of secular cubic equation

$$\begin{aligned} z_n(x) &= 2P(x) \cos \left[\phi(x) + 2\pi(n-1)/3 \right], \\ P^2(x) &= (J/3)^2 + 2(J\Delta)^2/3 + x^2/3, \\ Q(x) &= (J/3)^3 + J(J\Delta)^2/3 - Jx^2/3, \\ \phi(x) &= \frac{1}{3} \arctan \left[-\sqrt{P^6(x) - Q^2(x)}/Q(x) \right]. \end{aligned} \quad (5)$$

It is worthy to note that Eq.(2) constitutes the basic result of our calculation, since it enables simple calculation of a large number of quantities.

To understand the behaviour of ferromagnetic Ising-Heisenberg model ($J > 0, J_1 > 0$) more deeply, we have examined in detail magnetizations, correlations and quadrupolar momentum

$$\begin{aligned} m_i^\mp &\equiv \frac{1}{2} \langle \hat{\mu}_{k1}^z \mp \hat{\mu}_{k2}^z \rangle, & m_h^\mp &\equiv \frac{1}{2} \langle \hat{S}_{k1}^z \mp \hat{S}_{k2}^z \rangle, \\ C_{hh}^{xx} &\equiv \langle \hat{S}_{k1}^x \hat{S}_{k2}^x \rangle, & C_{hh}^{zz} &\equiv \langle \hat{S}_{k1}^z \hat{S}_{k2}^z \rangle, \\ C_{ii}^{zz} &\equiv \langle \hat{\mu}_{k1}^z \hat{\mu}_{k2}^z \rangle, & C_{ih}^{zz} &\equiv \frac{1}{2} \langle \hat{S}_{k1}^z \hat{\mu}_{k1}^z + \hat{S}_{k2}^z \hat{\mu}_{k2}^z \rangle, \\ \eta &\equiv \frac{1}{2} \langle (\hat{S}_{k1}^z)^2 + (\hat{S}_{k2}^z)^2 \rangle, \end{aligned} \quad (6)$$

where $\langle \dots \rangle$ denotes standard canonical averaging, subscript $i(h)$ marks the spin-1/2 (spin-1) atom, and superscript the spatial component of spin.

Ground-state phase diagram in the $J_1/J - \Delta$ plane is shown Fig.2. As one can see, the parameter space is separated by first-order transition lines into 3 regions: the usual ferromagnetic phase (FP), valence-bond ferromagnetic phase

(VBF) and antiferromagnetic phase (AP). FP can be characterized by saturated values of both sublattice magnetizations ($m_i^+ = 1/2, m_h^+ = 1$) and the spin order well-known from pure Ising model.

Contrary to this, in VBF one finds: $m_i^+ = 1/2, m_h^+ = 1/2, C_{hh}^{zz} = 0, C_{hh}^{xx} = 1/2, C_{ih}^{zz} = 1/4, C_{ii}^{zz} = 1/4$ and $\eta = 1/2$, indicating that each couple of decorating spins comprises of a spin '01' pair (one spin in '0' state, another one in spin '1' state). Moreover, nonzero C_{hh}^{xx} clearly implies an influence of quantum fluctuations. Hence, it is reasonable to assume that both decorating spin-1 atoms interchange their spin states and thus, they effectively act as the spin-1/2 atoms [2].

Finally, there also appears the unusual AP with a perfect antiferromagnetic spin alignment in the spin-1/2 sublattice ($m_i^- = 1/2, C_{ii}^{zz} = -1/4$). On the other hand, in the spin-1 sublattice we found the coexistence of an antiferromagnetic long-range order along z axis ($m_h^- < 1, C_{hh}^{zz} \ll 0$) together with a ferromagnetic short-range ordering in xy plane ($C_{hh}^{xx} \gg 0$). To the best of our knowledge, the existence of such antiferromagnetic long-range ordering has not been reported in any pure ferromagnetic system before.

Acknowledgement: This work was supported by APVT grant No. 20-009902 and VEGA grant No. 1/9034/02

References

- [1] M. E. Fisher, Phys. Rev. 113 (1959) 969.
- [2] J. Strečka and M. Jaščur, Phys. Stat. Sol. (b) 233 (2002) R12.
- [3] J. Lu *et al*, Chem. Commun. (1996) 399.

FIGURE CAPTIONS:

Fig.1: Typical example of the doubly decorated planar lattice (decorated square lattice). Full and empty circles denote the spin-1/2 and spin-1 atoms, respectively. The ellipse demarcates all interaction terms involved in $\hat{\mathcal{H}}_k$ (1).

Fig.2: Ground-state phase diagram in $J_1/J - \Delta$ plane.



