

SPIN FRUSTRATION IN AN EXACTLY SOLVABLE ISING-HEISENBERG DIAMOND CHAIN

Michal Jaščur and Jozef Strečka

Department of Theoretical Physics and Astrophysics, Faculty of Science,
P.J. Šafárik University, Moyzesova 16, 041 54 Košice, Slovak Republic
E-mail: jascur@kosice.upjs.sk, jozkos@pobox.sk

Abstract

Exact results of a mixed spin-1/2 and spin-1 Ising-Heisenberg diamond chain are obtained applying a generalized decoration-iteration transformation. Depending on the values of external field, exchange parameters and anisotropy there exist six different phases in the ground state of the system. The magnetic order of these phases is briefly described in this work.

Keywords: Ising-Heisenberg diamond chain; Exact results

During last two decades low-dimensional magnetic materials have become important subject in the research of magnetism. Among systems exhibiting interesting quantum magnetic phenomena one should mention those having a diamond-chain structure that have been recently intensively studied both theoretically and experimentally (see [1] and references therein). In order to describe these materials, the quantum anisotropic Heisenberg model has been frequently adopted. Though this model proves to be very useful, it brings mathematical complexities preventing one to obtain exact results even for 1D systems. For this reason, the main purpose of this work is to introduce a simplified version of the Heisenberg model which is exactly solvable for the diamond chain. We will investigate here the diamond chain consisting of Ising- (gray circles) and Heisenberg-type atoms (black circles), as it is depicted in Fig. 1. In order to obtain exact expression for the partition function of the model, we express the total Hamiltonian $\hat{\mathcal{H}}$ as a sum of the bond Hamiltonians, i.e. $\hat{\mathcal{H}} = \sum_{k=1}^N \hat{\mathcal{H}}_k$, where N denotes the total number of the Ising-type atoms and the bond Hamiltonian $\hat{\mathcal{H}}_k$ is given by (see Fig. 1.)

$$\begin{aligned} \hat{\mathcal{H}}_k &= J_1(\hat{S}_{k1}^z + \hat{S}_{k2}^z)(\hat{\mu}_{k1}^z + \hat{\mu}_{k2}^z) \\ &+ J_2[\Delta(\hat{S}_{k1}^x \hat{S}_{k2}^x + \hat{S}_{k1}^y \hat{S}_{k2}^y) + \hat{S}_{k1}^z \hat{S}_{k2}^z] \\ &- H_A(\hat{\mu}_{k1}^z + \hat{\mu}_{k2}^z)/2 - H_B(\hat{S}_{k1}^z + \hat{S}_{k2}^z) \end{aligned} \quad (1)$$

In this equation, $\hat{S}_{k\alpha}^\gamma$ and $\hat{\mu}_{k\alpha}^z$ ($\gamma = x, y, z$; $\alpha = 1, 2$) represent relevant components of the standard spin-1 and spin-1/2 operators, respectively, the exchange interaction J_1 couples the nearest-neighbouring Ising and Heisenberg atoms, J_2 couples nearest-neighbouring Heisenberg pairs and Δ represents the XXZ anisotropy parameter. Finally, the terms including H_A and H_B describe interactions of relevant atoms with an external magnetic field. Next, introducing a generalized decoration-iteration transformation [2, 3] and utilizing the commutability of the bond Hamiltonians, we easily find the following relation

$$\mathcal{Z}(\beta, J_1, J_2, \Delta, H_A, H_B) = A^N \mathcal{Z}_0(\beta, R, H_0), \quad (2)$$

where \mathcal{Z} represents the partition function of our model and \mathcal{Z}_0 is the well-known partition function of a simple spin-1/2 Ising linear chain and $\beta = 1/(k_B T)$. For unknown transformation parameters A , R and H_0 , we obtain relatively simple expressions [2, 3]:

$$A^4 = (W_0 + W_1)(W_0 + W_2)(W_0 + W_3)^2 \quad (3)$$

$$\beta R = \ln \frac{(W_0 + W_1)(W_0 + W_2)}{(W_0 + W_3)^2} \quad (4)$$

$$\beta H_0 = \beta H_A - \ln(W_0 + W_1) + \ln(W_0 + W_2) \quad (5)$$

$$W_0 = \exp(t_2) + f(t_2/2, a, 0, 0) - 4 \quad (6)$$

$$W_1 = f(-t_2, 2t_1 + 2h_B, t_1 + h_B, t_2\Delta) \quad (7)$$

$$W_2 = f(-t_2, 2t_1 - 2h_B, t_1 - h_B, t_2\Delta) \quad (8)$$

$$W_3 = f(-t_2, 2h_B, h_B, t_2\Delta) \quad (9)$$

with $a = \sqrt{1 + 8\Delta^2}$, $t_1 = \beta J_1$, $t_2 = \beta J_2$, $h_B = \beta H_B$, and $f(x_1, x_2, x_3, x_4) = 2e^{x_1} \cosh x_2 + 4 \cosh x_3 \cosh x_4$. Eqs. (2)-(9) enable one to calculate all physical quantities of interest.

In order to illustrate some interesting properties of the system, we have depicted the ground-state phase diagram in the $J_2 - H$ space for the case of the antiferromagnetic isotropic model i.e., $J_1 > 0$, $J_2 > 0$, $H_A = H_B = H$ and $\Delta = 1$. To describe different phases, we have calculated and analysed the reduced magnetization of the Ising (m_i^z) and Heisenberg sublattice (m_h^z) and various nearest-neighbour correlation functions (C_{hh}^{xx} , C_{hh}^{zz} , C_{ih}^{zz} and C_{ii}^{zz}). All these quantities are defined in the standard way, for instance, $m_i^z \equiv \langle \hat{\mu}_{k\alpha}^z \rangle$, $m_h^z \equiv \langle \hat{S}_{k\alpha}^z \rangle$, $C_{hh}^{xx} \equiv \langle \hat{S}_{k_1}^x \hat{S}_{k_2}^x \rangle$, where $\langle \dots \rangle$ means standard canonical average, subscripts $i(h)$ refer to the Ising (Heisenberg) atoms and superscripts represent spatial components of the spin. As one can see from Fig.2, there exist six different phases separated by first-order phase transition boundaries. With the help of above introduced quantities we can characterize these phases as follows:

Ordered phase 1 (OP₁):

$$m_i^z = -1/2, m_h^z = 1, C_{hh}^{zz} = 1, C_{hh}^{xx} = 0, C_{ih}^{zz} = -1/2, C_{ii}^{zz} = 1/4;$$

Ordered phase 2 (OP₂):

$$m_i^z = -1/2, m_h^z = 1/2, C_{hh}^{zz} = 0, C_{hh}^{xx} = -1/2, C_{ih}^{zz} = -1/4, C_{ii}^{zz} = 1/4;$$

Ordered phase 3 (OP₃):

$$m_i^z = 1/2, m_h^z = 0, C_{hh}^{zz} = -2/3, C_{hh}^{xx} = -2/3, C_{ih}^{zz} = 0, C_{ii}^{zz} = 1/4;$$

Ordered phase 4 (OP₄):

$$m_i^z = 1/2, m_h^z = 1/2, C_{hh}^{zz} = 0, C_{hh}^{xx} = -1/2, C_{ih}^{zz} = 1/4, C_{ii}^{zz} = 1/4;$$

Saturated paramagnetic phase (SP):

$$m_i^z = 1/2, m_h^z = 1, C_{hh}^{zz} = 1, C_{hh}^{xx} = 0, C_{ih}^{zz} = 1/4, C_{ii}^{zz} = 1/4;$$

Frustrated phase (FP):

$$m_i^z = 0, m_h^z = 0, C_{hh}^{zz} = -2/3, C_{hh}^{xx} = -2/3, C_{ih}^{zz} = 0, C_{ii}^{zz} = 0.$$

As one can see the OP₁ and SP represent standard phases that are usually observed also in the pure Ising models. On the other hand, the remaining four phases exhibit apparently a nontrivial spin ordering related to strong quantum fluctuations in the system. Due to this effect magnetic properties of the OP₂

and OP_4 can be explained within the valence-bond-solid picture as suggested in [4]. Moreover, the stepwise magnetization curves exhibit plateaux at relevant ordered phases OP_1 , OP_2 , OP_3 or OP_4 . In addition to the above mentioned phases, there appears for $H/J = 0$ and $J_2 > J_1$ an interesting frustrated phase (FP) having the nonzero entropy, namely, $S/Nk_B = \ln 2$. Even without presenting further details here, it is clear that the system under consideration is very interesting and it will exhibit nontrivial properties also at finite temperatures.

References

- [1] K. Okamoto and Y. Ichikawa, J. Phys. Chem. Solids 63 (2002) 1575.
- [2] M. E. Fisher, Phys. Rev. 113 (1959) 969.
- [3] J. Strečka and M. Jaščur, Phys. Rev. B 66 (2002) 174715.
- [4] J. Strečka and M. Jaščur, Phys. Stat. Sol. (b) 233 (2002) R12.

Acknowledgment: This work was supported under grants APVT 20-009902 and VEGA 1/9034/02

FIGURE CAPTIONS:

Fig.1: Part of the of the mixed-spin diamond chain. Gray circles represent spin-1/2 Ising atoms and black ones spin-1 Heisenberg atoms. The ellipse demarcates k th bond (see text)

Fig.2: Ground-state phase diagram of the Ising-Heisenberg diamond chain in the $J_2 - H$ space. OP_1 , OP_2 , OP_3 , OP_4 , SP and FP denote different phases described in the text



