## No dinosaurs in microcanonical gravitation: No special "microcanonical phase transitions"

D.H.E. Gross

<sup>1</sup> Hahn-Meitner-Institut, Bereich Theoretische Physik, Glienickerstr. 100, 14109 Berlin (Germany)

It is shown that the recently introduced singularities of the microcanonical entropy like "microcanonical phase transitions", and exotic pattern of the microcanonical caloric curve T(E) like multi-valuednes or the appearance of "dinosaur's necks" are inconsistent with Boltzmann's fundamental definition of entropy  $S = \ln[W(E)]$  for a system at equilibrium even for extremely large systems as astro-physical ones.

In a recent paper Bouchet and Barré[1] introduced a special class of "microcanonical" phase transitions. These are defined by singularities of the microcanonical entropy or the microcanonical temperature. However, such singularities do not exist in the microcanonical thermodynamics, i.e. of a system at equilibrium, based on Boltzmann's principle [2] neither for small systems nor for the real large self-gravitating astrophysical ones.

The entropy as defined by Boltzmann's principle

$$S(E) = \ln[W(E)] \tag{1}$$

as also the temperature

$$\beta(E) = T^{-1}(E) = \frac{dS}{dE} \tag{2}$$

are single valued, smooth, and multiple differentiable for any finite N. Here W(E) is the (6N-1) folded integral which measures the geometric area of the microcanonical many-fold of points with the energy E in the 6N-dim.

phase space. T(E) is a single valued differentiable function at all energies as long as  $\beta>0$ , i.e. E is finite. This clearly excludes all microcanonical "hysteretic cyclings", -zones, or "dinosaur's necks" as introduced by [3].

Chavanis argues in private communication that the "hysteretic cycles" reflect possible meta-stable configurations which are important in astro-physics. This might well be so but is outside of any controllable and fundamental conception of statistics. The microcanonical ensemble  $\mathcal{E}$  is per definition the ensemble of all possible configuration in the 6N-dim phasespace with the same energy. Its geometrical size is  $W = e^{S(E)}$ . Of course it is interesting and important to consider only those configurations  $\mathcal{M} \subseteq \mathcal{E}$  which represent e.g. stars at densities up to the ignition of their hydrogen burning, when further gravitational collapse is halted for a while. This can be done by adding an upper limit to the local density as done e.g by ref. [4, 5, 6].  $\mathcal{M}(E)$  is then still a compact manifold and similarly single valued multiple differentiable in E. Again there are no singularities and no "dinosaurs".

<sup>[1]</sup> F. Bouchet and J. Barré. Classification of phase transitions and ensemble inequivalence, in systems with long range interactions. *preprint*, pages cond–mat/0303307, 2003.

<sup>[2]</sup> D.H.E. Gross. Microcanonical thermodynamics: Phase transitions in "Small" systems, volume 66 of Lecture Notes in Physics. World Scientific, Singapore, 2001.

<sup>[3]</sup> P.H. Chavanis. Phase transitions in self-gravitating systems: Self-gravitating fermions and hard-sphere models. *Phys. Rev. E*, 65:056123–1–21, 2002.

<sup>[4]</sup> E.V. Votyakov, H.I. Hidmi, A. De Martino, and D.H.E. Gross. Microcanonical mean-field thermodynamics of

self-gravitating and rotating systems. *Phys.Rev.Lett.*, 89:031101–1–4; http://arXiv.org/abs/cond-mat/0202140, (2002).

<sup>[5]</sup> E.V. Votyakov, A. De Martino, and D.H.E. Gross. Asymmetric binaries and hierarchy of states in self-gravitating systems. *Eur.Phys.Lett*, 62:14–20; http://arXiv.org/abs/cond-mat/0210707, 2003).

<sup>[6]</sup> D.H.E. Gross. A new thermodynamics, from nuclei to stars ii. pages cond—mat/0306499, 2003).