## Effect of magnetic field in hybrid nanostructures

N. Stefanakis

Universität Tübingen, Tübingen Germany (October 28, 2018)

We examine the effect of the magnetic field on the proximity effect in nanostructures self consistently using the Bogoliubov-deGennes formalism within the two dimensional extended Hubbard model. We calculate the local density of states, and the pair amplitude. We study several nanostructures: superconductor - two dimensional electron gas, superconductor - ferromagnet. In these structures the magnetic field can be considered as a modulation parameter for the proximity effect.

### I. INTRODUCTION

When a magnetic field is applied in a two dimensional electron gas (2DEG) leads to a fractal energy spectrum known as the Hofstadter's butterfly, where small changes in magnetic field change completely the spectrum [1].

Recently conductance measurements have been performed on a superconductor 2DEG structure where an external magnetic field is applied perpendicular to the 2DEG [2]. In this case the magnetic field is sufficient large to induce Landau quantization in 2DEG, however is smaller than the upper critical field of the superconductor. More recently experiments have established the connection between the Hofstadter's spectrum observed in the 2DEG in the presence of a magnetic field and the quantum Hall effect [3].

On the other hand phase sensitive experiments can be used to probe the anisotropy of the pair amplitude in high temperature superconductors [4,5]. These experiments report the existence of a zero bias conduction peak (ZBCP) [6] which originates from the zero energy states (ZES) formed near the [110] surfaces of d-wave superconductors [7,8].

Recently the proximity effect has been probed as decaying oscillations of the density of states in s-wave superconductor ferromagnet hybrid structures [9] and a phase shift of half flux quantum in the diffraction pattern of a ferromagnetic  $0-\pi$  SQUID [10]. Similar effects have been observed in d-wave [11,12] superconductor ferromagnet hybrid structures. Theoretical explanation has been given in the framework of the quasiclassical theory for s-wave [13] and d-wave case [14]. In these structures the exchange field modulates the period of the pair amplitude oscillations. Moreover much interest has been focused recently on the manipulation of entangled states which are formed by extracting Cooper pairs from the superconductor. For example a beam splitter has been proposed [15] and also several experiments that involve ferromagnetic electrodes connected to superconductors [16–18]. These structures have acquired considerable interest the last years due to the possibility to use the  $\pi$  states in solid state qubit implementation.

In this paper our goal is to explore several new aspects related to the control of the proximity effect in nanostructures. We study superconductor - two dimensional electron gas, superconductor - ferromagnet. The basic quantities which we calculate are the local density of states and the pair amplitude as a function of several relevant parameters: the distance from the surface, the magnetic field, the barrier strength, and the symmetry of the pair potential. The method is based on exact diagonalizations of the Bogoliubov-de Gennes equations associated to the mean field solution of an extended Hubbard model.

Our principal result is that for superconductor 2DEG interface the LDOS shows a composite picture of energy bands due to Landau quantization and gaps due to the presence of superconductivity. In superconductor ferromagnet hybrid structures the magnetic field can modulate the period of the pair amplitude oscillations inside the ferromagnetic layer. Our predictions from the simulations of this model are of interest in view of future STM spectroscopy experiments on nanostructures.

The article is organized as follows. In Sec. II we develop the model and discuss the formalism. In Sec. III we discuss the effect of the magnetic field in pure 2DEG, superconductors. In Sec. IV we discuss the effect of the magnetic field and the effect of the strength of the barrier, in superconductor 2DEG heterostructures. In Sec. V we discuss the effect of the magnetic field in a superconductor ferromagnet hybrid structure. Finally summary and discussions are presented in the last section.

# II. BDG EQUATIONS, FOR THE SUPERCONDUCTOR-SEMICONDUCTOR JUNCTION WITHIN THE HUBBARD MODEL

The Hamiltonian of the extended Hubbard model on a two dimensional square lattice takes the form

$$H = -\sum_{\langle i,j \rangle} t_{ij} c_{i\sigma}^{\dagger} c_{j\sigma} + \mu \sum_{i\sigma} n_{i\sigma} + \sum_{i\sigma} \mu_i^I n_{i\sigma} + V_0 \sum_i n_{i\uparrow} n_{i\downarrow} + \frac{V_1}{2} \sum_{\langle ij \rangle \sigma\sigma'} n_{i\sigma} n_{j\sigma'} , \qquad (1)$$

where i, j are sites indices and the angle brackets indicate that the hopping is only to nearest neighbors. In the presence of a magnetic field the hopping integral  $t_{ij}$  is modified by the Peierls substitution, which account for the coupling of electrons to the magnetic field

$$t_{ij}(A) = t_{ij} \exp(\frac{ie}{\hbar c} \int_{r_j}^{r_i} Adl).$$
<sup>(2)</sup>

A is the vector potential which is chosen in a Landau gauge.  $n_{i\sigma} = c^{\dagger}_{i\sigma}c_{i\sigma}$  is the electron number operator at site  $i, \mu$  is the chemical potential.  $V_0, V_1$  are on site and nearest-neighbor interaction strength. Within the mean field approximation Eq. (1) reduces to the Bogoliubov deGennes equations [19]:

$$\begin{pmatrix} \hat{\xi} & \hat{\Delta} \\ \hat{\Delta}^* & -\hat{\xi} \end{pmatrix} \begin{pmatrix} u_n(r_i) \\ v_n(r_i) \end{pmatrix} = \epsilon_n \begin{pmatrix} u_n(r_i) \\ v_n(r_i) \end{pmatrix}, \tag{3}$$

such that

$$\hat{\xi}u_n(r_i) = -t\sum_{\hat{\delta}} u_n(r_i + \hat{\delta}) + (\mu^I(r_i) + \mu)u_n(r_i),$$
(4)

$$\hat{\Delta}u_n(r_i) = \Delta_0(r_i)u_n(r_i) + \sum_{\hat{\delta}} \Delta_\delta(r_i)u_n(r_i + \hat{\delta}),$$
(5)

where the gap functions are defined by

$$\Delta_0(r_i) \equiv V_0 < c_\uparrow(r_i)c_\downarrow(r_i) >, \tag{6}$$

$$\Delta_{\delta}(r_i) \equiv V_1 < c_{\uparrow}(r_i + \hat{\delta})c_{\downarrow}(r_i) >, \tag{7}$$

and where  $\hat{\delta} = \hat{x}, -\hat{x}, \hat{y}, -\hat{y}$ . Equation (3) is subject to the self consistency requirements

$$\Delta_0(r_i) = \frac{V_0(r_i)}{2} F_0(r_i) = \frac{V_0(r_i)}{2} \sum_n u_n(r_i) v_n^*(r_i) \tanh\left(\frac{\beta\epsilon_n}{2}\right),$$
(8)

$$\Delta_{\delta}(r_i) = \frac{V_1(r_i + \hat{\delta})}{2} F_{\delta}(r_i) = \frac{V_1(r_i + \hat{\delta})}{2} \sum_n (u_n(r_i)v_n^*(r_i + \hat{\delta}) + u_n(r_i + \hat{\delta})v_n^*(r_i)) \tanh\left(\frac{\beta\epsilon_{n\gamma_2}}{2}\right)).$$
(9)

We solve the above equations self-consistently. The numerical procedure has been described elsewhere [7,18]. We then compute the *d*-wave and the extended *s*-wave gap functions given by the expressions:

$$\Delta_d(r_i) = \frac{1}{4} [\Delta_{\hat{x}}(r_i) + \Delta_{-\hat{x}}(r_i) - \Delta_{\hat{y}}(r_i) - \Delta_{-\hat{y}}(r_i)],$$
(10)

$$\Delta_s^{ext}(r_i) = \frac{1}{4} [\Delta_{\hat{x}}(r_i) + \Delta_{-\hat{x}}(r_i) + \Delta_{\hat{y}}(r_i) + \Delta_{-\hat{y}}(r_i)].$$
(11)

The pair amplitude for the s-wave case is  $F_0(r_i)$ . The pair amplitude for the d-wave case is given by the expression

$$F_d(r_i) = \frac{1}{4} [F_{\hat{x}}(r_i) + F_{-\hat{x}}(r_i) - F_{\hat{y}}(r_i) - F_{-\hat{y}}(r_i)].$$
(12)

The local density of states (LDOS) at the *i*th site is given by

$$\rho_i(E) = -2\sum_n \left[ |u_n(r_i)|^2 f'(E - \epsilon_n) + |v_n(r_i)|^2 f'(E + \epsilon_n) \right],\tag{13}$$

f' is the derivative of the Fermi function,

$$f(\epsilon) = \frac{1}{\exp(\epsilon/k_B T) + 1}.$$
(14)

#### **III. EFFECT OF THE MAGNETIC FIELD**

We start our investigation of heterostructures by discussing the effect of the magnetic field on the local density of states for a 2DEG. We consider a two dimensional system of  $20 \times 20$  sites, and we suppose fixed boundary conditions by setting the impurity potential  $\mu^{I} = 100t$  at the surface. The temperature is  $k_{B}T = 0.1t$ .

The LDOS as a function of energy, for a two dimensional electron gas, for different values of the magnetic field is shown in Fig. 1 (we show only the E > 0 part of the spectrum due to symmetry). We see that when the magnetic flux through a plaquette is a rational number p/q the energy spectrum has q sub-bands. At each half filled sub-band there is a logarithmic singularity. The spectrum obtained corresponds to the Hofstadter's butterfly.

We also study the influence of the magnetic field in a pure superconducting state. We assume that the penetration depth is very large so that the magnetic field penetrates inside the superconductor. For the calculation we have chosen a Landau gauge for the vector potential, which neglects the effects of diamagnetic screening supercurrents. In a more realistic calculation the effect of the magnetic field should be calculated self-consistently. The LDOS for an *s*-wave superconductor in the presence of magnetic field shows additional gap at energy equal to zero (see Fig. 2(a)). However the above gap LDOS is not modified. For the case of *d*-wave superconductor the energy spectrum changes for the region inside the gap and also the overall shape of the Hofstadter's spectrum is modified qualitatively compared to the *s*-wave superconductor (see Fig. 2(b)).

#### IV. HOFSTADTER'S SPECTRUM IN A SUPERCONDUCTOR 2DEG HETEROSTRUCTURE

We now discuss the modification of the Hofstadter's spectrum in a superconductor - 2DEG heterostructure shown in Fig. 3 due to the magnetic field. We note that the magnetic field is applied only in the 2DEG. When the magnetic field is absent, the pair potential decays exponentially inside the 2DEG both for *s*-wave and *d*-wave superconductor (see Fig. 4). As we increase the exchange field the pair amplitude is not really modified inside the 2DEG. However inside the superconductor the pair amplitude increases for *s*-wave and does not change for *d*-wave.

The proximity effect is even more obvious in the LDOS (see Fig. 5). We see that for magnetic field equal to f = 1/2, the site inside the superconductor (-2) shows a gap structure as a signature of superconductivity and also a peak at E = 2 which is due to the effect of the magnetic field. As we move to the interior of the 2DEG the gap disappears while the peak the at E = 2 becomes more pronounced. The effect is similar to the *s*-wave and the *d*-wave case. So we may conclude that the Hofstadter's spectrum can also be observed even in the absence of external magnetic field, but only due to the proximity effect with a superconductor, and also that the same spectrum is strongly modified in the presence of superconducting correlations.

In general the increase of the inter-facial barrier potential suppresses the proximity effect because the leakage of Cooper pairs from the superconductor to the 2DEG is reduced if the tunnel amplitude is reduced. In this case the magnetic field is screened by the barrier. As a consequence for both s-wave and d-wave the pair amplitude is reduced on the 2DEG side by the increase of the barrier strength as seen in Fig. 6. In the s-wave case (see Fig. 6(a)) the pair amplitude on the superconducting side increases by the increase of the barrier strength since the barrier decouples the superconductor from the region where the magnetic field is applied, and the influence of the magnetic field in the superconductor is of reduced strength. In the d-wave case on the other hand the pair amplitude close to the interface does not change much, since the interface becomes pair breaking.

#### V. EFFECT OF THE MAGNETIC FIELD IN SUPERCONDUCTOR FERROMAGNET HYBRID STRUCTURES

We demonstrate in this section that the magnetic field can act as a control parameter for the proximity effect in hybrid structures. We study first the effect of the magnetic field in a pure ferromagnetic state. Due to the presence of the exchange field in the ferromagnet the energy bands in the LDOS that are formed due to the presence of the magnetic field are spitted (see Fig. 7).

In hybrid structures in the absence of magnetic field, there exists oscillations of the pair amplitude in the ferromagnetic layer. The magnetic field changes the period of the oscillations of the pair amplitude as seen in Fig. 8. This happens because the magnetic field favors single spin orientation and therefore modifies the exchange field of the ferromagnet to a different effective exchange field. When the magnetic field increases the effective exchange field increases and the period of oscillations of the pair amplitude decreases as seen in Fig. 8.

The proximity effect in hybrid structures appears also as oscillations in the local density of states with the distance from the interface. The external magnetic field modifies the period of these oscillations. We can see the effect of the magnetic field in the LDOS in Fig. 9. We see that for f = 0 the pic in the LDOS which is a signature of the  $\pi$  state appears after the x = 3 site from the interface, while for f = 1/2 it appear for smaller distance, because the effective period of the oscillations has been changed due to the presence of the magnetic field.

#### **VI. CONCLUSIONS**

We calculated the LDOS and the pair amplitude for several superconductor - 2DEG heterostructures, in the presence of an external magnetic field within the extended Hubbard model, self consistently. In the calculated quantities, like the pair amplitude and the LDOS, for the sites that are close to the interface, the gap which is a signature of superconductivity coexists with the bands which are formed due to the rational values of the magnetic field. We also demonstrated that the Hofstadter's spectrum is strongly modified due to the proximity effect with the superconductor. In superconductor ferromagnet structures the magnetic field can also be used as an external control parameter in order to modify the proximity effect.

- [1] D. Hofstadter, Phys. Rev. B 14, 2239 (1976).
- [2] T.D. Moore and D.A. Williams, Phys. Rev. B 59, 7308 (1999).
- [3] C. Albrecht, J.H. Smet, K. von Klitzing, D. Weiss, and H. Schwizer, Phys. Rev. Lett. 86, 147 (2001).
- [4] H. Hilgenkamp and J. Mannhart, Rev. Mod. Phys. 74, 485 (2002).
- [5] S. Kashiwaya, Y. Tanaka, Rep. Prog. Phys. 63, 1641 (2000).
- [6] H. Guillou, J. Chaussy, M. Charalambous, and M. Pissas, Physica C, 382 (2-3), 291 (2002).
- [7] N. Stefanakis, Phys. Rev. B 66, 024514 (2002).
- [8] J.-X. Zhu, B. Friedman, and C.S. Ting, Phys. Rev. B 59, 3353 (1999).
- [9] T. Kontos M. Aprili, J. Lesueur, and X. Grison, Phys. Rev. Lett. 86, 304 (2001).
- [10] W. Guichard, M. Aprili, O. Bourgeois, T. Kontos, J. Lesueur, and P. Gandit, Phys. Rev. Lett. 90, 167001 (2003).
- [11] M. Freamat and K.-W. Ng, cond-mat/0301081.
- [12] M. Freamat and K.-W. Ng, cond-mat/0305446.
- [13] M. Zareyan, W. Belzig, and Yu.V. Nazarov, Phys. Rev. Lett. 86, 308 (2001).
- [14] Z. Faraii and M. Zareyan, cond-mat/0304336.
- [15] G.B. Lesovik, T. Martin, and G. Blatter, Eur. Phys. J. B 24, 287 (2001).
- [16] R. Mélin and D. Feinberg, Eur. Phys. J. B 26, 101 (2002).
- [17] H. Jirari, R. Mélin and N. Stefanakis, Eur. Phys. J. B 31, 125 (2003).
- [18] N. Stefanakis and R. Mélin, J. Phys.: Condens. Matter 15, 3401 (2003).
- [19] P.G. de Gennes, Superconductivity of Metals and Alloys (Benjamin, New York, 1966).



FIG. 1. The LDOS as a function of energy, for a two dimensional electron gas, for different values of the magnetic field f = 0, 1/4, 1/3, 1/2.



FIG. 2. (a) The LDOS as a function of energy, for an s-wave superconductor, for different values of the magnetic field f = 0, 1/4, 1/3, 1/2. (b) The same for d-wave.



FIG. 3. The superconductor - 2DEG interface. Solid circles indicate interface sites. The LDOS is presented for sites along the dashed line. The 2DEG extends to the right of the interface.



FIG. 4. (a) The pairing amplitude as a function of position, for a s-wave superconductor - 2DEG interface, for different values of the magnetic field f = 0, 1/4, 1/3, 1/2. (b) The same but for a d-wave superconductor.



FIG. 5. (a) The LDOS as a function of energy, for an s-wave superconductor - 2DEG interface, for different values of distance from the interface x = -2, -1, 0, 1, for magnetic field equal to f = 1/2. (b) The same but for a d-wave superconductor.



FIG. 6. (a) The pair amplitude as a function of distance from the interface, for a s-wave superconductor - 2DEG interface, for different values of barrier strength 0, 5, 10, 15, for magnetic field equal to f = 1/2. (b) The same but for a d-wave superconductor.



FIG. 7. The LDOS as a function of energy, for a two dimensional electron gas, for different values of the magnetic field f = 0, 1/4, 1/3, 1/2, for exchange field equal to h = 0.5.



FIG. 8. The pairing amplitude as a function of position, for a superconductor - two dimensional electron gas interface, for different values of the magnetic field f = 0, 1/3, 1/2, for exchange field equal to h = 0.5.



FIG. 9. (a) The LDOS as a function of energy, for a superconductor - two dimensional electron gas interface, for different values of the distance from the interface x = 0, 1, 2, 3 for exchange field equal to h = 0.5, and magnetic field f = 0. (b) The same as in (a) but for f = 1/2.