Coulomb drag in longitudinal magnetic field in quantum wells

V. L. Gurevich and M. I. Muradov

Solid State Physics Division, A.F.Ioffe Institute, 194021 Saint Petersburg, Russia

(November 15, 2018)

The influence of a longitudinal magnetic field on the Coulomb drag current created in the ballistic transport regime in a quantum well by a ballistic current in a nearby parallel quantum well is investigated. We consider the case where the magnetic field is so strong that the Larmour radius is smaller than the width of the well. Both in Ohmic and non-Ohmic case, sharp oscillations of the drag current as a function of the gate voltage or chemical potential are predicted. We also study dependence of the drag current on the voltage V across the driving wire, as well as on the magnetic field B.

Studying the Coulomb drag one can make conclusions about the electron spectrum and and electron-electron interaction in quantum wells.

I. INTRODUCTION

The influence of a magnetic field on the Coulomb drag are investigated in different geometries. The Coulomb drag between two two-dimensional (2D) quantum wells in a strong magnetic field perpendicular to the planes of the wells and in the presence of disorder has been investigated in Ref. [1]. In magnetic field perpendicular to the planes the Hall voltage can be induced in the drag quantum well in the direction perpendicular to both direction of the magnetic field and of the current in the drive well [2], [3]. These two geometries can be called transverse.

The purpose of the present paper is to study the influence of an in-well magnetic field B on the Coulomb drag current in the course of ballistic (collisionless) electron transport in a quantum well due to a ballistic drive current in a parallel quantum well. In other words, we consider the longitudinal geometry, i.e. the case where the magnetic field is parallel to the applied electric field E and to the plane of the well itself.

We will concern ourselves with the case of a strong magnetic field that makes the motion of the carriers along the field one dimensional and alters the density of electron states. Moreover, we restrict ourselves with the quantum limit when only the ground Landau oscillator states are occupied by electrons in the two quantum wells, so that

$$
\hbar\omega_B \gtrsim \mu \tag{1.1}
$$

Here ω_B is the cyclotron frequency while μ is the chemical potential. A theory of electronic transport through three-dimensional ballistic microwires in longitudinal magnetic fields at low temperatures has been developed in Ref. [4]. Our geometry is similar to that considered in Ref. [4]. However, in the present paper we consider much simpler situation of a very strong magnetic field satisfying Eq. (1.1). Later on we hope to return to a more general case of a weaker magnetic field where several Landau levels may be involved.

The magnetic field making the motion of the electrons in the transverse direction one dimensional maps the problem under consideration onto the Coulomb drag problem in two one-dimensional wires already considered by the authors in Ref. [5] in the Fermi liquid approach. Therefore, our final formulae for the Coulomb drag current appear to be similar to those obtained in [5]. Physically the magnetic field may play the following important role. It will suppress the tunneling of electrons between the quantum wells that, if present, would impede observation of the Coulomb drag.

The magnetic field may change the electron quasimomentum relaxation time. Scattering of electrons by ionized impurities in sufficiently strong magnetic fields may be even weaker than for $B = 0$. As for the relaxation due to the phonon scattering, a strong magnetic field can alter the density of electron states and in the quantum limit the relaxation rate may be bigger than for $B = 0$. We, however, will assume the temperature to be so low that the transport remains ballistic even in the presence of magnetic field.

We consider the case where the magnetic length

$$
a_B = \sqrt{\frac{\hbar c}{|e|B}}\tag{1.2}
$$

is much smaller than the distance W between the quantum wells of the width $L_x \sim W$ each

$$
W/a_B \gg 1. \tag{1.3}
$$

This inequality establishes lower bound for the values of the magnetic field for a given distance between the quantum wells. For instance, for $W \sim 80$ nm the inequality requires magnetic fields of the order of $B \sim 1$ T, or bigger.

It is convenient to break our calculations into several parts. In the first part we will give the principal equations of our theory based on the Boltzmann treatment of the transport. We will consider a linear response in Sec. III. Next we will discuss a non-Ohmic case in Sec. IV. Comparison of our results with the 1D Coulomb drag results in the longitudinal geometry and 2D Coulomb drag results for $B = 0$ will be given in Summary.

II. BOLTZMANN EQUATION

We consider two parallel quantum wells perpendicular to x axis. The eigenfunctions and eigenvalues for a one electron problem in a magnetic field along the z axis in the ith quantum well is [we use the gauge $\mathbf{A} = (0, Bx, 0)$

$$
\psi_{0p_yp_z} = \frac{1}{\sqrt{L_y L_z}} \varphi_0 \left(\frac{x - x_{p_y}}{a_B}\right) \exp(i p_y y/\hbar + i p_z z/\hbar),\tag{2.1}
$$

$$
\varepsilon_{0p_z} = U_i + \frac{\hbar\omega_B}{2} + \frac{p_z^2}{2m}.
$$
\n(2.2)

Here m is the effective electron mass, $\varphi_0 \left[(x - x_{p_y})/a_B \right]$ is the wave function of a harmonic oscillator in the ground state oscillating about the point $x_{p_y} = -a_B^2 p_y/\hbar = -p_y/m\omega_B$. The wave function $\psi_{0p_y p_z}$ describes a state for which the electron probability distribution is large only within the slab of the width $\approx a_B$ symmetrically situated about the plane $x = x_{p_y}$ and falls off exponentially outside the slab. As we consider the case $L_x \gg a_B$ we will assume the wave function to be equal $\psi_{0p_y p_z}$ if x_{p_y} is within the quantum well and zero otherwise. In what follows we will need the matrix elements of the functions $\exp(\pm i\mathbf{qr})$ between two stationary states. We have

$$
\langle 0p'_{y}p'_{z}|e^{\pm i\mathbf{qr}}|0p_{y}p_{z}\rangle = e^{\pm iq_{x}(x_{p_{y}}+x_{p'_{y}})/2}
$$

$$
\times e^{-a_{B}^{2}q_{x}^{2}/4}e^{-(x_{p_{y}}-x_{p'_{y}})^{2}/4a_{B}^{2}}\delta_{p'_{z},p_{z}\pm\hbar q_{z}}\delta_{p'_{y},p_{y}\pm\hbar q_{y}}.
$$
(2.3)

The diagram representing Coulomb drag effect is illustrated in Fig. 1. The external driving force enters

FIG. 1. Coulomb drag diagram. Here the labels 2, (1) stand for the drive (drag) quantum wells.

the diagram through nonequilibrium distribution function represented by the solid lines marked by the symbol 2 indicating that they represent the drive quantum well.

Now we embark on analysis of the conservation laws for the collisions of electrons belonging to two different quantum wells. We have

$$
\varepsilon_{0p_{z}}^{(1)} + \varepsilon_{0p'_{z}}^{(2)} = \varepsilon_{0p_{z} + \hbar q_{z}}^{(1)} + \varepsilon_{0p'_{z} - \hbar q_{z}}^{(2)}
$$
\n(2.4)

where $\varepsilon_{0p_z}^{(1,2)} = U_{1,2} + \hbar\omega_B/2 + p_z^2/2m$.

The solution of Eq. (2.4) is

$$
\hbar q_z = p'_z - p_z. \tag{2.5}
$$

The δ -function describing energy conservation can be recast into the form

$$
\delta(\varepsilon_{np_z}^{(1)} + \varepsilon_{lp'_z}^{(2)} - \varepsilon_{np_z + \hbar q_z}^{(1)} - \varepsilon_{lp'_z - \hbar q_z}^{(2)}) = \frac{m}{\hbar |q_z|} \delta[\hbar q_z - (p'_z - p_z)]. \tag{2.6}
$$

Therefore, the initial quasimomenta p_z and p'_z after the collision become $p_z + \hbar q_z = p'_z$ and $p'_z - \hbar q_z = p_z$, i.e. the electrons swap their quasimomenta as a result of collision.

Following [5,6] we assume that the drag current in the quantum well 1 is much smaller than the drive ballistic current in the quantum well 2 and calculate it by solving the Boltzmann equation for the quantum well 1. We have

$$
v_z \frac{\partial \Delta F_{0p_y}^{(1)}(p_z, z)}{\partial z} = -I^{(12)} \{ F^{(1)}, F^{(2)} \},\tag{2.7}
$$

where $F^{(1,2)}$ are the electron distribution functions in the quantum wells 1 and 2 respectively, and the collision integral $I^{(12)}\lbrace F^{(1)}, F^{(2)}\rbrace$ takes into account the interwell electron-electron scattering

$$
I^{(12)}\{F^{(1)}, F^{(2)}\} = \sum_{p'_z p'_y q'_x \mathbf{q}} W^{1p_z + \hbar q_z, 2p'_z - \hbar q_z}_{1p_z, 2p'_z} (q'_x, q_x, q_y, p_y, p'_y) \mathcal{S}. \tag{2.8}
$$

In this expression the sum over p'_y should be determined by the requirement that the x-center of the oscillator function is within the second quantum well. The requirement imposes the constraint

$$
\frac{\hbar}{a_B^2} \left(W + L_x - \frac{L_x}{2} \right) < p_y' < \frac{\hbar}{a_B^2} \left(W + L_x + \frac{L_x}{2} \right),\tag{2.9}
$$

and the product of distribution functions S is

$$
S = F_{0p_z}^{(1)} F_{0p'_z}^{(2)} \left(1 - F_{0p_z + \hbar q_z}^{(1)} \right) \left(1 - F_{0p'_z - \hbar q_z}^{(2)} \right) - F_{0p_z + \hbar q_z}^{(1)} F_{0p'_z - \hbar q_z}^{(2)} \left(1 - F_{0p_z}^{(1)} \right) \left(1 - F_{0p'_z}^{(2)} \right). \tag{2.10}
$$

$$
F_{0p_z}^{(1)} = \theta[v_z]f(\varepsilon_{0p_z} - \mu_B^{1L}) + \theta[-v_z]f(\varepsilon_{0p_z} - \mu_B^{1R}) + \Delta F_{0p_z}^{(1)}
$$
\n(2.11)

where

$$
\theta[v_z] = \begin{cases} 1 & \text{for } v_z > 0 \\ 0 & \text{for } v_z < 0 \end{cases}.
$$

Here we assume that the electrons move ballistically within the quantum well and the electrons moving from the left and right reservoirs have the chemical potentials $\mu_B^{1L} = \mu_B - eV_d/2$ and $\mu_B^{1R} = \mu_B + eV_d/2$ respectively. We introduce also the drag voltage V_d induced across the drag quantum well due to the quasimomentum transfer from the driving quantum well, i.e. we assume an open circuit for the drag quantum well.

The solution of $Eq.(2.7)$ is (here we omit the equilibrium part)

$$
\Delta F_{0p_z}^{(1)} = -\left(z \pm \frac{L}{2}\right) \frac{1}{v_z} I^{(12)} \{F^{(1)}, F^{(2)}\}, \quad \text{for} \quad \frac{p_z > 0}{p_z < 0.} \tag{2.12}
$$

Using the particle conserving property of the scattering integral

$$
\sum_{p_z p_y} I^{(12)} \{F^{(1)}, F^{(2)}\} = 0
$$
\n(2.13)

we get for the total current in the drag quantum well defined as

$$
J = \frac{e}{L_z} \sum_{p_y p_z} v_z F_{0p_z}^{(1)},\tag{2.14}
$$

the result

$$
J = -e \sum_{p_y, (p_z > 0)} I^{(12)} \{F^{(1)}, F^{(2)}\} + e \frac{1}{L_z} \sum_{p_y, (p_z > 0)} v_z [f(\varepsilon_{0p_z} - \mu_B^{1L}) - f(\varepsilon_{0p_z} - \mu_B^{1R})]. \tag{2.15}
$$

In these equations the sum over p_y is restricted by the requirement that the x-center of Landau oscillator must be within the quantum well, so that $-\hbar L_x/2a_B^2 < p_y < \hbar L_x/2a_B^2$. Introducing the density of states (including spin) per unit quasimomentum interval

$$
N(p_z)dp_z = 2\frac{L_z}{(2\pi\hbar)^2} \frac{\hbar L_x L_y}{a_B^2} dp_z
$$
\n(2.16)

we have

$$
J_{\text{Ohm}} = -e \frac{2eV_{\text{d}}}{(2\pi\hbar)^2} \frac{\hbar L_x L_y}{a_B^2} \int_{U_1 + \hbar \omega_B/2}^{\infty} d\varepsilon \left(-\frac{\partial f(\varepsilon - \mu_B)}{\partial \varepsilon} \right). \tag{2.17}
$$

For the degenerate electron gas this expression can be written as

$$
J_{\text{Ohm}} = -\frac{e^2}{\pi \hbar} \frac{L_x L_y}{2\pi a_B^2} V_{\text{d}}.\tag{2.18}
$$

Here the number of Larmour circles covering the cross section of the quantum well $L_xL_y/2\pi a_B^2$ appears instead of the number of open channels in the 1D situation.

We assume that only the ground Landau oscillator state is occupied, so that

$$
U_1 + \frac{1}{2}\hbar\omega_B < \mu = U_1 + \frac{3}{2}\hbar\omega_B. \tag{2.19}
$$

Taking into account Eq. (2.6) we obtain for the Coulomb scattering probability Eq. (2.8)

$$
W_{1p_z,2p'_z}^{1p_z + hq_z,2p'_z - hq_z} (q'_x, q_x, q_y, p_y, p'_y) = \frac{m}{\hbar|q_z|} \delta[\hbar q_z - (p'_z - p_z)] \frac{2\pi}{\hbar} U_q U_{q'_z q_y q_z}
$$

\n
$$
\times \langle 0p_z p_y | e^{-iq'_x x - iq_y y - iq_z z} | 0p_z + \hbar q_z p_y + \hbar q_y \rangle \langle 0p_z + \hbar q_z p_y + \hbar q_y | e^{iq_x x + iq_y y + iq_z z} | 0p_z p_y \rangle
$$

\n
$$
\times \langle 0p'_z p'_y | e^{iq'_x x + iq_y y + iq_z z} | 0p'_z - \hbar q_z p'_y - \hbar q_y \rangle \langle 0p'_z - \hbar q_z p'_y - \hbar q_y | e^{-iq_x x - iq_y y - iq_z z} | 0p'_z p'_y \rangle.
$$
 (2.20)

Here we use the unscreened Coulomb potential postponing discussion as to when this approximation can be justified until the last section.

To calculate the drag current we iterate the Boltzmann equation in the interwell collision term that we assume to be small. Therefore one can choose the distribution functions in the collision term to be equilibrium ones, e.g. $F_{0p}^{(1)} = f(\varepsilon_{0p}^{(1)} - \mu_B)$ for the first quantum well.

We assume, in the spirit of the approach developed by Landauer [7], Imry [8] and Büttiker [9] the drive quantum well to be connected to reservoirs which we call 'left' l and 'right' r . Each of them is in independent equilibrium described by the shifted chemical potentials $\mu_B^l = \mu_B - eV/2$ and $\mu_B^r =$ $\mu_B + eV/2$, where μ_B is the equilibrium chemical potential in the magnetic field. Therefore, the electrons entering quantum well from the 'left'('right') and having quasimomenta $p'_z > 0$ ($p'_z < 0$) are described by $F_{0p'_z}^{(2)} = f(\varepsilon_{0p'_z}^{(2)} - \mu_B^1)$ [$F_{0p'_z}^{(2)} = f(\varepsilon_{0p'_z}^{(2)} - \mu_B^r)$] and we see that the collision integral Eq.(2.8) is identically zero if the initial p'_z and final $p'_z - q$ quasimomenta in the drive quantum well are of the same sign. This means that only the *backscattering processes* contribute to the drag current.

Due to Eq.(2.6) we are left only with $p'_z < 0$ (since we are restricted according to Eq.(2.15) by the constraint $p'_z - \hbar q_z = p_z > 0$ and obtain in view of the δ -function in Eq.(2.20) the following product of distribution functions in the collision term

$$
\mathcal{P} = F_{0p_z}^{(1)} F_{0p'_z}^{(2)r} \left(1 - F_{0p'_z}^{(1)} \right) \left(1 - F_{0p_z}^{(2)l} \right) - F_{0p'_z}^{(1)} F_{0p_z}^{(2)l} \left(1 - F_{0p_z}^{(1)} \right) \left(1 - F_{0p'_z}^{(2)r} \right),\tag{2.21}
$$

or

$$
\mathcal{P} = f(\varepsilon_{0p_z}^{(1)} - \mu_B) f(\varepsilon_{0p'_z}^{(2)} - \mu_B^r) [1 - f(\varepsilon_{0p'_z}^{(1)} - \mu_B)][1 - f(\varepsilon_{0p_z}^{(2)} - \mu_B^l)] - f(\varepsilon_{0p'_z}^{(1)} - \mu_B) f(\varepsilon_{0p_z}^{(2)} - \mu_B^l) [1 - f(\varepsilon_{0p_z}^{(1)} - \mu_B)][1 - f(\varepsilon_{0p'_z}^{(2)} - \mu_B^r)].
$$
\n(2.22)

This equation will be analyzed in the following sections.

III. LINEAR RESPONSE

In this case $eV/T \ll 1$ (we assume the Boltzmann constant to be equal 1) and Eq.(2.22) can be recast into the form

$$
\mathcal{P} = \frac{eV}{T} f(\varepsilon_{0p_z}^{(1)} - \mu_B) f(\varepsilon_{0p'_z}^{(2)} - \mu_B) [1 - f(\varepsilon_{0p'_z}^{(1)} - \mu_B)] [1 - f(\varepsilon_{0p_z}^{(2)} - \mu_B)]. \tag{3.1}
$$

Shifting the integration variable $p'_y \to p'_y + \hbar (W + L_x)/a_B^2$ we have for the drag current

$$
J_{\text{drag}} = -e \frac{eV}{T} \frac{2\pi}{\hbar} \left(\frac{4\pi e^2}{\kappa}\right)^2 \frac{1}{2\pi\hbar} \int_0^\infty \frac{2L_z dp_z}{2\pi\hbar} \int_0^\infty \frac{dp'_z}{2\pi\hbar} \frac{m}{(p_z + p'_z)}
$$

$$
\times f(\varepsilon_{0p_z}^{(1)} - \mu_B) f(\varepsilon_{0p'_z}^{(2)} - \mu_B) [1 - f(\varepsilon_{0p'_z}^{(1)} - \mu_B)][1 - f(\varepsilon_{0p_z}^{(2)} - \mu_B)]
$$

$$
\times \int_{-\hbar}^{\hbar L_x/2a_B^2} \frac{2L_y dp_y}{2\pi\hbar} \frac{dp'_y}{2\pi\hbar} g_{00} [(p_z + p'_z)/\hbar, (p_y - p'_y)/\hbar]
$$
(3.2)

where

$$
g_{00}(k_z, k_y) = \int_{-\infty}^{\infty} \frac{dq_y}{2\pi} e^{-a_B^2 q_y^2} A^2(k_z, k_y, q_y), \qquad (3.3)
$$

$$
A(k_z, k_y, q_y) = \int_{-\infty}^{\infty} \frac{dq_x}{2\pi} \frac{e^{-ia_B^2 q_x [k_y - (W + L_x)/a_B^2 + q_y]} e^{-a_B^2 q_x^2/2}}{(k_z^2 + q_\perp^2)}
$$
(3.4)

 $q_{\perp}^2 = q_x^2 + q_y^2$. The last integral over the centers of the Larmour circles (since $x_p = -a_B^2 p_y/\hbar$) in Eq.(3.2) plays the role of an effective Coulomb interaction potential between the electrons freely moving along the direction of applied magnetic field.

$$
\frac{\hbar L_x / 2a_B^2}{\int_{-\hbar L_x / 2a_B^2} 2L_y dp_y dp'_y}{\int_{-\hbar L_x / 2a_B^2} 2\pi \hbar} \frac{2L_y dp_y dp'_y}{2\pi \hbar} g_{00} \left[\frac{p_z + p'_z}{\hbar}, \frac{p_y - p'_y}{\hbar} \right]
$$

$$
= \frac{2L_y}{(2\pi)^2} \int_0^{L_x / a_B^2} dk_y k_y \left\{ g_{00} \left[\frac{p_z + p'_z}{\hbar}, \frac{L_x}{a_B^2} - k_y \right] + g_{00} \left[\frac{p_z + p'_z}{\hbar}, k_y - \frac{L_x}{a_B^2} \right] \right\} \tag{3.5}
$$

We keep only the first term in this expression since the second term includes a faster oscillating exponent $\sim \exp(i q_x (W + L_x))$ as compared to the oscillating exponent in the first term $\sim \exp(i q_x W)$.

As $W/a_B \gg 1$ we can sufficiently simplify the expression for g_{00} . We obtain

$$
g_{00}(k_z, k_y) = e^{a_B^2 k_z^2} \int_{-\infty}^{\infty} \frac{dq_y}{2\pi} A^2(k_z, k_y, q_y).
$$
 (3.6)

$$
A(k_z, k_y, q_y) \simeq \int \frac{dq_x}{2\pi} \frac{e^{iq_x[(W+L_x) - a_B^2 k_y]}}{q_x^2 + q_y^2 + k_z^2} = \frac{e^{-|W+L_x - a_B^2 k_y|} \sqrt{q_y^2 + k_z^2}}{2\sqrt{q_y^2 + k_z^2}}
$$
(3.7)

Finally, the interaction term acquires the form

$$
\int_{-\hbar L_x/2a_B^2}^{\hbar L_x/2a_B^2} \frac{2L_y dp_y}{2\pi\hbar} \frac{dp'_y}{2\pi\hbar} g_{00} \left[k_z, (p_y - p'_y)/\hbar\right] = \frac{L_y}{4a_B(2\pi a_B k_z)^3} e^{a_B^2 k_z^2} \Phi(2Wk_z)
$$
(3.8)

where

$$
\Phi(\alpha) = \int_1^{\infty} d\xi \frac{e^{-\alpha \xi}}{\xi^3 \sqrt{\xi^2 - 1}},
$$
\n(3.9)

For $\alpha \gg 1$

$$
\Phi(\alpha) \simeq \sqrt{\frac{\pi}{2\alpha}} e^{-\alpha}.
$$
\n(3.10)

This result for the effective interaction (3.8) can be explained as follows: Larmour circles within the quantum wells of the width $L_x \cdot 1/(k_zL_x)$ near the surfaces contribute to the interaction. The number of interacting circles from two quantum wells is

$$
\left(\frac{L_y}{a_B}\frac{L_x \cdot 1/(k_z L_x)}{a_B}\right)^2.
$$

The sum over $q_x, q'_x \sim k_z, q_y \sim \sqrt{k_z/W}$ leads to a factor $(k_z L_x)^2 \cdot L_y \sqrt{k_z/W}$. The exponential decay of the drag with the distance between the quantum wells W is a consequence of one-dimensional character of the drag in the strong longitudinal magnetic field. Combining all these factors and multiplying the result by $U^2 \sim (4\pi e^2)^2 / (L_x L_y L_z)^2 k_z^4$ we arrive at Eq.(3.8).

The product of the distribution functions in Eq. (3.2) is a sharp function of p_z and p'_z at small temperatures, acquiring nonzero values only at $p_z, p'_z \sim p_F^B \pm T/v_F^B$. We assume that the quasimomentum interval T/v_F^B is much smaller than \hbar/W

$$
T \ll \frac{\hbar v_F^B}{W}.\tag{3.11}
$$

Here we wish to note that the Boltzmann treatment of transport phenomena requires that the uncertainty in longitudinal momentum must be smaller than the same momentum interval $\hbar/L_z \ll T/v_F^B$. These two requirements automatically lead to the inequality $W \ll L_z$. We assume that the last inequality holds.

According to our assumptions we can regard the interaction term in Eq. (3.2) as the slowly varying function and obtain

$$
J_{\text{drag}} = J_0 \frac{eV}{4\varepsilon_F^B} \frac{T}{\varepsilon_F^B} \left(\frac{U_{12}}{2T}\right)^2 \left[\sinh\left(\frac{U_{12}}{2T}\right)\right]^{-2} \tag{3.12}
$$

where

$$
J_0 = -\frac{e^5 m}{\kappa^2 (4\pi \hbar)^3} \frac{L_z L_y}{a_B^2} \frac{1}{(a_B k_F^B)^2} e^{(2a_B k_F^B)^2} \Phi\left(4W k_F^B\right)
$$
(3.13)

Here we introduced notations $U_{12} = U_1 - U_2$ and $mv_F^B = p_F^B = \sqrt{2m[\mu_B - U_1 - \hbar\omega_B/2]}$, $k_F^B = p_F^B/\hbar$. We assume that the electrons remain degenerate in the magnetic field

$$
\varepsilon_F^B \equiv \mu_B - U_1 - \frac{\hbar \omega_B}{2} \gg T. \tag{3.14}
$$

We consider the quantum limit, i.e. the case when all electrons belong to the first Landau level

$$
\varepsilon_F^B < \hbar\omega_B. \tag{3.15}
$$

Since the electron concentration N_B under this condition is related to the chemical potential by the equation

$$
N_B = \frac{m\hbar\omega_B p_F^B}{\pi^2 \hbar^3} \tag{3.16}
$$

Eq. (3.14) and Eq. (3.15) lead to

$$
T \ll \frac{(p_F^B)^2}{2m} < \hbar\omega_B, \ \ p_F^B = \frac{\pi^2 \hbar^3}{m} \frac{N_B}{\hbar\omega_B}.\tag{3.17}
$$

The first inequality in this relation is weaker than Eq.(3.11) if $\varepsilon_F \sim \hbar \omega_B$ and $W k_F \geq 1$. Introducing the electron concentration N and the chemical potential μ for $B = 0$ given by

$$
N = \frac{(2m\varepsilon_F)^{3/2}}{3\pi^2\hbar^3}, \quad \varepsilon_F = \mu - U_1 \tag{3.18}
$$

one can rewrite Eq.(3.17) as

$$
T \ll \frac{4}{9} \left(\frac{N_B}{N}\right)^2 \left(\frac{\varepsilon_F}{\hbar \omega_B}\right)^2 \varepsilon_F < \hbar \omega_B.
$$
\n(3.19)

Note that the second inequality in this expression does not depend on the electron mass and can require magnetic fields stronger than Eq.(1.3) [thus imposing a constraint on the electron concentration, or, if the latter is given the inequality may require stronger magnetic fields than is required by the Eq.(1.3)]. For instance, in a magnetic field of the order of $B \sim 10$ T the electron concentration N must be smaller than $2.7 \cdot 10^{17}$ cm⁻³.

Considering the case of the aligned quantum wells, so that $U_1 = U_2$ [otherwise the effect is exponentially small, cf. Eq.(3.12)] and putting $N = N_B$ we obtain

$$
J_{\rm drag} = J_0 \frac{eV}{4T} \left(\frac{T}{\varepsilon_F}\right)^2 \left(\frac{3\hbar\omega_B}{2\varepsilon_F}\right)^4,\tag{3.20}
$$

$$
J_0 = -\frac{e^5 m L_y L_z k_F^2}{9\kappa^2 (4\pi\hbar)^3} \left(\frac{3\hbar\omega_B}{2\varepsilon_F}\right)^4 e^{12(2\varepsilon_F/3\hbar\omega_B)^3} \Phi\left(4W k_F \frac{2\varepsilon_F}{3\hbar\omega_B}\right). \tag{3.21}
$$

The drag current is a rapidly increasing function of the applied magnetic field, as the latter increases the density of states and decreases the transferred Fermi momentum.

FIG. 2. Drag current versus dimensionless magnetic field $b = \hbar \omega_B/\varepsilon_F$ for two values of the interwell distances $W = 40$ nm (1) and $W = 50$ nm (2). Other parameters are given in the text.

To make an estimate of the current we put $m = 0.07m_e$, $\hbar\omega_B \sim \varepsilon_F = 14$ meV, $\kappa = 13$, $L_z \sim L_y = 1 \mu$ m, $W = 40$ nm.

$$
J_{\rm drag} \sim 10^{-11} \text{ A}
$$

In the linear response regime we can introduce a drag resistance, i.e. we can introduce the coefficient that depends only on the quantum wells parameters and relates the drive current J_{drive} in the quantum well 2 to the induced voltage in the drag quantum well $J_{\text{drive}}R_{\text{D}} = V_d$. Here the drive current in the quantum well 2 is

$$
J_{\text{drive}} = -V \frac{e^2}{2\pi\hbar} \frac{L_x L_y}{\pi a_B^2} \tag{3.22}
$$

(cf. Eq.(2.18)) and V_d is determined by the condition of zero total current $J = J_{\text{drag}} + J_{\text{Ohm}} = 0$ in the drag quantum well in Eq.(2.15).

$$
R_{\mathcal{D}} = \frac{\pi \hbar}{e^2} \frac{E_B}{\varepsilon_F} \frac{T}{\varepsilon_F} \frac{L_z}{L_y} \frac{1}{(k_F L_x)^2} \left(\frac{3\hbar\omega_B}{4\varepsilon_F}\right)^6 e^{12(2\varepsilon_F/3\hbar\omega_B)^3} \Phi\left[4Wk_F \frac{2}{3} \frac{\varepsilon_F}{\hbar\omega_B}\right],\tag{3.23}
$$

where we introduced effective Bohr energy $E_B = me^4/\kappa^2\hbar^2$ and

$$
p_F^B = \frac{2}{3} \frac{N_B}{N} \frac{\varepsilon_F}{\hbar \omega_B} p_F, \ \ p_F = \sqrt{2m\varepsilon_F}.
$$
\n(3.24)

With the given above parameters we have the following estimate for the transresistance

$$
R_{\rm D} \sim 0.4 \,\mathrm{m}\Omega.
$$

Now let us discuss when one can neglect the screening. Since the transferred momenta are $q_z \sim 2p_F^B/\hbar$ we may not take into account the screening of the Coulomb potential if the inverse screening length is much smaller than the transferred momentum. We estimate the screening length at a transferred energy $\sim\,T$ as

$$
\frac{1}{r_s} \sim \sqrt{\frac{\pi e^2 N_B}{\kappa \varepsilon_F^B}} \ln \frac{\varepsilon_F^B}{T}.\tag{3.25}
$$

The required inequality can be written as (we put $N_B = N$)

$$
\frac{N^{1/3}e^2}{\kappa} \ln \left[\frac{\varepsilon_F}{T} \left(\frac{\varepsilon_F}{\hbar \omega_B} \right)^2 \right] \ll \varepsilon_F \left(\frac{\varepsilon_F}{\hbar \omega_B} \right)^4. \tag{3.26}
$$

We will assume this inequality to be satisfied.

IV. NON-OHMIC CASE

The product of distribution functions $Eq.(2.22)$ can be recast into the form

$$
\mathcal{P} = 2\sinh\left(\frac{eV}{2T}\right)\exp\left\{\frac{\varepsilon_{p_z}^{(1)} - \mu_B}{T}\right\}\exp\left\{\frac{\varepsilon_{p_z}^{(2)} - \mu_B}{T}\right\}
$$
\n
$$
\times f\left(\frac{\varepsilon_{p_z}^{(1)} - \mu_B}{T}\right)f\left(\frac{\varepsilon_{p_z}^{(2)} - \mu_B - eV}{2}\right)f\left(\frac{\varepsilon_{p_z}^{(1)} - \mu_B}{T}\right)f\left(\frac{\varepsilon_{p_z}^{(2)} - \mu_B + eV}{2}\right)
$$
\n(4.1)

As P is a sharp function of p_z and p'_z one can take out of the integral all the slowly varying functions and get

$$
\int_0^\infty dp_z dp'_z \frac{\mathcal{P}}{(p_z + p'_z)} \int_{-\hbar L_x/2a_B^2}^{\hbar L_x/2a_B^2} \frac{2L_y dp_y}{2\pi\hbar} \frac{dp'_y}{2\pi\hbar} g_{00} \left[(p_z + p'_z)/\hbar, (p_y - p'_y)/\hbar \right]
$$

$$
= \frac{L_y m^2 a_B^2 T^2 \exp\left(2a_B k_F^B\right)^2}{4(4\pi\hbar)^3 (a_B k_F^B)^6} \Phi(4W k_F^B) \sinh\left(\frac{eV}{2T}\right) \frac{\frac{eV}{4T} - \frac{U_{12}}{2T}}{\sinh\left(\frac{eV}{4T} - \frac{U_{12}}{2T}\right)} \cdot \frac{\frac{eV}{4T} + \frac{U_{12}}{2T}}{\sinh\left(\frac{eV}{4T} + \frac{U_{12}}{2T}\right)} \tag{4.2}
$$

The drag current is

$$
J_{\text{drag}} = J_0 \frac{1}{2} \left(\frac{T}{\varepsilon_F^B}\right)^2 \sinh\left(\frac{eV}{2T}\right) \frac{\frac{eV}{4T} - \frac{U_{12}}{2T}}{\sinh\left(\frac{eV}{4T} - \frac{U_{12}}{2T}\right)} \cdot \frac{\frac{eV}{4T} + \frac{U_{12}}{2T}}{\sinh\left(\frac{eV}{4T} + \frac{U_{12}}{2T}\right)} \tag{4.3}
$$

For $eV \ll T$ one gets from Eq. (4.3) the result of Eq. (3.20). Let us consider the opposite case $eV \gg T$. In this case one gets a nonvanishing result for Eq.(4.3) only if $|U_{12}| < eV/2$ and one obtains the following equation for the drag current

$$
J_{\text{drag}} = J_0 \left[\left(\frac{eV}{4\varepsilon_F} \right)^2 - \left(\frac{U_{12}}{2\varepsilon_F} \right)^2 \right] \left(\frac{3\hbar\omega_B}{2\varepsilon_F} \right)^4 \tag{4.4}
$$

Thus the drag current vanishes unless $eV > 2|U_{12}|$.

V. SUMMARY

We have developed a theory of the Coulomb drag between two quantum wells in a strong longitudinal magnetic field. We have considered a comparatively simple limiting case where only the lowest Landau level is occupied. The strong magnetic field makes transverse motion of an electron one dimensional. These one dimensional electron states can be visualized as quantum "tubes" or "wires". Therefore, the Coulomb drag problem in this situation becomes similar to the Coulomb drag problem between two parallel nanowires.

It is interesting to compare our results with two different geometries of experiment. First, let us consider the influence of magnetic field on 1D Coulomb drag for the longitudinal geometry. In this case the magnetic field is directed along z axis and is parallel to 1D nanowires. For simplicity, we assume that the confining potential in the absence of the magnetic field is

$$
U(x,y) = \frac{m\Omega^2}{2}(x^2 + y^2).
$$
\n(5.1)

The applied magnetic field shortens the radius of the state a_B so that it becomes

$$
a_B^2 = \frac{a_0^2}{\sqrt{1 + (B/B_c)^2}}, \quad B_c = 2\frac{\Omega mc}{|e|}, \quad a_0 = \sqrt{\frac{\hbar}{2m\Omega}}
$$
(5.2)

where a_0 is the radius in the absence of the magnetic field. For the lowest Landau level we have

$$
\phi = \frac{1}{\sqrt{2\pi}} \frac{1}{a_B} \exp\left(-\rho^2 / 4a_B^2\right), \ \ \varepsilon_p = \frac{\hbar^2}{2ma_B^2} + \frac{p_z^2}{2m}.\tag{5.3}
$$

The wave function of the electron in the second wire can be obtained by a gauge transformation of the wave function in the first one. Since the interaction term q is not phase sensitive we are left only with a shift by the distance W between the centers of the wires in the argument of the wave function (5.3) . As a result, one gets for the interaction

$$
g(2p_F^B) = 4e^{-W^2/2a_B^2} \left[\int_0^\infty d\rho \rho e^{-\rho^2} I_0 \left(\frac{W}{a_B} \rho \right) K_0 \left(4 \frac{p_F^B a_B}{\hbar} \rho \right) \right]^2, \tag{5.4}
$$

where $I_0(x)$, $K_0(x)$ are the modified Bessel functions. The quasimomentum

$$
p_F^B = \frac{1}{2}\pi\hbar N_L^B
$$

must satisfy the inequality

$$
T \ll (p_F^B)^2 / 2m < \frac{\hbar^2}{2ma_B^2},\tag{5.5}
$$

since we have assumed that only the lowest Landau level is occupied. Here N_L^B is the electron density per unit length in magnetic field. The expression (5.4) demonstrates that provided the magnetic field goes up the localization radius a_B of the wave functions suppresses the probability of the backscattering processes. Note that if one assumes $N_L^B = N_L$, where N_L is the electron density per unit length for $B = 0$ then the effective interaction depends on the magnetic field only via a_B . Therefore in this case the magnetic field does not change the magnitude of transferred momentum, in contrast with the previous case where such a change leads to a rapid increase of the drag current in a strong magnetic field. The drag current is

$$
J_{\text{drag}} = J_{01} \frac{eV}{T} \left(\frac{T}{\varepsilon_F^B}\right)^2 \left(\frac{U_{12}}{2T}\right)^2 \left[\sinh\left(\frac{U_{12}}{2T}\right)\right]^{-2} \tag{5.6}
$$

$$
J_{01} = -\frac{e^5 m}{2\pi^2 \kappa^2 \hbar^3} L_z k_F^B \, g(2p_F^B). \tag{5.7}
$$

Second, we can compare our results with the drag between two two-dimensional quantum wells [10] in the field-free case. In this case the transresistance ρ_{12} is proportional to

$$
\rho_{12} \sim T^2 \frac{1}{(k_S d)^2} \frac{1}{(k_F d)^2},\tag{5.8}
$$

where k_S is the single-quantum well (two-dimensional) Thomas-Fermi screening wave vector of the order of the inverse effective Bohr radius, d is the interwell distance. First we note that the temperature dependence of the Coulomb drag between two (three-dimensional) quantum wells in the strong magnetic fields is weaker than for the drag in two dimensions. Second, we note that in the latter case the contribution from the backscattering processes can be neglected as compared to the small angle scattering contribution with transferred momenta $0 < q < 1/d \ll k_S$ while in our case only the backscattering processes are important (this is again a consequence of one-dimensionality of the Coulomb drag problem in the quantum limit).

ACKNOWLEDGMENTS

The authors acknowledge support for this work by the Russian National Fund of Fundamental Research (Grant No 03-02-17638).

- [1] M. C. Bonsager, K. Flensberg, B. Y.-K. Hu and A.-P. Jauho, Phys. Rev. Lett. 77, 1366 (1996),
- M. C. Bonsager, K. Flensberg, B. Y.-K. Hu and A.-P. Jauho, Phys. Rev. B 56, 10314 (1997).
- [2] A. Kamenev and Y. Oreg, Phys. Rev. B 52, 7516, (1995).
- [3] B. Y. K. Hu, Phys. Scripta T69, 170 (1997).
- [4] E. N. Bogachek, M. Jonson, R. I. Shekhter and T. Swahn, Phys. Rev. B 50, 18341 (1994).
- [5] V. L. Gurevich, V. B. Pevzner, and E. W. Fenton J. Phys.: *Condens. Matter* 10, 2551, (1998).
- [6] V. L. Gurevich, M. I. Muradov, Pis'ma v ZhETF, 71, 164, (2000) [JETP Letters 71, 111 (2000)].
- [7] R. Landauer, IBM J. Res. Develop. 1, 233 (1957); 32(3), 306 (1989).
- [8] Y. Imry , Directions in Condensed Matter Physics, ed. G. Grinstein and G. Mazenko, 1986 (Singapore: World Scientific), 101.
- [9] Büttiker M., Phys.Rev. Lett., 57, 1761 (1986).
- [10] T. J. Gramila, J. P. Eisenstein, A. H. MacDonald, L. N. Pfeiffer and K. W. West, Phys. Rev. Lett., 66, 1216 (1991).