

Stability and correlations in dilute two-dimensional boson systems

Han Guangze

Department of Applied Physics, South China University of Technology, Guangzhou 510640, China

O. Sørensen,* A. S. Jensen, and D. V. Fedorov

Department of Physics and Astronomy, University of Aarhus, DK-8000 Aarhus C, Denmark

(Dated: October 29, 2019)

The hyperspherical adiabatic expansion method is used to describe correlations in a symmetric boson system rigorously confined to two spatial dimensions. The hyperangular eigenvalue equation turns out to be almost independent of the hyperradius, whereas the solutions are strongly varying with the strength of the attractive two-body potentials. Instability is encountered in hyperangular, hyperradial, and mean-field equations for almost identical strengths inversely proportional to the particle number. The derived conditions for stability are similar to mean-field conditions and closely related to the possible occurrence of the Thomas and Efimov effects. Renormalization in mean-field calculations for two spatial dimensions is probably not needed.

PACS numbers: 21.45.+v, 31.15.Ja, 05.30.Jp

Introduction. Lower dimensions than three are necessary in several branches of physics, for instance surface physics, semiconductor physics, artificial atoms, and quantum dots. The advanced tools of well-controlled external fields, used in atomic and molecular physics to manipulate the effective interactions, allow confinement of the systems to lower dimensions [1, 2]. The experimental investigations employ continuous variation of the dimension by use of tunable deformed external fields.

The basic properties vary dramatically with the dimensionality of the system, as highlighted in two dimensions (2D) where the centrifugal s -wave barrier is negative for two particles and even an infinitesimally small attraction provides a bound state [3, 4]. The stability is strongly dependent on the deformation or the effective dimension of the confining potential [5, 6, 7]. The simplest N -body structures are the Bose-Einstein condensates for identical bosons. These systems are dilute, weakly interacting, and well described by mean-field models [8, 9] with an interaction strength adjusted to reproduce the low-energy scattering cross section in the Born approximation. The interaction in 2D is then obtained from short- or zero-range three-dimensional (3D) potentials restricted to the mean-field Hilbert space [10, 11].

Inclusion of correlations in the wave function prohibits this renormalization. Instead a finite short-range potential with the correct scattering length should be used. If the correlations are appropriately accounted for, the large-distance behavior must come out correctly with the realistic interaction [12, 13]. The consequences for lower dimensions are not yet investigated. The huge difference between two- and three-body properties in two and three dimensions is most likely more pronounced for N -body systems. Experimental results from varying dimensionality are easier to interpret if the limits are known.

The purpose of this letter is to investigate the possibly correlated structure and establish stability conditions under strict 2D confinement for N weakly-interacting identical bosons in an external harmonic field. We also provide a conceptual link between the successfully renormalized mean-field models and the correlated solutions with the bare (effective) interaction [13].

Theoretical method. We shall briefly sketch the method to establish the notation and the pertinent formulae for two dimensions. We follow the derivation for three dimensions given in [14]. The N identical bosons have masses m and coordinates \vec{r}_i . We use the hyperspherical adiabatic expansion method where the only length coordinate is the hyperradius ρ defined by

$$\rho^2 \equiv \frac{1}{N} \sum_{i < j}^N r_{ij}^2 = \sum_{i=1}^N r_i^2 - NR^2, \quad (1)$$

where $\vec{r}_{ij} = \vec{r}_i - \vec{r}_j$ and $\vec{R} = \sum_i \vec{r}_i / N$ is the center-of-mass coordinate. The remaining $2N - 3$ relative coordinates in two dimensions are angles where we define α_{ij} related to the size of \vec{r}_{ij} by $r_{ij} \equiv \sqrt{2} \rho \sin \alpha_{ij}$. If permitted in the context, we shall omit the indices ij .

The center of mass separates out and we only need to deal with relative coordinates. The related volume element is $\rho^{2N-3} d\rho \sin \alpha \cos^{2N-5} \alpha d\alpha d\vartheta d\Omega_{N-2}$, where $\alpha = \alpha_{ij}$, ϑ describes the direction of \vec{r}_{ij} and Ω_{N-2} denotes the remaining angular part of the volume element corresponding to the last $N - 2$ relative vector coordinates.

An external harmonic potential $m\omega^2 \sum_i r_i^2 / 2$ of angular frequency ω is by use of eq. (1) divided into a center-of-mass part and a hyperadial part. The relative Hamiltonian is then separated into a hyperradial part and a hyperangular part, \hat{h}_Ω , i.e.

$$\hat{H} = \frac{\hbar^2}{2m} \left(-\frac{1}{\rho^{2N-3}} \frac{\partial}{\partial \rho} \rho^{2N-3} \frac{\partial}{\partial \rho} + \frac{\rho^2}{b_t^4} + \hat{h}_\Omega \right), \quad (2)$$

*Electronic address: oles@phys.au.dk

$$\hat{h}_\Omega = \hat{\Pi}^2 + D_{angle} + \frac{2m\rho^2}{\hbar^2} \sum_{i<j}^N \hat{V}(r_{ij}), \quad (3)$$

$$\hat{\Pi}^2 = -\frac{\partial^2}{\partial\alpha^2} + \frac{2N-6-(2N-4)\cos 2\alpha}{\sin 2\alpha} \frac{\partial}{\partial\alpha}, \quad (4)$$

where all differential angular dependence, except α , is collected in D_{angle} . The trap length b_t is given by $b_t^2 \equiv \hbar/(m\omega)$. The two-body interaction \hat{V}_{ij} is of short range, e.g. a Gaussian $V_0 \exp(-r_{ij}^2/b^2)$ or a square well $V_0\Theta(r_{ij} < b)$, where Θ is the truth function.

The relative wave function $\Psi(\rho, \Omega)$ obeys the Schrödinger equation

$$\hat{H}\Psi(\rho, \Omega) = E\Psi(\rho, \Omega), \quad (5)$$

where E is the energy. We write Ψ as an adiabatic expansion [3, 14] where the first term is

$$\Psi(\rho, \Omega) = \rho^{-(2N-3)/2} f(\rho) \Phi(\rho, \Omega), \quad (6)$$

with the hyperradial volume element explicitly extracted. The angular wave function $\Phi(\rho, \Omega)$ is for fixed ρ an eigenfunction of \hat{h}_Ω with the eigenvalue $\lambda(\rho)$, i.e.

$$\hat{h}_\Omega \Phi(\rho, \Omega) = \lambda(\rho) \Phi(\rho, \Omega). \quad (7)$$

The corresponding radial equation is then

$$\left(-\frac{\hbar^2}{2m} \frac{d^2}{d\rho^2} + U(\rho) - E \right) f(\rho) = 0, \quad (8)$$

$$\frac{2mU(\rho)}{\hbar^2} = \frac{\lambda(\rho)}{\rho^2} + \frac{(2N-3)(2N-5)}{4\rho^2} + \frac{\rho^2}{b_t^4}, \quad (9)$$

where the adiabatic potential U is a function of the hyperradius consisting of three terms, i.e. the angular average λ of the interactions and kinetic energies, the generalized centrifugal barrier, and the external field.

For large particle distances only relative s waves contribute. With a Faddeev decomposition of the angular wave function only the dependence on distance α_{ij} is left, i.e.

$$\Phi(\rho, \Omega) = \sum_{i<j}^N \tilde{\phi}(\rho, \alpha_{ij}) \equiv \sum_{i<j}^N \frac{\phi(\rho, \alpha_{ij})}{\sin^{\frac{1}{2}} \alpha_{ij} \cos^{N-\frac{5}{2}} \alpha_{ij}}, \quad (10)$$

where we again explicitly extracted the square root of the volume element.

The integro-differential equation for $\phi(\rho, \alpha)$ is obtained from eq. (7) by integrating over all other angles than $\alpha_{12} = \alpha$, denoted by τ [14], i.e.

$$\left(-\frac{\partial^2}{\partial\alpha^2} + v_c + v + v_1 + v_2 - \lambda \right) \phi(\rho, \alpha) = \int d\tau G, \quad (11)$$

where $G = G(\tau, \alpha)$ is linear in both ϕ and V [14], and the reduced potentials are given by

$$v_c \equiv \frac{(2N-5)(2N-7)}{4} \tan^2 \alpha - \frac{\cot^2 \alpha}{4} - \frac{4N-9}{2}, \quad (12)$$

$$v(\alpha) = 4a_B \left(\frac{\rho}{b}\right)^2 \exp(-2(\rho \sin \alpha/b)^2), \quad (13)$$

$$v_1(\alpha) \equiv 2a_B \left(\frac{\rho}{b}\right)^2 (N-3)^2 (N-2) \times \int dx x^{N-4} \exp[2(x-1)\left(\frac{\rho}{b} \cos \alpha\right)^2] \quad (14)$$

$$\approx \frac{(N-3)^2 (N-2) a_B}{\cos^2 \alpha}, \quad (15)$$

$$v_2(\alpha) \approx \frac{4}{3} v_1(\alpha) \left(1 - \frac{1}{3} \tan^2 \alpha\right)^{N-4} \Theta(\alpha < \pi/3), \quad (16)$$

$$a_B \equiv \frac{m}{2\pi\hbar^2} \int V(r) d^3r = \frac{mV_0 b^2}{2\hbar^2}. \quad (17)$$

The last expression for a_B is valid both for Gaussian and square-well potentials V . The approximations for v_1 and v_2 are very accurate for $\rho \cos \alpha \gg b$. For $\alpha = \pi/2$ we get exactly $v_1(\pi/2) = 2(N-3)(N-2)a_B(\rho/b)^2$, and v_2 is very small.

For a short-range interaction the right-hand side of eq. (11) is independent of ρ as well as v_1 and v_2 when α is not too close to $\pi/2$. The only ρ dependence is then through v , which approaches a zero-range interaction in α as ρ increases. The eigenvalue $\lambda(\rho)$ is therefore expected to be constant in large ranges of ρ . These features are unique for two dimensions.

Stability conditions. For comparison we first consider the mean-field approximation in two dimensions. For an interaction of short range, i.e. small b the differential equation is in dimensionless quantities given by

$$\left(-\frac{\partial^2}{\partial x^2} - \frac{1}{x} \frac{\partial}{\partial x} + \left(\frac{b}{b_t}\right)^4 x^2 + 2a_B(N-1)|f_m(x)|^2 - \epsilon \right) f_m(x) = 0, \quad (18)$$

where $x \equiv r/b$ and $\epsilon \equiv 2mE_m b^2/\hbar^2$ are measures of the single particle mean-field coordinate r and energy E_m . The radial wave function f_m is approximated by a Gaussian, i.e. $f_m = \exp[-x^2/(2d^2)]/(\sqrt{\pi}d)$ normalized as $\int d^3x dx |f_m(x)|^2 = 1$. The corresponding energy per particle ϵ is then as a function of d given by

$$\epsilon = \left(\frac{b}{b_t}\right)^4 d^2 + \frac{1}{d^2} [1 + a_B(N-1)], \quad (19)$$

which only has a minimum when

$$a_B(N-1) > -1. \quad (20)$$

Then the energy and the Gaussian width are

$$d = \frac{b_t}{b} [1 + (N-1)a_B]^{1/4}, \quad (21)$$

$$\epsilon = 2\left(\frac{b}{b_t}\right)^2 \sqrt{1 + (N-1)a_B}. \quad (22)$$

Thus only repulsive or very weakly attractive potentials provide stable mean-field solutions. The reason is that even infinitesimally small attractions bind two particles in two dimensions. The corresponding two-body

Schrödinger equation with wave function f_2 and energy E_2 is

$$\left[-\frac{\partial^2}{\partial x^2} - \frac{1}{x} \frac{\partial}{\partial x} + 2a_B \left(\frac{V(x) - E_2}{V_0} \right) \right] f_2(x) = 0, \quad (23)$$

which for a weakly attractive interaction has the bound-state energy $E_2 = -4\hbar^2/(mb^2) \exp(2/a_B - 2\gamma)$ and the mean-square radius $\langle r^2 \rangle = 2\hbar^2/(3m|E_2|)$, where γ is Euler's constant, see [3].




FIG. 1: The centrifugal terms v_c and v_1 from eqs. (12) and (14) as function of α . The thick lines show $v_c + v_1$ for $N = 20$, various values of a_B , and $\rho/b = 10^3$ unless otherwise indicated. The thin, solid lines show the contributions from v_1 and v_c for $a_B = -0.02$.

The radial potential in eq. (9) depends crucially on λ determined from eq. (11) which in turn is dominated by the terms v_c and v_1 shown in fig. 1. When α approaches $\pi/2$ the approximation in eq. (15) and v_c eq. (12) both diverge as $\cos^{-2} \alpha \approx (\pi/2 - \alpha)^{-2}$. Thus when $a_B < -(N - 5/2)(N - 7/2)/[(N - 2)(N - 3)^2] \approx -1/(N - 2)$ an attractive pocket inevitably appears for large ρ . This divergence, corresponding to the Thomas effect [3], at the point $\alpha = \pi/2$ disappears when the exact expression in eq. (14) is used for v_1 . Then the lowest eigenvalue λ would be finite and proportional to ρ^{-2} . These solutions correspond to many particles close together which probably violates our assumption of s -wave dominance in the wave function. To avoid this divergence at large ρ the strength of the attraction must be limited by $a_B(N - 2) > -1$, which is almost identical to eq. (20).

For small α the angular centrifugal term v_c from eq. (12) diverges as $-\alpha^2/4$. This is the limit rigorously separating attractions leading to either no bound states or infinitely many bound states of Thomas or Efimov character [3]. Thus, a small two-body attraction v

is for large ρ sufficient to bind a state in the pocket at small α . This is seen by substituting $x \equiv \sqrt{2}\rho\alpha/b$ in eq. (11) when only the potential v and v_c are included. The energy and the mean-square radius of the solution becomes $\lambda(\rho) \rightarrow 2m\rho^2 E_2/\hbar^2$ and $\langle \alpha^2 \rangle \rightarrow 2/(3|\lambda|) = 1/12(b/\rho)^2 \exp(2\gamma - 2/a_B)$. If the size in α space has to be smaller than unity, ρ/b must exceed $\exp(-1/a_B)$ which is huge when $-1/N < a_B < 0$, i.e. the interaction is attractive but allows physical solutions in agreement with eq. (20). Thus the diverging λ corresponding to the bound two-body state is never encountered because either the interaction is too attractive leading to solutions in the pocket at large α , or the interaction is too weak to bind at small α for ρ values less than the trap length. Therefore diatomic recombination is unlikely.

Radial solutions. To solve the hyperradial eq. (9) we need λ . For $\rho = 0$ only v_c is present in eq. (11). The free angular solutions are

$$\tilde{\phi}_\nu = P_\nu^{(0, N-3)}(\cos 2\alpha), \quad \lambda_\nu^{(f)} = 4\nu(\nu + N - 2), \quad (24)$$

where $P_\nu^{(0, N-3)}$ is the Jacobi function [3] and $\nu = 0, 1, 2, 3, \dots$ is a non-negative integer. For very small ρ perturbation then gives $\lambda \approx \lambda_\nu^{(f)} + 2a_B N(N - 1)(\rho/b)^2$.

Increasing ρ for a repulsive or weakly attractive potential leaves approximately the free wave function, but the energies are shifted from the contribution of the interaction. In first-order perturbation with the free wave functions the eigenvalues are denoted $\lambda_\nu^{(\delta)}$, where the lowest for $\rho \gg b$ is found to be $\lambda_{\nu=0}^{(\delta)} = a_B N(N - 1)(N - 2)$. When the attraction is stronger, both wave function and energy change. However, the only ρ dependence in the angular equation is in v and in v_1 when $\alpha \approx \pi/2$. Still, the eigenvalues λ are essentially ρ independent when $b \ll \rho \ll \exp(-1/a_B)$, but perhaps lower than λ_ν . The eigenvalues must be obtained by numerical calculations.

The behavior of the angular eigenvalues in the interesting range of ρ values are illustrated in fig. 2. They can be parametrized rather well by

$$\lambda(\rho) \approx \frac{\lambda_\nu^{(f)} - \lambda_\nu^{(\delta)}}{1 + (\rho/\rho_0)^2} + \lambda_\nu^{(\delta)}, \quad (25)$$

$$\left(\frac{\rho_0}{b} \right)^2 = -\frac{2a_B N(N - 1)}{\lambda_\nu^{(f)} - \lambda_\nu^{(\delta)}} = \frac{2}{N - 2}, \quad (26)$$

where the last equality only holds for $\nu = 0$. Thus ρ_0 is independent of a_B and inversely proportional to N . In these expressions we must for somewhat stronger attraction replace $\lambda_\nu^{(\delta)}$ by the solution obtained numerically. The curve for $a_B = -0.0559$ in fig. 2 suddenly decreases dramatically when $\rho/b \approx 3000$. This reflects the pronounced effect of the attractive pocket for large α appearing in fig. 2 for large ρ .

The radial potential in eq. (9) has the same simple structure as the mean-field energy in eq. (19). Analogously stable solutions only exist for constant λ when $\lambda > -(N - 3/2)(N - 5/2)$. Using $\lambda_{\nu=0}^{(\delta)}$ this implies collapse when $a_B < -(N - 3/2)(N - 5/2)/[N(N - 1)(N -$




FIG. 2: The lowest angular eigenvalues as functions of ρ for different values of a_B and N . The thick, solid line is λ_0 for $N = 20$ and $a_B = -0.02$. The parameters for the other curves are given directly on the figure.

2)] $\approx -1/(N + 1)$. This condition is again very similar to the mean-field condition in eq.(20), but now less surprising since the hyperradial and the mean-field radial equations both describe the same overall size dependence.

For ρ values confined by the external field, we can insert the constant $\lambda_\nu^{(\delta)}$ given by eq. (25) in eq. (9) which then is the radial harmonic-oscillator equation for an effective angular-momentum quantum number l^* defined by $l^*(l^* + 1) \equiv \lambda + (N - 3/2)(N - 5/2)$ or equivalently $l^* = -0.5 + [\lambda + (N - 3/2)(N - 5/2) + 1/4]^{1/2}$. When $l^*(l^* + 1) > -1/4$ the solutions are well defined and characterized by the corresponding real values of $l^* > -1/2$. The eigenfunctions and eigenvalues are $(\rho/b_t)^{l^*} L_n^{(l^*-1/2)}[(\rho/b_t)^2] \exp[-0.5(\rho/b_t)^2]$ and $\hbar\omega(2n +$

$l^* + 3/2)$, where L_n are generalized Laguerre polynomials and n is a non-negative integer. The ground-state energy is then $\hbar\omega(l^* + 3/2)$ and the level spacing is precisely $2\hbar\omega$ as for harmonic-oscillator excitations maintaining parity.

In mean-field 3D calculations the realistic short- and zero-range interactions must be renormalized to reproduce the correct scattering length in the Born approximation [13]. This is equivalent to $\lambda_\nu \rightarrow \lambda_\nu^{(\delta)}$ for large ρ . Thus, to the extent the large- ρ behavior of λ_ν is well described by $\lambda_\nu^{(\delta)}$, this renormalization is not needed in 2D. Averaging the Gaussian interaction over the z direction confined by a typical length scale b_z , we find for a weak confinement, i.e. $b \ll b_z$ that $a_B \propto a_B^{(3)}/b_z$, where $a_B^{(3)}$ is the Born approximation to the scattering length in three dimensions [9]. The stability condition $Na_B > -1$ then becomes $Na_B^{(3)}/b_z > -1$, which coincides with the condition for a deformed trap [15] when the 3D scattering length is substituted by its Born approximation. Again this indicates that renormalization is not needed for 2D calculations. When the 3D scattering length is small compared to b_z , the 2D and 3D coupling strengths for a zero-range interaction are related by $g_{2D} \approx g_{3D}/(\sqrt{2\pi}b_z)$. When the axial confinement is strong, g_{2D} becomes density dependent [10].

Conclusions. We investigated symmetric N -boson systems in the rigorous two-dimensional limit for attractive two-body interactions. We derive the stability condition where the product of particle number, strength, mass, and square of the range has to be sufficiently small. The condition is the same for both mean-field and correlated solutions and independent of the external field. Diatomic recombination into two-body bound states is not likely, since the two-body bound state would extend beyond the trap length for potentials sufficiently weak to allow stable solutions. Thus these weakly attractive potentials, inevitably binding two particles, are able to support stable condensates. These features are completely different from three-dimensional properties. These rigorous two-dimensional results cannot be compared to other calculations where the effective interactions are repulsive.

-
- [1] D. Rychtarik, B. Engeser, H.-C. Nägerl, and R. Grimm, E-print arXiv cond-mat/0309536 (2003).
 - [2] A. Görlitz *et al.*, Phys. Rev. Lett. **87**, 130402 (2001).
 - [3] E. Nielsen, D. V. Fedorov, A. S. Jensen, and E. Garrido, Phys. Rep. **347**, 373 (2001).
 - [4] N. N. Khuri, A. Martin, and T.-T. Wu, Few-Body Syst. **31**, 83 (2002).
 - [5] J. L. Roberts *et al.*, Phys. Rev. Lett. **86**, 4211 (2001).
 - [6] A. Gammal, T. Frederico, and L. Tomio, Phys. Rev. A **64**, 055602 (2001).
 - [7] S. K. Adhikari, Phys. Rev. E **65**, 016703 (2002).
 - [8] C. J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases* (Cambridge University Press, Cambridge, 2002).
 - [9] L. P. Pitaevskii and S. Stringari, *Bose-Einstein Condensation* (Clarendon Press, Oxford, 2003).
 - [10] M. D. Lee, S. A. Morgan, M. J. Davis, and K. Burnett, Phys. Rev. A **65**, 043617 (2002).
 - [11] K. K. Rajagopal, P. Vignolo, and M. P. Tosi, E-print arXiv cond-mat/0309366 (2003).
 - [12] O. Sørensen, D. V. Fedorov, and A. S. Jensen, Phys. Rev. A **68**, 063618 (2003).
 - [13] O. Sørensen, D. V. Fedorov, and A. S. Jensen, J. Phys. B: At. Mol. Opt. Phys. **37**, 93 (2004).
 - [14] O. Sørensen, D. V. Fedorov, and A. S. Jensen, Phys. Rev. A **66**, 032507 (2002).
 - [15] G. Baym and C. J. Pethick, Phys. Rev. Lett. **76**, 6 (1996).