## Optimal Traffic Organisation in Ants under Crowded Conditions.

3 Audrey Dussutour<sup>\*‡</sup>, Vincent Fourcassié<sup>\*</sup>, Dirk Helbing<sup>#</sup>, Jean-Louis Deneubourg<sup>‡</sup>

4 <sup>\*</sup>Centre de Recherches sur la Cognition Animale, UMR CNRS 5169, Université Paul

5 Sabatier, 118 Route de Narbonne, F-31062 Toulouse Cedex 4, France

<sup>‡</sup>Centre d'Études des Phénomènes Non-linéaires et des Systèmes Complexe, Université Libre
de Bruxelles, CP231, Boulevard du Triomphe, 1050 Bruxelles, Belgium

8 <sup>#</sup>Institute for Economics and Traffic, Dresden University of Technology, D-01062 Dresden,

9 Germany

Efficient transportation, a hot topic in nonlinear science<sup>1</sup>, is most essential for 10 modern societies and the survival of biological species. As biological evolution has 11 generated a rich variety of successful solutions<sup>2</sup>, nature can inspire optimisation 12 strategies<sup>3,4</sup>. Foraging ants, for example, form attractive trails which support the 13 exploitation of initially unknown food sources in almost the minimum possible time<sup>5,6</sup>. 14 15 However, can this strategy cope with bottleneck situations, when interactions cause 16 delays, which reduce the overall flow? Here, we present an experimental study of ants 17 confronted with two alternative ways. We find that pheromone-based attraction 18 generates one trail at low densities, while at a high level of crowding, another trail is 19 established before the traffic volume is affected, which guarantees that an optimal rate 20 of food return is maintained. This bifurcation phenomenon is explained by a non-linear 21 modelling approach. Surprisingly, the underlying mechanism is based on inhibitory 22 interactions. It implies capacity reserves, a limitation of the density-induced speed 23 reduction, and a sufficient pheromone concentration for reliable trail perception. The

## balancing mechanism between cohesive and dispersive forces appears to be generic in natural, urban, and transportation systems.

Animals living in groups<sup>7,8</sup> often display collective movement along well-defined lanes 3 or trails<sup>9-13</sup>. This behaviour emerges through self-organisation resulting from the action of 4 individuals on the environment<sup>14-16</sup>. In ants<sup>17,18</sup> for example, mass recruitment allows a colony 5 6 to make adaptive choices solely based on information collected at the local level by individual 7 workers. When a scout ant discovers a food source, it lays an odour trail on its way back to 8 the nest. Recruited ants use the trail to find the food source and, when coming back to the 9 nest, reinforce it in turn. Without significant crowding, mass recruitment generally leads to the use of only one branch, i.e. to asymmetrical traffic<sup>19</sup>, because small initial differences in 10 pheromone concentration between trails are amplified. This also explains why ants use the 11 shorter of two branches leading to the same food source<sup>20</sup>, if it does not constitute a 12 bottleneck. 13

14 Here, we investigate the regulation of traffic flow during mass recruitment in the black garden 15 ant Lasius niger. Ants were forced to move on a diamond-shaped bridge with two branches 16 between their nest and a food source (Fig. 1). We have studied to what extent the traffic on 17 the bridge remains asymmetrical and, therefore, limited by the capacity of one branch, when 18 an increased level of crowding is induced by using branches of reduced widths (w = 10.0, 6.0,19 3.0 and 1.5 mm). The temporal evolution of the flow of ants on the bridge shown in Fig. 2 is typical of a trail recruitment process<sup>21</sup>. Surprisingly, the recruitment dynamics was not 20 influenced by the branch width w and the traffic volumes were the same (ANOVA, 21  $F_{3,44} = 0.500$ , P = 0.690). However, for w=10 mm, the majority of ants used the same branch 22 (Fig. 3a), while for  $w \le 6$  mm most experiments led to symmetrical traffic ( $\chi_2 = 1.686$ , 23 d.f. = 2, P = 0.430). This suggests that there is a transition from asymmetrical to symmetrical 24 traffic between 10.0 and 6.0 mm ( $\chi_2$ = 12.667, d.f. = 3, *P* = 0.005). 25

1 The mechanism for this "symmetry restoring transition" has been identified by experiments, 2 analytical calculations, and Monte Carlo simulations. Figure 3b supports indeed that the 3 proportion of experiments producing symmetrical flows on narrow bridges increases with the 4 total number of ants crossing the bridge ( $\chi_2 = 4.6$ , d.f. = 2, P = 0.04). Moreover, Fig. 3c 5 shows that both branches were equally used by the opposite flow directions. Therefore, in contrast to army ants<sup>6</sup> and pedestrians<sup>1</sup>, separation of the opposite flow directions was not 6 7 found. This could be explained by a high level of disturbance, e.g. a large variation in the 8 speeds of  $ants^{22}$ .

9 The flow of ants over the bridge can be understood analytically: The concentration of the trail 10 pheromone  $C_{ij}$  on branch *i* (*i* = 1, 2) immediately behind each choice point *j* (*j* = 1, 2) changes 11 in time *t* according to

12

$$dC_{ij}/dt = q\Phi_{ij}(t) + q\Phi_{ij'}(t-\tau) - \nu C_{ij}(t) \quad \text{with} \quad j' = 3 - j, \tag{1}$$

14

where  $\Phi_{i1}(t)$  represents the overall flow of foragers from the nest to the food source choosing branch *i* behind the choice point 1,  $\Phi_{i2}(t)$  the opposite flow on branch *i* behind the other choice point j' = 3 - j = 2,  $\tau$  the average time required for an ant to get from one choice point to the other, *q* the quantity of pheromone laid on the trail, and v the decay rate of the pheromone. Moreover, if the density is low enough,

20 21

$$\Phi_{ij}(t) = \phi_j(t) F_{ij}(t),$$

22

where  $\phi_1$  is the outbound flow of foragers from the nest to the food source and  $\phi_2$  the opposite, nestbound flow. The function  $F_{ij}$  describes the relative attractiveness of the trail on branch *i* at choice point *j*. For *Lasius niger* it is given by<sup>19</sup>

26

(2)

1 
$$F_{ij} = \frac{(k+C_{ij})^2}{(k+C_{1j})^2 + (k+C_{2j})^2} = 1 - F_{i'j} \text{ with } i'=3-i,$$
(3)

2

3 where k denotes a concentration threshold beyond which the pheromone-based choice of a
4 trail begins to be effective.

5 Equation (2) describes the flow dynamics without interactions between ants. However, when 6 traffic was dense, we observed that ants that had just engaged on a branch were often pushed 7 to the other branch when colliding frontally with another ant coming from the opposite 8 direction. This behaviour will turn out to be essential in generating symmetrical traffic on 9 narrow bridges. While pushing was practically never observed on a 10 mm bridge, on narrow 10 bridges the frequency of pushing events was high and proportional to the flow (Fig. 3d).

When pushing is taken into account, the overall flow of ants arriving at choice point *j*and choosing branch *i* can be expressed by the following formula:

13 
$$\Phi_{ii}(t) = \phi_i(t)F_{ii}(t)[1 - \gamma a \Phi_{ii'}(t - \tau)/w] + \phi_i(t)F_{i'i}(t)\gamma a \Phi_{i'i'}(t - \tau)/w$$
(4)

14 The first term on the right-hand side of (4) represents the flow of ants engaged on branch i  $[\phi_i(t)F_{ii}(t)]$ , diminished by the flow of ants pushed towards the other branch i' by ants 15 arriving from the opposite direction.  $a\Phi_{ii'}(t-\tau)/w$  is the proportion of ants decelerated by 16 17 interactions. The factor *a* is proportional to the interaction time period and the lateral width 18 of ants. Up to the symmetry restoring transition it can be considered approximately constant. 19  $\gamma \approx 0.57$  denotes the probability of being pushed in case of an encounter (Fig. 3d). The 20 second term on the right-hand side of Eq. (4) represents the flow of ants that were engaged on 21 branch *i*' and were pushed towards branch *i*.

22 The stationary solutions of this model are defined by the conditions  $dC_{ij}/dt = 0$ ,

23 
$$F_{ij}(t-\tau) = F_{ij}(t) = F_{ij}, \ \Phi_{ij}(t-\tau) = \Phi_{ij}(t) = \Phi_{ij}, \ \text{and} \ \phi_j(t-\tau) = \phi_j(t) = \phi_{j'}(t) = \phi$$
 (as the

- 24 nestbound flow and the outbound flow should be equal). These imply
- 25  $C_{ij} = q(\Phi_{ij} + \Phi_{ij'}) / \nu = C_{ij'}, \ F_{ij} = F_{ij'} = 1 F_{i'j} = 1 F_{i'j'}$

1 and 
$$C_{ij} = C_{ij'} = \frac{q\phi}{v} + D$$
,  $C_{i'j} = C_{i'j'} = \frac{q\phi}{v} - D$  (5)

2 with 
$$D = 0$$
 (6)

3 or

$$D^{2} = \frac{q^{2}\phi^{2}}{v^{2}} - \frac{k^{2} + \gamma a\phi(k + 2q\phi/v)^{2}/w}{1 + \gamma a\phi/w}$$
(7)

We can distinguish two cases: i) When  $D^2 > 0$ , the stable stationary solution corresponds to 4 asymmetrical traffic with  $C_{ij} = \frac{q\phi}{v} \pm \sqrt{D^2}$  and  $C_{i'j} = \frac{q\phi}{v} \mp \sqrt{D^2}$ . If  $\gamma = 0$ , this situation is 5 given for  $q\phi/v > k$ . If  $q\phi/v \le k$ , i.e. if the pheromone concentration is too low, the ants 6 7 choose both branches at random, corresponding to a symmetrical distribution (Fig. 4a). ii) 8 When pushing is taken into account with  $\gamma > 0$ , the organization of traffic changes considerably: the asymmetrical solution cannot be established anymore as soon as  $D^2 < 0$  for 9 10 large traffic volumes  $\phi$  (Fig. 4b). In this case, symmetrical traffic is expected with  $C_{ij} = q\phi/\nu$  and  $\Phi_{ij} = \phi/2$  for both branches *i* and choice points *j*. Therefore, high traffic 11 volumes can be maintained and none of the branches is preferred despite of the competitive 12 13 effect due to the accumulation of pheromone on both branches. Moreover, the model implies that the outbound flow  $\Phi_{i1}$  and the nestbound flow  $\Phi_{i2}$  are equal, indicating that one-way 14 flows are not required to maintain a high traffic volume<sup>23</sup>. These analytical results have been 15 16 confirmed by Monte-Carlo simulations (see Fig. 4c,d). Our simulations have also 17 demonstrated that, if pushed ants, instead of moving to the other branch, made a U-turn in the direction they were coming from, the overall flow of ants crossing the bridge was affected by 18 19 the branch width and no shift to symmetrical traffic was observed.

- 20 The traffic organisation in ants can be called optimal. The overall flow on branch *i* behind
- 21 choice point *j* is given by  $\Phi_{ij} = w \rho_{ij} V_{ij} \le \phi$ , where  $\rho_{ij}$  denotes the density of ants. Their
- 22 average speed is theoretically estimated by  $V_{ii} \approx V_m (1 a \Phi_{ii'} / w)$ , where  $V_m$  denotes the

(7)

average maximum speed and  $a\Phi_{ii'}/w$  again the proportion of decelerated ants. For the 1 2 symmetry restoring transition with  $D^2 = 0$ , Eq. (7) requires  $a\gamma\phi/w < 1/3$ , which implies  $V_{ii} > V_m [1 - 1/(3\gamma)] \approx 0.42 V_m$ . However, according to the *empirical* speed-density relation by 3 Burd et al.<sup>23</sup>, the maximum flow (the capacity) is only reached at the smaller speed 4  $V_{ii} = V_m [1 - 1/(n+1)] \approx 0.39 V_m$ . Although the empirical value  $n \approx 0.64$  was determined for 5 6 another ant species, the values for *L. niger* should be comparable. This shows that the 7 symmetry restoring transition occurs before the maximum flow is reached. The strict 8 inequality also implies capacity reserves and a limitation of the density-related speed 9 reduction. A marginal reduction in speed, however, would not be in favour of symmetrical 10 traffic because of benefits by using a single trail: First, a more concentrated trail provides a better orientation guidance and a stronger arousal stimulus<sup>24</sup>. Second, a higher density of ants 11 enhances information exchange and supports a better group defence $^{25}$ . 12

13 To conclude, this study demonstrates the surprising functionality of collisions among ants to keep up the desired flow level by generation of symmetrical traffic. Pushing behaviour may 14 be considered as an optimal behaviour to maintain a high rate of food return to the nest. It 15 would not be required in most models of Ideal Free Distribution<sup>26</sup> (IDF), as they neglect 16 17 effects of inter-attraction. The balancing between cohesive and dispersive forces avoids a 18 dysfunctional degree of aggregation and supports an optimal accessibility of space at minimal 19 costs allowing an efficient construction, maintenance and use of infrastructures. This 20 mechanism appears to be generic in nature, in particular in group living organisms. For 21 example, inhibitory interactions at overcrowded building sites in termites allow a smooth growth of the nest structure<sup>27</sup>. In the development of urban agglomerations they are important 22 to keep up the co-existence of cities<sup>28</sup> and in vehicle traffic, they determine the choice of 23 24 longer, less congested routes. The mechanism also suggests algorithms for the routing of data 25 traffic in networks.

- 1 Competing interests statement: The authors declare that they have no competing financial interests.
- 2 Correspondence and requests for materials should be addressed to dussutou@cict.fr

## 1 Figure Legends

2 Figure 1 Experimental set-up. Five queenless colonies of *Lasius niger* 3 (Hymenoptera, Formicidae) each containing 500 workers were used. During an 4 experiment, ants had access from the nest to a source of sucrose (2mL of 1M 5 solution) placed at the end of a diamond-shaped bridge. The solution was spread 6 over a large surface so that all ants arriving at the end of the bridge could have 7 access to the food. j = 1 and j = 2 indicate the choice points for branch selection. The 8 whole set-up was surrounded by white curtains to prevent any bias in branch choice 9 due to the use of visual cues. The colonies were starved for four days before each 10 experiment. The traffic on the two-branch bridge was recorded by a video camera for one hour. Nestbound and outbound ants were then counted and aggregated over 11 1-minute intervals. Counting began as soon as the first ant climbed the bridge. Four 12 13 different bridge widths w were used: 10.0, 6.0, 3.0 and 1.5mm. This set-up mimics 14 many natural situations in which the geometry of the trails is influenced by physical 15 constraints of the environment such as the diameter of underground galleries or of the branches in the vegetation along which the ants are moving. 16

Figure 2 a-d Average number of ants per minute crossing the two branches of the bridge within intervals of five minutes. The observed traffic volumes  $2\phi(t)$  and flow dynamics agreed for the four different branch widths, i.e. the bottleneck for smaller branches was compensated for by usage of both branches. *N* = 12 experiments were carried out for each bridge width. Error bars indicate the standard deviation.

Figure 3 Experimental results. (a) Experimental frequency distributions of the proportion  $F_{1j}$  of ants using the right branch for different branch widths *w*. We considered traffic to be asymmetrical when more than 2/3 of the cumulated traffic of ants at the end of the experiment had used a single branch. All experiments have been pooled. (b) Experimental frequency distributions similar to (a), but for different

1 total numbers of ants crossing the bridge. All experiments on bridges of 1.5, 3.0 and 2 6.0 mm width have been pooled. (c) Proportion of outbound ants on the left and right 3 branches of the bridge to the total number of ants that have passed the bridge at the 4 end of the experiment. The results contradict opposite one-way flows on both 5 branches. (d) Number of pushing events as a function of the total number of ant 6 encounters on the initial part of each branch of the bridge. For each branch width, 7 pushing events were evaluated at both choice points, for outbound and nestbound 8 ants, during the first ten and last ten minutes of a random sample of two experiments 9 characterised by symmetrical and two experiments characterised by asymmetrical 10 traffic. This yielded a total of 2 branches x 4 experiments x 2 time intervals = 16 points per branch width. When the number of encounters was too low ( $\leq$  3) the points 11 12 were not taken into account in the regression. The slope of the linear regression is 13 equal to 0.571  $\pm$  0.057 (Cl<sub>95%</sub>). This corresponds to the probability  $\gamma$  of an ant to be 14 pushed and redirected towards the other branch after encountering another ant 15 coming from the opposite direction (see **e-q**).

**Figure 4 Top:** Analytical results for the parameter values q = 1, k = 6, v = 1/40 min<sup>-1</sup>, 16 17 and a= 0.1 mm.min, which have been adjusted to experimental measurements. The 18 curves show the proportion  $F_{ii}$  of the overall flow  $\phi$  on each branch in the stationary 19 state. (a) In the absence of pushing ( $\gamma = 0$ ) the model predicts asymmetrical traffic 20 above very low values of the overall flow of ants, independently of the traffic volume. (b) When the proportion of pushed ants is high ( $\gamma = 0.6$ ), traffic is asymmetrical for 21 moderate values of the overall flow of ants, but stable symmetrical traffic with  $F_{ii}$  =0.5 22 23 is expected above a critical value of traffic flow which is an increasing function of 24 branch width w. The symmetry-restoring transition from asymmetrical to symmetrical traffic flow corresponds to an inverse pitchfork bifurcation. Bottom: Results of Monte-25 26 Carlo simulations. At time *t* = 0, the pheromone concentration and the number of ants on each branch are set to zero. Ants arrive at choice point j at a rate  $\phi_i(t) = \phi_i$ . The 27

9

1 probability of choosing the right or left branch at a choice point is governed by the 2 choice function  $F_{ij}$  [see Eq. (3)]. However, as soon as an ant has engaged on a 3 branch, it may be pushed to the other branch with probability  $\gamma$  by an oppositely 4 moving ant coming from the second choice point. The pushed ant then continues its 5 course on the alternative branch and lays a trail. For each value of  $\gamma$ , the simulations 6 are averaged over 1000 runs. The graphs show the relative frequency of simulations 7 in which a certain proportion of traffic was supported by the right branch. (c) 8 Assuming a pushing probability equal to zero ( $\gamma$  =0), most simulations ended with 9 asymmetrical traffic. However, when we used the experimental value  $\gamma = 0.6$ , most 10 simulations for narrow branches resulted in symmetrical traffic. (d) The proportion of 11 simulations with  $\gamma$  =0.6 in which asymmetrical traffic emerged is a function of the total 12 number of ants crossing the bridge, just as in our experiments (cf. Fig. 3c).

- 1 <sup>1</sup> Helbing, D. Traffic and related self-driven many-particle systems. *Reviews of Modern*
- 2 *Physics* **73**, 1067-1141 (2001).
- 3 <sup>2</sup> Camazine, S. et al., Self-organized Biological Systems (Princeton Studies in Complexity,
- 4 2001).
- 5 <sup>3</sup> Bonabeau, E., Dorigo, M. & Theraulaz, G. Inspiration for optimization from social insect
- 6 behaviour. *Nature* **406**, 39-42 (2000).
- 7 <sup>4</sup> Bonabeau, E., Dorigo, M. & Theraulaz, G. Swarm Intelligence. From Natural to Artificial
- 8 Systems (Oxford University Press, 1999).
- 9 <sup>5</sup> Burd, M. & Aranwela, N. Head-on encounter rates and walking speed of foragers in leaf-
- 10 cutting ant traffic. *Insectes soc.* **50**, 3-8 (2003).
- 11 <sup>6</sup> Couzin, I. D. & Franks, N. R. Self-organized lane formation and optimized traffic flow in
- 12 army ants. Proc. R. Soc. Lond. B 270, 139-146 (2003).
- <sup>7</sup> Parrish, J. K. & Hamner, W. K. *Animal Groups in three Dimensions* (Cambridge University
   Press, Cambridge, 1997)
- <sup>8</sup> Krause, J. & Ruxton, D. *Living in Groups* (Oxford Series in Ecology and Evolution, Oxford
- 16 University Press, 2002)
- <sup>9</sup> Galef, B. G. Jr. & Buckley, L. L. Using foraging trails by Norway rats. *Anim. Behav.* **51**,
- 18 765-771 (1996).
- <sup>10</sup> Fitzgerald, T. D. *The Tent Caterpillars* (Cornell University Press, Ithaca, N.Y., 1995).
- 20 <sup>11</sup> Erlandson, J. & Kostylev, V. Trail following, speed and fractal dimension of movement in a
- 21 marine prosobranch, *Littorina littorea*, during a mating and a non-mating season. *Marine*
- 22 Biol. 122, 87-94 (1995)
- 23 <sup>12</sup> Traniello, J. F. A. & Robson S. K. in: Chemical Ecology of Insects (eds. Cardé, R. T. &
- 24 Bell, W.) 241-286 (Chapman & Hall, 1995).
- 25 <sup>13</sup> Miura, T. & Matusmoto, T. Open-air litter foraging in the nasute termite *Longipeditermes*
- 26 longipes (Isoptera: Termitidae). J. Insect Behav. 11, 179-189 (1998).
- <sup>14</sup> Helbing, D., Keltsch, J. & Molnár, P. Modelling the evolution of human trail systems. *Nature* 388, 47-50 (1997).
- 29 <sup>15</sup> Edelstein-Keshet, L. W., Ermentrout, J. & Bard., G. Trail following in ants: individual
- 30 properties determine population behaviour. *Behav. Ecol. and Sociobiol.* **36**, 119-133 (1995).

- 1 <sup>16</sup> Schweitzer, F. *Brownian Agents and Active Particles. Collective Dynamics in the Natural*
- 2 *and Social Sciences* (Springer, Berlin, 2003).
- <sup>17</sup> Wilson, E. O. Chemical communication among workers of the fire ant *Solenopsis saevissima* (Fr. Smith). 1. The organization of mass-foraging. *Anim. Behav.* 10, 134-147
  (1962).
- 6 <sup>18</sup> Hölldobler, B. & Wilson E. O. *The Ants* (The Belknap Press of Harvard University Press,
- 7 Cambridge, MA, 1990).
- 8 <sup>19</sup> Beckers, R., Deneubourg, J. L. & Goss, S. Trails and U-turns in the selection of a path by
- 9 the ant Lasius niger. J. theor. Biol. **159**, 397-415 (1992).
- <sup>20</sup> Goss, S., Aron, S., Deneubourg, J. L. & Pasteels J. M. Self-organized shortcuts in the 11 Argentine ant. *Naturwissenschaften* **76**, 579-581 (1989).
- 12 <sup>21</sup> Pasteels, J. M., Deneubourg, J. L. & Goss, S.in: From Individual to Collective Behaviour in
- 13 Social Insects, Experientia supplementum, Vol. 54 (eds Pasteels, J. M. & Deneubourg, J. L.)
- 14 155-175 (Birkhäuser, Basel, 1987).
- 15 <sup>22</sup> Helbing, D. & Platkowski., T. Drift- or fluctuation-induced ordering and self-organization
- 16 in driven many-particle systems. *Europhys. Lett.* **60**, 227-233 (2002).
- <sup>23</sup> Burd, M. *et al.* Traffic dynamics of the leaf-cutting ant, *Atta cephalotes. Am. Nat.*, **159**, 283293 (2002).
- <sup>24</sup> Calenbuhr, V. *et al.* A model for osmotropotactic orientation. II. *J. theor. Biol.*, **158**, 395407 (1992).
- 21 <sup>25</sup> Feener, D.H. & Moss, K.A.G. Defense against parasites by hitchhikers in leaf-cutting ants:
- 22 a quantitative assessment. *Behav. Ecol. Sociobiol.* **49**, 348-356.
- 23 <sup>26</sup> Fretwell, S. D. & Lucas, H. L. On territorial behavior and other factors influencing habitat
- 24 distribution in birds. I. Theoretical development. Acta Biotheor. 19, 16–36 (1970).
- 25 <sup>27</sup> O'Toole, D.V., Robinson, P.A. & Myerscough, M.R. Self-organized criticality in termite
- architecture: a role for crowding in ensuring ordered nest expansion. J. theor. Bio., **198**, 305-
- 27 327.
- <sup>28</sup> Fujita, M., Krugman, P. & Venables, A. *The Spatial Economy Cities, Regions, and International Trade* (MIT Press, Cambridge, MA, 1999).







