Superfluid vs Ferromagnetic Behaviour in a Bose Gas of Spin-1/2 Atoms

S. Ashhab

Department of Physics, Ohio State University, 174 West Eighteenth Avenue, Columbus, Ohio 43210 August 31, 2004

Abstract

We study the thermodynamic phases of a gas of spin-1/2 atoms in the Hartree-Fock approximation. Our main result is that, for repulsive or weakly-attractive inter-component interaction strength, the superfluid and ferromagnetic phase transitions occur at the same temperature. For strongly-attractive inter-component interaction strength, however, the ferromagnetic phase transition occurs at a higher temperature than the superfluid phase transition. We also find that the presence of a condensate acts as an effective magnetic field that polarizes the normal cloud. We finally comment on the validity of the Hartree-Fock approximation in describing different phenomena in this system.

In recent years, studies of multi-component Bose-Einstein Condensates (BEC) have revealed a variety of interesting phenomena that reflect qualitatively different ideas from spinless condensates [1, 2, 3, 4, 5, 6, 7, 8, 9]. Gases of spin-1/2 [1], spin-1 [2, 3, 4, 5, 6, 7, 8] and spin-2 [8, 9] atoms have been studied in considerable detail, both experimentally and theoretically. New physical phenomena include fragmented states [5], coreless vortices [6] and complex spin dynamics [7, 8, 9]. Truely spin-1/2 particles are fermions. However, a gas of bosonic atoms each of which can occupy two different internal states can be treated as a gas of spin-1/2 atoms that obey Bose statistics (note that in nonrelativistic systems, there is no *a priori* connection between spin and statistics). Examples of such gases are mixtures of ⁸⁷Rb atomic gases in two hyperfine states and spin-polarized Hydrogen [1, 10].

It is well known that in a gas of noninteracting spin-1/2 bosonic atoms, the emergence of superfluid order is accompanied by the emergence of ferromagnetic order [11, 12]. A natural question that arises in that context is whether these two orders would remain connected if we include the effects of interatomic interactions. Perhaps a clearer way to pose the question is as follows: is it possible to have one of those two orders (i.e. superfluid or ferromagnetic) without having the other? One can give a number of elementary arguments that hint one way or the other. On the one hand, in the noninteracting system the connection between the two orders follows from the symmetry of the wave function and is not related to any thermodynamic arguments (except for saying that the normal cloud is unpolarized). That would suggest a rather robust connection between superfluid and ferromagnetic behaviours. On the other hand, if one considers the form of the order parameters or the symmetry of the Hamiltonian, there is no reason to believe that the two orders must be related. Keeping in mind certain caveats, the superfluid order parameter can be chosen as $|\langle \hat{\psi}_{\uparrow}(\mathbf{r}) \rangle|^2 + |\langle \hat{\psi}_{\downarrow}(\mathbf{r}) \rangle|^2$, whereas the ferromagnetic order parameter can be chosen as $\sum_{ij} \langle \hat{\psi}_i^{\dagger}(\mathbf{r}) \vec{\sigma}_{ij} \hat{\psi}_j(\mathbf{r}) \rangle$, where $\vec{\sigma}_{ij}$ is the vector of Pauli spin matrices, and the creation and annihilation operators used above will be explained when we give the Hamiltonian of the system below. The noninteracting Hamiltonian obeys SU(2) symmetry, whereas the interacting Hamiltonian obeys $U(1) \otimes U(1)$ symmetry. Furthermore, it has been shown recently by Yang and Li that the gound state of this system with no spin-dependent interactions is ferromagnetic, even if the interactions are strong enough to completely suppress superfluidity [13]. The energies of low-lying excitations, however, also drops to zero for such strong interactions. It is therefore not obvious, at first sight, whether or not one order can exist without the other at finite temperature. A similar problem for a Bose gas of spin-1 atoms with ferromagnetic interactions has been considered by Gu and Klemm [14]. One clear difference between spin-1/2 and spin-1 atoms is that with the latter one has a parameter that determines whether interatomic interactions are ferromagnetic or antiferromagnetic, even in a rotation-symmetric system. With spin-1/2atoms, apart from some obvious cases that we shall explain shortly, it is in general not immediately clear whether a certain set of interaction parameters describes ferromagnetic interatomic interactions. In fact, as explained above, a rotation-symmetric gas of spin-1/2atoms always exhibits ferromagnetic behaviour.

In the present Paper, we shall try to obtain the answer to the above-posed question in the Hartree-Fock (i.e. mean-field) approximation. We do that by calculating the free energy as a function of the macroscopic thermodynamic variables and minimizing the free energy with respect to those variables. Since similar approximations, as well as a number of systematic field-theoretic calculations, are not reliable in predicting the superfluid transition temperature in a spinless system (see e.g. [15, 16], also see Appendix A), our results are necessarily plagued by the same type of unreliability, and they cannot be considered conclusive. The advantage of using the the Hartree-Fock approximation, however, is that whenever it gives correct results, it gives them with a simple physical explanation. At the end of our treatment, we shall give our intuitive assessment of which results we believe describe real physical phenomena and which results we believe are merely artifacts of the Hartree-Fock approximation.

One rather trivial phenomenon in the context of spin-1/2 atoms occurs when the two internal states have different internal energies. In that case, the gas can be almost completely polarized at microkelvin temperatrues, assuming chemical equilibrium between the two spin states is reached. Although that phenomenon can be considered one form of ferromagnetic behaviour, we are not interested in it. Another ferromagnetic behaviour that we do not wish to consider here occurs when the two intra-component interaction strengths are sufficiently different. In that case it can, under certain conditions, be favourable for the majority of the atoms to occupy a single spin state. The system we shall consider in this Paper, which is also the most experimentally-relevant system of spin-1/2 atoms, is the one where the total spin along some axis is conserved, but the total spin perpendicular to that axis is not conserved. In the language of spin systems, one says that the longitudinal spin-relaxation time T_1 is much longer than the time scale of performing the experiment, whereas the transverse spinrelaxation time T_2 is shorter than the time scale of performing the experiment. We take the longitudinal axis to be the z-axis. Note that in the situation described above, and assuming that one starts with no net magnetization along the z-axis, the macroscopic magnetization expected for a condensate of noninteracting atoms lies entirely in the xy-plane.

Let us take a Bose gas of spin-1/2 atoms in a three dimensional box. The Hamiltonian of the system can be expressed as:

$$\dot{H} = \dot{H}_{\rm kin} + \dot{H}_{\rm int},\tag{1}$$

where

$$\hat{H}_{\rm kin} = -\frac{\hbar^2}{2m} \int d\mathbf{r} \sum_{\sigma=\uparrow,\downarrow} \hat{\psi}^{\dagger}_{\sigma}(\mathbf{r}) \nabla^2 \hat{\psi}_{\sigma}(\mathbf{r}), \qquad (2)$$

$$\hat{H}_{\text{int}} = \int d\mathbf{r} \sum_{\sigma,\sigma'=\uparrow,\downarrow} \frac{g_{\sigma\sigma'}}{2} \hat{\psi}^{\dagger}_{\sigma}(\mathbf{r}) \hat{\psi}^{\dagger}_{\sigma'}(\mathbf{r}) \hat{\psi}_{\sigma'}(\mathbf{r}) \hat{\psi}_{\sigma}(\mathbf{r}), \qquad (3)$$

m is the atomic mass, $\hat{\psi}^{\dagger}_{\sigma}(\mathbf{r})$ is an operator that creates an atom at position \mathbf{r} in spin state σ (the spin states \uparrow and \downarrow correspond to the z-component of atomic spin being equal to $\pm 1/2$, respectively), $\hat{\psi}_{\sigma}(\mathbf{r})$ is its Hermitian conjugate, $g_{\sigma\sigma'} = 4\pi\hbar^2 a_{\sigma\sigma'}/m$, and $a_{\sigma\sigma'}$ is the scattering length between two atoms in spin states σ and σ' . We assume that the number of atoms in the σ and σ' spin states is individually conserved, whereas the total spin in the xy-plane is not conserved. Furthermore, we shall assume that there is an equal number of atoms in the two spin states \uparrow and \downarrow [17]. In order to avoid dealing with the possibility of mechanical collapse of the gas or phase separation of the two components, we shall assume that both $g_{\uparrow\uparrow}$ and $g_{\downarrow\downarrow}$ are positive and that $g^2_{\uparrow\downarrow} < g_{\uparrow\uparrow}g_{\downarrow\downarrow}$.

We now derive an expression for the free energy of the above-described system as a function of the macroscopic thermodynamic variables and use it to determine the state of the system for a given temperature. The macroscopic variables are the number of atoms in the condensate N_o , the total spin of the condensate \mathbf{S}_C , the number of atoms in the normal cloud N_N , and the total spin of the normal cloud \mathbf{S}_N . We use the canonical ensemble, where the constraint of fixed total number of atoms $N_o + N_N$ is imposed explicitly. Since we are dealing with a macroscopic system, we treat the above variables as classical variables. A condensate of spin-1/2 atoms is ferromagnetic as a result of Bose symmetry [11, 12]. Its total spin S_C is equal to $N_o/2$ [i.e. $\hat{S}_C^2 = N_o/2(N_o/2+1)$]. On the other hand, the only constraint on the total spin of the normal cloud is that $|\mathbf{S}_N| \leq N_N/2$. Therefore, we express the x, y and z components of the total spin of the condensate as $(N_o/2\sin\theta_o\cos\varphi_o, N_o/2\sin\theta_o\sin\varphi_o, N_o/2\cos\theta_o)$ and those of the normal cloud as $(N_N s_{N\perp} \cos\varphi_N, N_N s_{N\perp} \sin\varphi_N, -N_o/2\cos\theta_o)$. Note that with the above values of condensate and normal-cloud spins, we have taken into account our assumption that the z-component of the total spin must vanish. For any given set of values of the thermodynamic variables, the free energy is given by:

$$F = \langle \hat{H}_{\rm kin} \rangle + \langle \hat{H}_{\rm int} \rangle - TS \tag{4}$$

where T is the temperature and S is the entropy of the system. The entropy S and the expectation values are calculated by considering all the different microscopic configurations corresponding to the given values of the macroscopic variables. We first consider the first and third terms of the free energy, which we denote by F_{ideal} . The condensate does not contribute to those terms in the free energy. Therefore we only need to evaluate F_{ideal} for a normal cloud of N_N atoms with total spin S_N . As we shall see below, the interaction energy is constant to leading order for all the different microscopic configurations in the Hartree-Fock thermodynamic ensemble for given N_N and S_N . Therefore, F_{ideal} is given by the same expression that it takes in the noninteracting system. The free energy of a spinless noninteracting uniform Bose gas is given by [18]:

$$F_{\text{ideal}} \equiv \langle \hat{H}_{\text{kin}} \rangle - TS \tag{5}$$

$$= N_N k_B T \left(\ln z - \frac{g_{5/2}(z)}{g_{3/2}(z)} \right), \tag{6}$$

where k_B is Boltzmann's constant and z is given by

$$g_{3/2}(z) = n_N \lambda_t^3, \tag{7}$$

$$g_j(z) \equiv \sum_{l=1}^{\infty} \frac{z^i}{l^j}, \qquad (8)$$

$$\lambda_t \equiv \sqrt{\frac{2\pi\hbar^2}{mk_B T}},\tag{9}$$

 $n_N = N_N/V$, and V is the volume of the sample. F_{ideal} of the system at hand is given by the sum of two terms of the above form of F_{ideal} for two independent Bose gases, one with $N_N(1/2 + s_N)$ atoms and the other with $N_N(1/2 - s_N)$ atoms, where $s_N = \sqrt{s_{N\perp}^2 + N_o^2 \cos^2 \theta_o/4N_N^2}$:

$$\frac{F_{\text{ideal}}}{N_N k_B T} = \left(\frac{1}{2} + s_N\right) \left(\ln z_+ - \frac{g_{5/2}(z_+)}{g_{3/2}(z_+)}\right) + \left(\frac{1}{2} - s_N\right) \left(\ln z_- - \frac{g_{5/2}(z_-)}{g_{3/2}(z_-)}\right),\tag{10}$$

where z_{\pm} are given by

$$g_{3/2}(z_{\pm}) = n_N \lambda_t^3 \left(\frac{1}{2} \pm s_N\right).$$
 (11)

Note that since z_+ cannot exceed the value 1, s_N must obey the condition $n_N \lambda_t^3 (1/2 + s_N) \leq g_{3/2}(1)$. Also note that F_{ideal} is a monotonically increasing function of s_N . If we take the limit $s_N \to 0$, we find that:

$$\frac{F_{\text{ideal}}}{N_N k_B T} = \alpha + \gamma s_N^2 + \eta s_N^4 + O(s_N^6), \qquad (12)$$

where

$$\alpha = \ln z - \frac{g_{5/2}(z)}{g_{3/2}(z)},\tag{13}$$

$$\gamma = 2\frac{g_{3/2}(z)}{g_{1/2}(z)},\tag{14}$$

$$\eta = \frac{2g_{3/2}^3(z)}{3} \left(\frac{3g_{-1/2}^2(z)}{g_{1/2}^5(z)} - \frac{g_{-3/2}(z)}{g_{1/2}^4(z)} \right), \tag{15}$$

and z is evaluated from Eq. (11) with $s_N = 0$. When z = 0, $\gamma = 2$ and $\eta = 4/3$, and as $z \to 1$, $\gamma \propto \text{const.}\sqrt{1-z}$ whereas η decreases slightly from the value 4/3 and remains finite. We now calculate the interaction energy in the Hartree-Fock approximation. In that approximation we assume that there is no coherence between states of different relative momentum of a pair of interacting atoms (i.e. $\langle a_{k_1\sigma_1}^{\dagger} a_{k_2\sigma_2}^{\dagger} a_{k_3\sigma_3} a_{k_4\sigma_4} \rangle$ vanishes unless the momenta of the creation operators match those of the annihilation operators, not just the sum of the momenta). The two-particle correlation functions needed to evaluate the interaction energy can then be straightforwardly calculated to give:

$$\langle \hat{\psi}_{\uparrow}^{\dagger}(\mathbf{r}) \hat{\psi}_{\uparrow}^{\dagger}(\mathbf{r}) \hat{\psi}_{\uparrow}(\mathbf{r}) \hat{\psi}_{\uparrow}(\mathbf{r}) \rangle = n_o^2 \cos^4 \frac{\theta_o}{2} + \frac{n_N^2}{2} \left(1 - \frac{n_o}{n_N} \cos \theta_o \right)^2 + 2n_o n_N \cos^2 \frac{\theta_o}{2} \left(1 - \frac{n_o}{n_N} \cos \theta_{\downarrow} \right)$$

$$\langle \hat{\psi}_{\downarrow}^{\dagger}(\mathbf{r}) \hat{\psi}_{\downarrow}^{\dagger}(\mathbf{r}) \hat{\psi}_{\downarrow}(\mathbf{r}) \hat{\psi}_{\downarrow}(\mathbf{r}) \rangle = n_o^2 \sin^4 \frac{\theta_o}{2} + \frac{n_N^2}{2} \left(1 + \frac{n_o}{n_N} \cos \theta_o \right)^2 + 2n_o n_N \sin^2 \frac{\theta_o}{2} \left(1 + \frac{n_o}{n_N} \cos \theta_{\downarrow} \right)$$

$$\langle \hat{\psi}_{\uparrow}^{\dagger}(\mathbf{r}) \hat{\psi}_{\downarrow}^{\dagger}(\mathbf{r}) \hat{\psi}_{\downarrow}(\mathbf{r}) \hat{\psi}_{\downarrow}(\mathbf{r}) \rangle = \left(\frac{n_o + n_N}{2} \right)^2 + n_N^2 s_{N\perp}^2 + n_o n_N s_{N\perp} \sin \theta_o \cos(\varphi_o - \varphi_N)$$

$$(18)$$

where $n_o = N_o/V$. By direct substitution of the above correlation functions, one can find the expression for the interaction energy $\langle \hat{H}_{int} \rangle$ of the system. The state of the system can now be determined by minimizing the free energy:

$$\frac{F}{V} = f_o - g_- n_o^2 \cos \theta_o - \frac{g_+}{2} n_o^2 \cos^2 \theta_o + g_{\uparrow\downarrow} n_N^2 s_{N\perp}^2 + g_{\uparrow\downarrow} n_o n_N s_{N\perp} \sin \theta_o \cos(\varphi_o - \varphi_N) \\
+ \frac{F_{\text{ideal}}(N_N, k_B T, s_{N\perp}, \cos \theta_o)}{V},$$
(19)

where

$$f_{o} = \frac{\bar{g}}{2}n_{o}^{2} + \tilde{g}n_{N}^{2} + 2\tilde{g}n_{o}n_{N}$$

$$= \tilde{g}(n_{o} + n_{N})^{2} - \frac{g_{+}}{2}n_{o}^{2}$$
(20)

$$\bar{g} = \frac{g_{\uparrow\uparrow} + g_{\downarrow\downarrow} + 2g_{\uparrow\downarrow}}{4} \tag{21}$$

$$\tilde{g} = \frac{g_{\uparrow\uparrow} + g_{\downarrow\downarrow} + g_{\uparrow\downarrow}}{4} \tag{22}$$

$$g_{\pm} = \frac{g_{\uparrow\uparrow} \pm g_{\downarrow\downarrow}}{4} \tag{23}$$

Note that the fourth and fifth terms in Eq. (19), among others, result from exhangeinteractions. Those two terms describe scattering processes where an atom with spin \uparrow and another with spin \downarrow exchange their momenta. In the expression for the free energy (Eq. 19), the second and third terms favour maximizing $|\cos\theta_o|$ (i.e. having finite values of the z-component of the condensate and normal cloud), whereas the fifth and sixth terms favour taking $\cos \theta_o = 0$. The second term can be understood quite intuitively as follows. Since the interaction term is magnified in the normal cloud as compared to the condensate (because of the exchange terms), the normal atoms tend to accumulate in the spin state with lessrepulsive interactions. For example, if we take $g_{\uparrow\uparrow} > g_{\downarrow\downarrow}$, we find that the normal cloud will have an excess of atoms in the \downarrow spin state, leaving the condensate with an excess of atoms in the \uparrow spin state. Since in this Paper we are not interested in that phenomenon, we eliminate it by taking $g_{\uparrow\uparrow} = g_{\downarrow\downarrow}$. The third term in Eq. (19) can also be understood by considering the absence of exchange interaction terms within the condensate. Since we are requiring that half the atoms have spin \uparrow and the other half have spin \downarrow , it is straightforward to see that in order to gain the greatest reduction in interaction energy, the condensate atoms tend to accumulate in the same spin state (either \uparrow or \downarrow). The competition between the third term and last two terms of Eq. (19) determines whether the condensate and normal cloud will have any net magnetization along the z-axis. Clearly, at low enough temperatures, the third term will win, and one will have a finite z-component of the polarization of the condensate and normal cloud (note that the net polarization of the entire cloud must vanish, which is one of our main assumptions). That result is quite interesting in its own right. However, since the main question addressed in this Paper concerns the phase transitions, we focus our attention on those relatively-high temperatures. We minimize the free energy with respect to n_o , $\cos\theta$ and $s_{N\perp}$ at any given combination of the parameters $g_{\uparrow\uparrow}$, $g_{\uparrow\downarrow}$ and T to find the thermodynamic phases of the system. The results of a numerical calculation of the order of the phase transition are shown in Fig. 1. It is also worth making some analytical remarks about the bahaviour of the system. For clarity we address the following two cases separately (in the following remarks, we shall implicitly use a result that we found from our numerical calculation, namely that $\cos\theta$ remains negligibly small close to the transition temperatures): Case 1: $g_{\uparrow\downarrow} > 0$. If $n_o \neq 0$, the free energy is minimized by taking $\varphi_o - \varphi_N = \pi$ (i.e the polarizations of the condensate and normal cloud point in opposite directions) and a finite value of $s_{N\perp}$. In the limit $n_o/n_N \ll 1$,

$$s_{N\perp} = \frac{g_{\uparrow\downarrow}n_o}{2(\gamma k_B T + g_{\uparrow\downarrow}n_N)} + O\left(\frac{k_B T n_o^3}{(\gamma k_B T + g_{\uparrow\downarrow}n_N)^4}\right)$$
(24)

Note that above the BEC critical temperature T_c , i.e. when $n_o = 0$, the normal cloud is not

polarized at all. Below T_c the physics can be understood in terms of the condensate acting as an effective magnetic field that partially polarizes a paramagnetic normal cloud. Since the superfluid transition occurs at a temperature where $\gamma k_B T \sim g_{\uparrow\uparrow} n$ (see Appendix A), we find that:

$$n_N s_{N\perp} < \frac{n_o}{2} \tag{25}$$

Therefore, the polarization of the normal cloud is smaller than that of the condensate, and the net polarization does not vanish. Note that if we calculate the exact expression for $n_N s_{N\perp}$, substitute it in Eq. (19) and minimize F with respect to n_o (keeping $n_o + n_N$ fixed), we find that the global minimum of F jumps discontinuously from a point with $n_o = 0$ to a point with a finite value of n_o . That would suggest a first-order phase transition to the superfluid phase, which is also ferromagnetic. However, the Hartree-Fock approximation predicts a first-order phase transition in a spinless Bose gas [15] (also see Appendix A), and we therefore suspect that this result must be an artifact of the approximation.

Case 2: $g_{\uparrow\downarrow} < 0$. In this case the free energy is minimized by taking $\varphi_o - \varphi_N = 0$ (i.e the polarizations of the condensate and normal cloud point in the same direction). The term $g_{\uparrow\downarrow}n_N^2 s_{N\perp}^2$ favours a polarized normal cloud. That suggests that the gas might exhibit ferromagnetic behaviour even if $n_o = 0$, as we shall see shortly. Using the small $s_{N\perp}$ expression for the ideal-gas free energy (Eq. 12) we find that the free energy is minimized by choosing

$$s_{N\perp} = \frac{|g_{\uparrow\downarrow}|n_o}{2(\gamma k_B T - |g_{\uparrow\downarrow}|n_N)} + O\left(\frac{k_B T n_o^3}{(\gamma k_B T - |g_{\uparrow\downarrow}|n_N)^4}\right),\tag{26}$$

if $\gamma k_B T - |g_{\uparrow\downarrow}| n_N \gg |g_{\uparrow\downarrow}| n_o$. As in case 1, the condensate acts as an effective magnetic field that polarizes the normal cloud. One can also immediately see that the normal gas exhibits ferromagnetic behaviour when $\gamma k_B T - |g_{\uparrow\downarrow}| n_N$ becomes negative. Note that since $\gamma = 0$ when $T/T_c^o \leq 1$, the ferromagnetic phase transition must occur at a temperature higher than that of the ideal-gas BEC phase transition T_c^o . Assuming that $n_o = 0$ just below the ferromagnetic phase transition, we find that that transition occurs at a temperature where:

$$2\frac{g_{3/2}(z)}{g_{1/2}(z)}k_B T_{\text{ferro}} = |g_{\uparrow\downarrow}|n_N,$$
(27)

where z is given by:

$$\frac{g_{3/2}(z)}{g_{3/2}(1)} = \left(\frac{T_c^o}{T_{\rm ferro}}\right)^{3/2}.$$
(28)

Just below the transition temperature, the spin grows as:

$$s_N = \sqrt{\frac{\gamma k_B (T_{\text{ferro}} - T)}{2\eta}}.$$
(29)

Note that if $|g_{\uparrow\downarrow}|$ is smaller than a certain value, the ferromagnetic transition temperature can be smaller than the superfluid transition temperature, which is also higher than T_c^o for positive $g_{\uparrow\uparrow}$. In that case there would be a single (first-order) phase transition, just as in the case $g_{\uparrow\downarrow} > 0$. However, as can be seen from Fig. 1 and as explained in Appendices A and B, there is a region in parameter space where the ferromagnetic transition temperature is higher than the superfluid transition temperature.

We now summarize our results and comment on them. We found that when $g_{\uparrow\downarrow} > 0$, there is a single transition to a phase that has both superfluid and ferromagnetic order, and the

phase transition is first order. We believe that the first-order nature of the phase transition is an artifact of the Hartree-Fock approximation. However, since the physical mechanisms favouring an unpolarized gas (interaction energy) and those favouring an unpolarized normal gas (entropy) are both real physical mechanisms, we suspect that as soon as there is a finite fraction of atoms in the condensate, the induced polarization of the normal cloud will be smaller than the polarization of the condensate, and there will in fact be a single phase transition to a state with both superfluid and ferromagnetic order. In the case $g_{\uparrow\downarrow} <$ 0, we found that it is possible, for large enough $|g_{\uparrow\downarrow}|$, to have two phase transitions. At temperature T_{ferro} , we found a second-order phase transition to a ferromagnetic phase with no superfluid order. We suspect that that result will persist even beyond the Hartree-Fock approximation, because that transition occurs above the superfluid transition temperature, and therefore fluctuations in the superfluid order parameter are expected to be negligible. At a lower termperature we found a first-order phase transition to a superfluid phase (with ferromagnetic order). As above, we believe that the superfluid phase transition will also be second-order. Based on similar elementary arguments alone, we cannot comment on the location of the boundary separating the single-transition and double-transition regimes.

In conclusion, we have performed a Hartree-Fock (mean-field) calculation to study the phase transitions in a gas of spin-1/2 bosonic atoms. We found that it is possible to have ferromagnetic order with no superfluid order, but not vice versa. We also found that the phase transition to the ferromagnetic non-superfluid phase is second-order, whereas any transition to a superfluid phase is first order. We suspect that the result of a nonsuperfluid ferromagnetic phase describes a real physical phenomenon. Judging from the Hartree-Fock results in the spinless case, however, we believe that in the real system, all phase transitions will turn out to be second order in nature. Due to the unreliability of the Hartree-Fock approximation near the superfluid transition temperature, further study is required to confirm or refute the results of this Paper.

The author would like to thank T.-L. Ho for useful discussions and Q. Gu for useful correspondence about the results of Ref. [14]. This work was supported by the National Science Foundation through Grant Nos. DMR-0071630 and DMR-0109255 and by NASA through Grant Nos. NAG8-1441 and NAG8-1765.

Appendix A:

In this Appendix we show that the Hartree-Fock approximation predicts a first-order phase transition in a weakly-interacting spinless Bose gas [15] (Here we are assuming repulsive interactions). Using similar arguments to those used in the main text, we find that the free energy density as a function of condensate number density n_o and normal cloud number density n_N is given by:

$$\frac{F}{V} = n_N \left(\ln z - \frac{g_{5/2}(z)}{g_{3/2}(z)} \right) - \frac{g}{2} n_o^2 + g(n_o + n_N)^2, \tag{30}$$

where

$$g_{3/2}(z) = n_N \left(\frac{2\pi\hbar^2}{mk_BT}\right)^{3/2},$$
 (31)

with the constraint that the right-hand side of Eq. (31) cannot be greater than $g_{3/2}(1)$. The dependence of $(F(n_o)-F(0))/V$ on n_o (keeping the total density fixed) is shown schematically in Fig. A1. Clearly this calculation predicts a first-order phase transition at a temperature

 $T_c > T_c^o$, where T_c^o is the transition temperature of the noninteracting system. By expanding F/V in powers of n_o , we find that, to leading order in na^3 , the shift in transition temperature is given by:

$$\frac{T_c - T_c^o}{T_c^o} = 1.08(na^3)^{1/3}.$$
(32)

Note that as long as g > 0, we find that z < 1 even below the transition temperature. Also note that when applying the above analysis to the situation discussed in this Paper, e.g. by taking $a_{\uparrow\downarrow} = 0$, some additional factors of 2 appear that lead to replacing the factor 1.08 in Eq. (32) by 0.86, with that equation now relating $(T_c - T_c^o)/T_c^o$ to $(na_{\uparrow\uparrow}^3)^{1/3}$. Appendix B:

In this Appendix we calculate, to leading order the transition temperature to the ferromagnetic nonsuperfluid phase. As explained in the main text, we treat only the case $g_{\uparrow\downarrow} < 0$, and we assume that $|g_{\uparrow\downarrow}|$ is large enough that such a phase exists for a certain range of temperatures. We take Eq. (27), and use the asymptotic behaviour of the functions $g_{3/2}(z)$ and $g_{1/2}(z)$ as follows:

$$\lim_{z \to 1^{-}} \frac{g_{3/2}(z) - g_{3/2}(1)}{\sqrt{1-z}} = -3.545,$$
(33)

which gives the asymptotic function:

$$g_{3/2}(z) - g_{3/2}(1) \approx -3.545\sqrt{1-z}.$$
 (34)

The asymptotic limits of the functions $g_{1/2}(z), g_{-1/2}(z), ...$ can be derived using the above approximation for $g_{3/2}(z)$ and the relation $dg_j(z)/dz = g_{j-1}(z)/z$. Using the asymptotic forms of $g_{3/2}(z)$ and $g_{1/2}(z)$ in Eq. (27), we find that

$$\frac{T_{\text{ferro}} - T_c^o}{T_c^o} = 1.84 \left(n |a_{\uparrow\downarrow}|^3 \right)^{1/3}$$
(35)

Comparing the results of Appendices A and B, one would expect the boundary between the two regions, i.e. those corresponding to a single and double phase transitions, to occur when $|a_{\uparrow\downarrow}|/a_{\uparrow\uparrow} = 0.47$. The value of the ratio $|a_{\uparrow\downarrow}|/a_{\uparrow\uparrow}$ for the actual boundary will be higher than 0.47 if one takes into account the fact that a negative value of $a_{\uparrow\downarrow}$ reduces T_c , in agreement with the results of our numerical calculation shown in Fig. 1.

Figures

Fig. 1: Regions of different behaviour in a gas of spin-1/2 atoms. The x and y axes are given by $g_1 \equiv g_{\uparrow\uparrow} n/k_B T_c^o$ and $g_2 \equiv g_{\uparrow\downarrow} n/k_B T_c^o$. In region I we find a single phase transition to a superfluid ferromagnetic phase. In region II we find two phase transitions, with the ferromagnetic transition occurring at a higher temperature than the superfluid transition. Regions III and IV were not considered in this Paper because of possible instabilities: the former being susceptible to phase separation between the two spin species, and the latter susceptible to the whole cloud imploding because the net interatomic forces are attractive in that region.

Fig. A1: Schematic plot of the free energy $F(n_o) - F(0)$ of a spinless Bose gas as a function of condensate fraction n_o for $T > T_c$ (solid line), $T = T_c$ (dashed line), $T_c > T > T_c^o$ (dash-dotted line) $T = T_c^o$ (dash-double-dotted line). The dotted line is the x axis and serves as a guide to find the temperature of the first-order phase transition. As the temperature

is lowered, the point that minimizes F jumps discontinuously from 0 to a finite value at temperature T_c that is higher than the noninteracting transition temperature T_c^o .

References

- C. J. Myatt, E. A. Burt, R. W. Ghrist, E.A. Cornell, and C. E. Wieman, Phys. Rev. Lett. 78, 586 (1997).
- [2] J. Stenger, S. Inouye, D.M. Stamper-Kurn, H.-J. Miesner, A.P. Chikkatur, and W. Ketterle, Nature 396, 345 (1998).
- [3] T. Ohmi and K. Machida, J. Phys. Soc. Jpn. 67, 1822 (1998).
- [4] T.-L. Ho, Phys. Rev. Lett. **81**, 742 (1998).
- [5] C. K. Law, H. Pu, and N. P. Bigelow, Phys. Rev. Lett. 81, 5257 (1998).
- [6] A. E. Leanhardt, Y. Shin, D. Kielpinski, D. E. Pritchard, and W. Ketterle, Phys. Rev. Lett. 90, 140403 (2003).
- [7] H. Pu, C. K. Law, S. Raghavan, J. H. Eberly, and N. P. Bigelow, Phys. Rev. A 60, 1463 (1999).
- [8] H. Schmaljohann, M. Erhard, J. Kronjager, M. Kottke, S. van Staa, L. Cacciapuoti, J. J. Arlt, K. Bongs, and K. Sengstock, Phys. Rev. Lett. 92, 040402 (2004).
- [9] M.-S. Chang, C. D. Hamley, M. D. Barrett, J. A. Sauer, K. M. Fortier, W. Zhang, L. You, and M. S. Chapman, cond-mat/0309164.
- [10] I. J. Bonalde, T. M Brill, W. Yao, B. J. Verhaar, and I.F. Silvera, J. Low Temp. Phys. 113, 211 (1998).
- [11] E.D. Siggia and A.E. Ruckenstein, Phys. Rev. Lett. 44, 1423 (1980).
- [12] T.-L. Ho and L. Yin, Phys. Rev. Lett. 84, 2302 (2000).
- [13] K. Yang and Y.-Q. Li, Int. J. Mod. Phys. B 17, 1027 (2003).
- [14] Q. Gu and R. A. Klemm, Phys. Rev. A 68, 031604(R) (2003).
- [15] G. Baym and G. Grinstein, Phys. Rev. D 15, 2897 (1977).
- [16] See e.g. A. L. Fetter and J. D. Walecka, Quantum Theory of Many-Particle Systems (McGraw-Hill, New York, 1971).
- [17] The reason why we are restricting our analysis to states where the numbers of atoms in the \uparrow and \downarrow states are equal is to simplify the algebra. In the regimes of interest to us, the state of the system that minimizes the free energy will satisfy that condition even if it is not imposed externally.
- [18] See e.g. K. Huang, *Statistical Mechanics* (John Wiley & Sons, New York, 1963).