Possible Force-Entropy Correlation

Enrique Canessa[∗]

The Abdus Salam International Centre for Theoretical Physics, Trieste, Italy

Abstract

A statistical thermodynamic approach of moving particles forming an elastic body is presented which leads to reveal molecular-mechanical properties of classical and nonextensive dynamical systems. We derive the Boltzmann-Gibbs (BG) entropy form and relate it to Newton's law of motion in relation to a distinct tensile force acting on the systems at constant volume and number of particles. Tsallis generalization of the BG entropy is deduced assuming the thermal energy of the particles to be proportional to their energy states by the nonextensivity factor $q - 1$.

PACS numbers: 05.70.Ce; 05.70.Ln; 05.70.-a

Keywords: Statistical Physics and Thermodynamics; Probability theory; Nonlinear dynamics; Nonextensive statistical mechanics

[∗] E-mail: canessae@ictp.trieste.it

I. OVERVIEW

It was pointed out long ago by Einstein that the foundations of statistical mechanics should lie on the underlying dynamics [\[1\]](#page-5-0). All statistical mechanics results should derive from the dynamics if the systems considered are systems consisting of moving particles.

Within the classical Boltzmann-Gibbs (BG) statistics, it is still unkown how to derive from say Newton's law of motion the remarkable expression for the thermodynamic entropy [\[2](#page-5-1)]

$$
S_{BG}/k_B = -\sum_{i=1}^{W} p_i \ln p_i = \left\langle \ln \frac{1}{p_i} \right\rangle , \qquad (1)
$$

where mean values $\langle \cdots \rangle$ > correspond to $\sum_i (\cdots) p_i$. Even if this functional form has been profusely verified on isolated systems in thermal equilibrium, a rigorous derivation on how S_{BG} descends from dynamics has been limited to a dilute gas only [\[1](#page-5-0)]. It is usually approximated via the statistics of a large ensemble of identical replicas (see, $e.g., [3]$ $e.g., [3]$ $e.g., [3]$). Only recently, there has been computational evidence on the fact that BG thermal equilibrium distribution descends from Newton law [\[4](#page-6-0)].

Also unclear at present [\[1](#page-5-0), [2\]](#page-5-1), is the connection between the dynamics of a particle system and the Tsallis nonextensive generalization of the BG entropy

$$
S_q/k_B = \left\langle \ln_q \frac{1}{p_i} \right\rangle = \sum_{i=1}^W p_i \ln_q \frac{1}{p_i} = \frac{1 - \sum_{i=1}^W p_i^q}{q - 1} , \qquad (2)
$$

where

$$
\ln_q p_i \equiv \frac{p_i^{1-q} - 1}{1-q} \quad , \tag{3}
$$

which for $q = 1$, it coincides with S_{BG} giving positive Lyapunov exponents [\[5\]](#page-6-1). The nonextensivity is given by the pseudo-additivity property $S_q(A+B) = S_q(A) + S_q(B) + \frac{1-q}{k_B} S_q(A) S_q(B)$; A and B being two independent systems.

In order to shed some light on this physics dilemma, we propose a simple statistical thermodynamic model where moving particles of known type are assumed to form an elastic specimen (or body). It is shown that this type of approach leads to reveal intrinsic molecularmechanical properties on classical and nonextensive dynamical systems in relation to a distinct tensile force acting on these systems at constant volume and number of particles. In particular we derive the entropy of Eq.[\(1\)](#page-1-0) and relate it to Newton's second law $\vec{F} = m\vec{a}$. A new scenario for the entropic index q in Tsallis statistics in terms of the energy of the system is proposed.

II. FORMALISM

Let us consider a system of non-interacting moving particles and let p_i be the probability that the system is in the microstate $i(= 1, \dots, W)$ given by the product of two positive functions u_i and w_i and satisfying the normalization condition

$$
\sum_{i=1}^{W} p_i = \sum_{i=1}^{W} u_i w_i = 1 \quad . \tag{4}
$$

Derivative of this relation with respect to changes in particle displacements x , considering particles of mass m to move at velocity $v = \frac{\partial x}{\partial t}$ and acceleration $a = \frac{\partial v}{\partial t}$, gives

$$
\sum_{i=1}^{W} \frac{\partial p_i}{\partial x} = \sum_{i=1}^{W} p_i \frac{\partial}{\partial x} \ln \ p_i = 0 \tag{5}
$$

or, alternatively,

$$
\sum_{i=1}^{W} \left(\frac{\partial u_i}{\partial x}\right) \frac{p_i}{u_i} = -\sum_{i=1}^{W} \left(\frac{\partial w_i}{\partial x}\right) \frac{p_i}{w_i} \quad . \tag{6}
$$

Summing up both sides by $\sum_{i=1}(\frac{\partial p_i}{\partial x})$ ln u_i it is straightforward to show that

$$
\frac{\partial}{\partial x} < \ln u_i \rangle = -\frac{\partial}{\partial x} < \ln w_i \rangle + \sum_{i}^{W} \left(\frac{\partial p_i}{\partial x}\right) \ln p_i \quad . \tag{7}
$$

Let us also consider the *purely* thermodynamic relation for the Helmholtz free energy $A = E - TS$, assuming the temperature T and volume V to be independent variables with E the internal energy, A the Helmholtz free energy and S the entropy of the system. Therefore from Eq.[\(7\)](#page-2-0) we can readily identify (or, alternatively, define) three quantities

$$
A/k_B T \rightarrow \langle \ln u_i \rangle = \sum_{i=1}^{W} p_i \ln u_i , \qquad (8)
$$

$$
E/k_B T \to -\langle \ln w_i \rangle = -\sum_{i=1}^{W} p_i \ln w_i , \qquad (9)
$$

$$
-\frac{\partial}{\partial x}(S/k_B) \to \sum_{i=1}^{W} (\frac{\partial p_i}{\partial x}) \ln p_i , \qquad (10)
$$

with k_B a constant added to have reduced units. The choice of the sign for E is to approximate the system energy as the sum of individual contributions (or mean energy) $E \to E \to E \geq \sum_i p_i \epsilon_i$, where $\epsilon_i / k_B T = -\ln w_i$ represents their energy states, and thus obtain

$$
w_i = \exp(-\epsilon_i/k_B T) \ge 0 \quad . \tag{11}
$$

This relation is also found to be the solution of the differential equation: $\frac{\partial \ln (p_i/u_i)}{\partial (\epsilon_i/k_BT)} = -1$.

Furthermore from Eqs.[\(5\)](#page-2-1) and [\(10\)](#page-2-2) it follows immediately that the entropy satisfies

$$
\frac{\partial}{\partial x}(S/k_B) = -\frac{\partial}{\partial x}\left(\sum_{i=1}^{W} p_i \ln p_i\right) , \qquad (12)
$$

which is the basic result stated by the BG entropy of Eq.[\(1\)](#page-1-0).

Since the moving particles are assumed to form an elastic body, then the total work along an infinitesimal elongation $d\ell$ is given by $d\mathcal{W} = dA = f_{\ell} d\ell$ for a change at constant T, with f a (compressive < 0 , or expansive > 0) tensile force in the direction of ℓ . This relation is obtainable directly from the first and second laws of thermodynamics in a reversible isothermal process [\[6\]](#page-6-2), assuming that there is not another force acting on the system in addition to f_{ℓ} as for example due to hydrostatic pressure $P(dW) = -PdV \approx 0$; recall volume V here is an independent variable). Therefore, the dynamics of our particle system relating thermodynamic quantities reduces to

$$
F = \left(\frac{\partial A}{\partial x}\right)_T = k_B T \frac{\partial}{\partial x} \left(\sum_{i=1}^W p_i \ln u_i\right)_T = \left(\frac{\partial E}{\partial x}\right)_T - T \left(\frac{\partial S}{\partial x}\right)_T \quad . \tag{13}
$$

Thus through Eqs.[\(8\)](#page-2-2) and [\(9\)](#page-2-2) the sytem entropy can be correlated to Newton's law of motion in relation to a distinct tensile force acting on the system.

III. REMARKS

At a first glance the choice for p_i in Eq.[\(4\)](#page-2-3) may be seen quite arbitrary, but in view of the results derived for S, A and E starting only from this quantity it seems to be reasonable. As shown below, this choice can further be justified in relation with the measure for an inverse temperature for the system. Using Eqs. (4) , (9) , (11) and (12) , we obtain at once

$$
\frac{\Delta(S/k_B)}{\Delta(E/k_B T)} = 1 + \frac{\ln u_w}{\ln w_w} = 1 - \left(\frac{k_B T}{\epsilon_w}\right) \ln u_w \quad , \tag{14}
$$

where Δ represents functions difference between consecutive microstates fluctuations.

Although we do not know how each of our u_i and w_i functions may depend explicitly on the displacement x , we can still estimate all thermodynamic relations from given average values of either u or w at each state i. For example, interesting to note are the results obtained when setting $u_i = 1/Z$, a sort of mean value for $u(\forall i)$ and w_i variable. From Eqs.[\(4\)](#page-2-3) and [\(11\)](#page-2-4), it follows then that $p_i = \frac{e^{-\epsilon_i/k_B T}}{Z}$ $\frac{Z}{Z}$, which corresponds to the Gibbs distribution in

thermal equilibrium and satisfies $\frac{dp_i}{d\epsilon_i} = -\frac{p_i}{k_B}$ $\frac{p_i}{k_B T}$. Using the normalization condition for p_i we derive in turn the partition function $Z = \sum_i e^{-\epsilon_i/k_B T}$, where the summation is over all microstates of the system. From this result and that of the energy states ϵ_i plus Eq.[\(9\)](#page-2-2), we obtain a relation for the mean energy of a system at temperature T as $\lt E \gt = -\left(\frac{\partial \ln Z}{\partial (1/k_B)}\right)$ $\partial (1/k_BT)$. For the Helmholtz free energy of Eq.[\(8\)](#page-2-2) we can write $A/k_BT = -\ln Z$, which is also a wellknown classical statistical mechanics result [\[3\]](#page-5-2).

Let us analyse next a second case, *i.e.*, the case of having an averaged $w(\forall i)$ and u_i as the variable. To derive Tsallis statistics we assume the thermal energy of the particles k_BT to be proportional to their energy states ϵ by the nonextensivity (integer) factor $q-1$ for all the *i*-microstates. From Eq.[\(11\)](#page-2-4) and $q \neq 1$ this is equivalent to consider

$$
\ln w_i = \frac{1}{1-q} \quad . \tag{15}
$$

Using the ln_q definition of Eq.[\(3\)](#page-1-1) in conjunction with Eqs.[\(4\)](#page-2-3) and [\(15\)](#page-4-0), we then get the relation $-\ln_q(1/p_i) - \ln(p_i/u_i) = \frac{p_i^{q-1}}{q-1}$. For $q \to 1$ and by Taylor expansion [\[2](#page-5-1)] we can also approximate $p_i^q = p_i p_i^{-(1-q)} = p_i exp(\ln p_i^{-(1-q)})$ $\binom{n}{i} \approx p_i[1 + (1 - q) \ln p_i^{-1}]$. Thus in this limit $\ln_q p_i^{-1} \approx \ln p_i^{-1}$. These results imply that

$$
\ln u_i \approx \frac{p_i^{q-1}}{q-1} \quad . \tag{16}
$$

Therefore within our statistical thermodynamic approach it follows that

$$
A/k_B T \approx \sum_{i=1}^{W} p_i \left(\frac{p_i^{q-1}}{q-1}\right) , \quad E/k_B T \approx -\sum_{i=1}^{W} p_i \left(\frac{1}{1-q}\right) = \epsilon/k_B T , \qquad (17)
$$

whence

$$
S/k_B = E/k_B T - A/k_B T \approx \frac{1 - \sum_{i=1}^{W} p_i^q}{q - 1} \quad , \tag{18}
$$

which corresponds to the entropy term of $Eq.(2)$ $Eq.(2)$ introduced by Tsallis. As in Ref.[\[2\]](#page-5-1), if we assume equiprobability $p_i \equiv 1/W$ ($\forall i$) we also obtain $S = \ln_q W$.

Possible values of the q index are constrained by the logarithm functions of Eqs.[\(15\)](#page-4-0) and [\(16\)](#page-4-1). The condition $0 \le p_i \equiv u_i w_i \le 1$ ($\forall i$) implies that $0 \le exp(1/(1-q)) exp(p_i^{q-1}/(q-1))$ 1)) \leq 1. Hence we can deduce that $\Theta_i(q) \equiv (p_i^{q-1} - 1)/(q - 1) \leq 0$, due to the positive and continue nature of the exponential function. Then the limiting behaviour of this function as $q \to 1$ is obtained by L'Hôpital's rule because this is an indeterminate form of type $\frac{\infty}{\infty}$. We found $\lim_{q\to 1} \Theta_i(q) = \ln p_i$, hence $0 \leq exp(\ln p_i) \leq 1$. From this result we also

have $0 \le p_i^{q-1} \le 1$ ($\forall i, \forall q$), which via Eq.[\(15\)](#page-4-0) it implies that $\ln w_i \le \Theta_i(q) \le 0$. This, in turn, implies a positive w_i in agreement with Eq.[\(11\)](#page-2-4) such that $0 < w_i \leq 1$. Therefore we can conclude that the range of validity of our statistical thermodynamic model for Tsallis statistics is $q \in (\pm \infty, 1]$.

On the other hand, using Eqs.[\(15\)](#page-4-0) and [\(16\)](#page-4-1) for (u,w) at the upper microstate $i = W$, Eq.[\(14\)](#page-3-1) for $q \neq 1$ takes the form

$$
\frac{\Delta(S/k_B)}{\Delta(E/k_B T)} \approx 1 - p_w^{q-1} \quad . \tag{19}
$$

This implies that the thermodynamic definition for the temperature at constant volume and number of particles $T = \left(\frac{\partial E}{\partial S}\right)_{V,N}$ follows for $q \gg 1$ since, in physical terms, $p_w \leq 1$ for large *subextensive* systems. For *superextensive* systems with integers $q < 1$, this relation becomes negative. For $q = 1$, from Eq.[\(14\)](#page-3-1) we readily found $\frac{\Delta(S/k_B)}{\Delta(E/k_B T)} = 1$ as should be expected since $S_{q=1}$ recovers S_{BG} .

Finally for the force on the system within Tsallis statistics we have

$$
F = k_B T \frac{\partial}{\partial x} \left(\frac{\sum_{i=1}^{W} p_i^q}{q - 1} \right)_T = ma \quad , \tag{20}
$$

which gives a physical significance to the q-values representing complex systems and correlates them to the system dynamics.

In summary, the new insights gained by our statistical thermodynamic approach of moving particles forming an elastic body are on the molecular-mechanical properties derived for classical and nonextensive dynamical systems including a possible correlation between force and entropy in the system. Our initial factorization for $p_i = u_i w_i$ may still be seen as rather arbitrary, but we have shown that this simple multiplicative form leads to interesting connections between an applied stretching force (tension) and thermodynamics quantities of dynamical systems [\[7\]](#page-6-3). Such class of normalized product of positive functions appears formally, e.g. in the analysis of stochastic processes on graphs according to the Hammersley-Clifford Theorem.

- [1] E.G.D. Cohen, Phys. A 305, 19 (2002).
- [2] C. Tsallis, [arXiv:cond-mat/0312500.](http://arxiv.org/abs/cond-mat/0312500)
- [3] F. Mandl, "Statistical Physics" (John Wiley & Sons, 2nd Edition, 1988).
- [4] F. Baldovin, L.G. Moyano and C. Tsallis, [arXiv:cond-mat/0402635.](http://arxiv.org/abs/cond-mat/0402635)
- [5] C. Tsallis, J. Stat. Phys. 52, 479 (1988).
- [6] L.R.G. Treloar, "The Physics of Rubber Elasticity" (Clarendon Press, Oxford, 1975).
- [7] This work was inspired on preliminary ideas discussed by the Author in Phys. A 328, 44 (2003), where set of numbers played the role of classical particles.