Is a "homogeneous" description of dynamic heterogeneities possible?

Grzegorz Szamel

Department of Chemistry, Colorado State University, Fort Collins, CO 80525

(Dated: February 8, 2020)

We study the simplest model of dynamic heterogeneities in glass forming liquids: one-spin facilitated kinetic Ising model introduced by Fredrickson and Andersen [G.H. Fredrickson and H.C. Andersen, Phys. Rev. Lett. **53**, 1244 (1984); J. Chem. Phys. **83**, 5822 (1985)]. We show that the low-temperature, long-time behavior of the density autocorrelation function predicted by a scaling approach can be obtained from a self-consistent mode-coupling-like approximation.

PACS numbers: 64.70.Pf, 64.60.Ht, 64.60.Cn, 05.50+q

Recently, a new general paradigm for dynamics of supercooled liquids has been proposed¹. It focuses directly on the spatial and temporal structure of dynamic heterogeneities, *i.e.* regions of higher-than-average mobility. The essence of this new approach is a mapping of a microscopic model of a supercooled liquid onto an effective model describing dynamics of the heterogeneities². It has been argued on physical grounds that very simple effective models can capture the most important features of the heterogeneities' dynamics. In the initial work¹ two models were used: a one-dimensional one-spin facilitated kinetic Ising model, *i.e.* the Fredrickson-Andersen (FA) $model^3$, and its asymmetric version, *i.e.* the East $model^4$. In the subsequent work⁵, more complicated (and more realistic) models were used. However, it is the simplest model (*i.e* the FA model) that is most often used^{6,7} to illustrate the main points of the new paradigm and to contrast it with other approaches to the glass transition problem like the mode-coupling theory⁸ and the landscape paradigm⁹.

The simple models describing dynamic heterogeneities have trivial static properties. However, they exhibit interesting dynamics. Hence, there arises a new problem: how to study the dynamics of the effective models. One should notice that solving this problem will be, most likely, considerably simpler than solving the original, microscopic model of a supercooled liquid: the effective models used so far are lattice-based and thus have a greatly reduced number of degrees of freedom; even more importantly, these models usually involve a small parameter (*i.e.* density of heterogeneities (mobile regions)) which can facilitate theoretical analysis.

The original work¹ concentrated on the analysis of the structure of trajectory spaces of the FA and East models; it was argued that the glass transition coincides with an entropy crisis in trajectory space rather than in configuration space. This is an ingenious suggestion because it focuses directly on the system's dynamics and thus it should be applicable to other glass-like systems with trivial equilibrium properties^{10,11}.

In Ref.¹, in order to get explicit results for dynamics of the FA and East models, the random walk theory was combined with scaling arguments. Subsequently² a dynamic field-theoretical model has been derived and analyzed using the renormalization group (RG) methods. It was argued that the *d*-dimensional FA model belongs to the same universality class as the field-theoretical model. RG analysis showed that, for d > 2, there is no finite temperature phase transition and the low temperature scaling properties are determined by the directed percolation critical point.

In our opinion, the simplicity of the effective models suggests that it might be possible to analyze their dynamics using well-known methods of non-equilibrium statistical mechanics. Such an approach was utilized by Fredrickson and Andersen³: the time-dependent density correlation function was expressed in terms of a memory function; the latter function was then analyzed using an exact diagrammatic expansion¹². The resulting self-consistent equation for the density correlation function predicted a power law dependence of the relaxation time on the density of heterogeneities. However, neither Ref.³ nor later work on the FA model¹³ reproduced correct scaling exponent for the relaxation time. This was, perhaps, the reason for the recent claim⁷ that no "homogeneous", mean-field¹⁴ approach is able to reproduce scaling predictions for the FA model.

In this note we present a simple theory that does reproduce previously predicted scaling behavior¹⁵ of the FA model density correlation function. The theory relies upon a self-consistent factorization approximation of a pair-density correlation function into a product of two density correlations. This approximation is analogous to one used in the mode-coupling theory⁸; the important difference is that we use the factorization approximation for pair-density involving non-nearest-neighbors. We argue that while this makes the structure of the theory somewhat complicated, it is necessary from a physical point of view since the dynamics of nearest-neighbor up-spins are drastically different from that of individual spins.

The theory presented here shows that it might be possible to use established "homogeneous" methods to analyze dynamics of effective models of heterogeneities.

The FA model consists of a chain of N non-interacting Ising spins in a unit magnetic field. We use density variables n_i ($n_i = 1$ corresponds to *i*th spin pointing up; physically, it corresponds to a more mobile region located near site *i*). In terms of n_i the Hamiltonian is $H = \sum_i n_i$. The average density is $\langle n_i \rangle \equiv c = 1/(1 + \exp(1/T))$ where T is the temperature.

We consider the normalized time-dependent density correlation function, $C_{i,j}(t)$,

$$C_{i,j}(t) = \left\langle \delta n_i \exp\left(\Omega t\right) \delta n_j \right\rangle / \left\langle \left(\delta n_i\right)^2 \right\rangle \tag{1}$$

where δn_i is the deviation of the density at site *i* from its equilibrium average, $\delta n_i = n_i - c$; note that $C_{i,j}(t = 0) = \delta_{i,j}$. In Eq. (1) Ω is the evolution operator; time evolution consists of single spin-flip dynamics with rate that is proportional to the number of nearest neighbors in the spin up state. Explicitly,

$$\Omega = -\sum_{i=1}^{N} m_i \left(1 - S_i \right) w_i \tag{2}$$

where S_i is the spin-flip operator, $S_i n_i = -n_i + 1$, m_i is the number of nearest neighbor up-spins, $m_i = n_{i+1} + n_{i-1}$, and w_i is the factor maintaining detailed balance, $w_i = c + n_i(1 - 2c)$.

Some earlier simulational studies¹⁶ and recent computational and theoretical works^{1,6,7} used the so-called persistence function to characterize dynamics of the FA model. The persistence function is defined as a fraction of spins that remain unflipped after a given time. While this function is easier to determine from simulations than the density correlation function, it is somewhat awkward to analyze theoretically because it is defined via a timenonlocal constraint.

The starting point of our analysis is the memory function representation of the density correlations function,

$$\sum_{j} \left(\delta_{i,j} + M_{i,j}^{1irr}(z) \left\langle \delta n_{j} m_{j} \delta n_{j} \right\rangle^{-1} \right) \left(z C_{j,k}(z) - \delta_{j,k} \right)$$
$$= \left\langle \delta n_{i} \Omega \delta n_{i} \right\rangle \left\langle \delta n_{i} \delta n_{i} \right\rangle^{-1} C_{i,k}(z) \tag{3}$$

Here $C_{i,j}(z)$ is the Laplace transform of $C_{i,j}(t)$, $C_{i,j}(z) = \int_0^\infty dt \exp(-zt)C_{i,j}(t)$, and $M_{i,j}^{1irr}(z)$ is the Laplace transform of the irreducible^{17,18} memory function. The memory function can be expressed in terms of the pairdensity correlation function involving nearest neighbor spins,

$$M_{i,j}^{1irr}(t) = A_{i,j}(t) + A_{i-1,j}(t) + A_{i,j-1}(t) + A_{i-1,j-1}(t).$$
(4)

Note that if the $M_{i,j}^{1irr}$ is neglected, the density correlation function $C_{i,j}$ is diagonal; motion of the isolated upspins is possible only due to the coupling to pair-densities.

The pair density correlation function $A_{i,j}(t)$ evolves with 1-spin irreducible evolution operator Ω^{1irr} ,

$$A_{i,j}(t) = \left\langle \delta n_i \delta n_{i+1} \exp\left(\Omega^{1irr} t\right) \delta n_j \delta n_{j+1} \right\rangle \tag{5}$$

where

$$\Omega^{1irr} = \hat{Q}_1 \left[\Omega + \sum_i m_i \delta n_i \rangle \left\langle \delta n_i m_i \delta n_i \right\rangle^{-1} \left\langle \delta n_i m_i \right] \hat{Q}_1.$$
(6)

In Eq. (6) $\hat{Q}_1 = 1 - \hat{P}_1$, and \hat{P}_1 is a projection operator on the one-spin density subspace,

$$\hat{P}_{1} = \sum_{i} \delta n_{i} \rangle \left\langle (\delta n_{i})^{2} \right\rangle^{-1} \langle \delta n_{i} \tag{7}$$

It should be noted at this point that the standard sequence of approximations leading to a mode-coupling theory⁸ would start from Eq. (5). Briefly, the pairdensity correlation function would be factorized into a product of two density correlations, and the irreducible operator Ω^{1irr} would be replaced by Ω^{19} . However, a closer investigation shows that the time dependence of $A_{i,j}$ is very different from that of the product of $C_{i,j}$ s. In particular, if in an equation of motion for $A_{i,j}$ one neglects the memory function term, one gets a (lattice) diffusion equation. This is in contrast to Eq. (3) that, upon neglecting the memory function term, leads to a site-diagonal density correlation function.

The strategy followed in this note is different: we propose to continue using the memory function approach until we generate a function that can be expressed in terms of pair-densities involving non-nearest neighbors, in terms of e.g. $\delta n_i \delta n_{i+l}$, $l \geq 2$. With this goal in mind, we derive a memory function representation for the pair-density correlation function, $A_{i,j}$. It turns out that the 2-spin irreducible memory function that enters the equation of motion for $A_{i,j}$ can be expressed in terms of the triple-density correlation function, $B_{i,j}$,

$$M_{i,j}^{2irr}(t) = B_{i,j}(t) + B_{i-1,j}(t) + B_{i,j-1}(t) + B_{i-1,j-1}(t),$$
(8)

where $B_{i,j}$ is defined as

$$B_{i,j}(t) = \left\langle \delta n_i \delta n_{i+1} \delta n_{i+2} \exp\left(\Omega^{2irr} t\right) \delta n_j \delta n_{j+1} \delta n_{j+2} \right\rangle,$$
(9)

and the 2nd order irreducible evolution operator Ω^{2irr} is given by a formula analogous to (6). It should be noted that Ω^{2irr} involves operators \hat{Q}_2 projecting on the space orthogonal to the density and to the pair-density involving *nearest neighbor* sites, $\hat{Q}_2 = 1 - \hat{P}_1 - \hat{P}_2$. Here \hat{P}_1 is given by Eq. (7) and \hat{P}_2 is defined as

$$\hat{P}_2 = \sum_{i} \delta n_i \delta n_{i+1} \rangle \left\langle \left(\delta n_i \delta n_{i+1} \right)^2 \right\rangle^{-1} \langle \delta n_i \delta n_{i+1}. \quad (10)$$

Next, we derive the memory function representation for $B_{i,j}$. Its irreducible memory function is given by the following expression

$$M_{i,j}^{3irr}(t) = \left\langle \delta n_i \delta n_{i+1} \delta n_{i+2} \left(m_{i+2} + m_i \right) \hat{Q}_3 \right.$$
$$\left. \exp\left(\Omega^{3irr} t \right) \, \hat{Q}_3 \left(m_{j+2} + m_j \right) \delta n_j \delta n_{j+1} \delta n_{j+2} \right\rangle, (11)$$

where $\hat{Q}_3 = 1 - \hat{P}_1 - \hat{P}_2 - \hat{P}_3$, and \hat{P}_3 is defined as

$$\hat{P}_{3} = \sum_{i} \delta n_{i} \delta n_{i+1} \delta n_{i+2} \left\langle \left(\delta n_{i} \delta n_{i+1} \delta n_{i+2} \right)^{2} \right\rangle^{-1} \\ \left\langle \delta n_{i} \delta n_{i+1} \delta n_{i+2} \right\rangle.$$
(12)

Finally, the 3rd order irreducible evolution operator Ω^{3irr} in Eq. (11) is given by a formula analogous to (6).

At this point we note that $M_{i,j}^{3irr}$ is a correlation function of a quantity that has a non-zero overlap with the pair-density involving non-nearest neighbor sites:

$$\left\langle \delta n_i \delta n_{i+2} \hat{Q}_3 \left(m_{i+2} + m_i \right) \delta n_i \delta n_{i+1} \delta n_{i+2} \right\rangle$$

= 2 (c(1-c))³. (13)

Thus, it is at this stage that we propose to follow the usual sequence of approximations⁸. As a result we get the following expression for the memory function $M_{i,j}^{3irr}$

in terms of the density correlations $C_{i,j}$ s:

$$M_{i,j}^{3irr}(t) \approx 4 \left(c(1-c) \right)^4 C_{i,j}(t) C_{i+2,j+2}(t).$$
 (14)

The final self-consistent equation for the density correlation function is easiest to present in terms of the Laplace-Fourier transform, C(q; z),

$$C(q;z) = \sum_{k} \exp(-iqk) C_{k,0}(z).$$
 (15)

It reads

$$C(q;z) = \frac{1}{z + \frac{2c}{1 + \frac{2c}{1 + \frac{2(1-c)\cos^2(q/2)}{z + 2(1-c)\sin^2(q/2) + \frac{2c}{1 + \frac{2(1-c)\cos^2(q/2)}{z + 2(1-c)\cos^2(q/2) + \frac{2}{1 + 2c(1-c)\phi_2(q;z)}}}},$$
(16)

where $\phi_2(q; z)$ is the Fourier-Laplace transform of $C_{i,j}(t)C_{i+2,j+2}(t)$.

To investigate the low temperature (*i.e.* low c) behavior of the density correlation function we consider the scaling function $\tilde{C}(q^*; z^*)$,

$$\tilde{C}(q^*; z^*) = \lim_{c \to 0} c^3 C(q^* c; z^* c^3) = \int_0^\infty dt^* \exp(-z^* t^*) \tilde{C}(q^*; t^*) \quad (17)$$

where $\tilde{C}(q^*;t^*) = \lim_{c\to 0} C(q^*c;t^*c^{-3})$. It can be easily shown that in the limit $c \to 0$ the self-consistent equation (16) reduces to the following *c*-independent equation for the scaling function $\tilde{C}(q^*;z^*)$:

$$\tilde{C}(q^*; z^*) = \frac{1}{z^* + q^{*2}/2 + 1/\tilde{\phi}_2(q^*; z^*)}.$$
 (18)

Here $\tilde{\phi}_2(q^*; z^*)$ is the Laplace transform of $\tilde{\phi}_2(q^*; t^*)$,

$$\tilde{\phi}_{2}(q^{*};t^{*}) = \lim_{c \to 0} c^{-1} \phi_{2}(q^{*}c;t^{*}c^{-3})$$
$$= \int_{-\infty}^{\infty} \frac{dk^{*}}{2\pi} \tilde{C}(k^{*};t^{*}) \tilde{C}(q^{*}-k^{*};t^{*}) \quad (19)$$

The scaling (17) and *c*-independence of Eq. (18) shows that, in the low *c* limit, up-spins (*i.e.* mobile regions) undergo diffusive motion with a diffusion constant $D \propto c$; up-spin diffusion is cut-off at long times by a longest relaxation time $\tau \propto c^{-3}$. These results agree with known low *c* scaling predictions¹⁵. In order to get quantitative results for the diffusion constant and the relaxation time one would have to solve the scaling equation (18) combined with Eq. (19). The numerical solution of Eqs. (18-19) and a comparison of the resulting density correlation function with computer simulations are left for a future study. Here we note that a single iteration of Eqs. (18-19) starting from a zeroth order approximation $\tilde{C}^{(0)}(q^*; z^*) = 1/(z^* + aq^{*2}/2 + b)$ leads to

$$\tilde{C}^{(1)}(q^*; z^*) = \frac{1}{z^* + q^{*2}/2 + (4az^* + 8ab + a^2q^{*2})^{1/2}}.$$
(20)

Whereas the zeroth order approximation $\tilde{C}^{(0)}$ has a simple pole, the first order approximation $\tilde{C}^{(1)}$ has a pole and a branch cut. The simplest self-consistency condition that allows one to estimate a and b (and thus the diffusion constant and the longest relaxation time, respectively) is to identify the pole in $\tilde{C}^{(0)}$ and the one in $\tilde{C}^{(1)}$. This leads to the following approximate results for the diffusion constant²⁰ and the longest relaxation time: $D = 0.4c, \tau = 0.3125c^{-3}$.

One should note that the structure of Eq. (18) is similar to that of the self-consistent equation of the modecoupling theory of the glass transition⁸. The very important difference is the $q^{*2}/2$ term in the denominator. It is this term that cuts off the ergodicity-breaking transition that is generic for mode-coupling theories.

In summary, we have shown here that a simple selfconsistent, mode-coupling-like approximation reproduces known scaling predictions for the density correlation function of the FA model. It would be of interest to investigate whether a similar approach can be applied to more complicated models like the East model or the Kob-Andersen lattice gas model²¹, which recently have attracted renewed interest^{1,22}.

Note added: After this paper had been submitted a preprint presenting an RG approach to the onedimensional FA model appeared²³.

Support by NSF Grant No. CHE-0111152 is gratefully acknowledged.

- ¹ J.P. Garrahan and D. Chandler, Phys. Rev. Lett. **89**, 035704 (2002).
- ² S. Whitelam, L. Berthier, and J.P. Garrahan, Phys. Rev. Lett. **92**, 185705 (2004).
- ³ G.H. Fredrickson and H.C. Andersen, Phys. Rev. Lett. 53, 1244 (1984); J. Chem. Phys. 83, 5822 (1985).
- ⁴ J. Jäckle and S. Eisinger, Z. Phys. B **84**, 115 (1991).
- ⁵ J.P. Garrahan and D. Chandler, Proc. Natl. Acad. Sci. 100, 9710 (2003).
- ⁶ L. Berthier and J.P. Garrahan, J. Chem. Phys. **119**, 4367 (2003).
- ⁷ L. Berthier and J.P. Garrahan, Phys. Rev. E 68, 041201 (2003).
- ⁸ W. Götze, in *Liquids, Freezing and the Glass Transition*, J.P. Hansen, D. Levesque, and J. Zinn-Justin, eds. (North-Holland, Amsterdam, 1991).
- ⁹ P.G. Debenedetti and F.H. Stillinger, Nature (London) 410, 259 (2001).
- ¹⁰ C. Renner, H. Löwen, and J.-L. Barrat, Phys. Rev. E 52, 5091 (1995).
- ¹¹ R. Schilling and G. Szamel, Europhys. Lett. **61**, 207 (2003).
- ¹² Diagrammatic approach has been also used to analyze dynamics of the East model: S.J. Pitts and H.C. Andersen, J. Chem. Phys. **114**, 1101 (2001).
- $^{13}\,$ J. Reiter and J. Jäckle, Physica A $\mathbf{215},\,311$ (1995).
- ¹⁴ Note that "mean-field" is often meant to be synonymous with "mode-coupling". This identification is based on re-

sults obtained for fully connected spin systems. See, e.g., L.F. Cugliandolo, in *Slow Relaxations and Nonequilibrium Dynamics in Condensed Matter*, J.-L. Barrat *et al.*, eds. (EDP Sciences-Springer Verlag, 2003), and references therein.

- ¹⁵ For a recent review see, *e.g.*, F. Ritort and P. Sollich, Adv. Phys. **52**, 219 (2003).
- ¹⁶ S. Butler and P. Harrowell, J. Chem. Phys. **95**, 4454 (1991).
- ¹⁷ K. Kawasaki, Physica A **215**, 61 (1995).
- ¹⁸ S. Pitts and H.C. Andersen, J. Chem. Phys. **113**, 3945 (2000).
- ¹⁹ Such a procedure was applied to the East model in Ref.¹⁷. For an extended discussion of two different mode-coupling approximations for the East model see S.J. Pitts, T. Young, and H.C. Andersen, J. Chem. Phys. **113**, 8671 (2000) and Ref.¹².
- ²⁰ It is has been claimed¹³ that the exact low density result for the diffusion constant is D = c/2. We believe that a more sophisticated self-consistent approximation may reproduce the exact value. This point is left for a future study.
- ²¹ W. Kob and H.C. Andersen, Phys. Rev. E **48**, 4364 (1993).
- ²² C. Toninelli, G. Biroli, and D. Fisher, Phys. Rev. Lett. **92**, 185504 (2004).
- 23 S. Whitelam and J.P. Garrahan, cond-mat/0405647.