

# Bose-Einstein condensation

V.I. Yukalov<sup>1,2</sup>

<sup>1</sup>*Bogolubov Laboratory of Theoretical Physics,  
Joint Institute for Nuclear Research, Dubna 141980, Russia*

<sup>2</sup>*Institut für Theoretische Physik,  
Freie Universität Berlin, Arnimallee 14, D-14195 Berlin, Germany*

Bose-Einstein condensation is the *occupation of a single quantum state by a large number of identical particles*. This implies that particles are assumed to be bosons, satisfying the Bose-Einstein statistics allowing for many particles to pile up in the same quantum state. This is contrary to fermions, satisfying the Fermi-Dirac statistics, for which the Pauli exclusion principle does not allow the occupation of any single quantum state by more than one particle.

The role of quantum correlations, caused by the Bose-Einstein statistics, is crucial for the occurrence of BEC. This statistics was advanced by Bose (1924) for photons, having zero mass, and generalized by Einstein (1924) to particles with nonzero masses. Einstein (1925) also described the phenomenon of condensation in ideal gases. The possibility of BEC in weakly nonideal gases was theoretically demonstrated by Bogolubov (1947). The wave function of Bose-condensed particles in dilute gases satisfies the Gross-Pitaevskii equation, suggested by Gross (1961) and Pitaevskii (1961). Its mathematical structure is that of the nonlinear Schrödinger equation. Experimental evidence of BEC in weakly interacting confined gases was achieved, seventy years after the Einstein prediction, almost simultaneously by three experimental groups (Anderson *et al.*, 1995; Bradley *et al.*, 1995; Davis *et al.*, 1995).

To say that many particles are in the same quantum state is equivalent to saying that these particles display the *state coherence*. That is, BEC is a particular case of *coherence phenomena* related to the arising state coherence. For the latter to occur, the particles are to be strongly correlated with each other. The expected conditions, when such a strong correlation could appear, may be qualitatively understood applying the de Broglie duality arguments to an ensemble of atoms in thermodynamic equilibrium at temperature  $T$ . Then the thermal energy of an atom is given by  $k_B T$ , where  $k_B$  is the Boltzmann constant. This energy defines the thermal wavelength

$$\lambda_T \equiv \sqrt{\frac{2\pi\hbar^2}{m_0 k_B T}} \quad (1)$$

for an atom of mass  $m_0$ , with  $\hbar$  being the Planck constant. Thus, an atom can be associated with a matter wave characterized by the wavelength (1). Atoms become correlated with each other when their related waves overlap, which requires that the wavelength (1) be larger than the mean interatomic distance,  $\lambda_T > a$ . The average atomic density  $\rho \equiv N/V$  for  $N$  atoms in volume  $V$  is related to the mean distance  $a$  through the equality  $\rho a^3 = 1$ . Hence, condition  $\lambda_T > a$  may be rewritten as  $\rho \lambda_T^3 > 1$ . With the thermal wavelength (1), this yields the inequality

$$T < \frac{2\pi\hbar^2}{m_0 k_B} \rho^{2/3}, \quad (2)$$

which tells that state coherence may develop if temperature is sufficiently low or the density of particles is sufficiently high.

An accurate description of BEC for an ideal gas is based on the *Bose-Einstein distribution*

$$n(\mathbf{p}) = \left[ \exp\left(\frac{\varepsilon_p - \mu}{k_B T}\right) - 1 \right]^{-1}, \quad (3)$$

describing the density of particles with a single-particle energy  $\varepsilon_p = p^2/2m_0$  for a momentum  $\mathbf{p}$  and with a chemical potential  $\mu$ . The latter is defined from the condition  $N = \sum_p n(\mathbf{p})$  for the total number of particles. Assuming the *thermodynamic limit*

$$N \rightarrow \infty, \quad V \rightarrow \infty, \quad \frac{N}{V} \rightarrow \text{const}$$

allows the replacement of summation over  $\mathbf{p}$  by integration. Then the fraction of particles, condensing to the state with  $\mathbf{p} = 0$ , is

$$n_0 \equiv \frac{N_0}{N} = 1 - \left(\frac{T}{T_c}\right)^{3/2} \quad (4)$$

below the *condensation temperature*

$$T_c = \frac{2\pi\hbar^2 \rho^{2/3}}{m_0 k_B \zeta^{2/3}}, \quad (5)$$

where  $\zeta \approx 2.612$ , while  $n_0 = 0$  above the critical temperature (5). The latter is about twice smaller than the right-hand side of inequality (2).

The condensate fraction (4) is derived for an ideal, that is, noninteracting, Bose gas. A weakly nonideal, i.e., weakly interacting Bose gas, also displays Bose-Einstein condensation, though particle interactions deplete the condensate, so that at zero temperature the condensate fraction is smaller than unity,  $n_0 < 1$ . A system is called weakly interacting if the characteristic interaction radius  $r_{int}$  is much shorter than the mean interparticle distance,  $r_{int} \ll a$ . This inequality can be rewritten as  $\rho r_{int}^3 \ll 1$ , because of which such a system is termed dilute.

Superfluid liquids, such as liquid  $^4\text{He}$ , are far from being dilute. Nevertheless, one commonly believes that the phenomenon of superfluidity is somehow connected with BEC, although an explicit relation between the superfluid and condensate fractions is not known. Theoretical calculations and experimental observations for superfluid helium estimate the condensate fraction at  $T = 0$  as  $n_0 \approx 0.1$ .

A strongly correlated pair of fermions can approximately be treated as a boson. This is why the arising superfluidity in liquid  $^3\text{He}$  can be interpreted as the condensation of coupled fermions. Similarly, superconductivity is often compared with the condensation of the Cooper pairs that are formed by correlated electrons or holes. However, one should understand that superconductivity of fermions is analogous but not identical to BEC of bosons.

An ideal object for the experimental observation of BEC is a dilute atomic Bose gas confined in a trap and cooled down to temperatures satisfying condition (2). Such experiments with different atomic gases have been recently realized, BEC explicitly observed, and

a variety of its features carefully investigated. It has been demonstrated that the system of Bose-condensed atoms displays a high level of state coherence.

There exist different types of traps, magnetic, optical, and their combinations, which makes it possible to confine atoms for sufficiently long times of up to 100 s. There are single-well and double-well traps. Employing a standing wave of laser light, one forms a multi-well periodic effective potentials called *optical lattices*. The use of the latter has allowed the demonstration of a number of interesting effects, as Bloch oscillations, Landau-Zener tunneling, Josephson current, Wannier-Stark ladders, Bragg diffraction, and so on.

An ensemble of Bose-condensed atoms, displaying a high level of state coherence, forms a matter wave that is analogous to a coherent electromagnetic wave from a laser. Therefore a device emitting a coherent beam of Bose atoms is named *atom laser*.

The realization of BEC of dilute trapped gases is of great importance because of several reasons. First of all, this explicitly proved the actual existence of the phenomenon predicted by Einstein many years ago. It is worth emphasizing that a direct observation of BEC in superfluid helium, despite enormous experimental efforts, have never been achieved. This is why such an unambiguous observation of BEC of trapped atoms is of great value. Second, dilute atomic gases are sufficiently simple statistical systems, which can serve as a touchstone for testing different theoretical approaches. Finally, Bose-condensed trapped gases display such an enormous variability of their properties that they will certainly find diverse practical applications.

*See also* Coherence phenomena; Critical phenomena; Lasers; Nonequilibrium statistical mechanics; Nonlinear optics; Nonlinear Schrödinger equations; Order parameters; Pattern formation; Phase transitions; Quantum nonlinearity; Quantum theory; Superconductivity; Superfluidity

### Further Reading

Anderson, M.H., Ensher, J.R., Matthews, M.R., Wieman, C.E. and Cornell, E.A. 1995. Observation of Bose-Einstein condensation in a dilute atomic vapor. *Science*, 269: 198–201

Bogolubov, N.N. 1947. On the theory of superfluidity. *Journal of Physics* (Moscow), 11: 23–32

Bose, S.N. 1924. Plancks gesetz und lichtquantenhypothese. *Zeitschrift für Physik*, 26: 178–181

Bradley, C.C., Sackett, C.A., Tollett, J.J. and Hulet, R.G. 1995. Evidence of Bose-Einstein condensation in an atomic gas with attractive interactions. *Physical Review Letters*, 75: 1687–1690

Coleman, A.J. and Yukalov, V.I. 2000. *Reduced Density Matrices*, Berlin: Springer

Courteille, P.W., Bagnato, V.S. and Yukalov, V.I. 2001. Bose-Einstein condensation of trapped atomic gases. *Laser Physics*, 11: 659–800

Dalfovo, F., Giorgini, S., Pitaevskii, L.P. and Stringari, S. 1999. Theory of Bose-Einstein condensation in trapped gases. *Reviews of Modern Physics*, 71: 463–512

- Davis, K.B., Mewes, M.O., Andrews, M.R., van Druten, N.J., Durfee, D.S., Kurn, D.M. and Ketterle, W. 1995. Bose-Einstein condensation in a gas of sodium atoms. *Physical Review Letters*, 75: 3969–3973
- Einstein, A. 1924. Quantentheorie des einatomigen idealen gases. *Sitzungsberichte der Preussischen Akademie der Wissenschaften, Physik-Mathematik*, 1924: 261–267
- Einstein, A. 1925. Quantentheorie des einatomigen idealen gases. Zweite abhandlung. *Sitzungsberichte der Preussischen Akademie der Wissenschaften, Physik-Mathematik*, 1925: 3–14
- Gross, E.P. 1961. Structure of a quantized vortex in boson systems. *Nuovo Cimento*, 20: 454–477.
- Huang, K. 1963. *Statistical Mechanics*, New York: Wiley
- Klauder, J.R. and Skagerstam, B.S. 1985. *Coherent States*, Singapore: World Scientific
- Lifshitz, E.M. and Pitaevskii, L.P. 1980. *Statistical Physics: Theory of Condensed State*, Oxford: Pergamon
- Nozières, P. and Pines, D. 1990. *Theory of Quantum Liquids: Superfluid Bose Liquids*, Redwood: Addison-Wesley
- Parkins, A.S. and Walls, D.F. 1998. The physics of trapped dilute-gas Bose-Einstein condensates. *Physics Reports*, 303: 1–80
- Pitaevskii, L.P. 1961. Vortex lines in an imperfect Bose gas. *Journal of Experimental and Theoretical Physics*, 13: 451–455
- Ter Haar, D. 1977. *Lectures on Selected Topics in Statistical Mechanics*, Oxford: Pergamon
- Yukalov, V.I. and Shumovsky, A.S. 1990. *Lectures on Phase Transitions*, Singapore: World Scientific
- Ziff, R.M., Uhlenbeck, G.E. and Kac, M. 1977. The ideal Bose-Einstein gas revisited. *Physics Reports*, 32: 169–248