Rotational Brownian motion on the sphere surface and rotational relaxation

Ekrem Aydıner*

Dokuz Eylül University, Department of Physics 35160 İzmir, Turkey

(Dated: November 19, 2018)

The spatial components of the autocorrelation function of noninteracting dipoles are analytically obtained in terms of rotational Brownian motion on the surface of a unit sphere using multi-level jumping formalism based on Debye's rotational relaxation model, and the rotational relaxation functions are evaluated.

PACS numbers: 77.22.Gm, 05.40.Jc, 05.40.-a

Rotational Brownian motion and rotational relaxation phenomena has received considerable interest in the literature for a long time. Historically, the problem of isotropic rotational Brownian motion arose in connection with the work of Purcell on spin relaxation in systems with more than two spins in the same molecule. The dynamics of rotational Brownian motion of a sphere around a given axis was discussed briefly by Einstein [1] in one of his early papers. Later, Debye [2] proceeded by extending Einsteins treatment of the translational Brownian motion to rotational Brownian motion of noninteracting permanent dipoles in the presence of an external timevarying field and solving the appropriate Fokker-Planck equation. Here that equation is the Smoluchowski equation, which is an approximate Fokker-Planck equation in the space of angular coordinates for the distribution function of the orientations of the dipoles on the surface of the unit sphere when the influence of the inertia of the molecules on the relaxation process is ignored. Following years many generalizations of Debye model have been extensively studied [3, 4, 5, 6, 7, 8, 9, 10, 11]. However, Debye theory cannot explain relaxation phenomena in strongly interacting systems. The relaxation behavior in strongly interacting systems such as amorphous polymers, spin glasses, glass forming liquids, etc., deviates considerably from the exponential (Debye) pattern $\Phi \sim \exp\left[-t/\tau\right]$ and is characterized by a broad distribution of relaxation times. The relaxation in strongly interacting systems obey to the stretched exponential relaxation function experimentally $\Phi \sim \exp\left[-(t/\tau)^{\alpha}\right]$ with $0 < \alpha < 1$, which often referred to as the Kohlrausch-William-Watts (KWW) function [12, 13].

In this Letter our aim is to obtain rotational relaxation function in terms of autocorrelation function of rotational Brownian motion using multi-level jumping process (MJP) formalism based on historical Debye rotational relaxation scenario. The MJP is the generalization of the two-level jumping process (TJP) formalism. The TJP and MJP have been used to obtain the relaxation function [14, 15, 16, 17]. Recently, the non-exponential relaxation function in complex systems has been obtained using MJP formalism [18]. Autocorrelation function ϕ is simply given by

$$\phi\left(t\right) = \left\langle \mu\left(0\right)\mu\left(t\right)\right\rangle \tag{1}$$

where $\mu(0)$ and $\mu(t)$ denote values of fluctuation variable μ , i.e. dipole, at time zero and the same variable at a later time t, respectively. Physically, it measures the time over which the variable μ retains its own memory until this memory is averaged out by statistical randomness.

If it is assumed that the external field has been applied in the z-direction, the spatial components of the autocorrelation function of dipoles can be written in terms of the Brownian motion on the surface of the unit sphere in a simple form

$$\phi_x(t) = \langle \mu_x(0) \, \mu_x(t) \rangle = \langle \mu_0 \sin \theta \cos \varphi \rangle \qquad (2a)$$

$$\phi_z(t) = \langle \mu_y(0) \, \mu_y(t) \rangle = \langle \mu_0 \sin \theta \sin \varphi \rangle \qquad (2b)$$

$$\phi_z(t) = \langle \mu_z(0) \, \mu_z(t) \rangle = \langle \mu_0 \cos \theta \rangle \tag{2c}$$

where μ_0 denotes magnitude of the dipole ($\mu_0 = 1$), θ and φ are stochastic variables which define the spatial orientation of a dipole, in other words, position of the Brownian particle on the sphere surface. For simplicity, it can be considered that both of θ and φ are discrete stochastic variables. The θ takes random values in between 0 and π , and on the other hand, φ takes random values in between 0 and 2π , where π 's value is defined in terms of degree. Hence, stochastic variable μ can take randomly integer or non-integer values depending on θ and φ . In a such process, fluctuation variable μ jumps from one value of μ to another stochastically. As it can be seen in Eqs. (2) only z-component of the autocorrelation function depends on θ , however, x and y components depend on two variables i.e. θ and φ . We will see below that all components of the autocorrelation function are carried out using MJP after a little bit of algebra.

All components of the autocorrelation function in Eqs. (2) can be derived by using stochastic formalism. For example, the z-component $\phi_z(t)$ of the autocorrelation function is given by

$$\phi_{z}(t) = \sum_{\mu_{z0}} \sum_{\mu_{z}} p(\mu_{z0}) \mu_{z0} P(\mu_{z0} \mid \mu_{z}, t) \mu_{z} \qquad (3)$$

^{*}Electronic address: ekrem.aydiner@deu.edu.tr

where the quantities μ_{z0} and μ_z are the values of stochastic variable μ_z at times 0 and t, respectively. Having started at the initial state μ_{z0} with the statistical weight $p(\mu_{z0})$, and $P(\mu_{z0} | \mu_z, t)$ measures the probability of propagation from μ_{z0} to μ_z in time t, which is known as conditional probability.

The conditional probability $P(\mu_{z0} \mid \mu_z, t)$ can be given by the matrix representation associate a stochastic state $|\mu_z\rangle$. In the operator formalism (See Refs. [14, 19]) the conditional probability $P(\mu_{z0} \mid \mu_z, t)$ is simply written with the matrix elements of operators as

$$P(\mu_{z0} \mid \mu_z, t) \equiv \langle \mu_z \mid \vec{P}(t) \mid \mu_{z0} \rangle \tag{4}$$

in the matrix form where $\widehat{P}(t)$ is the conditional probability operator, and $|...\rangle$ indicates vector space of stochastic states. Note that the conditional probability satisfy

$$P(\mu_{z0} \mid \mu_z, 0) \equiv \langle \mu_z \mid \widehat{P}(0) \mid \mu_{z0} \rangle = \delta(\mu_z - \mu_{z0})$$
 (5)

for t = 0.

Similarly, the fluctuation variable μ_z can be also given by matrix form, the eigenvalues of which correspond to stochastic variables μ_0 , $\mu_1,...,\mu_\pi$, respectively. The matrix representation the non-interacting fluctuation variable μ_z can be represented associated with operator \hat{Z} as

$$\widehat{Z} = \begin{pmatrix} \mu_0 & 0 & 0 & \dots & 0 \\ 0 & \mu_1 & 0 & \dots & 0 \\ 0 & 0 & \mu_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \vdots & \vdots & \mu_\pi \end{pmatrix}.$$
 (6)

Eigenvalues of operator \widehat{Z} are given by $\mu_z = \cos\theta$ where $z = 0, 1, 2, ..., \pi$.

Thus, the autocorrelation function in Eq.(3) can be transformed to the matrix representation

$$\phi_{z}(t) = \sum_{\mu_{z},\mu_{z0}} p(\mu_{z0}) \langle \mu_{z0} | \hat{Z} | \mu_{z0} \rangle \langle \mu_{z} | \hat{P}(t) | z_{z0} \rangle \langle \mu_{z} | \hat{Z} | \mu_{z} \rangle$$
(7)

using above definitions.

As yet, the autocorrelation function (3) has been given by matrix representation as seen in Eq. (7). The Eq. (7) can be solved easily using matrix algebra and with more convenient basis vector notation if the operator form of the conditional probability is known.

The operator form of the conditional probability can be evaluated from master equation. The conditional probability $P(\mu_{z0} \mid \mu_z, t)$ can be given associated with master equation

$$\frac{\partial}{\partial t}P\left(\mu_{z0} \mid \mu_{z}, t\right) = \sum_{\mu'_{z}} W\left(\mu'_{z} \mid \mu_{z}\right) P\left(\mu_{z0} \mid \mu'_{z}, t'\right) \quad (8)$$

where $W(\mu'_z \mid \mu_z)$ is the jump rate from μ_z to μ'_z of the Brownian particle i.e. dipole.

The jumping rate $W(\mu'_z \mid \mu_z)$ can be given by matrix form

$$W(\mu'_{z} \mid \mu_{z}) = \langle \mu_{z} \mid \widehat{W} \mid \mu'_{z} \rangle \tag{9}$$

where \widehat{W} is the jumping rate operator. Thus Eq. (8) can be rewritten completely by a matrix form as

$$\frac{\partial}{\partial t} \langle \mu_z | \, \widehat{P}(t) \, | \mu_{z0} \rangle = \sum_{\mu'_z} \langle \mu_z | \, \widehat{W} \, | \mu'_z \rangle \, \langle \mu'_z | \, \widehat{P}(t) \, | \mu_{z0} \rangle$$
(10)

using Eq. (4) and (9).

The stochastic states taken to form an orthonormal set which provide *closure* property is as

$$\sum_{\mu_z} |\mu_z\rangle \langle \mu_z| = 1 \tag{11}$$

for discrete variables. Using closure property, Eq. (10) is reduced to

$$\langle \mu_z | \frac{\partial \widehat{P}(t)}{\partial t} | \mu_{z0} \rangle = \langle \mu_z | \widehat{W} \widehat{P}(t) | \mu_{z0} \rangle \quad . \tag{12}$$

This equation can be simplified as

$$\frac{\partial}{\partial t}\widehat{P}\left(t\right) = \widehat{W}\widehat{P}\left(t\right) \quad . \tag{13}$$

As a result, the solution of Eq. (13) is

$$\widehat{P}(t) = \exp\left(\widehat{W}t\right) \tag{14}$$

where \widehat{W} is the jump matrix and can be written in terms of collision and unit matrix elements as

$$\widehat{W} = \lambda \left(\widehat{J} - \mathbf{1} \right) \tag{15}$$

where λ is defined as the relaxation rate, \hat{J} is the collision matrix and **1** is a unit matrix. The relaxation rate λ for π -level jumping is given by

$$\lambda = \pi w \tag{16}$$

and collision matrix \widehat{J} is presented by

where w is the jump rate from one value of μ_z to another.

It is helpful to write down the explicit form of the jump matrix \widehat{W} in terms of w. If Eqs. (16) and (17) are inserted into Eq. (15), the jump matrix \widehat{W} can be constructed as

$$\widehat{W} = \begin{pmatrix} -(\pi - 1)w & w & \dots & w \\ w & -(\pi - 1)w & \dots & w \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ w & w & \dots & -(\pi - 1)w \end{pmatrix}.$$
(18)

The usefulness of the decomposition in Eq. (15) comes out from the fact that the \widehat{J} matrix has a very simple property

$$\widehat{J^2} = \widehat{J}, \quad \widehat{J^3} = \widehat{J}, \quad ..., \quad \widehat{J^k} = \widehat{J}$$
 (19)

for any integer k > 0. This property allows us to immediately construct the conditional probability matrix as

$$\widehat{P}(t) = \exp\left[\lambda\left(\widehat{J} - \mathbf{1}\right)t\right]$$
 (20)

or, alternatively, it can be given in the matrix form

where η is defined by

$$\eta = \frac{1}{\pi} \left(1 - e^{-\lambda t} \right) \quad . \tag{22}$$

The conditional probability operator $\widehat{P}(t)$ can be expressed in useful form as

$$\widehat{P}(t) = \exp(-\lambda t) \left[1 - \widehat{J} + \widehat{J} \exp(\lambda t) \right]$$
(23)

using direct power series expansion and relation which is given by Eq. (19).

On the other hand, it is convenient to introduce the more general notation for basis vectors to solve Eq. (7). Therefore, let us associate a stochastic state $|\theta\rangle$ ($\theta = 0, 1, 2, ..., \pi$ for multi-level jumping process) instead $|\mu_z\rangle$ of π values

$$|0\rangle = \begin{pmatrix} 1\\0\\0\\.\\.\\.\\0 \end{pmatrix}, |1\rangle = \begin{pmatrix} 0\\1\\0\\.\\.\\.\\0 \end{pmatrix}, ..., |\pi\rangle = \begin{pmatrix} 0\\0\\0\\.\\.\\.\\1 \end{pmatrix}$$
(24)

The *closure* property for new stochastic states $|\theta\rangle$ (and similarly $|\theta'\rangle$) is presented as

$$\sum_{\theta} \left| \theta \right\rangle \left\langle \theta \right| = 1 \tag{25}$$

in a discrete form.

Now, Eq. (7) can be rewritten using new basis vectors as

$$\phi_{z}(t) = \sum_{\theta,\theta'} p_{\theta} \langle \theta | \, \widehat{Z} \, | \theta \rangle \, \langle \theta' | \, \widehat{P}(t) \, | \theta \rangle \, \langle \theta' | \, \widehat{Z} \, | \theta' \rangle \qquad (26)$$

with apriori occupation probability

$$p_{\theta} = \frac{1}{\pi}, \quad \theta = 0, 1, 2, ..., \pi$$
 (27)

The expectation value of conditional probability matrix $\hat{P}(t)$ in Eq. (26) is obtained as

$$\langle \theta | \hat{P}(t) | \theta' \rangle = \frac{1}{\pi} + \left(\delta_{\theta\theta'} - \frac{1}{\pi} \right) \exp\left(-\lambda t\right)$$
 (28)

using Eq. (23) and new basis vectors in Eq. (24), and on the other hand, the expectation value of \hat{Z} is given by

$$\langle \theta | \, \widehat{Z} \, | \theta' \rangle = Z_{\theta} \delta_{\theta \theta'} \tag{29}$$

with $\theta, \theta' = 0, 1, 2, ..., \pi$, where Z_{θ} are the allowed values of the stochastic variables.

If Eqs. (27)-(29) are inserted into Eq. (26), the autocorrelation function (26) can be reduced to a simple form

$$\phi_z(t) = \langle \mu_z \rangle^2 + \left(\langle \mu_z^2 \rangle - \langle \mu_z \rangle^2 \right) \exp\left(-\lambda t\right)$$
(30)

where the deterministic quantities are weighted averages over the available states of the corresponding variable. The average values of $\langle \mu_z \rangle$ and $\langle \mu_z^2 \rangle$ in the stationary state are given by

$$\langle \mu_z \rangle = \sum_{\theta=0}^{\pi} p_\theta \left\langle \theta \right| \widehat{Z} \left| \theta \right\rangle = \frac{1}{\pi} \sum_{\theta=0}^{\pi} Z_\theta \tag{31}$$

and

$$\left\langle \mu_{z}^{2}\right\rangle = \sum_{\theta=0}^{\pi} p_{n} \left\langle \theta \right| \widehat{Z^{2}} \left| \theta \right\rangle = \frac{1}{\pi} \sum_{\theta=0}^{\pi} Z_{\theta}^{2}$$
(32)

respectively. The diagonal elements of \widehat{Z} matrix (6) are given as $\mu_z = \cos \theta$, i.e., $Z_{\theta} = \cos \theta$. Thus, it is expected that $\langle \mu_z \rangle = 0$ and $\langle \mu_z^2 \rangle = 1/2$.

Finally, z-component of the autocorrelation function for the Brownian motion on the spherical surface can be obtained simply as

$$\phi_z\left(t\right) = \frac{1}{2}\exp\left[-\lambda t\right] \quad . \tag{33}$$

As it can be seen in Eq. (33) z-component of the rotational autocorrelation function has exponential form which indicates that the same component of rotational relaxation must has exponential form.

The relaxation function for z-component can be immediately obtained using linear response theory

$$\Phi(t) = \Psi(t = \infty) - \Psi(t) \tag{34}$$

where $\Psi(t)$ is well known response function which can be associated to the famous response-relaxation function as

$$\Psi(t) = \frac{1}{kT} \left(\left\langle \mu_z^2 \right\rangle - \left\langle \mu_z(0) \, \mu_z(t) \right\rangle \right) \tag{35}$$

where k is the Boltzmann constant, and T is the temperature. If Eq. (35) is inserted into Eq. (34), as a final result, the rotational relaxation function can be obtained as

$$\Phi_z\left(t\right) = \frac{1}{2kT} \exp\left[-\lambda t\right] \quad . \tag{36}$$

We have shown that the z-component of the autocorrelation function and relaxation function can be obtained using the multi-level jumping formalism. Similarly, the other components of the autocorrelation function, that is, $\phi_x(t)$ and $\phi_y(t)$ can be also derived using this technique. However, to obtain the components ϕ_x and ϕ_y , we should eliminate the φ in ϕ_x and ϕ_y .

At this point, firstly, it can be suggested that components of the autocorrelation function in Eq. (2) must satisfy the relation

$$\left\langle \phi_x^2(t) \right\rangle + \left\langle \phi_y^2(t) \right\rangle + \left\langle \phi_z^2(t) \right\rangle = 1$$
 (37)

which is caused from a simple trigonometric identity as

$$\left\langle \sin^2 \theta \cos^2 \varphi \right\rangle + \left\langle \sin^2 \theta \sin^2 \varphi \right\rangle + \left\langle \cos^2 \theta \right\rangle = 1$$
 (38)

Therefore, the components ϕ_x and ϕ_y can be written as

$$\left\langle \phi_x^2\left(t\right) + \phi_y^2\left(t\right) \right\rangle = 1 - \left\langle \cos^2\theta \right\rangle \quad .$$
 (39)

On the other hand, it is expected that the component $\phi_y(t)$ should be equal to $\phi_x(t)$. Thus, $\phi_x(t)$ is written as

$$\phi_x(t) \equiv \langle \phi_x(t) \rangle = \frac{1}{\sqrt{2}} \langle \sin \theta \rangle$$
 (40)

in simple form.

As it can be seen in Eq. (40), using mathematical trick, we have got rid of φ in the components $\phi_x(t)$ and $\phi_y(t)$. Now, to calculate the component $\phi_x(t)$ we need only values $\langle \mu_x \rangle$ and $\langle \mu_x^2 \rangle$. The averages can be given as

$$\langle \mu_x \rangle = \sum_{\theta=0}^{\pi} p_\theta \langle \theta | \, \widehat{X} \, | \theta \rangle = \frac{1}{\pi} \sum_{\theta=0}^{\pi} X_\theta \tag{41}$$

and

$$\left\langle \mu_x^2 \right\rangle = \sum_{\theta=0}^{\pi} p_\theta \left\langle \theta \right| \widehat{X^2} \left| \theta \right\rangle = \frac{1}{\pi} \sum_{\theta=0}^{\pi} X_\theta^2 \tag{42}$$

respectively. The matrix \widehat{X} has the same form with (6). But elements of \widehat{X} matrix are equal to $\mu_x = \sin \theta$, i.e., $X_{\theta} = \sin \theta$. Therefore, required averages can be carried out as $\langle \mu_x \rangle = \sqrt{2}/\pi$ and $\langle \mu_x^2 \rangle = 1/4$. Hence, the relaxation function for x and y components are given by

$$\Phi_x(t) = \Phi_y(t) = \frac{1}{kT} \left(\frac{1}{4} - \frac{2}{\pi^2}\right) \exp\left[-\lambda t\right] \quad . \tag{43}$$

The Eq. (43) clearly show that the components $\phi_x(t)$ and $\phi_y(t)$ of rotational relaxation function have also exponential form.

In this Letter, the spatial components of the autocorrelation function of noninteracting dipoles have been analytically obtained in terms of rotational Brownian motion on the surface of a unit sphere using multi-level jumping formalism based on Debye's rotational relaxation model, and the rotational relaxation functions have been evaluated. As it is expected, all components of the rotational relaxation function have exponential form when the influence of the inertia of the dipoles is ignored.

This work was partially supported by Dokuz Eylül University (Project No: 04.KB.FEN.098)

- [1] A. Einstein, Ann. Physik. **19**, 371 (1906).
- [2] P. Debye, *Polar Molecules* (Chemical Catalog, New York, 1929, reprinted by Dover Publications, New York, 1954).
 [2] W. H. E. D. D. D. Tor, 7 (1977).
- [3] W. H. Furry, Phys. Rev. **107**, 7 (1957).
- [4] D. Pines and C. P. Slichter, Phys. Rev. **100**, 1014 (1955).
- [5] E. Fatuzzo and P. R. Mason, Proc. Phys. Soc. 90, 741 (1967).
- [6] G. W. Ford, J. T. Lewis, and J. McConnell, Phys. Rev. A 19, 907 (1979).
- [7] L. D. Favro, Phys. Rev. **119**, 53 (1960).
- [8] L. Pauling, Phys. Rev. **36**, 430 (1930).
- [9] S. Dattagupta, Phys. Rev. A 14, 480 (1976).
- [10] J. R. McConnell, Rotational Brownian Motion and Dielectric Theory (Academic, New York, 1980).

- [11] W. F. Brown, J. Appl. Phys. **30**, 130 (1959).
- [12] R. Kohlrausch, Ann. Phys. (Leipzig) 12, 393 (1847).
- [13] G. Williams and D. C. Watts, Trans. Faraday Soc. 66, 80 (1970).
- [14] S. Dattagupta, Relaxation Phenomena in Condensed Matter Physics (Academic Press, Orlando, 1987).
- [15] S. Dattagupta and M. Blume, Phys. Rev. B 10, 4551 (1974).
- [16] R. Kubo, J. Phys. Soc. Jpn., 9, 935 (1954).
- [17] P. W. Anderson, J. Phys. Soc. Jpn., 9, 316 (1954).
- [18] E. Aydıner, E-print: cond-mat/0602461.
- [19] N. G. van Kampen, Stochastic Processes in Physics and Chemistry (North-Holland, Amsterdam, 1981).