# A Kinetic Approach to the Calculation of Surface Tension in a Spherical Drop

Vincenzo Molinari, Domiziano Mostacci<sup>\*</sup> and Margherita Premuda

INFM-BO and Laboratorio di Ingegneria Nucleare di Montecuccolino,

Università degli Studi di Bologna via dei Colli 16, I-40136 Bologna (ITALY)

# Abstract

In literature, surface tension has been investigated mainly from a thermodynamic standpoint, more rarely with kinetic methods. In the present work, surface tension in drops is studied in the framework of kinetic theory, starting from the Sutherland approximation to Van Der Waals interaction between molecules. Surface tension is calculated as a function of drop radius: it is found that it approaches swiftly an asymptotic value, for radii of several times the intermolecular distance. This theoretical asymptotic value is compared to experimental values for a few liquids, and is found in reasonable agreement.

 $<sup>*</sup> Electronic \ address: \ domiziano.mostacci@mail.ing.unibo.it; \ URL: \verb+http://tori.ing.unibo.it/DIENCA/ita/personale/assisted address: \ domiziano.mostacci@mail.ing.unibo.it; \ URL: \ http://tori.ing.unibo.it/DIENCA/ita/personale/assisted address: \ domiziano.mostacci@mail.ing.unibo.it; \ url address: \ domiziano.mostacci@mail.ing.unibo.it; \ url address: \ domiziano.mostacci@mail.ing.unibo.it; \ url address: \ url a$ 

## I. INTRODUCTION

The effect of curvature on surface tension has been investigated widely in the 50's, but mainly within the framework of Thermodynamics (see, e.g., [1, 2]). Some studies have taken the point of view of kinetic theory [3, 4]. More recent work investigates surface tension, in plane geometry, from the point of view of kinetic theory, making use of the Sutherland potential [5]; still much work is being done taking the Thermodynamics standpoint (see for instance [6, 7, 8]). In the present work the problem is studied through kinetic equations derived in the following simplifying assumptions:

1) Free volume method [1]. This approximation can also be bettered including the molecular exchange with the surrounding atmosphere [9]. This effect might form the object of subsequent work.

2) Uncorrelated molecules. This assumption is dictated by two considerations: first, the double distribution function (or pair distribution function, as it often referred to) is not well known as it can be calculated exactly only for extremely simplified situations. Albeit it is almost invariably called upon, still the assumptions made to calculate it are often such as to render its benefits very limited; secondly, the importance of correlation can be judged from the ratio  $\gamma$  between the potential energy at the average intermolecular distance and the average kinetic energy of molecules [10, 11]:

$$\gamma = \frac{\Phi\left(r_0\right)}{\frac{3}{2}K_BT}\tag{1}$$

The intermolecular distance can be estimated, as usual, from the number density n as  $n^{\frac{1}{3}}$ . As will be shown in section IV, where  $\gamma$  is calculated for several cases, this ratio is of order unity, as might have been expected: therefore, correlation between molecules might play some role. However, it will be neglected here as a first approximation.

3) **Constant density**. The number density will be considered constant with radius: this approximation might turn out questionable at the interface, particularly in view of the fact that surface tension is, after all, an interface effect. This assumption will be made here, as a first approximation, and then reviewed in commenting the results.

4) **Sutherland potential**. The intermolecular Van der Waals forces are modeled with the Sutherland potential. This assumption could be bettered using the full Lennard-Jones potential, however at the cost of greater mathematical difficulties; on the other hand, it can be argued that the error introduced using the Sutherland potential, with suitable parameters, is a reasonable price to pay for the mathematical simplification, which allows solution in closed form. As will be shown in the last section, the results obtained are in very reasonable agreement with experimental data.

In the following sections, a simple method will be derived within the approximations discussed above, to obtain a simple expression for the surface tension as a function of radius. As the radius becomes larger, the value of surface tension approaches that of a plane surface. Calculations presented for several different liquids, in the large radius limit, are consistent with experimental results for plane surfaces.

## **II. GOVERNING EQUATIONS**

Consider a spherical droplet of radius R centered in the point O, and a point P at a distance r from O, with  $r \in [0,R]$ . A system of spherical coordinates can be defined, with origin in P and the O-P direction as the polar axis, as depicted in Fig. 1. The  $\varphi=0$  half-plane can be chosen arbitrarily, due to the spherical symmetry of the system.



Figure 1: Geometry of the problem - section of the droplet on the half-plane  $\varphi=0$ 

Any point Q within the drop that lies on the  $\varphi=0$  half-plane can then be described

uniquely with the pair of coordinates  $\rho$  and  $\vartheta$  (or  $\mu = \cos \vartheta$ ) as defined in Fig. 1.

If forces between particles are purely central forces, i.e., depending only on a given power  $\alpha$  of the distance, then the force due to the attraction of a particle in Q on the particle in P can be written as

$$\mathbf{F}_{QP} = \frac{G}{\rho^{\alpha}}\hat{\rho} \tag{2}$$

as it is directed from P to Q. In spherical coordinates, the half circle on the  $\varphi=0$  half-plane depicted in Fig. 1 is described by the equation

$$\rho = \sqrt{(\mu r)^2 + (R^2 - r^2)^2} - \mu r \tag{3}$$

or equivalently

$$\mu = \frac{(R^2 - r^2) - \rho^2}{2r\rho} \tag{4}$$

therefore the domain corresponding to the intersection of the drop with the half-plane is defined as

$$\mu \in [-1,1] \quad ; \quad \rho \in \left[0, \sqrt{(\mu r)^2 + (R^2 - r^2)^2} - \mu r\right]$$
(5)

or equivalently

$$\rho \in [0, R+r] \quad ; \quad \mu \in \left[-1, \min\left\{1, \frac{(R^2 - r^2) - \rho^2}{2r\rho}\right\}\right]$$
(6)

If there are  $n(r_1)$  particles per unit volume (due to the spherical symmetry the density can only depend on the radial position  $r_1$ , and this can be expressed in terms of r,  $\rho$  and  $\mu$ ), in a volume element dV (expressed in spherical coordinates) at location ( $\rho, \mu, \varphi$ ) there are a number of particles given by

$$ndV = n\left(r_1\left[r,\rho,\mu\right]\right)\rho^2 d\rho d\mu d\varphi \tag{7}$$

exerting on the particle in P a force  $d\mathcal{F}$ 

$$d\mathcal{F} = \frac{G}{\rho^{\alpha}} n dV = Gn\left(r_1\right) \rho^{2-\alpha} d\rho d\mu d\varphi \tag{8}$$

The z-component of  $d\mathcal{F}$  is

$$d\mathcal{F}_z = d\mathcal{F}\cos\vartheta = Gn\left(r_1\right)\rho^{2-\alpha}\mu d\rho d\mu d\varphi \tag{9}$$

and in the same vein the component perpendicular to z:

$$d\mathcal{F}_{\perp} = d\mathcal{F}\sin\vartheta = Gn\left(r_{1}\right)\rho^{2-\alpha}\sqrt{1-\mu^{2}}d\rho d\mu d\varphi$$
(10)

The first can be integrated readily with respect to  $\varphi$ , yielding

$$\int_{0}^{2\pi} d\mathcal{F}_{z} = 2\pi G n\left(r_{1}\right) \rho^{2-\alpha} \mu d\rho d\mu \tag{11}$$

As for the second, some further discussion is appropriate. To evaluate the surface tension one can proceed as follows: imagine cutting the drop in two halves, each one exerting a certain amount of attraction on the other. Consider now a molecule that is lying right on the edge of one half drop, say at the intersection of the polar axis with the drop surface. The component perpendicular to the axis, and lying on on the  $\varphi = 0$  plane, of the force acting on this molecule that is due to an element of volume at location ( $\rho, \vartheta, \varphi$ ) is given by

$$d\mathcal{F}_{\perp}\cos\varphi = Gn\left(r_{1}\right)\rho^{2-\alpha}\sqrt{1-\mu^{2}}d\rho d\mu\cos\varphi d\varphi$$
(12)

The tangential force  $dF_T$  acting on this molecule due to all elements with coordinates  $(\rho, \vartheta)$ , is given by the integral over  $\varphi$  of  $d\mathcal{F}_{\perp} \cos \varphi$  in one of the halves of the drop, e.g., for  $\varphi \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ :

$$dF_T = \int_{-\frac{\pi}{2}}^{+\frac{\pi}{2}} \left[ Gn(r_1) \rho^{2-\alpha} \sqrt{1-\mu^2} d\rho d\mu \right] \cos\varphi d\varphi = 2Gn(r_1) \rho^{2-\alpha} \sqrt{1-\mu^2} d\rho d\mu \qquad (13)$$

To calculate the total tangential force on the molecule, the above expression is to be integrated over  $\rho$  and  $\mu$ . In the following the density will be considered constant, as discussed in point 3) in the introduction. Recalling the expression for the domain in Eq. (6), noted that in this case r = R, the following integral is obtained:

$$\frac{F_T}{2nG} = \int_D^{2R} \rho^{2-\alpha} d\rho \int_{-1}^{\frac{-\rho}{2R}} \sqrt{1-\mu^2} d\mu = \frac{1}{2} \int_D^{2R} \rho^{2-\alpha} \left\{ \arcsin\left(\frac{-\rho}{2R}\right) - \frac{\rho}{2R} \sqrt{1-\left(\frac{\rho}{2R}\right)^2} + \frac{\pi}{2} \right\} d\rho$$
(14)

Here D is the distance of minimum approach, as will be discussed in the next section.

# III. SOLUTION WITH VAN DER WAALS TYPE FORCES

Van der Waals interactions are usually represented as a Lennard-Jones potential [12]

$$U_{LJ}(r) = 4\varepsilon \left[ \left(\frac{\sigma}{r}\right)^{12} - \left(\frac{\sigma}{r}\right)^6 \right]$$
(15)

This last can be approximated by a Sutherland potential [12], usually written as

$$U_S(r) = \begin{cases} \infty & r < D \\ -A\left(\frac{D}{r}\right)^{\omega} & r \ge D \end{cases}$$
(16)

with suitable choice of the (positive) constants  $\omega$ , A and D. A condition typically imposed is to conserve the asymptotic behavior of ULJ, which entails  $\omega = 6$  and  $A \cdot D^6 = 4\varepsilon \cdot \sigma^6$ ; a second condition can be chosen as follows: the rest position of a particle in the field of another particle is the bottom of the potential well, therefore it seems reasonable to choose the constants so that this position is the same in the two cases. Therefore,  $D = \sigma \cdot 2^{\frac{1}{6}}$ . With the above positions the Sutherland potential becomes

$$U_S(r) = \begin{cases} \infty & r < D = \sigma \cdot 2^{\frac{1}{6}} \\ -\varepsilon \left(\frac{D}{r}\right)^{\omega} & r \ge D = \sigma \cdot 2^{\frac{1}{6}} \end{cases}$$
(17)

This potential is depicted in Fig. 2. This corresponds to the present model for the force Eq. (2), with  $\varphi = 7$  and

$$G = 6\varepsilon D^6 \tag{18}$$



Figure 2: Schematized Sutherland potential

In the Sutherland approximation, D is the distance of minimum approach, the lower limit

of the integral in Eq. (14). Calculating that integral, and recalling the expression for G in Eq. (18), the following expression is obtained for the tangential force acting on a molecule:

$$F_T = \frac{3}{2}n\varepsilon D^2 \left\{ \arcsin\left(\frac{-D}{2R}\right) + \frac{D}{2R}\sqrt{1 - \left(\frac{D}{2R}\right)^2} \left[1 - 2\left(\frac{D}{2R}\right)^2\right] + \frac{\pi}{2} \right\}$$
(19)

The number of molecule per unit length can be estimated as  $n^{\frac{1}{3}}$ . Therefore the force per unit length can be calculated as

$$T_{sur}\left(R\right) = \frac{3}{2}n^{\frac{4}{3}}\varepsilon D^{2} \left\{ \arcsin\left(\frac{-D}{2R}\right) + \frac{D}{2R}\sqrt{1 - \left(\frac{D}{2R}\right)^{2}} \left[1 - 2\left(\frac{D}{2R}\right)^{2}\right] + \frac{\pi}{2} \right\}$$
(20)

The graph of the reduced surface tension  $\frac{T_{sur}}{\frac{3}{2}n^{\frac{4}{3}}\varepsilon D^2}$  as a function of  $\frac{2R}{D}$ , the drop diameter in units of D, is shown in Fig. 3



Figure 3: Reduced surface tension vs. drop diameter in units of D

As can be gathered from the graph, the surface tension approaches rapidly the limiting value of as the drop diameter grows to only a few tens of times D.

## IV. COMPARISON WITH EXPERIMENTAL DATA

Limiting values of surface tension for large radii have been calculated from Eq. (20) for several liquids, and compared with known experimental values for plane surfaces. The results are shown in Table 1. As can be gathered from the data reported, the agreement is quite good for such a qualitative approach, being always within a factor of 2 to 3 (3.4 for water). As the results depend on the parameters of the Sutherland potential, agreement might be improved by a different choice, the one choice made here being somewhat arbitrary albeit reasonable. In [15], for instance, detailed calculation and comparison with Lennard-Jones parameters is reported for several inert gases.

Table 1: comparison of calculated and experimental values of surface tension						
	Density [m <sup>-3</sup> ]	Lennard-Jones parameters <sup>[13]</sup>		$\gamma = \frac{\Phi(\mathbf{r}_0)}{\frac{3}{5}\mathrm{KT}}$	Calculated value	Experimental value
		σ[Å]	ε/K <sub>B</sub> [K]	2 ***	[N/m]	[N/m]
Water, 20°C	3.34×10 <sup>28</sup>	2.641	809.1	1.4	248×10 <sup>-3</sup>	72.75×10 <sup>-3</sup> <sup>[14]</sup>
Ethanol, 20°C	1.01×10 <sup>28</sup>	4.530	362.6	1.5	66.5×10 <sup>-3</sup>	22.75×10 <sup>-3</sup> <sup>[14]</sup>
Acetone, 20°C	0.671×10 <sup>28</sup>	4.600	560.2	0.57	61.5×10 <sup>-3</sup>	23.70×10 <sup>-3</sup> <sup>[14]</sup>
Benzene, 20°C	0.677×10 <sup>28</sup>	5.349	412.3	2.01	62.0×10 <sup>-3</sup>	28.88×10 <sup>-3 [13]</sup>
Liquid Oxigen, -183°C	2.14×10 <sup>28</sup>	3.467	106.7	1.26	31.2×10 <sup>-3</sup>	13.2×10 <sup>-3</sup> [14]
Liquid Nitrogen, -183°C	$1.72 \times 10^{28}$	3.798	71.4	0.94	18.8×10 <sup>-3</sup>	6.6×10 <sup>-3</sup> <sup>[14]</sup>

On the other hand, the density is indeed essentially constant throughout the drop except near the boundary, where it is generally found to decrease [1, 5, 16, 17, 18, 19, 20]. As the layers close to the boundary give the main contribution to the integral in Eq. (14), the actual density profile should be considered in the integration. This will form the object of a subsequent work, for the present the following view will be taken: in the literature, the density is always shown to decrease rather abruptly near the interface, taking a relative value at the interface of roughly one half the bulk value. Therefore, introducing the actual density in the integral will decrease this latter by a factor that is between 0.5 and 1. On the other hand, in calculating the number of molecules per unit length at the surface, the value of density at the surface must be considered; altogether then, adding the two effects, the result changes by a factor of between 0.4 and 0.8, which is the right magnitude to offer the correction needed to reconcile calculated values with those found experimentally.

## V. CONCLUSIONS

As anticipated in the introduction, aim of this work was to propose a simple equation for calculating surface tension in drops as a function of radius. The result is in very reasonable agreement with the experiment. This result was obtained in the simplifying assumptions discussed in the introduction. Accurate description of the density profile is needed to improve confidence in the results yielded by the method proposed.

- J.O. Hirschfelder, C.F. Curtiss, R.B. Bird, "Molecular Theory of Gases and Liquids", Wiley, New York (1954).
- [2] T.L. Hill, J. Chem. Phys. **19**, 1203 (1951).
- [3] J.G. Kirkwood and F.P. Buff, J. Chem. Phys. 17, 338 (1949).
- [4] F.P. Buff and J.G. Kirkwood, J. Chem. Phys. 18, 991 (1949).
- [5] A. Frezzotti, L. Gibelli, F. Lorenzani, Phys. Fl. 17, 012102-1 (2005).
- [6] L.K. Antanovskii, Phys. Rev. E 54, 6285 (1996).
- [7] A.J. Queimada et al., Fl. Phase Eq. **222-223**, 161 (2004).
- [8] J. Escobedo, G.A. Mansoori, AIChE J. 42, 1425 (1996).
- [9] D. Giusti, V. Molinari, D. Mostacci, Il Nuovo Cimento **21C**, 123 (1998).
- [10] R.L. Liboff, "Kinetic Theory", Prentice Hall, Englewood Cliffs, NJ (1990).
- [11] J.L. Delcroix, A. Bers, "Physique des plasmas", Interéditions, Paris (1994).
- [12] S. Chapman and T.Cowling, "The Mathematical Theory of Non-Uniform Gases", University Press, Cambridge, UK (1952).
- [13] R.C. Reid, J.M. Prausnitz, B.E. Poling, "The Properties of Gases and Liquids", McGraw-Hill, Singapore (1987).
- [14] A. Frohn, N. Roth, "Dynamics of Droplets", Springer Verlag, Berlin (2000).
- [15] H.W. Graben, R.D. Present, Rev. Mod. Phys. 36, 1025 (1964).
- [16] C.E. Upstill, R. Evans, J. Phys. C 10, 2791 (1977).
- [17] A. Laaksonen, R. McGraw, Europhys. Lett. 35, 367 (1996).
- [18] T.L. Hill, J. Chem. Phys. **19**, 261 (1951).
- [19] T.L. Hill, J. Chem. Phys. **20**, 141 (1952).

[20] A.J.M. Yang, P.D. Fleming, J.H. Gibbs, J. Chem. Phys. 64, 3732 (1976).