

Multiconfigurational Hartree-Fock theory for identical bosons in a double well

D. Masiello,^{1,*} S. B. McKagan,^{2,†} and W. P. Reinhardt^{1,2}

¹*Department of Chemistry, University of Washington, Seattle, Washington 98195-1700, USA*

²*Department of Physics, University of Washington, Seattle, Washington 98195-1560, USA*

(Dated: May 24, 2019)

Multiconfigurational Hartree-Fock theory is presented and implemented in an investigation of the fragmentation of a Bose-Einstein condensate made of identical bosonic atoms in a double well potential at zero temperature. Our approach builds in the effects of the condensate mean field and of atomic correlations by describing generalized many-body states that are composed of multiple configurations into which atomic interactions are incorporated. Nonlinear and linear optimization is utilized in conjunction with the variational and Hylleraas-Undheim theorems to find the optimal ground and excited states of the interacting system. The resulting energy spectrum and associated eigenstates are presented as a function of double well barrier height. Delocalized and localized single configurational states are found in the extreme limits of the simple and fragmented condensate ground states, while multiconfigurational states and exotic macroscopic quantum superposition states are revealed throughout the full extent of barrier heights. Comparison is made to existing theories that either neglect mean field or correlation effects and it is found that contributions from both interactions are essential in order to obtain a more complete microscopic understanding of the condensate's atomic structure throughout the fragmentation process.

PACS numbers:

I. INTRODUCTION

Recent experimental realization of a trapped atom interferometer using a coherently split Bose-Einstein condensate (BEC) [1, 2] as well as the direct observation of tunneling and self-trapping in weakly linked BECs [3, 4] have provided an impetus to formulate a comprehensive theoretical description of the zero temperature Bose gas confined to a trapping potential that can be continuously deformed from a single well into a double well with large barrier height. Such a deformation of the trapping potential impels the BEC from a single coherent entity to a fragmented condensate made up of two coherent and potentially correlated moieties. The quantum many-body physics governing this complex fragmentation process involves competition and balance between the effects of the condensate's mean field on the interacting atomic gas and the correlations that emerge between atoms in different Fock states.

Various theoretical descriptions of the simple and fragmented BEC exist in the literature today. At both low and high barrier heights, the many-body ground state of the condensate is well approximated in mean field theory by a single Fock state, which expresses a particular arrangement of atoms among one, two, or even many single-particle states [5, 6, 7]. Where mean field equations specify the underlying single-particle states, we call the single Fock state a single *configuration* [49]. If only one single-particle state is involved, then mean field theory reduces to the standard Gross-Pitaevskii theory [8, 9] provided

that the contact interaction approximation [10, 11] has been made. However, more complex many-body states do exist at all barrier heights that are made up of multiple configurations. Being that mean field theory describes only a single configuration, it necessarily lacks all correlation effects that arise between configurations. To this end, multiconfigurational approaches have been developed, which when applied to cold atomic gases in a double well trap geometry, are largely based on two-well limits of continuum models such as the Bose-Hubbard model. Both atomic correlation and, to some extent, mean field interactions arise automatically in these theories. Correlations emerge between configurations while mean field interactions occur in two places: first, directly through the interaction term in the many-body Hamiltonian, and second in the underlying single-particle states that make up the matrix elements in the Hamiltonian as well as each configuration. Previous multiconfigurational efforts found in the BEC literature include only the first type of mean field effect and therefore do not describe general many-body states of the system by superpositions of configurations but rather by superpositions of Fock states, where atom-atom interactions are neglected on the shape of the single-particle states. In these works, the underlying single-particle states have been chosen either to be solutions of the single-particle Schrödinger equation [12] or parameters have been introduced to replace the matrix elements in the Hamiltonian altogether [13, 14, 15, 16, 17]. In other words, mean field effects are not fully included in these treatments. Both theoretical approaches are, therefore, separate and complementary.

It is evident that a more complete description should characterize a general state of the system by a superposition of many configurations in which the underlying

*Electronic address: masiello@u.washington.edu

†Present address: JILA, University of Colorado, Boulder, Colorado 80309-0440, USA

single-particle states include the effects of the condensate mean field. Such a theory would be capable of illuminating the quantum many-body structure of the BEC *throughout* the fragmentation process as well as at the extreme simple and fragmented limits. It is the purpose of the present paper to provide such a formulation.

The challenge to formulate such a theory has already been partially fulfilled by several authors [5, 6, 7, 12, 13, 14, 15, 16, 17]. However, to our knowledge, no satisfactory work has been presented in the cold atom context that fully addresses this task. To this end, we formulate a new approach that variationally combines the Hartree-Fock mean field theory for N identical bosons in two single-particle states with a full diagonalization of the many-body Hamiltonian restricted to a basis of $N + 1$ generalized configurations stemming from each Hartree-Fock configuration. This allows for a description of the Bose gas at zero temperature where atomic correlations emerge between configurations into which mean field effects are built. Due to its composition, our approach incorporates both types of interaction and serves as a didactic device for elucidating where they become important in the fragmentation process as well as how these two intertwined but distinct effects change the system. By choosing the atomic interaction strength to be zero in the bosonic Hartree-Fock mean field equations, our approach recovers the Schrödinger based model developed in [12], while limiting our generalized many-body state to a single configuration recovers the mean field theories of [5, 6]. We acknowledge that a fermionic analog of our model, called multiconfigurational self-consistent field theory, is widely known in quantum chemistry and has been quite successful in accurately describing atomic and molecular electronic structure and geometry both at equilibrium and at dissociation [18, 19, 20]. In fact, a proper description of dissociation of polyatomic molecules in the Born-Oppenheimer approximation [18, 19, 20] is closely related to the problem of fragmentation of a BEC into two or more fragments. In this spirit, we refer to the work developed in this paper as the *multiconfigurational bosonic Hartree-Fock theory*.

The paper is organized as follows. In Section II, the many-body theory of a gas of N identical bosons restricted to a finite model space is discussed. Ground and excited eigenstates of the system are expanded onto a basis consisting of $N + 1$ Fock states made up of two single-particle states. Model calculations, which lack the effects of atomic interactions on the underlying single-particle states, are reviewed where the trapping potential is deformed from an initially single well to a double well geometry.

Section III is entirely devoted to a survey of Hartree-Fock mean field theory for N identical bosons in two single-particle states. It is emphasized that mean field theory describes only a single configurational state and therefore lacks the effects of correlation between multiple configurations. After making the standard contact interaction approximation [10, 11], we briefly discuss imagi-

nary time integration for efficient solution of the resulting coupled nonlinear differential equations. While imaginary time integration schemes are a standard method of solution for the bosonic mean field equations with one single-particle state, *i.e.*, for the Gross-Pitaevskii equation [21], or with multicomponent spinors [22], we highlight our method of maintaining spatial orthogonality between two single-particle states. Solutions of the mean field and Schrödinger equations are compared at various barrier heights in the strongly interacting limit.

Lastly, in the Section IV, we introduce the multiconfigurational bosonic Hartree-Fock theory. For each individual energy level, multiconfigurational Hartree-Fock states of the system are constructed from the variationally optimal linear combination of generalized configurations in which the underlying mean field states are chosen so that the partitioning of atoms between its two states allows the corresponding energy to be minimized. Systematic investigation of the atomic structure of the BEC, as a function of barrier height, throughout the fragmentation process is carried out. An Appendix is devoted to an informal statement of the Hylleraas-Undheim theorem, which justifies such a state-by-state use of the variational theorem. In particular, we discuss our application of this theorem to the optimization of BEC excited states and their corresponding energies.

II. EXACT SOLUTION OF THE MANY-BOSON THEORY IN A RESTRICTED BASIS

The many-body Hamiltonian for a gas of N identical spinless bosonic atoms of mass m at zero temperature interacting via a two-body potential $V(\mathbf{x}, \mathbf{x}') \equiv V(|\mathbf{x} - \mathbf{x}'|)$ is given by [23]

$$\begin{aligned} \hat{H} = & \int \hat{\Psi}^\dagger(\mathbf{x}) \{ (-\hbar^2/2m)\nabla^2 + V_{\text{ext}}(\mathbf{x}) \} \hat{\Psi}(\mathbf{x}) d^3x \\ & + (1/2) \int \hat{\Psi}^\dagger(\mathbf{x}) \hat{\Psi}^\dagger(\mathbf{x}') V(\mathbf{x}, \mathbf{x}') \hat{\Psi}(\mathbf{x}') \hat{\Psi}(\mathbf{x}) d^3x d^3x'. \end{aligned} \quad (1)$$

Here $\hat{\Psi}$ and $\hat{\Psi}^\dagger$ are boson field operators which satisfy the equal time commutation relations $[\hat{\Psi}(\mathbf{x}), \hat{\Psi}^\dagger(\mathbf{x}')] = \delta(\mathbf{x} - \mathbf{x}')$ and $[\hat{\Psi}(\mathbf{x}), \hat{\Psi}(\mathbf{x}')] = [\hat{\Psi}^\dagger(\mathbf{x}), \hat{\Psi}^\dagger(\mathbf{x}')] = 0$. The external trapping potential V_{ext} that we have in mind throughout this paper is one that can be continuously deformed from a single well to a double well geometry.

A. Restriction to Two Single-Particle States

The atomic structure of a BEC confined in a double well trapping potential at zero temperature can be reasonably described in a basis of restricted Fock states of the form [12, 14]

$$|N_1, N_2\rangle = (\hat{b}_1^\dagger)^{N_1} (\hat{b}_2^\dagger)^{N_2} |\text{vac}\rangle / \sqrt{N_1! N_2!}, \quad (2)$$

where N_k atoms are in each of the single-particle states $|\chi_k\rangle = \hat{b}_k^\dagger |\text{vac}\rangle$ for $k = 1, 2$ and the total number of

atoms is fixed at $N = N_1 + N_2$. Here, $|\text{vac}\rangle$ is the vacuum state in which no atoms are present. The operators \hat{b}_k^\dagger and \hat{b}_k are boson creation and annihilation operators that add and remove single atoms in the $|\chi_k\rangle$. They satisfy the basic commutation relations $[\hat{b}_k, \hat{b}_l^\dagger] = \delta_{kl}$ and $[\hat{b}_k, \hat{b}_l] = [\hat{b}_k^\dagger, \hat{b}_l^\dagger] = 0$ for $k, l = 1, 2$. While (2) is certainly not the most general eigenstate imaginable, we are interested mainly in the zero temperature properties of the condensate as its constituent atoms are exchanged between two macroscopically occupied single-particle states. Where the effects of finite temperature and of fragmentation into more than two condensates are negligible, the states (2) provide a rich basis in which to explore the many-body physics of the BEC fragmentation process.

The set of all Fock states of the form (2), with all possible numbers of atoms in each of the two single-particle states, exhausts the restricted N -boson Fock space. That is, the model space is spanned by the collection

$$\{|N, 0\rangle, |N-1, 1\rangle, |N-2, 2\rangle, \dots, |0, N\rangle\}, \quad (3)$$

which is taken as a complete set having $N+1$ elements.

Therefore, it is possible to expand an eigenstate of the many-body Hamiltonian (1) as a linear combination of $N+1$ individual Fock states according to [12, 14]

$$|\Psi^N\rangle_\nu = \sum_{N_1=0}^N C_{N_1}^\nu |N_1, N_2 = N - N_1\rangle, \quad (4)$$

where the expansion coefficient $C_{N_1}^\nu$ expresses the probability amplitude for the ν th excited state of the system to be in $|N_1, N_2\rangle$.

The many-body Hamiltonian that is associated with this model space may be derived from (1) by substitution of the two-state expansion of the boson field operator

$$\hat{\Psi}(\mathbf{x}) = \chi_1(\mathbf{x})\hat{b}_1 + \chi_2(\mathbf{x})\hat{b}_2, \quad (5)$$

where the expansion coefficients $\chi_k(\mathbf{x}) = \langle \mathbf{x} | \chi_k \rangle$ are coordinate space single-particle wavefunctions that are, as yet, unspecified. After some standard algebra, one obtains [23]

$$\begin{aligned} \hat{H} = & \hat{N}_1 h_{11} + \hat{N}_2 h_{22} + [h_{12} + V_{1112}(\hat{N}_1 - 1) + V_{2221}\hat{N}_2]\hat{b}_1^\dagger \hat{b}_2 + [h_{21} + V_{1112}\hat{N}_1 + V_{2221}(\hat{N}_2 - 1)]\hat{b}_2^\dagger \hat{b}_1 \\ & + (1/2)[V_{1111}(\hat{N}_1^2 - \hat{N}_1) + V_{2222}(\hat{N}_2^2 - \hat{N}_2)] + (1/2)V_{1122}[\hat{b}_1^\dagger \hat{b}_1^\dagger \hat{b}_2 \hat{b}_2 + \hat{b}_2^\dagger \hat{b}_2^\dagger \hat{b}_1 \hat{b}_1 + 4\hat{N}_1 \hat{N}_2], \end{aligned} \quad (6)$$

where $\hat{N}_k = \hat{b}_k^\dagger \hat{b}_k$ is the occupation number operator for each state $|\chi_k\rangle$, $\hat{N} = \hat{N}_1 + \hat{N}_2$ is the total particle number operator, and

$$\begin{aligned} h_{kl} &= \int \chi_k(\mathbf{x}) \{(-\hbar^2/2m)\nabla^2 + V_{\text{ext}}(\mathbf{x})\} \chi_l(\mathbf{x}) d^3x \\ V_{klmn} &= \int \chi_k(\mathbf{x}) \chi_l(\mathbf{x}') V(\mathbf{x}, \mathbf{x}') \chi_m(\mathbf{x}) \chi_n(\mathbf{x}') d^3x d^3x' \end{aligned} \quad (7)$$

with $k, l, m, n = 1, 2$ are matrix elements of the single-particle Hamiltonian $h(\mathbf{x})$ and two-body interaction potential $V(\mathbf{x}, \mathbf{x}')$. Throughout this paper, the wavefunctions $\chi_1(\mathbf{x})$ and $\chi_2(\mathbf{x})$ are real-valued functions so that

$$\begin{aligned} V_{klmn} &= V_{lknm} = V_{mnkl} = V_{nmlk} = V_{mlkn} \\ &= V_{lmnk} = V_{knml} = V_{nklm}. \end{aligned} \quad (8)$$

This Hamiltonian is more general than that, for example, of Spekksens and Sipe [12] in that the external potential V_{ext} is not assumed to be symmetric *a priori* since we anticipate the possibility for deformation of a single well into either a symmetric or asymmetric double well. Exclusion of certain nonlinear terms, reduces (6) to a two-state Bose-Hubbard Hamiltonian. In the thermodynamic limit, Bose-Hubbard theory provides, *inter alia*, the standard model for the description of quantum phase transitions [24, 25] in Bose gases confined to optical lattices [26].

With the Hamiltonian (6) and associated eigenstates (4), the many-body Schrödinger equation

$$\hat{H}|\Psi^N\rangle_\nu = E^\nu |\Psi^N\rangle_\nu, \quad (9)$$

which is a linear equation, may be represented in the restricted Fock state basis (3). This results in the set of $(N+1) \times (N+1)$ matrix equations

$$\begin{bmatrix} H_{00} & H_{01} & \cdots & H_{0N} \\ H_{10} & H_{11} & \cdots & H_{1N} \\ \vdots & \vdots & \ddots & \vdots \\ H_{N0} & H_{N1} & \cdots & H_{NN} \end{bmatrix} \begin{bmatrix} C_0^\nu \\ C_1^\nu \\ \vdots \\ C_N^\nu \end{bmatrix} = \begin{bmatrix} C_0^\nu \\ C_1^\nu \\ \vdots \\ C_N^\nu \end{bmatrix} E^\nu \quad (10)$$

with matrix elements and overlap

$$\begin{aligned} H_{N'_1 N_1} &= \langle N'_1, N - N'_1 | \hat{H} | N_1, N - N_1 \rangle \\ \delta_{N'_1 N_1} &= \langle N'_1, N - N'_1 | N_1, N - N_1 \rangle \end{aligned} \quad (11)$$

for $N'_1, N_1 = 0, \dots, N$. Once the wavefunctions χ_1 and χ_2 are specified, the matrix elements h_{kl} and V_{klmn} are defined and can be used to build $H_{N'_1 N_1}$ [50]. Then, the eigenvalue equations (10) can, in principle, be solved to give the expansion coefficients $C_{N_1}^\nu$ and energies E^ν of the ground state and first N excited states of the system. This computation may be repeated as the external

potential V_{ext} is deformed from a single well to a double well where the geometry encourages break up into two condensate fragments. Following Penrose and Onsager [27], a fragmented condensate may be characterized by the presence of at least two large eigenvalues of the single-particle reduced density matrix [28, 29]

$$\gamma^\nu(\mathbf{x}, \mathbf{x}') = \nu \langle \Psi^N | \hat{\Psi}^\dagger(\mathbf{x}) \hat{\Psi}(\mathbf{x}') | \Psi^N \rangle_\nu. \quad (12)$$

Within this model space, the full diagonalization of (10) has been found, in practice, to be achievable for systems with a large number of atoms. This is facilitated by the significant reduction of the Hamiltonian matrix in (10) to a pentadiagonal form [12]. Standard linear algebra routines [30] are well suited for the resulting banded eigenvalue problem.

B. Model Calculations

The energies of the ground and first N excited states of the Hamiltonian (6) have been mapped out as a function of the single dimensionless parameter α in correlation diagrams by Reinhardt and Perry [16] as illustrated in Figure 1. In this work, it has been assumed that the single-particle energies and matrix elements of the two-body potential V , in the standard contact interaction approximation $V(\mathbf{x}, \mathbf{x}') = g\delta(\mathbf{x} - \mathbf{x}')$, can be parameterized according to [51]

$$\begin{aligned} V_{kkkk} &= h_{kk} = \hbar\omega \\ V_{kkkl} &= h_{kl} = -\hbar\omega \exp(-\alpha) \\ V_{kkl} &= \hbar\omega \exp(-\alpha) \end{aligned} \quad (13)$$

in terms of the harmonic oscillator energy $\hbar\omega$ and length $\beta = \sqrt{\hbar/m\omega}$ for a symmetric double well trapping potential with $k \neq l = 1, 2$. Variation of the parameter α allows for a continuous change between strong tunneling (small α) and weak tunneling (large α) regimes. Within this simple ansatz, where none of the parameters depend upon the particle number N , the wavefunctions χ_1 and χ_2 are not specified. The ridge structure in Figure 1 shows the boundary between simple BEC for small α and fragmented BEC for large α . When working in the Fock basis (2) where N_1 and N_2 are the number of atoms localized in the left and right wells of a double well potential, it has been found in [17] that the distribution of Fock states contributing to the ground state below the ridge is binomial in form, while the distribution of Fock states above the ridge reveals the existence of macroscopic quantum superposition states. The study of the correlation diagrams in Figure 1 has led to the prediction that many interesting highly excited states, such as Schrödinger cats, exist in the weak tunneling regime and may be created by phase engineering [17, 31].

Other authors [12] opt to approximate the single-particle wavefunctions χ_1 and χ_2 for a given barrier height by solutions of the single-particle Schrödinger




FIG. 1: (Color online) Parameterized model ground and excited state energies for $g = 0.1 \hbar\omega \cdot \beta^3$ and $N = 20$ as a function of α . Note the energy level mergings and resulting ridge structure separating BEC (small α) and fragmented states (large α). The Fock states below the ridge are delocalized and nondegenerate while the states above the ridge are localized and doubly degenerate. The alternation of line style has been chosen to aid in visualization.




FIG. 2: (Color online) Schrödinger model ground and excited state energies for $g = 0.1 \hbar\omega \cdot \beta^3$ and $N = 20$ as a function of barrier height. Note the qualitative similarities to the parameterized model. The solid curve beginning at $20 \hbar\omega$ is the single-particle Schrödinger energy while the remaining curves are the eigenvalues of the Hamiltonian (6) built from the Schrödinger solutions at each barrier height. Note that the Schrödinger energy $N_1\varepsilon_1 + N_2\varepsilon_2$ differs from the many-body eigenvalues E^ν because it includes no atom-atom interaction. The alternation of line style has been chosen to aid in visualization.

equation

$$h(\mathbf{x})\chi_k(\mathbf{x}) = \varepsilon_k\chi_k(\mathbf{x}) \quad (14)$$

where $\varepsilon_k \equiv h_{kk}/\langle\chi_k|\chi_k\rangle$ is the single-particle Schrödinger energy for state k and where the wavefunctions are independent of the particle number N . The two-state Hamil-

tonian (6) in the contact interaction approximation is then diagonalized. Figure 2 presents the associated energy eigenvalues as a function of barrier height, where χ_1 and χ_2 are computed from (14). A qualitatively similar energy correlation diagram exists with ridge structure marking the phase transition between nondegenerate states and doubly degenerate states. This approach is well justified for weakly interacting atomic gases where the effect of atomic interactions on the shape of the wavefunctions is small, however, it may break down in the strong coupling regime where atom-atom interactions are important enough to affect the value of the parameters or matrix elements themselves. Here, mean fields effects dominate the shape of the atomic wavefunctions.

C. Lack of Mean Field Effects in the Single-Particle Wavefunctions

Diagonalization of the Hamiltonian (6) within the basis of restricted Fock states (3), furnishes a basis representation of the ground state and N excited states due to exchanges of atoms between the two single-particle states $|\chi_1\rangle$ and $|\chi_2\rangle$. As no equations have been derived for these states, their functional form is not specified by this approach and approximate models have been chosen that either define the χ_1 and χ_2 as solutions of the single-particle Schrödinger equation or parameterize the Hamiltonian matrix elements h_{kl} and V_{klmn} directly. Neither method takes into account the effects of the condensate mean field on the shape of the single-particle wavefunctions, which becomes more and more important as the interaction strength between the constituent atoms becomes larger and larger. We will demonstrate, in Section IV, how to build atomic interactions into the underlying single-particle states that enter the matrix elements in the many-body Hamiltonian (6). But first, it is important to discuss mean field theory by itself for identical bosons.

III. MEAN FIELD THEORY FOR IDENTICAL BOSONS

Single-particle wavefunctions χ_k were first introduced in the Fock space approach of the previous section. Within that model, no equations were developed to determine these functions. In this section, we derive a set of equations by minimizing the energy associated with (6) to dictate the functional form of the χ_k . The equations that arise are the bosonic Hartree-Fock equations.

A. Restriction to One Single-Particle State

The mean field theory in which all N atoms occupy the same single-particle state $|\chi\rangle = \hat{b}^\dagger|\text{vac}\rangle$ is known as the Gross-Pitaevskii (GP) theory. The many-body

wavefunction Ψ^N restricted to the Fock state $|N\rangle = (\hat{b}^\dagger)^N|\text{vac}\rangle/\sqrt{N!}$ is approximated by the product

$$\Psi^H(1, \dots, N) = \chi(1)\chi(2) \cdots \chi(N) \quad (15)$$

of single-particle wavefunctions χ . This particular type of product is also called a Hartree product since it involves no symmetrization whatsoever. Variation of the expectation value of the many-body Hamiltonian (1) with $\hat{\Psi} = \chi\hat{b}$ in $|\Psi^H\rangle \equiv |N\rangle$ with respect to χ and subject to the constraint that χ is normalized to unity leads to the GP equation

$$\{(-\hbar^2/2m)\nabla^2 + V_{\text{ext}} + g(N-1)|\chi|^2\}\chi = \mu\chi \quad (16)$$

provided that the contact interaction approximation has been made. The chemical potential μ enters (16) as a Lagrange multiplier which enforces the normalization $\langle\chi|\chi\rangle = 1$. While this approximation provides an excellent mean field description of the simple BEC, it is not flexible enough to characterize the break up of a single condensate into multiple fragments, where potentially several single-particle wavefunctions are macroscopically occupied.

B. Restriction to Two Single-Particle States

A mean field theory for bosons, which generalizes the GP (or Hartree) ansatz by adding a second single-particle state, is the bosonic Hartree-Fock (BHF) theory. The BHF ansatz rests on approximating the many-body wavefunction for N bosons as a *symmetric* product of the two single-particle wavefunctions χ_1 and χ_2 . That is

$$\begin{aligned} \Psi^{\text{BHF}}(1, \dots, N) \\ = \mathcal{S}\{\chi_1(1) \cdots \chi_1(N'_1)\chi_2(N'_1+1) \cdots \chi_2(N'_1+N'_2)\}, \end{aligned} \quad (17)$$

where $\mathcal{S} = (\sqrt{N!})^{-1} \sum_P P$ is the symmetrization operator and P is an operator that permutes the atomic coordinates. We place primes on the occupation numbers N'_1 and N'_2 for reasons that will become evident in Section IV. This BHF wavefunction is also called a single permanental wavefunction in contrast to the single determinantal wavefunction for fermions built from an *antisymmetric* product of single-particle wavefunctions. If we had omitted the symmetrization operator \mathcal{S} all together in (17), then we would have a two-single-particle state Hartree or GP theory. Two-state Hartree theory provides an alternative mean field theory that neglects the quantum-mechanical exchange interaction associated with identical particles.

The coupled BHF equations may be determined by variation of the expectation value of the functional

$$\hat{K}[\chi_1, \chi_2] = \hat{H} - \sum_{kl=1,2} \mu_{kl} \hat{N}'_k (\Delta_{kl} - \delta_{kl}) \quad (18)$$

in the BHF state

$$|\Psi^{\text{BHF}}\rangle \equiv |N'_1, N'_2\rangle = (\hat{b}_1^\dagger)^{N'_1} (\hat{b}_2^\dagger)^{N'_2} |\text{vac}\rangle / \sqrt{N'_1! N'_2!} \quad (19)$$

with respect to the two wavefunctions χ_1 and χ_2 , where \hat{H} is the many-body Hamiltonian (6) restricted to the model space of two single-particle states, $\hat{N} = \hat{N}'_1 + \hat{N}'_2$ is the total particle number operator, and $\Delta_{kl} \equiv \langle \chi_k | \chi_l \rangle$ are the matrix elements of the wavefunction overlap Δ . The second term on the right hand side of this equation adds Lagrange multipliers μ_{kl} whose purpose is to constrain the single-particle wavefunctions to be orthonormal. This leads to the coupled two-single-particle state BHF equations [52]

$$\begin{aligned} h\chi_1 + (N'_1 - 1)\Gamma_1\chi_1 + N'_2[\mathcal{J}_2 + \mathcal{K}_2]\chi_1 &= \mu_{11}\chi_1 + \mu_{12}\chi_2 \\ h\chi_2 + (N'_2 - 1)\Gamma_2\chi_2 + N'_1[\mathcal{J}_1 + \mathcal{K}_1]\chi_2 &= \mu_{21}\chi_1 + \mu_{22}\chi_2, \end{aligned} \quad (20)$$

where $h(\mathbf{x}) = (-\hbar^2/2m)\nabla^2 + V_{\text{ext}}(\mathbf{x})$ is the sum of kinetic energy and external trapping potential, Γ_k accounts for the interaction of one atom in the k th single-particle state with the mean field of $N'_k - 1$ other atoms in the same state, \mathcal{J}_l is the direct interaction between a single atom in the k th single-particle state and the mean field of N'_l atoms in the l th ($l \neq k$) single-particle state, and \mathcal{K}_l is the exchange interaction between states k and l which arises due to the symmetrization of the BHF wavefunction. These equations have already been derived by others [5, 6, 32] and have been further extended to treat identical bosons in arbitrarily many single-particle states [7].

The diagonal Lagrange multipliers μ_{kk} in (20) ensure the proper normalization of the single-particle wavefunctions while the off-diagonal μ_{kl} enforce their orthogonality. If the external potential V_{ext} is symmetric, then the off-diagonal Lagrange multipliers are not necessary as the wavefunctions are automatically spatially orthogonal by symmetry. Koopmans' theorem [19, 20, 33] is satisfied for the diagonal multipliers. Therefore $\mu_{11} = E^{\text{BHF}}[N'_1, N'_2] - E^{\text{BHF}}[N'_1 - 1, N'_2]$ and $\mu_{22} = E^{\text{BHF}}[N'_1, N'_2] - E^{\text{BHF}}[N'_1, N'_2 - 1]$ inherit the roles of chemical potentials [5, 6], where the BHF energy is given by

$$\begin{aligned} E^{\text{BHF}} &= \sum_{k=1,2} N'_k h_{kk} + (1/2) \sum_{k=1,2} N'_k (N'_k - 1) V_{kkkk} \\ &+ (1/2) \sum_{k \neq l=1,2} N'_k N'_l [V_{klkl} + V_{kllk}], \end{aligned} \quad (21)$$

and where h_{kl} and V_{klmn} are matrix elements of the single-particle Hamiltonian $h(\mathbf{x})$ and two-body interaction potential $V(\mathbf{x}, \mathbf{x}')$ in (7). The direct and exchange integrals in (20) are defined as

$$\begin{aligned} \Gamma_k(\mathbf{x})\chi_k(\mathbf{x}) &= \int_V [\chi_k(\mathbf{x}')V(\mathbf{x}, \mathbf{x}')\chi_k(\mathbf{x}')] \chi_k(\mathbf{x}) d^3x' \\ \mathcal{J}_l(\mathbf{x})\chi_k(\mathbf{x}) &= \int_V [\chi_l(\mathbf{x}')V(\mathbf{x}, \mathbf{x}')\chi_l(\mathbf{x}')] \chi_k(\mathbf{x}) d^3x' \\ \mathcal{K}_l(\mathbf{x})\chi_k(\mathbf{x}) &= \int_V [\chi_l(\mathbf{x}')V(\mathbf{x}, \mathbf{x}')\chi_k(\mathbf{x}')] \chi_l(\mathbf{x}) d^3x' \end{aligned} \quad (22)$$




FIG. 3: (Color online) The BHF energy reduces to that of the GP ground state when $N'_1/N = 1$ and to that of the first GP excited state when $N'_1/N = 0$. This corresponds to all particles in the symmetric GP ground state and the first antisymmetric GP excited state respectively. An interaction strength of $4\pi a/\beta = 0.4$ [see (27) below] for 100 atoms in a single well potential was used in this calculation of E^{BHF} .

for $k \neq l$ and $k, l = 1, 2$. Note that the potential Γ_k is a direct interaction that arises only for bosons. There is no analogous term for fermions due to Pauli exclusion.

The BHF equations (20) reduce to the GP equation (16) in the extreme limit of only one occupied single-particle state, *i.e.*, when $N'_1 = N$ and $N'_2 = 0$ or vice versa as demonstrated in Figure 3. However, unlike the GP equation, the BHF equations (20) offer additional freedom in that they accommodate the possibility of various numbers of atoms in each of two single-particle states. By choosing N'_1 atoms to be in the first state and N'_2 atoms to be in the second, a set of BHF equations corresponding to that particular arrangement is obtained. Each arrangement is associated with a symmetrized BHF single-permanent wavefunction like (17). Where BHF equations specify the underlying single-particle wavefunctions χ_1 and χ_2 , we call the Fock state $|\Psi^{\text{BHF}}\rangle \equiv |N'_1, N'_2\rangle$ a *single configuration*. These Fock states are distinguished from those in Section II, which were written without primes as $|N_1, N_2\rangle$, because the single-particle states that make up each single configuration are determined by solving BHF equations. It is then possible to vary the number of particles in each of the single-particle states to find the lowest energy single configuration for a particular trap geometry. This flexibility gives rise to two very different physical meanings for χ_1 and χ_2 . In the first, $N'_1 \sim N$ and there are a relatively small number of atoms in $|\chi_2\rangle$. In this regime, $|\chi_1\rangle$ is a condensate state and state two is a single-particle excited state. The energetic cost of making excitations from $|\chi_1\rangle$ to $|\chi_2\rangle$ is macroscopically large due to the effect of bosonic amplification [10]. In the second, fragmented, regime, N'_1 and N'_2 are *both* on the order of N . In this case, both single-particle states are condensate states. Single-atom excitations between

$|\chi_1\rangle$ and $|\chi_2\rangle$ also cost a macroscopic amount of energy, but now there is also a macroscopic gain of energy as the atom is added to a second macroscopically occupied state [16]. This distinction, which is quite important, will be elaborated on in Section IV.

C. Numerical Integration Method

We numerically solve the coupled BHF equations using a fast Fourier transform based pseudospectral grid method [34, 35] in quasi-one dimension [36]. Note that quasi-one dimension does not mean one dimension but rather that the variation of the BEC density is negligible in the two transverse dimensions and a separation of variables is permissible. Using this approach, the equations are expanded onto a discrete Fourier sine basis with 2^8 fixed grid points [53], which satisfies the proper boundary conditions, and the expansion coefficients are variationally optimized. Rather than solving the time-independent equations (20) self-consistently, we work with their time-dependent version, where μ_{11} and μ_{22} are replaced by $i\hbar(d/dt)$ [22]. We then solve the time-dependent BHF equations by the method of steepest descents in imaginary time [21]. That is, we employ the Wick time rotation $t \rightarrow \tau = it$, which takes $i\hbar(d/dt)$ to $-\hbar(d/d\tau)$, and integrate the pair of coupled nonlinear diffusion equations

$$\begin{aligned} -\hbar(d/d\tau)\chi_1 &= \mathcal{F}_1\chi_1 - \mu_{12}\chi_2 \\ -\hbar(d/d\tau)\chi_2 &= \mathcal{F}_2\chi_2 - \mu_{21}\chi_1 \end{aligned} \quad (23)$$

as an initial value problem in τ , where \mathcal{F}_k is the boson Fock operator for the k th single-particle state ($k = 1, 2$),

$$\mathcal{F}_k(\mathbf{x}) = h(\mathbf{x}) + (N'_k - 1)\Gamma_k(\mathbf{x}) + N'_l[\mathcal{J}_l(\mathbf{x}) + \mathcal{K}_l(\mathbf{x})]. \quad (24)$$

The resulting equations are well-defined once the two-body interaction potential V is identified. As was done in Section II, we make use of the contact potential $V(\mathbf{x}, \mathbf{x}') = (4\pi\hbar^2 a/m)\delta(\mathbf{x} - \mathbf{x}')$. Following the argument of Esry *et al.* [22], the renormalized S -wave scattering length a is taken from multichannel T -matrix calculations using *symmetrized* two-body wavefunctions. Thus, the contact interaction approximation effects the replacement of the symmetric combination of two-body matrix elements in the BHF energy (21) with a single contact potential and not each matrix element separately. In symbols that is

$$V_{klkl} + V_{kllk} \equiv \langle kl|V|kl + lk\rangle \rightarrow (4\pi\hbar^2 a/m)\delta(\mathbf{x} - \mathbf{x}'). \quad (25)$$

The resulting BHF equations (20) are [54]

$$\begin{aligned} \{h + g(N'_1 - 1)|\chi_1|^2 + gN'_2|\chi_2|^2\}\chi_1 &= \mu_{11}\chi_1 + \mu_{12}\chi_2 \\ \{h + g(N'_2 - 1)|\chi_2|^2 + gN'_1|\chi_1|^2\}\chi_2 &= \mu_{21}\chi_1 + \mu_{22}\chi_2. \end{aligned} \quad (26)$$

In practice, the physical coordinates in (26) are scaled by the harmonic oscillator length $\beta = \sqrt{\hbar/m\omega}$ according to $\mathbf{x} = \beta\mathbf{x}'$. In these units, the dimensionless parameter that characterizes the coupling strength is

$$4\pi a/\beta. \quad (27)$$

We choose as an initial condition for the single-particle wavefunction χ_1 the square root of the Thomas-Fermi density $\rho^{\text{TF}}(\mathbf{x}) = [\mu - V_{\text{ext}}(\mathbf{x})]/g(N - 1)$ at a particular barrier height, chemical potential, and value of coupling constant. For a symmetric trapping potential, the second wavefunction χ_2 is taken to be antisymmetric to χ_1 and the off-diagonal Lagrange multipliers are unnecessary. Otherwise, χ_2 need only be orthogonal to χ_1 but constraints are needed to maintain orthogonality. Both wavefunctions are initially normalized so that $\|\chi_1(0)\| = \|\chi_2(0)\| = 1$ and are real-valued. Thus, the overlap matrix elements $\Delta_{kl} = \delta_{kl}$ initially. We then employ the standard relaxation approach, *i.e.*, we subtract a guess $\bar{\mu}$ of the ground state chemical potential from the Fock operator and allow the system to time evolve. The wavefunctions relax according to

$$\chi_k(\tau) \approx \sum_{\mu} \exp(-[\mu - \bar{\mu}_{kk}]\tau/\hbar)\chi_k^{\mu}(0)c_{\mu} \quad (28)$$

for $k = 1, 2$, where the expansion coefficients c_{μ} are projections of the evolving state $|\chi_k(\tau)\rangle$ onto the stationary basis $|\chi_k^{\mu}(\tau)\rangle$. Eventually, all excited states decay away after repetition of this procedure together with intermittent renormalization. If the wavefunctions are initially orthogonal and share the symmetry of the trapping potential, then the wavefunctions that remain are the symmetric and antisymmetric solutions which minimize E^{BHF} for each configuration.

Whenever the external potential V_{ext} is asymmetric, off-diagonal Lagrange multipliers must be introduced to keep the single-particle wavefunctions orthogonal to each other throughout the evolution. The proper choice for the μ_{kl} ($k \neq l$) will ensure that $(d/d\tau)\Delta_{kl}(\mu_{kl}) = 0$ for all time τ . The appropriate μ_{kl} ($k \neq l$) are found by multiplying the first equation in (23) by χ_2 and the second equation in (23) by χ_1 , adding the two equations together, and then performing a volume integral. One then arrives at the following expression for the time derivative of the off-diagonal overlap

$$\begin{aligned} -\hbar(d/d\tau)\Delta_{12} &= \langle\chi_2|\mathcal{F}_1 - \bar{\mu}_{11}|\chi_1\rangle - \mu_{12}\langle\chi_2|\chi_2\rangle \\ &\quad + \langle\chi_1|\mathcal{F}_2 - \bar{\mu}_{22}|\chi_2\rangle - \mu_{21}\langle\chi_1|\chi_1\rangle, \end{aligned} \quad (29)$$

where the $\bar{\mu}_{kk}$ are intermediate guesses of the ground state chemical potentials associated with χ_1 and χ_2 . The time derivative of Δ_{21} is the same, as $\Delta^T = \Delta$. With (29) and the relation $N'_1\mu_{12} = N'_2\mu_{21}$ [6], values for μ_{12} and μ_{21} can be chosen so that the right hand side of (29)

equals zero. Those values are

$$\begin{aligned}\mu_{12} &= \frac{\langle \chi_2 | \mathcal{F}_1 - \bar{\mu}_{11} | \chi_1 \rangle + \langle \chi_1 | \mathcal{F}_2 - \bar{\mu}_{22} | \chi_2 \rangle}{(N'_1/N'_2) \langle \chi_1 | \chi_1 \rangle + \langle \chi_2 | \chi_2 \rangle} \\ \mu_{21} &= \frac{\langle \chi_2 | \mathcal{F}_1 - \bar{\mu}_{11} | \chi_1 \rangle + \langle \chi_1 | \mathcal{F}_2 - \bar{\mu}_{22} | \chi_2 \rangle}{\langle \chi_1 | \chi_1 \rangle + (N'_2/N'_1) \langle \chi_2 | \chi_2 \rangle}.\end{aligned}\quad (30)$$

With this choice, the time derivatives of the off-diagonal matrix elements of Δ are zero. Therefore, if the single-particle wavefunctions are initially orthogonal, then they will stay orthogonal for all time τ regardless of the trap symmetry. For each choice of BHF configuration $|N'_1, N'_2\rangle$, the relaxation algorithm will then find the associated two orthogonal wavefunctions which minimize the energy (21). This is the essence of our integration scheme.

D. Bosonic Hartree-Fock and Schrödinger Wavefunctions

Figure 4 displays the BHF single-particle wavefunctions χ_1 and χ_2 , which are obtained by solving (26) at four different barrier heights 0, 3, 9, 13 $\hbar\omega$. In each panel, χ_1 is associated with the BHF configuration $|N, 0\rangle$ and χ_2 is associated with the BHF configuration $|0, N\rangle$. In these extreme configurations, the single-particle BHF wavefunctions reduce to the GP ground and first excited wavefunctions. The configuration $|N, 0\rangle$ minimizes E^{BHF} at each barrier height. Figure 5 displays the solutions of the single-particle Schrödinger equation $\hbar\chi_k = \varepsilon_k\chi_k$ at the same four barrier heights 0, 3, 9, 13 $\hbar\omega$. Since the Schrödinger equation includes no atom-atom interaction, the Schrödinger solutions have no dependence on the number of particles in each single-particle state. For 100 atoms with an interaction strength of $4\pi a/\beta = 0.4$, it is seen that the BHF wavefunctions follow the Thomas-Fermi result quite well whereas the Schrödinger wavefunctions bear little resemblance to either BHF or Thomas-Fermi solutions.

E. Lack of Correlation Effects

The solutions χ_1 and χ_2 of the BHF equations (20) include the effects of the condensate mean field on their shape. An appropriate symmetrized product of these wavefunctions results in a BHF wavefunction like that in (17). Unlike in the single determinantal case where the wavefunction is defined only up to unitary transformations of its constituent single-particle wavefunctions [37], the BHF single permanental wavefunction seemingly corresponds uniquely to a single configuration. That is, the single permanent

$$\mathcal{S}\{\chi_1(1)\cdots\chi_1(N'_1)\chi_2(N'_1+1)\cdots\chi_2(N'_1+N'_2)\} \quad (31)$$

is in one-to-one correspondence with the single configuration $|N'_1, N'_2\rangle$, while another BHF permanent

$$\mathcal{S}\{\chi_1(1)\cdots\chi_1(N''_1)\chi_2(N''_1+1)\cdots\chi_2(N''_1+N''_2)\} \quad (32)$$

is in one-to-one correspondence with the single configuration $|N''_1, N''_2\rangle$, and so on. All such single configurations can be collected into the set

$$\{|N, 0\rangle, |N-1, 1\rangle, |N-2, 2\rangle, \dots, |0, N\rangle\}, \quad (33)$$

which, like (3), spans an $(N+1)$ -dimensional Fock space but, in addition, has states that are now specified by BHF equations.

Considering that the eigenstates of the many-body Hamiltonian (6) are not, in general, a single configuration but are rather composed of a linear combination of $N+1$ such configurations, it is evident that a single configurational description is quite limiting. In particular, it lacks all effects of correlation that arise between atoms in different configurations. To this end, we formulate a new approach in Section IV that combines the full diagonalization of the many-body Hamiltonian in a restricted basis with mean field theory for the underlying single particle states.

IV. MULTICONFIGURATIONAL BOSONIC HARTREE-FOCK THEORY

The many-boson theory restricted to a Fock basis consisting of two single-particle states and the two-single-particle state BHF mean field theory provide complementary descriptions of the BEC and its fragmentation into two condensates. For arbitrary interaction strength, the BHF approach is well justified whenever the state of the BEC can be described by a single configuration, but breaks down whenever a multiconfigurational description is appropriate. Alternatively, the finite basis representation of the many-body Schrödinger equation accounts for atomic correlation between each Fock state and includes the effects of the condensate mean field directly in the Hamiltonian. However, it does not provide equations that specify the underlying single-particle wavefunctions. The use of parameters or even single-particle Schrödinger wavefunctions may not capture certain properties of the BEC that are associated with strongly interacting atomic gases, where mean field effects on the shape of the single-particle wavefunctions are important.

To this end, we variationally combine the BHF mean field theory of Section III with the restricted Fock state representation of the many-body theory of Section II, allowing for both:

- (1.) The effects of the condensate mean field on the shape of the single-particle wavefunctions.
- (2.) The ability to describe states that are made up of multiple configurations.




FIG. 4: (Color online) BHF single-particle wavefunctions χ_1 and χ_2 versus coordinate x at four different barrier heights 0, 3, 9, 13 $\hbar\omega$. An interaction strength of $4\pi a/\beta = 0.4$ for 100 atoms was used. In each panel, χ_1 corresponds to the BHF configuration $|N, 0\rangle$, while χ_2 corresponds to the BHF configuration $|0, N\rangle$. The BHF energies associated with $|N, 0\rangle$ and $|0, N\rangle$ at zero barrier height appear in Figure 3. Both wavefunctions have been set at the chemical potentials μ_{11} and μ_{22} associated with each configuration. At each barrier height, we plot the corresponding Thomas-Fermi wavefunction as a solid black curve.

The former is necessary in the strongly interacting regime, while the latter is needed to describe condensate fragmentation within our approach. We refer to the union of these disjoint theories as the *multiconfigurational bosonic Hartree-Fock theory* or MCBHF [55]. This theory is rich enough to characterize the atomic structure of the simple BEC and its fragmentation into two condensates at zero temperature as its trapping potential is deformed from a single well to a double well with large barrier height [56].

A. General Theory

The basic idea behind our MCBHF approach is to diagonalize a representation of the many-body Hamiltonian

(6) in a set of basis functions of the form

$$|N_1, N_2; \{N'_1, N'_2\}\rangle, \quad (34)$$

where the total number of atoms $N = N_1 + N_2 = N'_1 + N'_2$, and $N_1, N'_1 = 0, \dots, N$. These basis states are a combination of the Fock states of Section II and the BHF configurations of Section III. We call the states (34) *generalized configuration states* or GCSs because for every underlying BHF configuration $|N'_1, N'_2\rangle$, which has been indicated in (34) by $\{N'_1, N'_2\}$, there are an additional $N + 1$ Fock states $|N_1, N_2\rangle$ that can be built off of this BHF reference configuration. For instance, the subset of GCSs stemming from the particular BHF configuration $|N'_1, N'_2\rangle$ is




FIG. 5: (Color online) Schrödinger single-particle wavefunctions χ_1 and χ_2 versus coordinate x at four different barrier heights 0, 3, 9, 13 $\hbar\omega$. These solutions have no dependence on the number of atoms in each single-particle state. Both wavefunctions have been set at the same BHF chemical potentials that were used in Figure 4. At each barrier height, we plot the corresponding Thomas-Fermi wavefunction with $4\pi a/\beta = 0.4$ as a solid black curve. Little resemblance is seen between the Schrödinger and Thomas-Fermi results.

$$\{|0, N; \{N'_1, N'_2\}\rangle, |1, N-1; \{N'_1, N'_2\}\rangle, |2, N-2; \{N'_1, N'_2\}\rangle, \dots, |N, 0; \{N'_1, N'_2\}\rangle\}. \quad (35)$$

However, there are $(N+1)$ -many underlying BHF configurations in total. The collection of all GCSs (34) can be organized into the set

$$\left\{ \begin{array}{cccc} |0, N; \{0, N\}\rangle, & |1, N-1; \{0, N\}\rangle, & |2, N-2; \{0, N\}\rangle, & \dots, |N, 0; \{0, N\}\rangle, \\ |0, N; \{1, N-1\}\rangle, & |1, N-1; \{1, N-1\}\rangle, & |2, N-2; \{1, N-1\}\rangle, & \dots, |N, 0; \{1, N-1\}\rangle, \\ |0, N; \{2, N-2\}\rangle, & |1, N-1; \{2, N-2\}\rangle, & |2, N-2; \{2, N-2\}\rangle, & \dots, |N, 0; \{2, N-2\}\rangle, \\ \vdots & \vdots & \vdots & \vdots \\ |0, N; \{N'_1, N'_2\}\rangle, & |1, N-1; \{N'_1, N'_2\}\rangle, & |2, N-2; \{N'_1, N'_2\}\rangle, & \dots, |N, 0; \{N'_1, N'_2\}\rangle, \\ \vdots & \vdots & \vdots & \vdots \\ |0, N; \{N, 0\}\rangle, & |1, N-1; \{N, 0\}\rangle, & |2, N-2; \{N, 0\}\rangle, & \dots, |N, 0; \{N, 0\}\rangle \end{array} \right\} \quad (36)$$

containing $(N+1) \times (N+1)$ -many elements of which (35) makes up just one row. These elements span a Fock space that contains the $(N+1)$ -dimensional Fock spaces of Sections II and III in certain limits. For example, the set of all GCSs (36) reduces to the set (3) associated with the Schrödinger model of Section II in the limit of vanishing atom-atom interaction within the BHF equations. This corresponds to the subset of all $N+1$ columns belonging to any

single row of (36), since all rows within each column are equivalent in the noninteracting limit of BHF. Alternatively, the subset of GCSs of the form $|N_1, N_2; \{N'_1 = N_1, N'_2 = N_2\}\rangle$ is identical to the set of BHF configurations (33) of Section III. The diagonal entries of (36) realize such a subset.

For each BHF configuration $|N'_1, N'_2\rangle$, which is determined by solving the BHF equations (20), the MCBHF state vector can be expanded onto the GCS basis (34) according to

$$|\Psi^N; \{N'_1, N'_2\}\rangle_\nu = \sum_{N_1=0}^N C_{N_1}^\nu [N'_1, N'_2] |N_1, N - N_1; \{N'_1, N'_2\}\rangle, \quad (37)$$

where the expansion coefficient $C_{N_1}^\nu [N'_1, N'_2]$ is the probability amplitude for the ν th excited state of the system to be in the GCS $|N_1, N_2; \{N'_1, N'_2\}\rangle$. With the MCBHF states, the many-body Schrödinger equation

$$\hat{H}|\Psi^N; \{N'_1, N'_2\}\rangle_\nu = E^\nu [N'_1, N'_2] |\Psi^N; \{N'_1, N'_2\}\rangle_\nu \quad (38)$$

is represented in the GCS basis (34). In a symmetric trapping potential V_{ext} , the Hamiltonian matrix elements $h_{kl} = \langle k|h|l\rangle$ and $V_{klmn} = \langle kl|V|mn\rangle$ are constructed out of the left- and right-localized states $(1/\sqrt{2})(|\chi_1\rangle \pm |\chi_2\rangle)$, which are obtained from the symmetric and antisymmetric BHF solutions $|\chi_1\rangle$ and $|\chi_2\rangle$ by unitary transformation. Such a transformation can always be performed within h_{kl} and V_{klmn} since they provide only matrix elements for the many-body Hamiltonian (6). Consequently, the Fock space occupation numbers N_1 and N_2 , which are the first two entries of the GCS (34), refer to the number of atoms in left- and right-localized single-particle states, while the BHF occupations numbers N'_1 and N'_2 , which appear in brackets within (34), refer to the number of atoms in symmetric and antisymmetric single-particle states. Full diagonalization in this basis gives rise to a set of energy eigenvalues $E^\nu [N'_1, N'_2]$ and associated eigenvectors $C_{N_1}^\nu [N'_1, N'_2]$ that depend upon the particular BHF reference configuration $|N'_1, N'_2\rangle$. Solving this eigenvalue problem for each of the $N + 1$ BHF configurations results in the set of all MCBHF energies

$$\{E^\nu [0, N], E^\nu [1, N - 1], E^\nu [2, N - 2], \dots, E^\nu [N'_1, N'_2], \dots, E^\nu [N, 0]\}, \quad (39)$$

where $\nu = 0, \dots, N$. The cardinality of this set is $(N + 1) \times (N + 1)$.

We then appeal to the variational principle [38] and Hylleraas-Undheim theorem [39, 40, 41, 42], which is presented in the Appendix without proof. The variational principle asserts that within the model space of MCBHF, the ground state energy E^0 stemming from any BHF configuration is an upper bound to the exact ground state energy and may be systematically reduced by adding more GCSs. Furthermore, by the Hylleraas-Undheim theorem, we know that ν th MCBHF excited state energy E^ν stemming from any BHF configuration is an upper bound to the exact ν th excited state energy, and may also be systematically reduced by adding more GCSs to the basis. Therefore, it is true in general, that the optimal BHF configuration at each energy level is the one that minimizes the energy at that particular level. By choosing the BHF configuration $|N'_1, N'_2\rangle$ that minimizes E^0 , and the BHF configuration $|N''_1, N''_2\rangle$ that minimizes E^1 , and so on, we generate a set of $N + 1$ optimal MCBHF energies

$$\{E^0 [N'_1, N'_2], E^1 [N''_1, N''_2], \dots, E^N [N'''_1, N'''_2]\}, \quad (40)$$

where each energy E^ν may stem from a different BHF reference configuration. Each of these optimal energies has an associated optimal eigenvector that is based off of the same BHF configuration that optimizes the energy. For example, the set of eigenvectors corresponding to the

set (40) is

$$\{C_{N_1}^0 [N'_1, N'_2], C_{N_1}^1 [N''_1, N''_2], \dots, C_{N_1}^N [N'''_1, N'''_2]\}, \quad (41)$$

where $N_1 = 0, \dots, N$ and where each of the individual BHF reference configurations $\{N'_1, N'_2\}$, $\{N''_1, N''_2\}$, and $\{N'''_1, N'''_2\}$ are the same in both (40) and (41). In this way, the MCBHF theory simultaneously optimizes both the underlying basis functions, *i.e.*, the BHF reference configurations, and the expansion coefficients of the GCSs. Therefore, each state vector $|\Psi^N; \{N'_1, N'_2\}\rangle_\nu$ is potentially made up of multiple GCSs between which arise correlations, and where the effects of the condensate's mean field are built into the Hamiltonian as well as into the shape of the underlying single-particle wavefunctions.

We call the MCBHF state vector that has the lowest MCBHF energy at the ν th energy level the optimal ν th MCBHF state vector and denote it by

$$|\Psi^N; \{N'_1^\bullet, N'_2^\bullet\}\rangle_\nu. \quad (42)$$

The bullets symbolize that the energy of the ν th state is minimized in the BHF configuration $|N'_1, N'_2\rangle$. The corresponding set of constituent GCSs forms an optimal $(N + 1)$ -dimensional subset of (36) that spans a Fock space supporting lower energies than either of those in Sections II and III. We now provide an algorithmic method by which to realize this subset.

B. Implementation Algorithm

Implementation of the MCBHF theory begins by solving the BHF equations (26) for a particular trap geometry, say a single well potential, and a particular BHF configuration, say the single configuration [57]

$$|N'_1 = 0, N'_2 = N\rangle \equiv |\Psi^{\text{BHF}}\rangle = (\hat{b}_1^\dagger)^0 (\hat{b}_2^\dagger)^N |\text{vac}\rangle / \sqrt{0!N!}, \quad (43)$$

where there are no atoms in the single-particle state $|\chi_1\rangle$ and N atoms in the single-particle state $|\chi_2\rangle$. We choose V_{ext} to be symmetric so that the solutions χ_1 and χ_2 of the BHF equations are symmetric and antisymmetric. With these single-particle wavefunctions we first calculate the BHF energy (21). We then unitarily transform χ_1 and χ_2 to the left- and right-localized functions $(1/\sqrt{2})[\chi_1 \pm \chi_2]$, and use them to build the matrix elements h_{kl} and V_{klmn} in the many-body Hamiltonian (6). With the MCBHF state vector

$$|\Psi^N; \{0, N\}\rangle_\nu = \sum_{N_1=0}^N C_{N_1}^\nu [0, N] |N_1, N - N_1; \{0, N\}\rangle, \quad (44)$$

full diagonalization of the Hamiltonian yields a basis representation of the ground and excited state energies and associated eigenvectors stemming from the single BHF configuration $|N'_1 = 0, N'_2 = N\rangle$.

Next, we solve the BHF equations (26) associated with the single configuration $|N'_1 = 1, N'_2 = N - 1\rangle = (\hat{b}_1^\dagger)^1 (\hat{b}_2^\dagger)^{N-1} |\text{vac}\rangle / \sqrt{1!(N-1)!}$, in which a single atom has been moved from the second single-particle state to the first. With the BHF solution for this configuration we again calculate E^{BHF} , make a unitary transformation to left- and right-localized states, and then build the matrix elements h_{kl} and V_{klmn} . With the MCBHF state vector

$$\begin{aligned} & |\Psi^N; \{1, N - 1\}\rangle_\nu \\ &= \sum_{N_1=0}^N C_{N_1}^\nu [1, N - 1] |N_1, N - N_1; \{1, N - 1\}\rangle \end{aligned} \quad (45)$$

the full diagonalization of the associated Hamiltonian yields the MCBHF ground and excited state energies and eigenvectors stemming from $|N'_1 = 1, N'_2 = N - 1\rangle$.

We then repeat this procedure for each BHF configuration until we reach the last BHF configuration $|N, 0\rangle = (\hat{b}_1^\dagger)^N (\hat{b}_2^\dagger)^0 |\text{vac}\rangle / \sqrt{N!0!}$ in which all N atoms are in the symmetric state $|\chi_1\rangle$ and no atoms are in the antisymmetric state $|\chi_2\rangle$. Mapping out the complete set of MCBHF energies versus all possible BHF reference configurations at this trap geometry results in a diagram like that displayed in Figure 6. Notice the detailed energy level mergings and splittings that occur in this figure. By appealing to the variational and Hylleraas-Undheim theorems, we find the variationally optimal MCBHF ground state and excited state energies and denote their location by a single bullet. The location




FIG. 6: (Color online) MCBHF ground and excited state energies and BHF ground state energy for $N = 100$ atoms in a single well potential (or zero barrier limit of a double well potential) with $4\pi a/\beta = 0.4$ versus BHF reference configuration. The solid black curve represents the BHF ground state energy. Bullets are placed at the minima of BHF ground state and MCBHF ground and excited energies. The line style is alternated to aid in visualization. This figure is a zoomed in and more detailed version of the first panel in Figure 9.

of these bullets also indicates which BHF configuration provides the optimal MCBHF ground and excited state energy. In this way, we are variationally optimizing both the shape of the single-particle wavefunctions and the expansion coefficients. That is, a nonlinear optimization determines the shape of the BHF single-particle wavefunctions for each BHF configuration while a second linear Hylleraas-Undheim optimization determines the expansion coefficients of the optimal MCBHF state vector $|\Psi^N; \{N'_1, N'_2\}\rangle_\nu$. We then repeat this algorithm as the trapping potential is deformed from a single to a double well geometry.

C. Trapping Potential

The external trapping potential V_{ext} that we use in these calculations is harmonic with a gaussian function centered at the oscillator's minimum. In quasi-one dimension, it has the dimensionless form

$$V_{\text{ext}}(x) = (1/2)[x - L_x/2]^2 + A \exp(-[x - L_x/2]^2/2), \quad (46)$$

where L_x is the spatial grid length and A is the amplitude of the gaussian in oscillator units. In practice, $A = 1 + \varepsilon B$ and B is varied from zero up to 150 in increments of $\varepsilon = 0.1 \hbar\omega$. As the amplitude is increased, the constant $1 + \ln A$ is subtracted off of the potential so that its minimum always lies at $0 \hbar\omega$. All energies in this paper are, therefore, reported as relative to the zero of the trapping potential. In this way, V_{ext} is deformed from an initially single well with zero barrier height to a




FIG. 7: (Color online) MCBHF expansion coefficients $C_{N_1}^{51}$ of the 51st excited state $|\Psi^N; \{56^\bullet, 44^\bullet\}\rangle_{51}$ versus the occupation number in each GCS. The label $\{56, 44\}$ in the state vector signifies that the minimal 51st MCBHF excited state energy has been attained in the BHF configuration $|N'_1 = 56, N'_2 = 44\rangle$. This probability amplitude corresponds to $N = 100$ atoms in a single well trapping potential with dimensionless coupling $4\pi a/\beta = 0.4$.

$$|\Psi^N; \{56^\bullet, 44^\bullet\}\rangle_{51} \approx C_{10}^{51}|10, 90; \{56, 44\}\rangle + C_{11}^{51}|11, 89; \{56, 44\}\rangle + \dots + C_{56}^{51}|56, 44; \{56, 44\}\rangle + \dots + C_{90}^{51}|90, 10; \{56, 44\}\rangle, \quad (47)$$

where $C_{N_1}^{51} \equiv C_{N_1}^{51}[56, 44]$ and where the label $\{56, 44\}$ in each ket signifies that the matrix elements of the many-body Hamiltonian (6) for the optimal 51st excited state have been built out of the single BHF configuration $|N'_1 = 56, N'_2 = 44\rangle$, where $N = 100 = N_1 + N_2$. While this BHF configuration minimizes the energy for the 51st excited state, another BHF configuration may be optimal for the 52nd, and so on. Furthermore, each optimal state may also have a different distribution of expansion coefficients $C_{N_1}^\nu$. In other words, every optimized MCBHF state is potentially derived from a different BHF reference configuration and is potentially made up of a unique superposition of GCSs.

E. Discussion

Allowing the interwell barrier to grow, we repeat the procedure of Section IV B for each and every barrier height. A panorama of MCBHF energies versus BHF configuration is presented in Figure 9 corresponding to four different barrier heights 0, 3, 9, and 13 $\hbar\omega$. The first panel is an enlarged version of Figure 6, where only every fifth line is plotted to aid in visualization. Bullets have again been placed at each of the variationally optimal energies. The structure of the MCBHF ground and ex-

double well with a barrier height of $\varepsilon B = 15 \hbar\omega$. Figure 6 corresponds to zero barrier height.

D. Example: MCBHF Excited State

Consider, for example, the optimal 51st excited state in Figure 6. It is built off of the $|N'_1 = 56, N'_2 = 44\rangle$ BHF reference configuration, which can be read off of Figure 6, and has expansion coefficients $C_{N_1}^{51}[56, 44]$ as shown in Figure 7. This particular optimal MCBHF excited state is not described by the single BHF configuration $|N'_1 = 56, N'_2 = 44\rangle$ displayed in Figure 8 but rather by almost all possible GCSs within this model space. It is roughly of the form

cited state energies versus BHF reference configuration is quite detailed and it is important to explain some of its features. For example, it can be seen in Figure 6 or in the upper two panels of Figure 9 that there are two energetic pathways that can support optimal solutions. One pathway provides the globally optimal solutions while the second pathway provides only locally optimal solutions. One might ask why this structure is present and why it is not symmetric about the $|N/2, N/2\rangle$ BHF configuration. The answer lies in the role of the two single-particle states $|\chi_1\rangle$ and $|\chi_2\rangle$, which are the symmetric and anti-symmetric BHF solutions, as various numbers of atoms are placed in each state.

Consider the BHF reference configuration $|0, N\rangle$ in which there are no atoms in the symmetric state and N atoms in the antisymmetric state. Excited MCBHF states $|\Psi^N; \{0, N\}\rangle_\nu$ are made up of single atom excitations out of the highest energy BHF reference configuration describable in the basis. Furthermore, these excitations move atoms to states that have zero occupation. Due to the effect of bosonic amplification, these excitations of an excited state cost a macroscopic amount of energy [10]. Alternatively, consider the MCBHF state $|\Psi^N; \{N, 0\}\rangle_\nu$, which is built out of the BHF reference configuration $|N, 0\rangle$, in which all atoms are in the symmetric ground state. Single atom excitations out of this state also cost a macroscopic amount of energy, due




FIG. 8: (Color online) BHF single-particle wavefunctions χ_1 and χ_2 versus coordinate x associated with the BHF configuration $|N'_1 = 56, N'_2 = 44\rangle$, which underlies the optimal 51st MCBHF excited state at a barrier height of $0 \hbar\omega$. A dimensionless coupling of $4\pi a/\beta = 0.4$ for 100 atoms was used. Both wavefunctions have been set at the chemical potentials μ_{11} and μ_{22} associated with this configuration. The corresponding Thomas-Fermi wavefunction is plotted as a solid black curve.

to bosonic amplification, as atoms are moved from a macroscopically filled state to empty states. However, these excitation energies lie lower than the previous energies because they correspond to excitations out of the ground state rather than excitations from an excited state. This explains why the energies on the left side of each panel, *i.e.*, $N'_1/N \sim 0$, are larger than the energies on the right, *i.e.*, $N'_1/N \sim 1$. Lastly, consider the MCBHF state $|\Psi^N; \{N'_1 \approx N/2, N'_2 \approx N/2\}\rangle_\nu$, which is approximately built out of the BHF reference configuration $|N/2, N/2\rangle$, in which both single-particle states are macroscopically occupied. Single atom excitations out of $|\Psi^N; \{N/2, N/2\}\rangle_\nu$ *exchange* atoms between two macroscopically occupied states. In distinction to the previous two cases, here there is a macroscopic energy cost to move a single atom out of $|\chi_1\rangle$, but there is also a macroscopic energy gain as the atom is added to $|\chi_2\rangle$. Therefore, the MCBHF energies in this region represent the smallest energy excitations possible within our model. The location of each minimum is biased to right because it is energetically easier to make excitations out of the ground state than out of excited states. This is why the energy pathway on the right provides the global minimum solutions while the energy pathway on the left supports only local minima.

By collecting the set of all optimal energies at each barrier height, *i.e.*, by collecting all the bulleted energies appearing in Figure 9 plus those at all other barrier heights, an energy level correlation diagram similar to those in Figures 1 and 2 can be made. The MCBHF correlation diagram associated with Figure 9 is displayed in Figure 10, where the atom-atom interaction strength

is characterized by $4\pi a/\beta = 0.4$. There are qualitative similarities to those of the parametrized and Schrödinger model calculations such as, for example, the pronounced ridge structure that marks the merging of nondegenerate energy levels to degenerate as the interwell barrier is raised. As the atom-atom interaction strength decreases, the similarity between the results of the MCBHF and the Schrödinger model calculation increases. To this end, the energy level correlation diagram for 100 atoms with $4\pi a/\beta = 0.1$ is presented in Figure 11, which already looks more like Figure 2. To aid in visualization, the second panels in Figures 10 and 11 have been zoomed in precisely at the phase transition between nondegenerate and degenerate energies.

Examining the distribution of GCSs in each of the optimal MCBHF states, we find that below the ridge the distribution is harmonic oscillator like in form, while exotic macroscopic superpositions of GCSs exist above the ridge. This behavior is generic across all barrier heights where ridge structures are present. In order to exemplify these characteristics, we present optimal MCBHF expansion coefficients in Figure 12 taken from Figures 10 and 11 at the particular barrier height of $3 \hbar\omega$. Coefficients are presented that correspond to the optimal MCBHF ground state and first and second excited states, which are below the ridge, as well as to the optimal MCBHF 63rd, 64th, and 65th excited states, which are above the ridge. The two left panels in Figure 12 correspond to an interaction strength of $4\pi a/\beta = 0.4$, while the two right panels correspond to $4\pi a/\beta = 0.1$.

Below the ridge, the optimal MCBHF states are not energetically degenerate. The ground state and first and second excited states at $4\pi a/\beta = 0.4$ are optimized in the BHF reference configurations

$$\begin{aligned} |N'_1 = 99, N'_2 = 1\rangle & \text{ ground state} \\ |N'_1 = 94, N'_2 = 6\rangle & \text{ first excited state} \\ |N'_1 = 93, N'_2 = 7\rangle & \text{ second excited state,} \end{aligned} \quad (48)$$

while the BHF reference configurations

$$\begin{aligned} |N'_1 = 100, N'_2 = 0\rangle & \text{ ground state} \\ |N'_1 = 97, N'_2 = 3\rangle & \text{ first excited state} \\ |N'_1 = 93, N'_2 = 7\rangle & \text{ second excited state} \end{aligned} \quad (49)$$

are appropriate for $4\pi a/\beta = 0.1$. In each case, the optimal MCBHF ground state is described by a binomial distribution of GCSs peaked around $N_1 = 50$ and $N_2 = 50$ in the basis where N_1 and N_2 are the occupation numbers of states that are left- and right-localized in the trapping potential. However, by making a unitary transformation directly on the coefficients of the GCSs taking them back to the symmetric and antisymmetric basis, the expansion coefficients are tightly peaked around a single GCS and the optimal MCBHF ground states become

$$\begin{aligned} |\Psi^N; \{99^\bullet, 1^\bullet\}\rangle_0 & \approx |100, 0; \{99, 1\}\rangle \quad \text{for } 4\pi a/\beta = 0.4 \\ |\Psi^N; \{100^\bullet, 0^\bullet\}\rangle_0 & \approx |100, 0; \{100, 0\}\rangle \quad \text{for } 4\pi a/\beta = 0.1. \end{aligned} \quad (50)$$




FIG. 9: (Color online) MCBHF ground and excited state energies and BHF ground state energy for $4\pi a/\beta = 0.4$ and $N = 100$ versus BHF configuration. Four different barrier heights 0, 3, 9, and 13 $\hbar\omega$ are displayed from left to right and top to bottom. The solid black curve in each panel represents the BHF ground state energy at each barrier height and for each BHF configuration. Bullets are placed at the minima of the BHF ground state and all MCBHF energies. These are the variationally optimal states at each energy level. To aid in visualization, we alternate line style and only display every fifth excited state. The box in the first panel corresponds to Figure 6.

The underlying BHF single-particle wavefunctions associated with (50) are displayed in the upper two panels of Figure 13. It is important to note that these BHF wavefunctions depend strongly on the number of atoms in each single-particle state as well as on the interaction strength and do look quite different from the solutions of the single-particle Schrödinger equation found in the second panel of Figure 5. Being that the MCBHF states (50) are made up of a single configuration with almost all $N = 100$ atoms in $|\chi_1\rangle$, it is not surprising that GP theory [8, 9] provides a good description of the ground state of the simple BEC.

A harmonic oscillator like distribution of expansion coefficients continues from the ground state up to and including the excited states that lie within the ridge at approximately 470 $\hbar\omega$ in Figure 10 and at approximately 190 $\hbar\omega$ in Figure 11. However, above the ridge, the distribution of GCSs making up each optimal MCBHF

state takes on a striking new form. Macroscopic quantum superpositions states emerge in the spectrum. These states, which are pairwise degenerate, are also known as entangled number states and are colloquially called Schrödinger cats. The lower two panels in Figure 12 depict the distribution of GCSs in the optimal 63rd, 64th, and 65th excited MCBHF states at a barrier height of 3 $\hbar\omega$ and corresponding to interaction strengths of $4\pi a/\beta = 0.4$ and $4\pi a/\beta = 0.1$. It is found that these three states are optimized in the BHF reference configurations

$$\begin{aligned}
 |N'_1 = 50, N'_2 = 50\rangle & \quad 63\text{rd excited state} \\
 |N'_1 = 50, N'_2 = 50\rangle & \quad 64\text{th excited state} \\
 |N'_1 = 50, N'_2 = 50\rangle & \quad 65\text{th excited state,}
 \end{aligned} \tag{51}$$




FIG. 10: (Color online) MCBHF energy level correlation diagram versus barrier height for $4\pi a/\beta = 0.4$ and $N = 100$ atoms. Every fifth energy level is plotted in the left panel and the line styles are alternated in both to aid in visualization. The solid black curve in the left panel denotes the BHF ground state energy. Energy level mergings, which lead to a pronounced ridge structure separating nondegenerate oscillator like states from doubly degenerate macroscopic superposition states, are focused on in the second panel. This panel is a zoomed and more detailed version of the boxed region in the first, where every line has been displayed.




FIG. 11: (Color online) MCBHF energy level correlation diagram versus barrier height for $4\pi a/\beta = 0.1$ and $N = 100$ atoms. Every fifth energy level is plotted in the left panel and the line styles are alternated in both to aid in visualization. The solid black curve denotes the BHF ground state energy. Qualitative similarities are evident in comparison of this figure to Figure 10. Differences exist in the shape and placement of the ridge structure. The second panel is a zoomed and more detailed version of the boxed region in the first, where every line has been displayed.

for $4\pi a/\beta = 0.4$, while

$$\begin{aligned}
 |N'_1 = 40, N'_2 = 60\rangle & \quad 63\text{rd excited state} \\
 |N'_1 = 40, N'_2 = 60\rangle & \quad 64\text{th excited state} \\
 |N'_1 = 39, N'_2 = 61\rangle & \quad 65\text{th excited state}
 \end{aligned} \tag{52}$$

provide the optimal BHF reference configurations for $4\pi a/\beta = 0.1$. The underlying BHF single-particle wavefunctions for the 63rd excited states are displayed in the lower two panels of Figure 13. Once again, it is important to note that these BHF wavefunctions depend strongly on the number of atoms in each single-particle state as well as on the interaction strength and do look quite different from the solutions of the single-particle Schrödinger

equation found in the second panel of Figure 5. The optimal MCBHF excited states are of the approximate form

$$\begin{aligned}
 |\Psi^N; \{N'_1, N'_2\}\rangle_\nu & \approx f_{15}^\nu |15, 85; \{N'_1, N'_2\}\rangle \\
 & \quad \pm g_{85}^\nu |85, 15; \{N'_1, N'_2\}\rangle
 \end{aligned} \tag{53}$$

where $\nu = 63, 64, 65$ and where the optimal BHF reference configurations $|N'_1, N'_2\rangle$ have already been given for both interaction strengths. The coefficients f_{15}^ν and g_{85}^ν are meant to indicate that these states are not made up of just two GCSs, but rather there is a distribution of GCSs peaked around $|15, 85; \{N'_1, N'_2\}\rangle$ and $|85, 15; \{N'_1, N'_2\}\rangle$ as is evident from the lower two panels of Figure 12.




FIG. 12: (Color online) Upper two panels: Expansion coefficients $C_{N_1}^\nu$ of the optimal ground state and first and second excited states $|\Psi^N; \{N_1^\bullet, N_2^\bullet\}\rangle_\nu$ versus occupation number in each GCS. Lower two panels: Expansion coefficients $C_{N_1}^\nu$ of the optimal 63rd, 64th, and 65th excited states $|\Psi^N; \{N_1^\bullet, N_2^\bullet\}\rangle_\nu$ versus occupation number in each GCS. The optimal BHF reference configurations $|N_1^\bullet, N_2^\bullet\rangle$ have been provided for each state separately within the text. These probability amplitudes correspond to $N = 100$ atoms at a barrier height of $3 \hbar\omega$. The left column has the dimensionless coupling $4\pi a/\beta = 0.4$, while the right column has $4\pi a/\beta = 0.1$.

These states are, therefore, multiconfigurational in nature. In Figure 12, the “+” combination is for $\nu = 64$ and the “−” combination is for $\nu = 63, 65$. Sharper and more extreme macroscopic superposition states do appear in the spectrum at even higher lying energies.

Furthermore, we should point out that in the large barrier height limit, the BHF reference configuration underlying the optimal MCBHF ground state and low lying excited states approaches $|N_1^\bullet = 50, N_2^\bullet = 50\rangle$ for both interaction strengths. This can be seen by comparing trends in the lower two panels of Figure 9, which correspond to barrier heights of $9 \hbar\omega$ and $13 \hbar\omega$. The distribution of GCSs at these barrier heights is also sharply peaked at $N_1 = 50$ and $N_2 = 50$. Therefore, the optimal MCBHF ground state takes the form

$$|\Psi^N; \{50^\bullet, 50^\bullet\}\rangle_0 \approx |50, 50; \{50, 50\}\rangle \quad (54)$$

in the large barrier height limit, which is a fragmented

BEC state. Since the optimal BHF reference configuration has approximately 50 atoms in each of two single-particle states, it is not surprising that a two-single-particle state BHF based mean field theory provides a realistic description of the fragmented BEC ground state [5, 6]. The MCBHF state (54) is, after all, a single configuration with about $N/2$ atoms in each of two single-particle states.

We have demonstrated that the optimal MCBHF ground states at both zero and large barrier heights are described by a single GCS. The low barrier limit is characterized by only one single-particle state. This is why ground state zero temperature properties of the simple BEC are well understood with GP theory alone. The large barrier height limit requires, at minimum, two single-particle states. GP theory, being a one-single-particle state theory fails in this case, however, the two-




FIG. 13: (Color online) BHF single-particle wavefunctions χ_1 and χ_2 versus coordinate x at a barrier height of $3 \hbar \omega$. The upper two panels correspond to the BHF configuration that optimizes the MCBHF ground state at both interaction strengths. That is, the first panel depicts the BHF configuration $|99, 1\rangle$ associated with $4\pi a/\beta = 0.4$, while the second panel depicts the BHF configuration $|100, 0\rangle$ associated with $4\pi a/\beta = 0.1$. The lower two panels correspond to the BHF configuration that optimizes the 63rd MCBHF excited state at both interaction strengths. That is, the third panel depicts the BHF configuration $|50, 50\rangle$ associated with $4\pi a/\beta = 0.4$, while the fourth panel depicts the BHF configuration $|40, 60\rangle$ associated with $4\pi a/\beta = 0.1$. All wavefunctions have been set at the chemical potentials μ_{11} and μ_{22} associated with each configuration. At each barrier height, we plot the corresponding Thomas-Fermi wavefunction as a solid black curve.

single-particle state BHF theory provides a good description here. In addition, it has been shown that states do exist at all barrier heights that can only be described by multiple configurations in which atom-atom interactions strongly influence the shape of the underlying single-particle states. Therefore, one must include the full effects of the condensate mean field as well as correlation in order to gain understanding of the entire process of fragmentation and not just an understanding of the simple or fragmented condensate ground state.

V. CONCLUSION

Multiconfigurational Hartree-Fock theory has been formulated for the many-body problem associated with a

gas of identical bosonic atoms trapped at zero temperature in potentials that can be continuously deformed from single to double well geometries. A didactic survey of our approach has been presented which clarifies many of the principles and approximations that are found in other relevant approaches from the literature. In the extreme limit of a single configuration, MCBHF theory recovers two-single-particle state mean field theory [5, 6], which includes the effect of the condensate mean field on the single-particle states but lacks the correlation that arises between configurations. In the opposite extreme limit, our approach recovers the exact diagonalization of the many-body Hamiltonian in a restricted two-state basis of Fock states [12]. Atomic correlation arises automatically in this case, but the full effects of atom-atom interaction are missing. In particular, the underlying single-particle

states have no dependence upon the mean field of the condensate. By incorporating both approaches, the MCBHF theory is capable of describing general many-body states of the system that are made up of a superposition of many configurations into which mean field effects are included.

MCBHF theory has been implemented in a systematic study of the many-body atomic structure of the BEC and its fragmentation in a double well trapping potential. Nonlinear and linear optimization has been utilized in conjunction with the variational and Hylleraas-Undheim theorems to find the optimal ground and excited states of the condensate throughout its break up. A variety of interesting delocalized and localized, single configurational and multiconfigurational states have been found to arise throughout the spectrum at all barrier heights. Contributions from the condensate mean field on the underlying single-particle states as well as correlation effects, which arise between configurations, have been emphasized to be essential in order to obtain a more complete microscopic understanding of the condensate's atomic structure throughout the fragmentation process. Future work will be devoted to the construction of an associated dynamical theory that incorporates the rich atomic structure of MCBHF theory.

Acknowledgments

This work was partially funded by the NSF grant PHY-0140091 (W.P.R.). Furthermore, D.M. would like to acknowledge helpful discussions with Prof. B.D. Esry and Prof. C.H. Greene as well as other participants at the Few- and Many-Body Physics in Quantum Liquids and Gases Conference held at the University of Washington's Institute for Nuclear Theory, regarding the contact po-

tential replacement in BHF theory.

APPENDIX: HYLLERAAS-UNDHEIM THEOREM AND OPTIMIZATION OF BOUNDS FOR EXCITED STATE EIGENVALUES

A nonrigorous statement of the Hylleraas-Undheim theorem is as follows: The N energy eigenvalues E_ν of the Hamiltonian $H_{kl} = \langle g_k | H | g_l \rangle$ represented in the orthonormal basis $\{g_1, \dots, g_N\}$ can be ordered in such a way that

$$E_1 \leq E_2 \leq \dots \leq E_N. \quad (\text{A.1})$$

Each E_ν is an upper bound to the exact ν th eigenvalue E_ν^{ex} of the same symmetry, *i.e.*,

$$E_1^{\text{ex}} \leq E_1, E_2^{\text{ex}} \leq E_2, \dots, E_N^{\text{ex}} \leq E_N. \quad (\text{A.2})$$

The bounds can be systematically improved as N is increased since the energy eigenvalues necessarily move downward or stay the same. For example, the eigenvalues E'_ν of the Hamiltonian represented in the orthonormal basis $\{g_1, \dots, g_N, g_{N+1}\}$, which contains one extra basis function, interlace the eigenvalues E_ν in such a way that

$$E_1^{\text{ex}} \leq E'_1 \leq E_1, E_2^{\text{ex}} \leq E'_2 \leq E_2, \dots, E_N^{\text{ex}} \leq E'_N \leq E_N. \quad (\text{A.3})$$

Continued interlacing occurs as more basis functions are added. In this way, bounds for excited states can be independently optimized. A proof of the theorem appears in [39, 41, 42], while the more casual reader is referred to [40].

-
- [1] Y. Shin, M. Saba, T. A. Pasquini, W. Ketterle, D. E. Pritchard, and A. E. Leanhardt, *Phys. Rev. Lett.* **92**, 050405 (2004).
 - [2] M. Saba, T. A. Pasquini, C. Sanner, Y. Shin, W. Ketterle, and D. E. Pritchard, *Science* **307**, 1945 (2005).
 - [3] T. Anker, M. Albiez, R. Gati, S. Hunsmann, B. Eiermann, A. Trombettoni, and M. K. Oberthaler (2004), *cond-mat/0410176*.
 - [4] M. Albiez, R. Gati, J. Fölling, S. Hunsmann, M. Cristiani, and M. K. Oberthaler (2004), *cond-mat/0411757*.
 - [5] L. S. Cederbaum and A. I. Streltsov, *Phys. Lett. A* **318**, 564 (2003).
 - [6] L. S. Cederbaum and A. I. Streltsov, *Phys. Rev. A* **70**, 023610 (2004).
 - [7] O. E. Alon and L. S. Cederbaum (2005), *cond-mat/0502187*.
 - [8] E. P. Gross, *Nuovo Cimento* **20**, 454 (1961).
 - [9] L. P. Pitaevskii, *Sov. Phys. JETP* **13**, 451 (1961).
 - [10] K. Huang and C. N. Yang, *Phys. Rev.* **105**, 767 (1957).
 - [11] K. Huang, *Statistical Mechanics* (John Wiley and Sons, 1987), 2nd ed.
 - [12] R. W. Spekkens and J. E. Sipe, *Phys. Rev. A* **59**, 3868 (1999).
 - [13] G. J. Milburn, J. Corney, E. M. Wright, and D. F. Walls, *Phys. Rev. A* **55**, 4318 (1997).
 - [14] M. J. Steel and M. J. Collett, *Phys. Rev. A* **57**, 2920 (1998).
 - [15] S. Raghavan, A. Smerzi, and V. M. Kenkre, *Phys. Rev. A* **60**, R1787 (1999).
 - [16] W. P. Reinhardt and H. Perry, in *Fundamental World of Quantum Chemistry*, edited by E. J. Brändas and E. S. Kryachko (Cambridge University Press, 2003), vol. 2, p. 385.
 - [17] K. W. Mahmud, H. Perry, and W. P. Reinhardt, *J. Phys. B.* **36**, L265 (2003).
 - [18] R. Shepard, *Adv. Chem. Phys.* **69**, 63 (1987).
 - [19] I. N. Levine, *Quantum Chemistry* (Prentice Hall, New Jersey, 2000), 5th ed.
 - [20] A. Szabo and N. S. Ostlund, *Modern Quantum Chemistry* (Dover Publications, Mineola, 1989).
 - [21] F. Dalfovo and S. Stringari, *Phys. Rev. A* **53**, 2477 (1996).

- [22] B. D. Esry, C. H. Greene, J. P. Burke, Jr., and J. L. Bohn, Phys. Rev. Lett. **78**, 3594 (1997).
- [23] A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems* (McGraw-Hill Book Company, New York, 1971).
- [24] M. P. A. Fisher, P. B. Weichman, G. Grinstein, and D. S. Fisher, Phys. Rev. B **40**, 546 (1989).
- [25] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, 1999).
- [26] D. Jaksch, C. Bruder, J. I. Cirac, C. W. Gardiner, and P. Zoller, Phys. Rev. Lett. **81**, 3108 (1998).
- [27] O. Penrose and L. Onsager, Phys. Rev. **104**, 576 (1956).
- [28] P.-O. Löwdin, Phys. Rev. **97**, 1474 (1955).
- [29] A. J. Leggett, Rev. Mod. Phys. **73**, 307 (2001).
- [30] E. Anderson, Z. Bai, C. Bischof, J. Demmel, J. Dongarra, J. DuCroz, A. Greenbaum, S. Hammarling, A. McKenney, S. Osrtouchov, et al., *LAPACK Users Guide* (Society for Industrial and Applied Mathematics, Philadelphia, 1992).
- [31] K. W. Mahmud, H. Perry, and W. P. Reinhardt, Phys. Rev. A **71**, 023615 (2005).
- [32] B. D. Esry, Ph.D. thesis, University of Colorado (Boulder) (1997).
- [33] T. Koopmans, Physica **1**, 104 (1934).
- [34] M. Frigo and S. G. Johnson, Proceedings of the IEEE **93**, 216 (2005).
- [35] D. Gottlieb and S. A. Orszag, *Numerical Analysis of Spectral Methods: Theory and Applications* (SIAM, Bristol, England, 1977).
- [36] L. D. Carr, C. W. Clark, and W. P. Reinhardt, Phys. Rev. A **62**, 063610 (2000).
- [37] D. J. Thouless, Nuclear Physics **21**, 225 (1960).
- [38] E. Merzbacher, *Quantum Mechanics* (John Wiley & Sons, Inc., New York, 1998), 3rd ed.
- [39] R. G. Newton, *Scattering Theory of Waves and Particles* (McGraw Hill Book Company, 1966).
- [40] R. Peierls, *Surprises in Theoretical Physics* (Princeton University Press, 1979).
- [41] E. A. Hylleraas and B. Undheim, Z. Phys. **65**, 759 (1930).
- [42] J. K. L. MacDonald, Phys. Rev. **43**, 830 (1933).
- [43] B. D. Esry, Phys. Rev. A **55**, 1147 (1997).
- [44] T.-L. Ho and V. B. Shenoy, Phys. Rev. Lett. **77**, 3276 (1996).
- [45] R. J. Ballagh, K. Burnett, and T. F. Scott, Phys. Rev. Lett. **78**, 1607 (1997).
- [46] R. Graham and D. Walls, Phys. Rev. A **57**, 484 (1998).
- [47] E. H. Lieb and W. Liniger, Phys. Rev. **130**, 1605 (1963).
- [48] S. McKinney (now McKagan), Ph.D. thesis, University of Washington (2004).
- [49] We choose to use this terminology, which is adapted from quantum chemistry, because we wish to emphasize the connection between the atomic structure theory of cold atomic gases and the well established theory of electronic structure in atoms and molecules. This relationship has already been promoted in the BEC literature [43].
- [50] The theory of diagonalizing the many-body Hamiltonian in a basis of N -electron determinants made up of the single determinantal ground state plus all possible excited state determinants is well known in quantum chemistry [19, 20]. Since every determinant specifies a configuration, this method is known as configuration interaction.
- [51] The specific parameterization of the matrix elements of h and V given in [16, 31] has been misprinted. Rather, the correct parameterization, which is associated with the energy level correlation diagram in Figure 1, is given in (13). We explicitly indicate the energy units $\hbar\omega$ in these formulae whereas [16, 17, 31] do not.
- [52] Similar equations have been derived for *distinguishable* multicomponent (spinor) BECs [22, 44, 45, 46], however, it is important to note that they are not the same as the BHF equations (20), which describe *identical* bosons.
- [53] We have also performed calculations with 2^9 and 2^{10} fixed grid points but our results have adequately converged with only 2^8 points.
- [54] In a one-dimensional model of interacting bosons [47], the two-body potential $V(x, x') = g\delta(x - x')$ is a real interaction potential and is *not* an effective potential. With this choice for V , the interaction terms $N'_l[\mathcal{J}_l + \mathcal{K}_l]\chi_k$ in the BHF equations (20) would become $2gN'_l|\chi_l|^2\chi_k$ [5, 6, 7] rather than $gN'_l|\chi_l|^2\chi_k$ as we have in (26). This is due to the fact that $\langle kl|V|kl + lk\rangle$ is no longer replaced by an effective potential, but rather each matrix element $\langle kl|V|kl\rangle$ and $\langle kl|V|lk\rangle$ contributes separately to the equations since V is a real scattering potential.
- [55] A fermionic analog of the bosonic MCBHF theory is well established in theoretical chemistry. In the fermionic context, this model is called multiconfigurational self-consistent field theory or MCSCF [18, 19, 20]. It combines self-consistent field theory (*i.e.*, fermionic Hartree-Fock) with configuration interaction. In MCSCF, both the expansion coefficients of the generalized configuration states and the shape of the single-particle wavefunctions enter the theory as variational parameters. The optimal MCSCF ground state and excited states may be variationally determined and are found to lie lower than those found from a separate self-consistent field plus configuration interaction calculation.
- [56] Preliminary work based on MCBHF can be found in [48].
- [57] In practice, we do not use the extreme BHF configurations $|0, N\rangle$ and $|N, 0\rangle$ as they are GP states and not BHF. Rather in these two limits we choose $|\varepsilon, N - \varepsilon\rangle$ and $|N - \varepsilon, \varepsilon\rangle$, where $\varepsilon = 0.01$. Note that solution of the BHF equations does not restrict N'_1 and N'_2 to be integers.