

# Beyond the Landau Criterion for Superfluidity

Sara Ianeselli<sup>1</sup>, Chiara Menotti<sup>1</sup> and Augusto Smerzi<sup>1,2</sup>

*1) Istituto Nazionale di Fisica per la Materia BEC-CRS and Dipartimento di Fisica, Università di Trento, I-38050 Povo, Italy*

*2) Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

(Dated: September 16, 2018)

According to the Landau criterion for superfluidity, a Bose-Einstein condensate flowing with a group velocity smaller than the sound velocity is energetically stable to the presence of perturbing potentials. We found that this is strictly correct only for vanishingly small perturbations. The superfluid critical velocity strongly depends on the strength and shape of the defect. We quantitatively study, both numerically and with an approximate analytical model, the dynamical response of a one-dimensional condensate flowing against an instantaneously raised spatially periodic defect. We found that the critical velocity  $v_c$  decreases by increasing the strength of the defect  $V_0$ , up to to a critical value of the defect intensity where the critical velocity vanishes.

*Introduction.* Dissipationless flow in homogeneous bosonic systems is one of the manifestations of the interplay between interatomic interaction and Bose-Einstein condensation. Bogoliubov ground-breaking calculation of the many-body properties of a dilute condensate showed that this interplay produces a linear dispersion of the low-energy excitation modes which, according to Landau, inhibits dissipation when the group velocity of the fluid is smaller than the sound velocity [1, 2, 3]. However, the question if a Bose-Einstein condensed dilute gas (BEC) really displays this basic characteristics of superfluidity remained open until its experimental realization in 1995. Eventually, several typical superfluidity phenomena like vortices with quantized circulation [4, 5], the scissor mode [6, 7] and Josephson dynamics [8] have been experimentally observed. Instead, the direct experimental verification of the Landau critical velocity has been more elusive. The MIT group studied the response of a trapped elongated condensate perturbed by a blue detuned laser beam [9]. They found a transition between the superfluid and the dissipative regime at a laser velocity smaller than the estimated sound velocity. The reason for this discrepancy is not fully understood, but it might be related to the topological creation of vortices [10]. It is worth noticing that even in superfluid  $^4\text{He}$  the observed superfluid critical velocity is smaller than Landau's prediction [11].

In this paper we study the response of a one dimensional homogeneous condensate flowing with momentum  $p$  against an instantaneously raised defect. The analysis is done numerically, solving the Gross-Pitaevskii equation (GPE), and analytically, with both a dynamical Bogoliubov approach and a nonlinear two-mode approximation. The Bogoliubov linear framework provides a clear dynamical analysis of the Landau criterion, predicting the transition from a superfluid to a dissipative regime at a condensate group velocity equal to the sound velocity. However, and this is the main result of this paper, we show that nonlinear corrections to the Landau criterion become very important even for small strengths of the defect.

We consider, as a defect, a spatially periodic potential. This choice is convenient for two reasons: first, in the linear regime it is possible to selectively excite a given pair of quasi-particle modes, and study their behaviour below and above the critical velocity; second, periodic potentials can be easily tailored and accurately controlled experimentally.

Our analysis has several analogies with the Bragg spectroscopy of condensates, which has been extensively investigated both theoretically and experimentally [12, 13, 14, 15, 16]. The crucial difference is that we focus on the signature of the energetic instability associated with the Landau criterion of superfluidity, and on the study of how the nonlinear dynamics modifies the Landau critical velocity.

In particular, for a spatially periodic defect, we find that the Landau critical velocity  $v_c$  rapidly decreases with increasing defect strength, up to a critical value where superfluidity disappears. This result might be related to the downwards shift of the excitations frequencies with respect to the Bogoliubov spectrum, as predicted in [15]. While the regimes investigated in Bragg spectroscopy experiments are still described with a good accuracy by linear Bogoliubov theory, the experimental investigation of our predictions would require Bragg experiments in highly non-linear regime, i.e. beyond the regimes investigated so far. On the other hand, the investigation of energetical instability in the linear regime of very weak defects, might require times inaccessible experimentally.

*Bogoliubov Analysis.* We describe the BEC dynamics by an effective one-dimensional GPE

$$i\hbar\psi(x, t) = \left[ -\frac{\hbar^2\partial_x^2}{2m} + gnL|\psi|^2 + \Theta(t)V_{def}(x) \right] \psi(x, t) \quad (1)$$

with  $g = 4\pi\hbar^2a/m$ ,  $m$  the atomic mass,  $a$  the  $s$ -wave interatomic scattering length,  $n$  the 3D density and  $L$  the length of the system. Moreover  $V_{def}(x)$  is the defect instantaneously raised at  $t = 0$  and  $\Theta(t)$  the Heaviside theta function. The GPE can be linearized by writing the

condensate wave-function, normalised to 1 in the interval  $L$ , as

$$\psi(x, t) = \frac{1}{\sqrt{L}} e^{i(px - \mu t)/\hbar} \times \left[ 1 + \sum_q \left( c_q(t) U_q e^{i(qx/\hbar - \omega t)} + c_q^*(t) V_q^* e^{i(qx/\hbar + \omega t)} \right) \right], \quad (2)$$

being  $\mu = p^2/2m + gn$  the chemical potential. The coefficients  $c_q(t)$  are time-dependent quasi-particle amplitudes and  $U_q, V_q$ , normalized to  $|U_q|^2 - |V_q|^2 = 1$ , satisfy the Bogoliubov equations with dispersion

$$\hbar\omega_q = \frac{pq}{m} + \sqrt{\frac{q^2}{2m} \left( \frac{q^2}{2m} + 2gn \right)}. \quad (3)$$

The momentum  $q$  represents the momentum of the quasi-particle excitation with respect to the momentum  $p$  of the condensate. The sound velocity in the condensate at rest ( $p = 0$ ) is given by  $c = \sqrt{gn/m}$ . When the condensate momentum  $p < p_B \equiv mc$ ,  $\omega_q \neq 0$  for all  $q$ , while for  $p > p_B$  there exists a value of the momentum  $q$  for which  $\omega_q = 0$ . The time evolution of the coefficients  $c_q(t)$  can be calculated analytically following the quasi-particle expansion approach developed in [16]. We obtain

$$|c_q(t)|^2 = \begin{cases} \sqrt{\frac{q^2}{q^2 + 4mgn}} |I_q|^2 \frac{\sin^2(\omega_q t)}{\hbar^2 \omega_q^2}; & \omega_q \neq 0 \\ \sqrt{\frac{q^2}{q^2 + 4mgn}} |I_q|^2 \frac{t^2}{\hbar^2}; & \omega_q = 0 \end{cases} \quad (4)$$

being  $I_q = \frac{1}{L} \int dx e^{-iqx/\hbar} V_{def}(x)$  the spatial Fourier transform of the defect. According to Eq.(4), when  $\omega_q \neq 0$ , the quasi-particles modes are *stable*: their amplitudes oscillate in time remaining finite. Instead, the amplitude of the *unstable* mode with  $\omega_q = 0$  explodes, leading to a strong dissipation of the condensate kinetic energy.

It is worth emphasizing here the main physical differences between energetic and dynamical instabilities. The latter is characterized by a complex dispersion relation so that, even in absence of perturbing potentials, small fluctuations in the population of the quasi-particle modes increase exponentially fast. On the contrary, when the system is energetically unstable, fluctuations remain small unless the condensate is perturbed by an external defect. In this case, Eq.(4) predicts an increase of the instable mode quadratically in time.

We notice that in the Bogoliubov linear approach, the Landau critical velocity is independent of the shape and the strength of the defect. With a  $\delta$ -like defect  $V_{def} = V_0 \delta(x/L)$ ,  $I_q = V_0$  for all  $q$ , so that all quasi-particle modes are excited. Instead, with a periodic potential  $V_{def} = V_0 \cos(\alpha x/\hbar)$ ,  $I_q = (V_0/2) \delta_{q=\pm\alpha}$  and it is possible to selectively excite the two modes at  $q = \pm\alpha$  [17]. In the Bogoliubov framework it is possible to predict some simple typical behaviours:

1) for  $p \neq 0$  the spectrum becomes asymmetric and the dynamics, involving the two modes at  $q = \pm\alpha$ , will be dominated by the the quasi-particle mode with smaller frequency, since its amplitude is enhanced by  $\omega_q$  at the denominator;

2) this quasi-particle excitation contains two momentum components at  $\pm q$ , the relative weight of which depends on  $\alpha$ : for small  $\alpha$  the phonon character of the excitation will be strong ( $U_q \sim V_q$ ) and both momentum components become important; for large  $\alpha$  the excitation is almost free-particle like and it mostly contains only one momentum component at  $q$ .

From the above considerations, one can deduce that for large  $p$  and large  $\alpha$  only a single momentum component other than the condensate will be considerably excited, permitting to consider a two-mode approximation, as we will discuss in the following.

To inquiry about the energetical stability of the system using a spatially periodic defect, it is necessary to scan the wavevector of the periodic potential  $\alpha$  over all possible values. The Landau criterion can be then stated in the following terms: *below the critical velocity, the depletion from the condensate oscillates with finite amplitude for all  $\alpha$ ; above the critical velocity, there exist a specific value of  $\alpha$  at which the depletion grows quadratically in time*. It is interesting to point out that a  $\delta$ -like defect exciting simultaneously all momenta plays the role of the scan over all possible  $\alpha$  mentioned just above. However, this is of course true only in the linear regime. In the non linear regime mode-coupling makes things much more complicated and of difficult interpretation, so that we restrict ourselves to consider in detail only the case of a spatially periodic defect.

*Numerical Simulations.* Bogoliubov's predictions can be verified numerically by integrating the full GPE. The numerical solutions of the GPE are well approximated by Bogoliubov theory until the amplitude of the depletion is of the order of few percents of the condensate. At longer times, the two evolutions differ dramatically. First of all, as expected, also the population of the unstable mode oscillates, since it cannot grow without limit. However stable modes might oscillate with larger amplitude than the unstable one, and in general they oscillate with periods and amplitudes different from those predicted by Bogoliubov and strongly depending on the value of  $V_0$ . The main consequence is that the clear signature of the onset of energetic instability present in Bogoliubov theory does not exists: for each value of the momentum of the condensate  $p$  and each value of the wavevector  $\alpha$  of the perturbing potential, the excited modes oscillate in time. It is not possible to claim whether for a given  $p$  and a given  $\alpha$  the system is stable or unstable, but a comparison of the results at different  $\alpha$  is needed.

In order to identify a general signature of the energetic instability, we performed several numerical simulations fixing the momentum of the condensate  $p$  and changing the wavevector  $\alpha$  and the strength  $V_0$  of the defect. In Fig. 1, we report the depletion from the condensate and

the population of the momentum components at  $q = \pm\alpha$  as a function of  $\alpha$  for different values of  $p$ . Comparing the results relative to different momenta  $p$ , we observe different situations. For large  $p$ , the depletion presents a discontinuity at a certain value of  $\alpha$ . Below a critical value  $p_c$  this discontinuity disappears. We define operationally the critical velocity for energetic instability as the smallest velocity of the condensate at which we observe such a discontinuity in the depletion. The critical value strongly depends on the strength of the defect  $V_0$ , and, decreasing  $V_0$ , the depletion asymptotically reproduces the Bogoliubov prediction.




FIG. 1: Depletion and maximum amplitude of oscillation of the momentum component at  $q = \pm\alpha$  for  $p = 0.36, 0.48, 0.6, 0.72 p_B$  and  $V_0 = 0.2 gn$ . The thick line is the depletion from the condensate, while the normal and thin lines below are the contributions to it from the momentum component at  $q = -\alpha$  and  $q = +\alpha$  respectively.

At this point we can remark the relation with the previous formulation of the Landau criterion: *below the critical velocity, the depletion varies smoothly over all  $\alpha$ ; above the critical velocity, there exist a specific value of  $\alpha$  at which the depletion makes a jump*, which we take as the signature of the energetic instability. In the limit  $V_0 \rightarrow 0$ , the region around the discontinuity becomes sharper and asymptotically approaches the Bogoliubov prediction.

From the experimental point of view, it would be easier to study the response of the system at a fixed  $\alpha$  as a function of the condensate momentum  $p$  as shown in Fig. 2. The jump in the depletion occurs at a momentum larger than the critical one, and it is a signature of the energetic instability of the system at such value of  $p$ .

Following our definition, the critical velocity can be studied as a function of  $V_0$ , as shown in Fig. 3. For small  $V_0$ , we recover the Landau critical velocity, indicated by the dotted line at  $v = c$ . The limit of very small defect ( $V_0 < 10^{-3} gn$ ) can not be investigated numerically due to the very long time scale of the dynamics. The critical velocity decreases with increasing  $V_0$  and practically vanishes around  $V_0 = gn/2$ . For  $V_0 > gn/2$ , being the strength of the defect larger than the interaction energy,




FIG. 2: Depletion and maximum amplitude of oscillation of the momentum component at  $q = \pm\alpha$  as a function of  $p$  for  $\alpha = 0.21, 0.35, 0.5 p_B$  and  $V_0 = 0.2 gn$ . The thick line is the depletion from the condensate, while the normal and thin lines below are the contributions to it from the momentum component at  $q = -\alpha$  and  $q = +\alpha$  respectively.

the dynamics involves a huge number of excitations, and it is difficult to establish numerically the presence or the absence of the jump.




FIG. 3: Critical velocity as a function of the strength of the defect as defined in the text. The blue dots are the values obtained by GP simulations, the diamonds are obtained from the two-mode effective potential and the dotted line indicates the sound velocity  $c$ .

*Nonlinear two-mode model:* As we have briefly mentioned above, there is a regime (large  $p$ , large  $\alpha$ ) in which a part from the condensate only one single momentum component is considerably populated and one can develop a two-mode model. We write the order parameter as

$$\psi(x, t) = \frac{1}{\sqrt{L}} \sum_k a_k(t) e^{ikx/\hbar}, \quad (5)$$

with  $\sum_k |a_k(t)|^2 = 1$ . The initial conditions are  $a_{k=p} = 1$  and  $a_{k \neq p} = 0$ . Replacing (5) in the GPE, we obtain a set of coupled equations for the amplitudes  $a_k(t)$ , the coupling between different modes being provided by the presence of the defect and being proportional to its strength  $V_0$ . The system of equations shows that the defect couples the condensate to the modes with momentum  $k = p \pm \alpha$ , and to higher harmonics, with momentum  $k = p \pm \ell \alpha$ , with  $\ell$  arbitrary integer. A drastic simplification in this set of coupled equations is obtained noticing that, when both the momentum of the condensate  $p$  and  $\alpha$  are large, only the momentum  $k = p - \alpha$  (for  $p > 0$ ) couples significantly to the condensate. This is consistent to the picture above, which predicted that, for  $p$  and  $\alpha$  large, the Bogoliubov mode with  $q = -\alpha$  lying in the free-particle region of the spectrum ( $U_q \simeq 1$  and  $V_q \simeq 0$ ) dominates the dynamics and gives rise to a single momentum component at  $p - \alpha$ .

The equation of motion for the relative population of the two modes  $z = (N_p - N_{p-\alpha})/N \in [-1, 1]$  is

$$\dot{z}(t)^2 + W[z(t)] = \text{const.}, \quad (6)$$

with

$$W(z) = \frac{-V_0^2(1-z^2) + [\Delta E(1-z) - gn(1-z^2)/2]^2}{\hbar^2} \quad (7)$$

where  $\Delta E = p^2/2m - (p-\alpha)^2/2m$ , and the initial condition are  $z_0 = z(0) = 1$  and  $\dot{z}(0) = 0$ . These equations are similar to the Josephson equations for the double well and many other two-mode nonlinear systems [19]. With a mechanical analogy, the dynamics of the system is described by the motion of a particle in the potential  $W(z)$  in Eq. (6). When the condensate has momentum  $p$  smaller than the critical value, the effective potential displays a double-well shape for every value of  $\alpha$ . The potential barrier remains higher than the initial energy of the system so that the motion is always localized in the right well. On the other hand if the momentum  $p$  of the condensate is larger than the critical value, by decreasing  $\alpha$  the potential  $W(z)$  displays a transition from a double-well to a single-well, i.e. the potential barrier becomes smaller than the initial energy of the system. In this case, the population of the excited mode suddenly increases, qualitatively reproducing the jumps obtained with the GP simulation shown in Figs.(1,2).

We have checked the capability of the two-mode model of reproducing the results of the GP simulations, and confirmed that it works well for large  $p$  and large  $\alpha$ , as shown in Fig. 4.

Close to the critical velocity, the two-mode model does not work quantitatively well and overestimates the critical velocity, as shown in Fig. 3. However it reproduces the decrease of the critical velocity as a function of  $V_0$  and gives correctly the point at which the critical velocity

becomes negligible. Based on this result, we can use the




FIG. 4: Comparison between the results of the two-mode model (green lines) with the results of the GP equation (blue lines with dots).

two-mode model to calculate the critical defect strength, at which the critical velocity vanishes. To that aim, we consider small  $p$  and small  $\alpha$ , i.e. small  $\Delta E$ . Within this assumption the effective potential can be expanded and becomes

$$W(z) \approx -V_0^2(1-z^2) + (gn/2)^2(1-z^2)^2 - gn\Delta E(1-z^2)(1-z). \quad (8)$$

When  $V_0 = gn/2$ , the effective potential undergoes a double to single-well transition for each value of  $p$ : for  $\alpha > 2p$ ,  $\Delta E < 0$  and we have that  $W(0) > W(1)$ ; for  $\alpha \in [0, 2p]$ ,  $\Delta E > 0$  and we obtain that  $W(0) < W(1)$ . This means that at  $\alpha \approx 2p$ ,  $W(z)$  undergoes a transition from double to single-well, and hence the population of the excited mode undergoes a jump, signature of the energetic instability.

*Conclusions.* The Landau criterion for superfluidity predicts that a weakly interacting Bose-Einstein condensate can flow without dissipation when its group velocity is smaller than the sound velocity. We found that the Landau criterion remains valid only in the linear (Bogoliubov) regime or, in other words, for vanishingly small perturbations. Indeed, the superfluidity critical velocity rapidly decreases while increasing the strength of the defect, up to a critical value of the perturbation where the flow is always dissipative. This finding can be tested experimentally with spatially periodic perturbations, and can qualitatively explain the critical velocities lower than sound observed with condensates flowing against large defects.

*Acknowledgements.* We thank F. Dalfovo and W.D. Phillips for useful discussions. This work has been partially supported by the DOE.

- 
- [1] Kerson Huang *Statistical Mechanics*. John Wiley & Sons, Second edition.
- [2] G. E. Astrakharchik and L. P. Pitaevskii, *Phys. Rev. A* **70**, 013608 (2004)
- [3] For a review see A.J. Leggett, *Rev. Mod. Phys.* **73**, 307 (2001)
- [4] J.R. Abo-Shaeer et al., *Science* **292**, 476 (2001).
- [5] E. Hodby et al., *Phys. Rev. Lett.* **88**, 010405 (2002).
- [6] K.B. Davis et al., *Phys. Rev. Lett.* **75**, 3969 (1995).
- [7] O. Maragò et al., *Phys. Rev. Lett.* **86**, (2001).
- [8] Cataliotti et al., *Science* **293** 843 (2001); M. Albiez et al., *Phys. Rev. Lett.* **95**, 010402 (2005)
- [9] R. Onofrio et al., *Phys. Rev. Lett.* **85**, 2228 (2000).
- [10] B.Jackson, J.F.McCann and C.S.Adams, *Phys. Rev. A* **61**, 013604 (2000), **61**, 051603(R) (2000).
- [11] D. R. Tilley and J. Tilley, *Superfluidity and Superconductivity* (Adam Hilger Ltd., Bristol/Boston,1991).
- [12] M. Kozuma et al., *Phys. Rev. Lett.* **82**, 871 (1999); G. Birkel et al., *Phys. Rev. Lett.* **75**, 2823 (1995).
- [13] J.Stengel et al., *Phys. Rev. Lett.* **82**, 4569 (1999).
- [14] J.Steinhauser et al., *Phys. Rev. Lett.* **88**, 120407 (2002).
- [15] C. Tozzo and F. Dalfovo, *New J. Phys.*, **5**, 54 (2003).
- [16] S.A. Morgan, S. Choi, K. Burnett, and M. Edwards, *Phys. Rev.* **A57**, 3818 (1998); P.B.Blakie, R.J.Ballagh and C.W. Gardiner, *Phys. Rev.* **A65**, 033602 (2002).
- [17] With respect to the experiments performed with the Bragg potential, where the condensate is at rest and the periodic potential  $V_B \cos(q_B x - \omega_B t)$  moves with velocity  $\omega_B/q_B$ , this is simply a change of reference frame: in our case the potential is at rest and the condensate moves with velocity  $v = p/m$ . Therefore the significant quantities,  $q_B$  and  $\omega_B$  in the Bragg experiments,  $\alpha$  and  $p$  in our system, are related one to the other in the following way  $\alpha = \hbar q_B$  and  $p = m\omega_B/q_B$ .
- [18] Compare this approach with the case of a defect on the nonlinear discrete Schroedinger equation on a ring, A. Trombettoni, A. Smerzi and A.R. Bishop, *Phys. Rev. Lett.* **88** 173902 (2002)
- [19] Raghavan et al., *Phys. Rev.* **A59**, 59 (1999) and refs therein.