## Anomalous hydrodynamics and "normal" fluids in rapidly rotating BECs

A. Bourne<sup>(1)</sup>, N. K. Wilkin<sup>(1,2)</sup> and J.M.F. Gunn<sup>(1,2)</sup>

<sup>(1)</sup>School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, U. K. and

<sup>(2)</sup> Laboratoire Physique Théorique et Modèles Statistiques,

Bâtiment 100, Université Paris-Sud, 91405 Orsay, France

In rapidly rotating bose systems we show that there is a region of anomalous hydrodynamics whilst the system is still condensed, which coincides with the mean field quantum Hall regime. An immediate consequence is the absence of a normal fluid in any conventional sense. However, even the superfluid hydrodynamics is not described by conventional Bernoulli and continuity equations. We show there are kinematic constraints which connect spatial variations of density and phase, that the positions of vortices are not the simplest description of the dynamics of such a fluid (despite their utility in describing the instantaneous state of the condensate) and that the most compact description allows solution of some illuminating examples of motion. We demonstrate, inter alia, a very simple relation between vortices and surface waves. We show the surface waves can form a "normal fluid" which absorbs energy and angular momentum from vortex motion in the trap. The time scale of this process is sensitive to the initial configuration of the vortices, which can lead to long-lived vortex patches - perhaps related to those observed at JILA.

PACS numbers: 03.75.Kk, 73.43.-f

The area of rapidly rotating Bose Einstein condensates was one of the first to produce predicted phenomena quite distinct to the analogous condensed matter system, <sup>4</sup>He. Initially an instability was found [1] to a noncondensed, Laughlin, state for repulsive effective interactions between the atoms with sufficiently high angular momentum, and the production[1] of a fragmented condensate in the rotating attractive case. Both of these novel features occur in a regime where the atoms reside in restricted set of single particle states, the "Lowest Landau level" (LLL), defined below. Subsequently it has been understood that when the atoms reside in the Lowest Landau level at intermediate amounts of angular momentum mean field theory provides a good description (Mean field quantum Hall regime or MFQHR). In that regime the ground state is a vortex lattice [3, 4] and it has been shown that other phases occur en route to the Laughlin state in the correlated domain [5, 6, 7] as the amount of angular momentum in increased. The nature of these transitions in the thermodynamic limit[8] and in finite systems remains a very active area of research. extending now into anisotropic traps[9, 10].

Following the pioneering studies [11, 12] of the vortex lattice, current experiments are in [13] or near [14, 15] the mean field quantum Hall regime. The correlated regime is still to be investigated. The experiments find that the vortex lattice becomes very soft with an increasingly lengthy period for recovery after disruption. In this Letter we show the underlying hydrodynamics in the LLL is very unconventional and will be a contributory factor to understanding these experimental observations. There has been considerable theoretical discussion [8, 16] about the conventional or unconventional nature of excitations in the vortex lattice; this Letter provides a formulation of the hydrodynamics underlying such excitations. The defining characteristic of the wave functions,  $\psi(x, y)$ , belonging to the lowest Landau level is that  $\psi(z, z^*) = f(z) e^{-|z|^2/2}$ , where z = x + iy and f(z) is analytic (or entire) in z.  $\psi(z, z^*)$  describes the system in the plane perpendicular to the axis of rotation. In general f(z, t) is time-dependent. The standard basis set, reflecting the analyticity of f(z), is  $\{(z^m/\sqrt{\pi m!}) e^{-|z|^2/2}\}$ , where  $m = 0, 1, \cdots$ .

To derive hydrodynamic equations from the Gross-Pitaevskii equation one conventionally represents the wave-function,  $\psi = \sqrt{\rho} e^{i\phi}$ , i.e. in terms of its modulus,  $\sqrt{\rho}$ , and argument  $\phi$ .  $\rho$  is interpreted as the density and  $\phi$  is the order parameter, and in the nonrotating case is the velocity potential of the fluid. In the case of LLL hydrodynamics this motivates a representation of f(z,t) as  $f(z,t) = \sqrt{\rho} e^{i\phi}$ , since the Gaussian factor is time-independent. The analyticity of f (i.e.  $\partial f/\partial z^* = 0$ ) leads to "polar" analogues of the Cauchy-Riemann conditions for the derivatives of  $\tilde{\rho}$  and  $\phi$ :

$$\frac{1}{2}\frac{\partial\tilde{\rho}}{\partial x} = \tilde{\rho}\frac{\partial\phi}{\partial y} \text{ and } \frac{1}{2}\frac{\partial\tilde{\rho}}{\partial y} = -\tilde{\rho}\frac{\partial\phi}{\partial x}$$
$$\Leftrightarrow \frac{\partial\ln\sqrt{\tilde{\rho}}}{\partial x} = \frac{\partial\phi}{\partial y} \text{ and } \frac{\partial\ln\sqrt{\tilde{\rho}}}{\partial y} = -\frac{\partial\phi}{\partial x}$$

The importance of these relations is that they constrain kinematically the spatial variation of the density (via  $\rho = \tilde{\rho} e^{-|z|^2}$ ) and phase unlike conventional superfluid hydrodynamics. In addition, non-locality due to projection of the interaction term to the LLL affects both the continuity and "Bernoulli" equations, making them difficult to use.

With these shortcomings in mind, we incorporate the analyticity of the wave-function from the start. There are clearly two complementary representations [2, 3, 4] which accomplish this. Firstly the Taylor expansion of f(z,t)

may be written as:

$$\psi(z, z^*, t) = f(z, t) e^{-|z|^2/2} = \sum_{m=0}^{\infty} a_m(t) z^m e^{-|z|^2/2}.$$

By the fundamental theorem of algebra, we may factorise the polynomial in terms of its roots  $\zeta_{\alpha}(t)$  (4):

$$\psi(z, z^*, t) = S^{-1/2} \prod_{\alpha=1}^{\infty} (\zeta_{\alpha}(t) - z) e^{-|z|^2/2}$$

where S is the normalisation integral. The points  $\zeta_{\alpha}(t) = X_{\alpha}(t) + iY_{\alpha}(t)$  are the vortex positions, or coordinates, at time t.

We immediately see that there is an important distinction between a condensate wave-function residing in the LLL and a general condensate wave-function. In the former case the wave-function is *completely specified* by the position of the vortices. In particular, the density is specified by the vortex positions - there are no independent sound waves (long wavelength Bogoliubov quasiparticles), unlike the general case. There is thus *no possibility* of the existence of a normal fluid component in a conventional sense.

Since the vortices uniquely determine the wavefunction in the LLL, it is natural to re-express the Hamiltonian in terms of the vortex coordinates,  $\zeta_{\alpha}(t)$ . We use the standard model Hamiltonian for N particles in a rotating reference frame:

$$\mathcal{H} = -\frac{1}{2} \sum_{n=1}^{N} \nabla_n^2 + \frac{1}{2} \sum_{n=1}^{N} r_n^2 + \frac{1}{2} \eta \sum_{n \neq n'=1}^{N} \delta(\mathbf{r}_n - \mathbf{r}_{n'}) - \omega \sum_{n=1}^{N} L_{\tau}^2$$

Units of length,  $\ell_0$ , and energy,  $\hbar\omega_0$ , are those provided by the harmonic trap; angular momenta,  $L_n^z$  are scaled by  $\hbar$ . There are two remaining dimensionless parameters. Firstly,  $\omega$ , is the angular velocity of the rotating frame divided by the natural frequency of the harmonic trap. Secondly the coupling constant,  $\eta = 4\pi \bar{n} a \ell_0^2$ .

Then the Hamiltonian takes the form (assuming M vortices):  $\mathcal{V}$ , inter-particle potential,  $\mathcal{U}$  energies and the rotating frame term,  $\mathcal{R}$ , are defined as (using  $\dot{=} d/dt$ ):

$$S = \pi N \sum_{m}^{M} |P_{M-m}^{*}(\zeta)|^{2} m!$$
$$\mathcal{K} = \mathcal{V} = \frac{1}{2} \pi N \sum_{m}^{M} |P_{M-m}^{*}(\zeta)|^{2} (m+1)!$$
$$\mathcal{R} = \pi N \omega \sum_{m}^{M} |P_{M-m}^{*}(\zeta)|^{2} mm!$$
$$= \frac{\lambda}{4} N (N-1) S^{-2} \sum_{m=n}^{M} n n q = 0 P_{M-m}^{*}(\zeta) P_{M-n}^{*}(\zeta)$$

$$\mathcal{U} = \frac{\lambda}{4} N(N-1) S^{-2} \sum_{m,n,p,q=0}^{M} P_{M-m}^{*}(\zeta) P_{M-n}^{*}(\zeta)$$
$$P_{M-p}(\zeta) P_{M-q}(\zeta) (p+q)! 2^{-(p+q)} \delta_{m+n,p+q}$$

where the quantity  $P_n(\zeta) = \sum_{i_1 < i_2 < \cdots < i_n} \zeta_{i_1} \zeta_{i_2} \cdots \zeta_{i_n}$  is the *n*th symmetric polynomial in the variables  $\zeta_{\alpha}$ .

It is instructive to contrast this expression with the corresponding expression for vortices in an incompressible medium in a container of size R:

$$\mathcal{H} = -\frac{1}{2}\Gamma^2 n_0 \sum_{\alpha \neq \beta = 1}^M \ln |\zeta_\alpha - \zeta_\beta| - \omega \Gamma n_0 (R^2 - |\zeta_\alpha|^2) + \text{images}$$

where  $\omega$  is the unscaled angular velocity of the rotating frame,  $\Gamma$  is the circulation of the vortices (assumed to be identical) and  $n_0$  is the constant density.

There are two striking differences. Firstly [2], the interaction between the vortices is analytic and is not pairwise, with multi (up to M-) vortex contributions. Secondly the expression for the angular momentum of the fluid in terms of the vortices (entering the term involving  $\omega$ ) is not a sum of single vortex contributions but is collective in nature in the LLL case.

To determine the equations of motion of the condensate constrained to reside in the LLL, we employ a variational formulation[17, 18] where the LLL wavefunction is a trial function leading to the Lagrangian,  $\mathcal{L} = \mathcal{T} - \mathcal{H}$ , where (assuming  $\psi(z, z^*, t)$  is normalised)

$$\mathcal{T} = \pi N \sum_{m=0}^{M} m! \frac{i}{2} (P_{M-m}^{*}(\zeta) \dot{P}_{M-m}(\zeta) - P_{M-m}(\zeta) \dot{P}_{M-m}^{*}(\zeta))$$

If the corresponding action is varied with respect to the  $\zeta_{\alpha}$  the resulting equations of motion for the vortex coordinates are rather complicated. However the form of  $\mathcal{L}$  shows that the vortex coordinates always enter in the form of the symmetric polynomials,  $P_n(\{\zeta_{\alpha}\})$ . This suggests that their numerical values should be used as the primary variables. Since, up to a sign, the  $P_n(\{\zeta_{\alpha}\})$  are the coefficients in the polynomial whose roots are the  $\zeta_{\alpha}$ , there is a one-to-one relation between the vortex coordinates and the symmetric polynomial values. The dynamics are most simply expressed in the variables  $\rho_m(t)$ and  $\theta_m(t)$  implicitly defined by:

$$(-1)^m P_{M-m}(\zeta) = a_m = \sqrt{\frac{\rho_m}{\pi m!}} e^{\mathrm{i}\theta_m}$$

In particular normalisation can be easily enforced via the  $\rho_m$ .

In terms of  $\rho_m$  and  $\theta_m$  we find ('indicates a time derivative and  $\bar{\lambda} = \lambda (N-1)/4\pi$ ):

$$\mathcal{L} = N \left\{ S^{-1} \sum_{m=0}^{M} \rho_m [-\dot{\theta}_m - (1 + m[1 - \omega])] - \bar{\lambda} \sum_{m,n,p,q=1}^{M} \sqrt{\rho_m \rho_n \rho_p \rho_q} e^{i[(\theta_m + \theta_n) - (\theta_p + \theta_q)]} \sqrt{\frac{(m+n)!}{2^{m+n} m! n!}} \sqrt{\frac{(p+q)!}{2^{p+q} p! q!}} \delta_{p+q,m+n} \right\}$$

where  $S = \sum_{m=0}^{M} \rho_m$ .

Differences between incompressible[22] or Thomas-Fermi[23] and LLL vortex dynamics are already apparent in the case of m vortices symmetrically arranged and equidistant with respect to the centre of the trap. Up to a phase, the vortices are in the directions of the mth roots of unity in the  $\zeta$ -plane, and hence the wave-function is of the form  $\phi(z) \sim (a_0(t) + a_m(t)z^m)$ . We will initially restrict ourselves to a reduced Lagrangian only involving the  $\rho_m$ ,  $\theta_0$  and  $\theta_m$  variables (as  $\rho_0$  can be eliminated by the normalisation constraint  $1 = \rho_0 + \rho_m$ ):

$$\begin{aligned} \frac{\mathcal{L}}{N} &= -\dot{\theta}_0 - 1 - \bar{\lambda} \\ &+ \rho_m \left( \left[ -\dot{\theta}_m - \dot{\theta}_0 \right] - m[1 - \omega] - \bar{\lambda} [2^{-(m-2)} - 2] \right) \\ &+ \rho_m^2 \bar{\lambda} \left( 2^{-(m-2)} - 1 - \frac{(2m)!}{(m!)^2} 2^{-2m} \right) \end{aligned}$$

 $\theta_0$  is not determined and its time derivative may be interpreted as the zero of energy in the original quantum problem.

The significant Euler-Lagrange equation is that associated with  $\rho_m$  (the equation for  $\theta_m$  expresses the unitarity of the original time evolution, in this case  $\dot{\rho}_m = 0$ ), yielding the excitation frequency  $\omega_m$ :

$$\begin{split} \omega_m &= -\dot{\theta}_m + \dot{\theta}_0 \\ &= m(1-\omega) - 2\bar{\lambda}(1-2^{-(m-1)}) \\ &+ \rho_m 2\bar{\lambda} \left( 1 + \frac{(2m)!}{(m!)^2} 2^{-2m} - 2^{-(m-2)} \right) \end{split}$$

This expression describes, in a unified manner, several physical phenomena: surface wave excitations, instabilities and anomalous vortex dynamics.

We may re-express the equation for  $\omega_m$  in terms of the radial distance of the vortices from the centre,  $\zeta$ , as  $\rho_m = 1/(1 + \zeta^{2m}/m!)$ . This implies a natural scale for the radial vortex positions of  $\zeta_m = (m!)^{1/(2m)} \sim \sqrt{m}$ , which accords with the extent of the condensate with mquanta of angular momentum. For  $\zeta \lesssim \zeta_m$ , the vortices are "inside" the condensate, and for  $\zeta \gtrsim \zeta_m$  the vortices are in the evanescent tail.

Thus in the limit that the vortices are very far from the centre of the trap (i.e.  $\rho_m \to 0$ ) we recover the linear surface excitation frequencies derived by Kavoulakis et al [19]. We are thus immediately provided with a representation of surface waves in the LLL of angular momentum m in terms of m vortices. A relationship between vortices and surface waves in the Thomas-Fermi limit has been discussed by Anglin [20] and Tsubota et al[21].

To obtain the equilibrium radial distance of the vortices,  $\zeta_{eq}(\omega)$ , we set  $\omega_m = 0$  so that the vortices are stationary in the rotating frame. We find  $\zeta_{eq}(\omega) = \infty$ for  $\omega$  less than a critical rotation frequency,  $\omega_c^{(m)} =$ 

$$1 - 2\overline{\lambda} \frac{1 - 2^{-(m-1)}}{m}$$
. For  $\omega > \omega_{c}^{(m)}$ , we see:

$$\frac{\zeta_{\rm eq}^{2m}(\omega)}{m!} = \frac{2\lambda}{m(\omega - \omega_{\rm c}^{(m)})} \left(1 + \frac{(2m)!}{(m!)^2} 2^{-2m} - 2^{-(m-2)}\right)$$

These critical frequencies do not have a simple physical interpretation as shape instabilities of the condensate - as, for example,  $\omega_c^{(2)}$  coincides with the frequency of a first order transition of a single vortex to the centre of the trap. This is unlike the Thomas-Fermi limit[24].

One might expect the dynamics of the vortices when they are "inside" the trap,  $\zeta \leq \zeta_m$ , to be reminiscent to incompressible vortex dynamics as the phases of the wave-functions are identical. However the LLL dynamics is completely different. For example as the separation of the vortices vanishes, the frequency  $\omega_m(\zeta = 0)$  obeys:

$$\omega_m(\zeta = 0) = m(1-\omega) + 2\bar{\lambda}2^{-2m} \left(\frac{(2m)!}{(m!)^2} - 2\right)$$
$$\underset{m \to \infty}{\sim} m(1-\omega) + \frac{2\bar{\lambda}}{\sqrt{\pi m}}$$

It does not diverge, in stark contrast with the incompressible case. There the frequency,  $\Omega$ , of rotation of such a polygonal array of m vortices, each with circulation  $\Gamma$ , a distance  $\zeta$  from the centre is[22] (in the laboratory frame)  $\Omega = \Gamma(m-1)/(4\pi\zeta^2)$ .

We will now show that energy and angular momentum can be deposited in surface waves (vortices at large distances) by vortices near the centre of the trap. In other words the surface waves can in part play the role of a "normal fluid". Consider *n* vortices which are all initially very near the centre of the trap. A linearised approximation (where  $\rho_n \simeq 1$ ) allows coupling to all the angular momentum channels which conserve angular momentum (i.e.  $m \leq 2n$ ). This implies we will have *n* additional vortices, which will turn out to be at large distances from the centre of the trap. Because of angular momentum conservation  $\rho_{n-m} - \rho_{n+m}$  is a constant, which we choose to be zero. Then the equations for  $R_m = \frac{1}{2}(\rho_{n-m} + \rho_{n+m})$ and  $\sigma_m = (\theta_{n-m} - \theta_n) + (\theta_{n+m} - \theta_n)$  are:

$$\dot{R}_m = -\Lambda_m R_m \sin \sigma_m \dot{\sigma}_m = \chi_m - \Lambda_m \cos \sigma_m$$

Where

$$\Lambda_m = 4\bar{\lambda} \frac{(2n)!}{2^{2n}n!\sqrt{(n+m)!(n-m)!}}$$
  

$$\chi_m = 4\bar{\lambda} \frac{1}{2^{2n}n!} \left[ \frac{(2n)!}{n!} - \frac{(2n-m)!}{2^{-m}(n-m)!} - \frac{(2n+m)!}{2^{+m}(n+m)!} \right]$$

These equations are readily integrated and one finds exponentially unstable growth for  $n + c\sqrt{n} \gtrsim m > n + 1$ and  $n - 1 > m \gtrsim n - c\sqrt{n}$ , in the limit where  $n \gg 1$ , where  $c = 2(-\ln[\sqrt{2}-1])^{1/2}$ . The extent of this unstable region is determined by the spatial overlap of the states with the state n. Oscillatory behaviour occurs for other values of m. The time constant,  $\tau_m$ , for the growth is  $\tau_m^{-1} = \frac{1}{2}\sqrt{\Lambda_m^2 - \chi_m^2}$ . For the most divergent cases,  $m* \simeq n \pm 2(\ln[2/(\sqrt{5}-1)])^{1/2}\sqrt{n}$ , we find  $\tau_{m*n \to \infty}^{-1} \bar{\lambda}/\sqrt{n}$ . If we assume that initial values of  $\rho_m$ ,  $\rho_m(0)$ , are of

If we assume that initial values of  $\rho_m$ ,  $\rho_m(0)$ , are of the order of  $\epsilon \ll 1$ , and then the unstable modes grow by time t to  $\rho_{m^*} \sim \delta$  where  $\epsilon \ll \delta \ll 1$ , then we can deduce the corresponding vortex positions from the crude representation of the polynomial:

$$z^{n} + \delta(z^{n-\sqrt{n}} + \dots + z^{n-1} + z^{n+1} + \dots + z^{n+\sqrt{n}}) + \epsilon(1 + z + \dots + z^{n-\sqrt{n-1}} + z^{n+\sqrt{n+1}} + \dots + z^{2n}) = 0$$

Noting that if z is a solution so is 1/z, let us look for solutions for large |z|. Keeping the largest terms multiplying each of the two small parameters, we obtain

$$1 + x + \frac{\epsilon}{\delta\sqrt{n}} x^{\sqrt{n}} \simeq 0$$

where  $z^{\sqrt{n}} = x/\delta$ . Balancing the middle term with either the final term or with unity, we find  $\sqrt{n}$  approximate solutions  $z^{\sqrt{n}} = -1/\delta$  and  $n - \sqrt{n}$  approximate solutions  $z^{\sqrt{n}(\sqrt{n-1})} = -\delta/\epsilon$ . (We assume  $\epsilon \ll \delta^{\sqrt{n}}$ .)

Geometrically, the roots form four concentric rings, two inside the trap and two outside. The closest and the most remote rings do not change their moduli significantly, but the rings with moduli  $|\delta|^{\pm 1/\sqrt{n}}$  change with time. The inner ring of  $\sqrt{n}$  vortices, moving outwards and the outer ring (with the same number) moving inwards. Each member of the inner ring has a corresponding member of the outer ring with the same argument. This demonstrates the exchange of angular momentum between the inner and outer rings.

This description assumes that one draws the initial conditions from the same probability distribution for each coefficient in the polynomial - which leads to rings of roots[25, 26]. However for initial vortex positions forming a cloud distributed normally around the origin in the complex plane, then the coefficients of the polynomial are very strongly peaked [25] at  $m \simeq n/2$ . Such a configuration will evolve much more slowly as it has negligible initial weight in the unstable modes,  $n - \sqrt{n} \lesssim m < n$ . A simple estimate using the expressions above yields time scales for evolution  $t_{\rm ring} \sim \sqrt{n} \ln n$  and  $t_{\rm cloud} \sim n^{3/2}$ . It is tempting to compare this distinction with the JILA experiments<sup>[27]</sup> where long-lived concentrations of vortices were observed. Although the experiments were not conducted under LLL conditions on average, in the lowdensity environment of the vicinity of the vortices the dynamics might be of a LLL nature.

NKW and JMFG would like to thank KITP, University of California at Santa Barbara, the Aspen Center for Physics and Gora Shlyapnikov for hospitality at Université Paris-Sud while parts of this work was performed and thank him, Nigel Cooper and Martin Long for several helpful discussions. This work was also supported by EPSRC grants GR/R80865 (NKW) and GR/R00920 (JMFG).

- N.K. Wilkin, J.M.F. Gunn and R.A. Smith, Phys. Rev. Lett. 80, 2265 (1998).
- [2] Z. Tesanovic and L. Xing, Phys. Rev. Lett. 67, 2729 (1991);
   Z. Tesanovic, Phys. Rev. B 44 12635 (1991).
- [3] D.A. Butts and D.S. Rokhsar, Nature (London), 397, 297 (1999).
- [4] T.-L. Ho. Phys. Rev. Lett. 87, 060403 (2001)
- [5] N.R. Cooper and N.K. Wilkin, Phys. Rev. B 60, R16279, (1999).
- [6] N.K. Wilkin and J.M.F. Gunn Phys. Rev. Lett. 84, 6 (2000).
- [7] N.R. Cooper, N.K. Wilkin and J.M.F. Gunn, Phys. Rev. Lett. 87, 120405 (2001).
- [8] J. Sinova, C.B. Hanna and A.H. Macdonald, Phys. Rev. Lett. 89, 030403 (2002).
- [9] S. Sinha and G.V. Shlyapnikov, Phys. Rev. Lett. 94, 150401 (2005)
- [10] M.I. Parke and N.K. Wilkin, (To be published).
- [11] K. W. Madison, F. Chevy, W. Wohlleben and J. Dalibard, Phys. Rev. Lett. 84, 806 (2000).
- [12] J.R. Abo-Shaeer, C. Raman, J.M. Vogels and W. Ketterle, Science **292**, 476 (2001).
- [13] V. Schweikhard, I. Coddington. P. Engels, V.P. Mogendorff and E.A. Cornell, Phys. Rev. Lett. 92, 040404 (2004).
- [14] V. Bretin, S. Stock, Y. Seurin and J. Dalibard, Phys. Rev. Lett. 92, 050403 (2004).
- [15] I. Coddington, P. Engels, V. Schweikhard and E.A. Cornell, Pys. Rev. Lett. **91**, 100402 (2003).
- [16] G. Baym, Phys. Rev. Lett. 91, 110402 (2003); G. Baym, Phys. Rev. A 69, 043618 (2004).
- [17] P. Kramer and M. Saraceno Geometry of the Time-Dependent Variational Principle in Quantum Mechanics, Lecture Notes in Physics 140 (Springer-Verlag, Berlin, 1981).
- [18] M. Linn and A.L. Fetter, Phys. Rev. A 61, 063603 (2000).
- [19] G. M. Kavoulakis, B. Mottelson and C. J. Pethick, Phys. Rev. A 62, 063605 (2000).
- [20] J.R. Anglin, Phys. Rev. Lett. 87, 240401 (2001).
- [21] M. Tsubota, K. Kasamatsu and M. Ueda, Phys. Rev. A 65, 023603 (2002); K. Kasamatsu, M. Tsubota and M. Ueda, Phys. Rev. A 67, 033610 (2003).
- [22] P. G. Saffman, Vortex dynamics (Cambridge University Press, Cambridge, 1995), p. 119.
- [23] J-K Kim and A.L. Fetter, Phys. Rev. A 70, 043624 (2004).
- [24] S. Sinha and Y. Castin, Phys. Rev. Lett. 87, 190402 (2001).
- [25] E.B. Bogomolny, O. Bohigas, P. Leboeuf, Phys. Rev. Lett. 68, 2726 (1992); E.B. Bogomolny, O. Bohigas, P. Leboeuf, J. Stat. Phys. 85, 639 (1996).
- [26] P.J. Forrester, G. Honner, J. Phys. A **32** 2961 (1999).
- [27] P. Engels, I. Coddington, P.C. Haljan, V. Schweikhard, E.A. Cornell, Phys. Rev. Lett. **90** 170405 (2003).