

Hamiltonian aspects of Bogoliubov quasiparticles

Z. Horváth*

Institute for Theoretical Physics, Eötvös University
H-1117 BUDAPEST (Hungary)

P. A. Horváthy†

Laboratoire de Mathématiques et de Physique Théorique
Université de Tours
F-37 200 TOURS (France)

L. Martina‡

INFN and Dipartimento di Fisica, Università di Lecce
I-73 100 LECCE (Italy)

November 20, 2018

Abstract

The Bogoliubov particle considered in [cond-mat/0507125] admits, contrarily to the claim of the authors, an interesting Hamiltonian structure.

cond-mat/0511099.

In [1] Zhang et al. derive, from the small deviation (Bogoliubov) equation for a superfluid condensate, a semiclassical model, where position and momentum satisfy

$$\dot{\mathbf{r}} = \frac{\partial h}{\partial \mathbf{k}}, \quad \dot{\mathbf{k}} = -\frac{\partial h}{\partial \mathbf{r}}, \quad (1)$$

where h , the energy [ω in their # (12)], is a rather complicated function of \mathbf{r} and \mathbf{k} , that also involves a Berry vector potential $\mathbf{A} = \mathbf{A}(\mathbf{q}, \mathbf{r})$, where $\mathbf{q} = \mathbf{k} - \mathbf{A}$. Eliminating \mathbf{k} in favor of \mathbf{q} and \mathbf{r} transforms (1) into a more complicated form, (3) below, which, Zhang et al. claim, would *no longer be Hamiltonian* and would lead to a *violation of Liouville's theorem* on the conservation of the phase space volume. These statements stem from a misinterpretation of Hamiltonian mechanics [2]. As we show below, the system admits in fact an interesting Hamiltonian structure, and the Liouville theorem is not violated : one simply has to use the correct phase-space volume.

Eqns. (1) are Hamiltonian : they correspond to the Hamiltonian $h = h(\mathbf{k}, \mathbf{r})$ and to the canonical Poisson brackets. But to be Hamiltonian is an intrinsic property of the system that can *not* be lost by a mere change of variables : the error is that the Authors of [1] transform the Hamiltonian but seem to forget about transforming simultaneously the Poisson structure.

The situation is conveniently explained working with the symplectic structure $\Omega = \frac{1}{2}\Omega_{\alpha\beta}d\xi^\alpha \wedge d\xi^\beta$, where ξ^α are coordinates on the phase space, and $\Omega_{\alpha\beta}$ is the inverse of the Poisson matrix

*e-mail: zalanh@ludens.elte.hu

†e-mail: horvathy@lmpt.univ-tours.fr

‡e-mail: Luigi.Martina@le.infn.it

$\Pi^{\alpha\beta} = \{\xi^\alpha, \xi^\beta\}$ [3]. The equations of motion are $\Omega_{\alpha\beta}\dot{\xi}^\alpha = \partial_\beta h$. In canonical coordinates, $\Omega = d\mathbf{k} \wedge d\mathbf{r}$ and we get (1). The change of variables $(\mathbf{k}, \mathbf{r}) \rightarrow (\mathbf{q}, \mathbf{r})$ takes this into

$$\Omega = \frac{1}{2}C_{ij}dq^i \wedge dr^j - \frac{1}{2}F_{ij}dr^i \wedge dr^j, \quad C_{ij} = \delta_{ij} - \frac{\partial A_i}{\partial q^j}, \quad F_{ij} = \frac{\partial A_j}{\partial r^i} - \frac{\partial A_i}{\partial r^j}. \quad (2)$$

C plays the role of an effective mass matrix; the equations of motion become Eq. (13) of [1],

$$C\dot{\mathbf{r}} = \frac{\partial h}{\partial \mathbf{q}} \quad - C\dot{\mathbf{q}} + F\dot{\mathbf{r}} = \frac{\partial h}{\partial \mathbf{r}}. \quad (3)$$

If \mathbf{A} was a function of \mathbf{r} alone, we would have $C = \mathbf{1}$, and for $h(\mathbf{q}, \mathbf{r}) = \mathbf{q}^2/2m + V(\mathbf{r})$ we would recover a particle with unit charge in an electromagnetic field. The \mathbf{q} -dependent case is more interesting, though. Assuming that the system is regular, $\det(\Omega_{\alpha\beta}) = [\det(C_{ij})]^2 \neq 0$, the matrix $C = (C_{ij})$ will be invertible, and the Poisson commutation relations become

$$\{q^i, q^j\} = (C^{-1}FC^{-1})_{ij}, \quad \{q^i, r^j\} = -(C^{-1})_{ij}, \quad \{r^i, r^j\} = 0. \quad (4)$$

The first of these generalizes the usual momentum-momentum relations in a magnetic field; the second modifies the ‘‘Heisenberg’’ commutation relations of momentum and position. The coordinates (Poisson-) commute. The equations of motion (3) are also obtained from the Hamiltonian framework as $\dot{x}^i = \{x^i, H\}$, $\dot{q}^i = \{q^i, H\}$ with $H(\mathbf{q}, \mathbf{r}) = h(\mathbf{k}(\mathbf{q}, \mathbf{r}), \mathbf{r})$.

When $\det \Omega = \det C^2 \equiv \det(\mathbf{1} - \partial\mathbf{A}/\partial\mathbf{q})^2 = 0$, the system becomes singular. This case, although spurious for the model of [1], is nevertheless interesting. The equations of motion (3) can indeed remain consistent when $\partial_{\mathbf{q}}h = 0$ (which fixes \mathbf{q} , which are no more dynamical), provided the motion follows a *generalized Hall law*,

$$\epsilon_{ijk}B^k\dot{r}^j = -E_i \quad (5)$$

where $E_i = -\partial h/\partial r_i$ and $B_i = \frac{1}{2}\epsilon_{ijk}F_{jk}$ are generalized electric and magnetic fields. The trajectories are defined by the vanishing of the ‘‘Lorentz’’ force. In this case, the Hamiltonian structure (4) blows up; Hamiltonian reduction [4] would yield a lower dimensional system.

Turning to the Liouville theorem, let us emphasise that the volume element of the phase space can *only* be defined through the symplectic form [3],

$$dV = \sqrt{\det(\Omega_{\alpha\beta})} \prod_{\alpha} d\xi^\alpha. \quad (6)$$

The pre-factor $\det(\mathbf{1} - \partial\mathbf{A}/\partial\mathbf{q})$ ‘‘discovered’’ by Zhang et al. is precisely the square-root of the determinant of the symplectic matrix. It is also the square-root of the Jacobian, J , of the transformation from canonical to arbitrary variables, and is hence always present whenever the change of variables is non-canonical. Now the Liouville theorem [3] says that the symplectic volume element is invariant w.r.t. the Hamiltonian flow.

In the singular case the Liouville volume form (6) becomes degenerate.

Analogous problems were studied before in [5, 2].

References

- [1] C-W. Zhang, A. M. Dudarev, Q. Niu, [cond-mat/0507125].

- [2] C. Duval, Z. Horváth, P. A. Horváthy, L. Martina, and P. Stichel, [cond-mat/0506051]; M. Stone, Urbana Lectures (2005), available from <http://w3.physics.uiuc.edu/~m-stone5/mmb/mmb.html>; K. Yu. Bliokh, [cond-mat/0507499]; S. Ghosh, P. Pal, [hep-th/0510038].
- [3] See e. g., R. Abraham and J. Marsden, *Foundations of Mechanics*. 2nd ed. Addison-Wesley, Reading, Mass (1978).
- [4] L. Faddeev and R. Jackiw, *Phys. Rev. Lett.* **60**, 1692 (1988).
- [5] C. Duval and P. A. Horváthy, *Phys. Lett.* **B 479**, 284 (2000) [hep-th/0002233]; *Journ. Phys.* **A 34**, 10097 (2001) [hep-th/0106089]; P. A. Horváthy, *Ann. Phys.* **299**, 128 (2002) [hep-th/0201007].