

Trapped ion chain as a neural network

M. Pons,¹ V. Ahufinger,² C. Wunderlich,³ A. Sanpera,⁴ and M. Lewenstein⁵

¹*Departamento de Física Aplicada I, Universidad del País Vasco, E-20600 Eibar, Spain*

²*ICREA and Grup d'Òptica, Departament de Física, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain.*

³*Fachbereich Physik, Universität Siegen, 57068 Siegen, Germany.*

⁴*ICREA and Grup de Física Teòrica, Departament de Física, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain.*

⁵*ICREA and Institut de Ciències Fotòniques, E-08034 Barcelona, Spain.*

We demonstrate the possibility of realizing a neural network in a chain of trapped ions with induced long range interactions. Such models permit to store information distributed over the whole system. The storage capacity of such network, which depends on the phonon spectrum of the system, can be controlled by changing the external trapping potential and/or by applying longitudinal local magnetic fields. The system properties suggest the possibility of implementing *robust distributed* realizations of quantum logic.

The Cirac-Zoller trapped ion computer [1] has become one of the paradigmatic models to implement a quantum computer [2]. Recently, spectacular experimental progress in both realization of simple algorithms and implementation of quantum logic has been achieved using a few ions qubits (c.f. [3]).

Still, serious experimental challenges remain on the way to a large-scale ion trap quantum computer. One should stress that while the achievement of an all-purpose quantum computer in the next years seems to be difficult, one can be quite optimistic about the applications of chains of trapped ions as quantum simulators. Recently, it has been shown that long range pairwise interaction between individual spins is induced in an ion trap when introducing an additional state-dependent force acting on the ions [4, 5]. The properties of such ion spin molecules can be designed by variation of the parameters characterizing the ion trap. Later, a state-dependent optical force has also been shown to evoke long range couplings and has been proposed to simulate general Heisenberg interactions and to investigate quantum phase transitions in ion traps [6]. Here we show that ion spin systems can serve to realize a neural network (NN) model. Neural networks are a prototype model of parallel distributed memory [7, 8], and have been intensively studied by physicists since the famous paper by Hopfield [9]. These disordered systems with long range interactions typically present a large number of metastable (free) energy minima, like in spin glasses [8]. These states can be used to store information distributed over the whole system. The patterns stored by the network have large basins of attraction in the thermodynamical sense which means that even fuzzy patterns are recognized as perfect ones. For this reason, attractive NN's can be used as associative memory. At the same time, NN are robust, so that destroying even a large part of the network does not necessarily diminish the network performance. The above listed properties make NN's interesting for *distributed* quantum information storing and processing where a quantum bit would correspond to two collective modes of the ion's chain instead of two internal states of each individual ion. Differ-

ent approaches to exploit the potential use of NN models for quantum information processing have been already discussed in the literature [10, 11]. In [11] we have shown that it is possible to create next neighbour entanglement in an ideal NN model. Here we propose, for the first time, a feasible implementation of NN using a chain of trapped ions.

Before explaining the details of NN implementation we first remind the readers the main features of the Hopfield model [9], and discuss the effective Hamiltonian derived in Ref.[5, 6]. The similarities between both models suggest the possibility of using a chain of trapped ions as a NN. To demonstrate this assumption we find the storage capacity of the system and its robustness which are the most appealing features of NN for distributed quantum information. Thus, the question of ergodicity and, therefore, the ability of the system to perform an associative memory is not considered here. We show that the storage capacity, which is determined by the phonon spectrum of the system, can be controlled by modifying the shape of the external axial trapping potential and/or by applying longitudinal magnetic fields. Although the storage capacity is a classical property of the network, spin ion systems permit also to study quantum neural networks by using a transverse magnetic field or an optical field that effectively simulates an additional transverse magnetic field. Transverse magnetic fields should permit tunneling processes between stored patterns and to realize for example of quantum stimulated annealing [8, 12]. This study is beyond the scope of this letter and will be treated elsewhere.

Following the models of Hopfield [9] and Little [13], a neuron can be viewed as an Ising spin with two possible states: an "up" position ($S = +1$) and a "down" position ($S = -1$) depending on whether the neuron has or has not fired an electromagnetic signal, in a given interval of time [14]. The state of the network of N neurons at a certain time is defined by the instantaneous configuration of all the spin variables $\{S_i\}$ at this time. The dynamic evolution of these states is determined by the interactions among neurons, J_{ij} , which are symmetric, so, for any pair

of neurons, $J_{ij} = J_{ji}$. Also full connectivity is assumed, that is, every neuron can receive an input from any other one, and can send an output to it. The Hamiltonian of the system can be written as:

$$H = -\frac{1}{2} \sum_{i,j} J_{ij} S_i S_j + h \sum_i S_i, \quad (1)$$

where h corresponds to an external magnetic field. The interactions are determined by the patterns or configurations of spins to be stored in the network. These patterns will be learned if the system is able to accommodate them as attractors, implying that a large set of initial configurations of the network will be driven dynamically (after sufficient enough time) to those patterns. A possible choice of the interactions is:

$$J_{ij} = \frac{1}{N} \sum_{\mu=1}^p \xi_i^\mu \xi_j^\mu, \quad (2)$$

with $i \neq j$. The p sets of $\{\xi_i^\mu\} = \pm 1$ are the patterns that we wish to store. The network will have the capacity of storage and retrieval of information if the dynamical stable configurations (local minima) reached by the system $\{S_i\}$ are correlated with the learned ones $\{\xi_i^\mu\}$. Despite the fact that the interactions have been constructed to guarantee that certain specified patterns be fixed points of the dynamics, the non-linearity of the dynamical process may induce additional attractors, the so called spurious states.

Recently it has been shown that the Hamiltonian describing a linear chain of harmonically trapped ions exposed to a magnetic field gradient [5] or interacting with convenient laser fields [6] can be transformed into an effective spin-spin Hamiltonian with long range interactions mediated by the collective motion of the ions:

$$H = -\frac{1}{2} \sum_{\alpha,i,j} J_{ij}^\alpha \sigma_i^\alpha \sigma_j^\alpha + \sum_{\alpha,i} B_i^\alpha \sigma_i^\alpha, \quad (3)$$

where:

$$J_{ij}^\alpha = \frac{(F_{ij}^\alpha)^2}{m} \sum_n \frac{M_{i,n}^\alpha M_{j,n}^\alpha}{\omega_{\alpha,n}^2}, \quad (4)$$

with $\alpha = x, y$ and z being the spatial directions, i, j labeling the ions, σ being the Pauli matrices, F_{ij}^α the forces in the α direction, m the mass of the ion, $\omega_{\alpha,n}$ the angular frequency of vibrational mode n . $M_{i,n}^\alpha$ are the unitary matrices that diagonalize the vibrational Hamiltonian:

$$M_{i,n}^\alpha \kappa_{i,j}^\alpha M_{j,m}^\alpha = \omega_{\alpha,n}^2 \delta_{nm}, \quad (5)$$

where $\kappa_{i,j}^\alpha$ are the elastic constants of the chain [15]. The coefficient $M_{i,n}^\alpha$ gives the scaled amplitude of the local oscillations of ion i around its equilibrium position when the collective vibrational mode n is excited. Thus, the

eigenvectors of M describe each ion's contribution to a given vibrational mode while the eigenvalues provide the oscillation frequencies, $\omega_{\alpha,n}$ of the collective modes.

The external trapping frequencies are chosen such that the laser cooled ions crystallize in a linear chain (*i.e.*, $\Omega_x = \Omega_y \gg \Omega_z$) and the external forces are such that on the z -axis $F_{ij}^x = F_{ij}^y = 0$ holds, and the index α is dropped. From now on we will consider the case of a chain of ions interacting with external laser fields [6] which allow us to consider site independent forces and magnetic field, $F_{ij} = F$, $B'_i = B'$ [16]. Thus, Eqs.(1) and (3) have the same form and the possibility to implement a classical neural network using a linear chain of ions arises. We substitute the Pauli matrices in Eq.(3) by Ising spins $S = \pm 1$, where the effective spin corresponds to the internal state of the ion. Nevertheless, there are some important differences between both models. First, in the Hopfield model, the interactions given by Eq.(2) are determined by the patterns to be stored $\{\xi_i^\mu\} = \pm 1$ while in the trapped ion chain, the interactions are fixed by the coefficients $M_{i,n}$ that do not necessarily equal ± 1 . Second, in Eq.(2), p corresponds to the number of patterns to be stored, which in the limit of large number of spins N is bounded from above by $p = 0.14N$ [7]. In our model, the sum (Eq.(4)) extends to the total number of vibrational modes which is larger than the total number of stored patterns. And finally, in the Hopfield model all the patterns have the same weight while in the ion chain each vibrational mode (Eq.(4)) is weighted by $1/\omega_n^2$. Due to this fact, to reproduce as close as possible a NN model, the vibrational modes should be ideally almost degenerate in energy. Moreover, the corresponding patterns must have large basins of attraction, *i.e.* they should correspond to sufficiently different configurations of spins so that each configuration is dynamically recovered even if several spins are randomly flipped. These requirements are not achievable with an harmonic trapping potential due to the fixed frequency ratio between the two lowest vibrational modes [17]. The desired degeneracy can be achieved either by engineering an adequate external magnetic field or by using a trapping potential of the form of $|x|^\gamma$ with $\gamma \leq 0.8$ (see Fig.1).

To check the feasibility of implementing a NN model in these ion spin systems, we calculate first the phonon spectrum, impose the learning rule (Eq.(4)), and probe the correlation between the dynamically stable configurations and the stored ones. Therefore, the first step consists in finding the equilibrium positions of the ions and calculate the vibrational modes and frequencies by using a standard diagonalization procedure. Once the complete vibrational spectrum of the system is known, we calculate the interactions among spins J_{ij} , and evaluate the energy of the system. The dynamical thermalization of the system is simulated using standard Monte Carlo techniques.

We fix the initial spin configuration by mimicking the signs of a given vibrational mode. Starting with this spin configuration, we flip randomly r spins and let the




FIG. 1: Ratio between the frequencies of the second and the first vibrational modes as a function of the exponent of the trapping potential, γ

system evolve towards the equilibrium situation assuming a noiseless scenario. If the system recovers the initial configuration, the configuration is stable under the flip of r spins. The number of initial spin flips determines the initial overlap, defined as $m_i = (N - r)/N$. After dynamical evolution, the final overlap is given by $m_f = (N - s)/N$ where s is the number of spins that differ from the initial configuration. We repeat this process over M initial different random r spin flips to evaluate statistically the final overlap with the initial configuration: $m_f = (\sum_{j=1}^N m_j n_j)/M$, n_j being the number of times that the system reaches the m_j configuration. The value of the initial overlap for which significant decrease of the final overlap occurs, is a good measure of the size of the basin of attraction of the corresponding pattern. For the harmonic trapping potential, the two lowest vibrational modes are the center of mass (all spins parallel, with frequency $\omega_1 = \Omega_z$) and the breathing mode (half up, half down, with $\omega_2 = \sqrt{3}\Omega_z$) [17]. We find that the center of mass mode is stable up to the flip of half of the spins, while the breathing mode is already unstable under the flip of a single spin. Therefore, the only configuration that can be recovered is the center of mass mode, corresponding to all spins aligned up or down. To increase the storage capacity of the network one can engineer the appropriate spatial dependence of a longitudinal magnetic field to stabilize one additional given pattern. We apply this technique to make the second pattern stable. The results are displayed in Fig.2 showing the basin of attraction of the two first modes for a chain of (a) 20 Ca^+ ions with $B = 7.7 \times 10^{-31} \text{Tm}^2\text{C/s}$ parallel to each spin of the second mode and (b) 40 ions for $B = 6.5 \times 10^{-31} \text{Tm}^2\text{C/s}$ parallel to each spin of the second mode. The results are averaged over 500 configurations and with random spin flip and random update. By increasing (diminishing) the external magnetic field the stability of the first pattern decreases (increases), while the second pattern exhibits the complementary behavior. In Fig.2(a) we show that




FIG. 2: Basin of attraction of the two lowest energy modes of an harmonically trapped ($\Omega_z = 1 \text{MHz}$) chain of ions in the presence of an external longitudinal magnetic field. (a) 20 Ca^+ ions with $B = 7.7 \times 10^{-31} \text{Tm}^2\text{C/s}$ parallel to each spin of the second mode and (b) 40 Ca^+ ions with $B = 6.5 \times 10^{-31} \text{Tm}^2\text{C/s}$ parallel to each spin of the second mode. Black squares (triangles) correspond to the lowest (second) mode.

with 20 ions, it is possible to stabilize completely the first and second pattern simultaneously up to the spin flip of 3 ions while the probability of recovering the initial pattern for four initial random spin flips is 99.2% for both patterns. Increasing the number of ions to 40, Fig.2(b), the basin of attraction increases up to the random flip of 9 ions with 100% of probability of recovering of the initial patterns, 99.7% (99.5%) of probability of recovering the first (second) pattern for the initial random flip of 10 spins and 98.5% (99%) of probability of recovering the first (second) pattern for the initial random flip of 11 spins.

The second approach used to increase the storage capacity of the network consists in modifying the spatial dependence of the trapping potential. We consider an external potential of the form $V(x) = A |x|^\gamma$ with an even number of ions in order to avoid any possible singularity of the potential at the origin. Fig.1 shows the ratio between the frequencies of the second and the first mode ω_2/ω_1 as a function of γ , for the case of 20 ions. For $N \geq 20$ this ratio does not depend on the number of ions neither on the value of A . For $\gamma = 1$, and assuming small oscillations around the equilibrium positions, the ratio ω_2/ω_1 is not well defined since the second mode does not exist because of the constant character of the force produced by the external potential. For $\gamma > 2$ severe limitations in the storage capacity of the network, similar to those encountered in the harmonic case, appear. The situation changes when $\gamma < 1$. In this case the two lowest modes become nearly degenerated. We obtain larger number of stored patterns for $\gamma = 0.5$ but




FIG. 3: Final overlap averaged over 500 initial configurations as a function of the initial overlap, for the two lowest vibrational modes of (a) 20 Ca^+ ions and (b) 40 Ca^+ ions in a potential $V = A|x|^{0.5}$ with $A = 6.6 \times 10^{-20} \text{J/m}^{1/2}$. The black squares (triangles) correspond to the first (second) pattern.

the desired degeneracy is only obtained for the first two collective modes. The results obtained with this optimal trapping potential are displayed in Fig.3 where the final overlap as a function of the initial overlap for the two lowest vibrational modes is depicted for (a) 20 Calcium ions and (b) 40 ions. The final overlap of the two lowest patterns and the corresponding ones with all the spins flipped is close to 1 up to two spin flips for $N = 20$. This shows that the system is able to recover the four patterns up to two spin flips. For $N = 40$ this holds up to

eight spin flips. Note that the system does not always recover the initial pattern but instead sometimes reaches a slightly deformed configuration, which differs only in one spin flip from the original one. For this reason m_f in Fig. 3(a) is not exactly 1 for one and two spin flips and up to eight in Fig. 3(b). In Fig. 3(a) the probability of recovering the initial pattern or one that differs in one spin flip with the initial one is 100% (96.9%) for the second (first) mode for the initial random spin flip of one spin and 98.7% (95.1%) for the second (first) mode for the initial random spin flip of two spins. In Fig. 3(b) the probability of recovering of the two modes is above 98% up to three initial random spin flips and above 97% up to eight initial spin flips.

Summarizing, we have shown that spin-ion systems can be used to implement NN models. We have calculated the storage capacity and robustness of such systems and showed how to control this storage capacity by changing the trapping potential and/or applying external longitudinal magnetic fields. Furthermore, such models should allow to implement quantum neural networks by applying transverse magnetic fields that permit tunneling processes between stored patterns. Finally, NN provide a clear example of robust distributed information which could be exploited as a quantum memory.

We thank I. Bloch, A. Bramon, H.-P. Büchler, G. Morigi, U. Sen, A. Sen (De), J. Wehr and P. Zoller for fruitful discussions. We acknowledge support from the Deutsche Forschungsgemeinschaft (SFB 407, SPP1078 and SPP1116), the RTN Cold Quantum Gases, ESF PESC BEC2000+, the Spanish Ministerio de Ciencia y Tecnología (MAT2002- 00699, FIS2005-04627), and the Alexander von Humboldt Foundation. V.A. acknowledges financial support from the Spanish Ministerio de Ciencia y Tecnología.

[1] J.I. Cirac and P. Zoller, Phys. Rev. Lett. **74**, 4091 (1995)
[2] D. Bouwmeester, A. Ekert, and A. Zeilinger (Eds.), *The Physics of quantum information* (Springer, Berlin, 2000).
[3] F. Schmidt-Kaler *et al.* Nature **422**, 408 (2003); J. Chiaverini *et al.* Science **308**, 997 (2005); M. Riebe *et al.* Nature **429**, 734 (2004); M. D. Barrett *et al.*, Nature **429**, 737 (2004).
[4] F. Mintert and C. Wunderlich, Phys. Rev. Lett. **87**, 257904 (2001); C. Wunderlich and C. Balzer, Adv. At. Mol. Opt. Phys. **49**, 293 (2003); D. Mc Hugh and J. Twamley, Phys. Rev. A **71**, 012315 (2005).
[5] C. Wunderlich, in *Laser Physics at the Limit* (Springer Verlag, Heidelberg, 2002), pp. 261-; also available as quant-ph/0111158.
[6] D. Porras and J.I. Cirac, Phys. Rev. Lett. **92**, 207901 (2004).
[7] D.J. Amit, *Modeling Brain Function*, (Cambridge University Press, 1989).
[8] M. Mézard, G. Parisi, and M.A. Virasoro, *Spin Glass and*

Beyond (World Scientific, Singapour, 1987).
[9] J.J. Hopfield, Proc. Natl. Acad. Sci. USA **81**,3088 (1984).
[10] P. Gralewicz quant-ph/0401127 and references therein.
[11] A. Sen (De) *et al.*, quant-ph/0507172.
[12] D.J. Amit, H. Gutfreund and H. Sompolinsky, Phys. Rev. A **32**, 1007 (1985).
[13] W.A. Little, Math. Biosci. **19**, 101 (1974).
[14] For an excellent introduction to NN see [7].
[15] H. Goldstein, *Classical Mechanics*, (Addison-Wesley, Reading, MA, 1980).
[16] In [6], the effective magnetic field B' in Eq. (3) includes a contribution from the pushing forces caused by the applied lasers: $B' = B + \frac{F^2}{m\Omega_z^2}$ This extra term due to the force is compensated by a constant external magnetic field. Notice also that F is such that it satisfies the validity of the derivation of the effective Hamiltonian: $\eta^2 = \frac{F^2}{2m\hbar\Omega_z^2}$ with $\eta = 0.1$.
[17] D.F.V. James, Appl. Phys. B **66**, 181 (1998).