

Towards Classification of Phase Transitions in Reaction–Diffusion Models

Vlad Elgart and Alex Kamenev

Department of Physics, University of Minnesota, Minneapolis, MN 55455, USA

(Dated: February 8, 2020)

Equilibrium phase transitions are associated with rearrangements of minima of a (Lagrangian) potential. Treatment of non–equilibrium systems requires doubling of degrees of freedom, which may be often interpreted as a transition from the “coordinate” to the “phase” space representation. As a result, one has to deal with the Hamiltonian formulation of the field theory instead of the Lagrangian one. We suggest a classification scheme of phase transitions in reaction–diffusion models based on the topology of the phase portraits of corresponding Hamiltonians. In models with an absorbing state such a topology is fully determined by intersecting curves of zero “energy”. We identify four families of topologically distinct classes of phase portraits stable upon RG transformations.

PACS numbers: 64.60.Ak, 05.40.-a, 64.60.Cn, 82.20.-w

I. INTRODUCTION

The last decade witnessed a rapid grow of interest in reaction–diffusion models^{1,2,3,4,5,6,7,8,9,10,11,12,13,14,15}. Such models are employed for description of phenomena ranging from kinetics of chemical reactions to evolution of biological populations. The subject of particular interest is description of dynamical phase transitions in reaction–diffusion kinetics. An important example is the absorbing phase transitions. Upon such a transition the system goes from an active (“living”) phase to an absorbing (“dead”) state with no escape from it.

Grassberger^{16,17} and Janssen¹⁸ realized that many of the absorbing state transitions belong to the same universality class as the directed percolation (DP) model. Since then the DP universality class was extensively studied both analytically and numerically (see Refs. [1,3,19,20,21,22,23,24]). The DP universality class is extremely robust. In fact, exceptions to the DP transitions are rare. However, if the microscopic dynamics possesses additional symmetries the universality class of the transition may be different. For example, the parity conservation (PC) is known to be driving the transition to a new distinct universality class^{1,3,5,25,26}. Recently another possible universality classes had been studied^{1,27,28,29,30,31,32,33,34,35,36,37,38,39,40}. Among them is pair contact process with diffusion (PCPD), which critical behavior was not yet described analytically^{8,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56}.

Despite of extensive accumulated knowledge, it seems that a guiding principle, allowing to distinguish between various types of the transitions, is still missing. The purpose of this paper is to suggest a simple scheme, providing with, at least, an educated guess regarding the universality class of a reaction–diffusion model at hand. The scheme is based on the topology of the phase portraits of the system’s Hamiltonian. Before elaborating on it, let us briefly remind the corresponding strategy for equilibrium systems.

An equilibrium system may be characterized by an ac-

tion (energy)

$$S = \int d^d x [D(\nabla q)^2 + V(q)] , \quad (1)$$

written in terms of the order parameter $q(x)$ (for simplicity we restrict ourselves to the one–component order parameter). The potential function $V(q)$ encompasses information about possible phase transitions. Specifically, one monitors behavior of the *minima* of $V(q)$ as a function of a control parameter to infer the existence and the type of the transition. For example, a wide class of models may be described with the potential of the form:

$$V(q) = h q + m q^2 + u q^4 . \quad (2)$$

For $m < 0$ the system exhibits the first order transition when h changes its sign (for $d > 1$) and the two minima interchange. In the symmetric case, $h = 0$, the system may undergo the second order transition when the parameter m is swept through zero, so a single minimum is split in two. Below the critical dimension $d_c = 4$ this second order transition is characterized by non–mean–field critical exponents. To find the exponents one typically employs the renormalization group (RG) technique. The RG treats the coefficients m and u of the potential (2) as functions of the running spatial scale. For $\epsilon = d_c - d > 0$ the potential scales towards the non–trivial fix–point potential, $V^*(q) \neq 0$, with $m^* \sim \epsilon^2$ and $u^* \sim \epsilon$. Notice that action (1) essentially specifies the Lagrangian field theory.

One may ask whether a similar strategy exists for non–equilibrium phase transitions in reaction–diffusion systems. To answer this question one has to recall that a description of non–equilibrium systems requires doubling of the degrees of freedom. There are various manifestations of this statement depending on the specific context. In quantum kinetics it is known as the Keldysh technique⁵⁷. It employs a time evolution along the closed contour, so one has to keep two copies of each field: one for the forward and another for the backward evolution. In classical context the Martin–Siggia–Rose^{58,59} method requires an additional set of fields to resolve the func-

tional delta-functions of Langevin equations. Most importantly for the present subject, in reaction-diffusion kinetics the Doi-Peliti^{60,61} operator technique deals with the creation and annihilation operators for each reagent. Thus it employs two variables (or one complex field) for every *real* physical degree of freedom. (For a discussion of connections between these techniques see e.g. Refs. [62,63].)

An important observation is that in all these examples the two sets of fields (being properly transformed) may be considered as canonically conjugated variables. As a result, instead of the equilibrium order parameter $q(x)$, one has to deal with the canonical pair: $q(x, t)$ and $p(x, t)$. Correspondingly a reaction-diffusion system may be described by the *Hamiltonian* action

$$S = \int dt \int d^d x [p \partial_t q + D \nabla p \nabla q - H_R(p, q)], \quad (3)$$

where the Hamiltonian $H_R(p, q)$ is determined by the set of reactions specific for a given model (see below).

Comparing Eqs. (3) and (1), one notices that the reaction Hamiltonian $H_R(p, q)$ plays the role similar to the effective potential $V(q)$ in the equilibrium statistical mechanics. Thus it is plausible that $H_R(p, q)$ may encode the information about possible non-equilibrium transitions in a way analogous to what $V(q)$ does. Specifically, one wants to know what is the Hamiltonian analog of the potential minima, given by $\partial_q V = 0$, in the Lagrangian formulation. The answer is that it is the *classical* equations of motion: $\partial_t q = \partial_p H_R$, $\partial_t p = -\partial_q H_R$. One is looking, therefore, for a geometric way to picture the Hamiltonian equations of motion. We argue below that the way to do it (at least for the one-component models) is to consider the phase space trajectories in the (p, q) plane. Indeed, the classical equations of motion conserve the “energy”. Thus the phase space trajectories are given by the curves $H_R(p, q) = \text{const}$. Moreover, for systems with absorbing states the only trajectories which may intersect correspond to zero energy. As a result, the set of curves:

$$H_R(p, q) = 0 \quad (4)$$

determines entirely the topology of the phase space.

The main message of this paper is that the curves specified by Eq. (4) and the corresponding topology of the phase portrait classify possible phase transitions in reaction-diffusion models. It is the web of the zero “energy” trajectories which plays the role of minima of $V(q)$ potential in the equilibrium statistical mechanics. A topological rearrangement of this web as a function of a control parameter signals the existence of a phase transition. The corresponding topology is in one-to-one correspondence with the universality classes.

Below we show that the number of distinct generic phase-space topologies (for one-component systems) is rather limited, indicating that all possible universality classes may be exhausted. Some of the topologies correspond to the first order transitions, while others to the

continuous ones (much the way potential (2) contained both). The latter class may develop non-trivial critical exponents below a certain critical dimension d_c . The way to find these exponents is to follow the RG flow of constants of the Hamiltonian $H_R(p, q)$ upon elimination of the small scale fluctuations. While the Hamiltonian itself may be complicated, it is only the topology of the phase space that matters, not a specific shape of the curves. Any given topology may be modeled by a simple polynomial of q and p , again much the way the simple polynomial (2) suffices to describe many equilibrium systems. Thus one must follow only changes of the topology of the phase portrait upon RG transformations. One should also verify that a given topology is stable upon RG transformations, i.e. it can’t be reduced to a more generic one by decimation. The resulting fix-point topologies and corresponding fix-point Hamiltonians, $H_R^*(q, p)$, provide the information about the universality classes.

We found that the DP universality class (represented by the simplest triangular structure on the phase plane) serves as a parent for a family of descending classes. Each subsequent class in the family is characterized by a minimal number k of particles needed to initiate reactions. We denote it as k -particle contact process with diffusion (kCPD). Here 1CPD is DP, while 2CPD is the pair contact process with diffusion (PCPD) (for a review see Ref. [8] and references therein). Above the upper critical dimension $d_c = 4/k$ the kCPD’s are characterized by the mean-field critical exponents, e.g. $\beta = 1$, $\nu_{\perp} = k/2$. For $d \leq d_c$ and $k = 2, 3$ we found that RG flows to a strong-coupling fix-point that can’t be accessed in the ϵ expansion (see also Refs [8,64]). We also discuss a possible nature of the strong-coupling fix-point for $k = 2$.

Similarly the parity conserving (PC) universality class generates a family of classes, characterized by a minimal number, k , of incoming particles required for *all* reactions. We call them k -particle parity conserving (kPC). Their upper critical dimension is $d_c = 2/k$. In addition to kCPD and kPC we identify two more families of universality classes. They both originate from reversible reactions which may go both directions with different rates. We call them k -particle reversible (kR) and k -particle reversible parity conserving (kRPC). In both cases k stays for minimal number of incoming particles. Their critical dimensions are $2/k$ and $2/(k+1)$ correspondingly.

These four families seem to exhaust all possible continuous transitions reachable by tuning of a *single* control parameter. That means that any phase portrait, topologically different from that of the four families, is unstable upon renormalization. In the large scale limit it flows towards one of the stable topologies. The latter are protected by certain symmetries of the action against deformations introduced by the RG.

The paper is organized as follows: in section II we introduce reaction Hamiltonians and their phase portraits. Section III is devoted to models, exhibiting the first order transitions and discuss topological structure of their phase portraits. In the section IV models of DP univer-

sality class and its derivatives, kCPD's, are considered. We demonstrate that triangular topology of the phase portrait is the typical feature of all universality classes of this kind. In section V we consider parity conserving model and its generalizations, kPC's. A rectangular phase portraits topology of the reversible reaction models, kR's and kRPC's, are discussed in section VI. Finally some conclusions and outlook are drawn in section VII.

II. REACTION HAMILTONIANS AND PHASE PORTRAITS

The standard way to introduce the “quantum” reaction Hamiltonian is by employing the creation, annihilation operator technique of Doi and Peliti^{10,11,60,61,65}. Here we choose to follow a different, though completely equivalent, strategy⁶⁶. Consider a generic reaction that transforms k particles into m equivalent ones with the probability λ :



The corresponding Master equation for the temporal evolution of the probability $\mathcal{P}_n(t)$ of a configuration with n particles has a form:

$$\partial_t \mathcal{P}_n(t) = \lambda \left[\binom{n+k-m}{k} \mathcal{P}_{n+k-m}(t) - \binom{n}{k} \mathcal{P}_n(t) \right]. \quad (6)$$

The two terms on the r.h.s. represent the probabilities of “in” and “out” processes correspondingly. The Master equation (6) is to be supplemented with an initial distribution, e.g. $\mathcal{P}_n(0) = e^{-n_0} n_0^n / n!$ – the Poisson distribution with the mean value n_0 , or $\mathcal{P}_n(0) = \delta_{n,n_0}$ – the fixed initial particle number.

Let us define now a *generating function* as:

$$G(p, t) \equiv \sum_{n=0}^{\infty} p^n \mathcal{P}_n(t). \quad (7)$$

Parameter p will play a role of the canonical momentum; so far it is introduced pure formally. The value $p = 1$ plays a special role. First, the conservation of probability implies the normalization condition:

$$G(1, t) \equiv 1. \quad (8)$$

Second, the moments of $\mathcal{P}_n(t)$, may be expressed through derivatives of the generating function at $p = 1$, e.g. $\langle n(t) \rangle \equiv \sum_n n \mathcal{P}_n(t) = \partial_p G(p, t)|_{p=1}$. Knowing the generating function, one may find a probability of having (integer) n particles at time t as $\mathcal{P}_n(t) = \partial_p^n G(p, t)|_{p=0} / n!$.

In terms of the generating function the Master equation (6) may be *identically* rewritten as

$$\partial_t G = \hat{H}_R(p, \hat{q}) G, \quad (9)$$

where the non-Hermitian normally-ordered operator, \hat{H}_R , stays for

$$\hat{H}_R(p, \hat{q}) = \frac{\lambda}{k!} (p^m - p^k) \hat{q}^k. \quad (10)$$

Here we have introduced the “coordinate” operator,

$$\hat{q} \equiv \frac{\partial}{\partial p}, \quad (11)$$

obeying the canonical commutation relation: $[\hat{q}, p] = 1$. Because of the obvious analogy with the Schrödinger equation, we shall refer to the operator \hat{H}_R as the Hamilton operator in the p -representation. From the normalization condition, Eq. (8), follows that

$$\hat{H}_R(p, \hat{q}) \Big|_{p=1} = 0. \quad (12)$$

Any Hamiltonian derived from a probability conserving Master equation necessarily satisfies this property.

One can easily generalize this construction for the case where many reactions with the rates λ_{km} take place at the same time. To this end one simply algebraically add the corresponding partial Hamiltonians to obtain the full reaction Hamiltonian. If there is no particle production from the vacuum, i.e. $k \neq 0$ for any m , the empty state with $n = 0$ is an absorbing state in a sense that the system can never leave it. According to Eq. (10) any Hamiltonian function describing a system with the empty absorbing state must satisfy

$$H_R(p, q)|_{q=0} = 0 \quad (13)$$

in addition to Eq. (12).

Before considering the full “quantum” problem, Eq. (9), let us analyze the corresponding classical dynamics. The classical equations of the motion are:

$$\partial_t q = \frac{\partial}{\partial p} H_R(p, q), \quad (14)$$

$$\partial_t p = -\frac{\partial}{\partial q} H_R(p, q). \quad (15)$$

Due to Eqs. (10) and (12), $p = 1$ is always one of the solutions of Eq. (15). Substituting $p = 1$ into Eq. (14), one obtains (for the Hamiltonian (10)):

$$\partial_t q = \frac{\lambda}{k!} (m - k) q^k. \quad (16)$$

This is nothing but the mean-field rate equation for the average particle number $\langle n(t) \rangle$, neglecting all fluctuation effects. Therefore one may identify the variable q as the reaction “coordinate” (in fact this notation is not precise, since it is true only along the line $p = 1$).

In order to proceed with the classical problem, Eqs. (14), (15), beyond the reaction rate approximation ($p = 1$), it is convenient to consider the phase space of the system. The classical equations of motion (14), (15)




FIG. 1: Phase portrait of the binary annihilation system, $2A \xrightarrow{\lambda} \emptyset$. The corresponding classical Hamiltonian is given by $H_R(p, q) = \frac{\lambda}{2}(1 - p^2)q^2$. Solid black lines show zero-energy trajectories: generic lines $p = 1$ and double degenerate $q = 0$ and the “accidental” line $p = -1$. Dashed colored curves indicate trajectories with non-zero energy. The arrows show the evolution direction.

conserve the energy. As a result, the phase space (p, q) evolution of the system takes place along the trajectories given by $H_R(p, q) = \text{const}$, where the constant energy is determined from the initial conditions. Among all possible trajectories the ones with $H_R = 0$ play a special role. For one thing, the evolution prescribed by the rate equation ($p = 1$), takes place along one of such lines (cf. Eq. (12)). More importantly, the trajectories with $H_R = 0$ may intersect each other. Indeed, the two zero-energy lines, guaranteed by Eqs. (12) and (13), i.e. $p = 1$ and $q = 0$, intersect in the point $(1, 0)$. Therefore the set of intersecting zero-energy curves plays the role of separatrix, i.e. it divides the entire phase space on the isolated sectors. All other trajectories cannot intersect the zero-energy ones and are confined to one of the sectors. The web of the zero-energy curves uniquely determines the topology of the phase portrait. An example of such a construction is given in Fig. 1.

Going back to the full “quantum” problem, one may write a formal solution of Eq. (9) as

$$G(p_f, t_f) = \int U(p_f, t_f; p_0, t_0) G_0(p_0) dp_0, \quad (17)$$

where the Green function $U(p_f, t_f; p_0, t_0)$ is given by the T-exponent: $T \exp\{\hat{H}_R t\}$. Dividing the time interval $[t_0, t_f]$ on $N \rightarrow \infty$ steps, and introducing the resolution of unity at each one of them, one obtains the Feynman representation

$$U(p_f, t_f; p_0, t_0) = \int \mathcal{D}p(t) \mathcal{D}q(t) e^{-S[p, q]}, \quad (18)$$

with the Hamiltonian action

$$S[p, q] = \int_{t_0}^{t_f} dt [p \partial_t q - H_R(p, q)]. \quad (19)$$

To combine reaction kinetics with the random walk on a lattice, one needs to modify the Master equation. The corresponding generating function becomes a function of

many variables p_i , where index i enumerates the lattice sites. One may also introduce the conjugated variables $\hat{q}_i = \partial/\partial p_i$. The resulting Hamiltonian takes the form:

$$\hat{H} = -\tilde{D} \sum_{\langle i, j \rangle} (p_i - p_j)(\hat{q}_i - \hat{q}_j) + \sum_i \hat{H}_R(p_i, \hat{q}_i), \quad (20)$$

where \tilde{D} is a hopping probability per unit time and the sum in the first term on the r.h.s. runs over nearest neighbors i, j . Taking the continuum limit^{60,61} and introducing the pair of canonically conjugated fields $p(x, t)$ and $q(x, t)$, one arrives at the quantum field theory with the Hamiltonian action (3). The diffusion constant in Eq. (3) is given by $D = \tilde{D}a^2$, where a is the lattice constant.

Unless the system is very close to the extinction, the functional integral in Eq. (18) may be evaluated in the saddle point approximation. In such a case the Green function is given by the exponentiated action of a classical trajectory, satisfying the proper boundary conditions⁶⁶. (Much the same way as the minima of the potential $V(q)$, Eq. (2), dominate the partition function away from an equilibrium phase transition.) A possible phase transition may be associated with a qualitative change in the behavior of the phase-space trajectories (c.f. with the rearrangement of minima of the potential upon an equilibrium transition). In other words, phase transitions lead to a change of topology of the phase space portrait. Since the latter is determined by the set of the zero-energy lines, it is the rearrangement of this set, upon variation of a control parameter, which must be associated with the phase transition.

If a system is close enough to a phase transition (or extinction) the saddle point approximation may lose its validity (below critical dimensionality d_c). One then employ the RG technique to progressively integrate out the small scale fluctuations. Upon such a procedure the constants and even the functional form of the reaction Hamiltonian flow. However, one needs to follow the topology of the phase space, rather than a specific form of the trajectories. Around the transition the topology may be fully encoded in a relatively simple polynomial, which in turn provides the full characterization of the transition (at least for small $\epsilon = d_c - d$). Considering all distinct topologies, stable upon RG transformations, one may classify the possible phase transitions.

We turn now to illustration of these ideas on specific examples.

III. MODELS WITH FIRST ORDER TRANSITIONS

Consider a set of reactions, given by







FIG. 2: Phase portraits of the system Eq. (21), exhibiting the first order phase transition. Thick solid lines represent the zero-energy trajectories. (a) extinction phase, (b) transition point, (c) unlimited proliferation phase, (d) reaction of the type of Eq. (23) with $k = 2$, $m = 3$ and $n = 0$ in the extinction phase.

The corresponding reaction Hamiltonian, according to Eq. (10), may be written as:

$$H_R(p, q) = \lambda(1-p)q + \mu(p^2 - p)q = (\lambda - \mu p)(1-p)q. \quad (22)$$

There are three zero-energy lines: two generic $p = 1$ and $q = 0$, following from Eqs. (12) and (13), and the additional “accidental” line $p = \lambda/\mu$. The phase portrait of the system is depicted in Fig. 2(a-c). Its topology is qualitatively different for $\mu < \lambda$ and $\mu > \lambda$. In the former case, Fig. 2(a) the flow along the mean-field line, $p = 1$, is directed towards $q = 0$ (extinction), while in the latter, Fig. 2(c) towards $q = \infty$ (unlimited proliferation). If the two reaction rates coincide, $\mu = \lambda$, the two of the zero-energy lines $p = 1$ and $p = \lambda/\mu$ degenerate, Fig. 2(b). This is the point of the first order transition^{51,52}.

This scenario may be generalized to a set



where $n < k < m$. This reaction set is characterized by the following reaction Hamiltonian

$$H_R = [\lambda(p^n - p^k) + \mu(p^m - p^k)] \frac{q^k}{k!} = h(p)(1-p)q^k, \quad (24)$$

where $h(p)$ is a polynomial the degree $m - 1$. The zero-energy lines are given by $p = 1$, the k times degenerate line $q = 0$ along with the lines $p = p_i$, where p_i are roots of the polynomial $h(p_i) = 0$, Fig. 2(d). It is easy to check that for $\lambda/\mu = (m - k)/(k - n)$ one of the roots of the polynomial is $p_1 = 1$. In this situation the $p = 1$ zero-energy line is double degenerate, corresponding to the first order transition.





FIG. 3: Phase portrait of the DP system in the active phase. Thick solid lines represent zero-energy trajectories which divide the phase space on a number of disconnected regions. Point $B = (1, 2(\mu - \lambda)/\sigma)$ represents the active mean-field point. The system is brought to the phase transitions if points A, B, C coalesce.

IV. DIRECTED PERCOLATION AND ITS GENERALIZATIONS

A. DP models

Consider a reaction set which includes death, branching and coalescence reactions:



The corresponding reaction Hamiltonian takes the form:

$$\begin{aligned} H_R &= \lambda(1-p)q + \mu(p^2 - p)q + \frac{\sigma}{2}(p - p^2)q^2 \\ &= \left(\lambda - \mu p + \frac{\sigma}{2} p q \right) (1-p)q. \end{aligned} \quad (26)$$

The phase portrait of the DP system is depicted in Fig. 3. The lines of zero energy are generic $p = 1$ and $q = 0$ trajectories along with the “accidental” trajectory $q = 2(\mu p - \lambda)/\sigma p$. According to the mean-field (classical equations (14), (15) with $p = 1$), there is an active phase with the average density

$$\langle n \rangle = 2 \frac{\mu - \lambda}{\sigma} \quad (27)$$

for $\mu > \lambda$. The active phase corresponds to the point B in Fig. 3. The system may be brought to the extinction by driving the control parameter $m = \mu - \lambda$ through zero. Therefore the $\mu = \lambda$ point corresponds to the continuous phase transition. The transition is represented by the phase portrait with the three zero-energy trajectories intersecting at the *single* point $(1, 0)$. According to Eq. (27) the mean-field order-parameter exponent is $\beta = 1$. The other mean-field exponents¹ are $\nu_{\perp} = 1/2$, $\nu_{\parallel} = 1$.

To go beyond the mean-field picture one needs to investigate the immediate vicinity of the phase transition.




FIG. 4: Generic phase portrait of DP models in the vicinity of the phase transition (after the shift Eq. (28)). (a) active phase, $m > 0$; (b) transition point, $m = 0$; (c) extinction phase, $m < 0$. The plus/minus signs show the sign of energy in each sector. (d) The one loop diagram renormalizing u -vertex (vertices m and v are renormalized in a similar way).

To focus on this regime it is convenient to shift the momentum variable

$$p - 1 \rightarrow p. \quad (28)$$

Moreover, close to the transition the phase portrait and thus the Hamiltonian may be modelled by the three intersecting *straight* lines, Fig. 4(a-c). This way one arrives to the model Hamiltonian, applicable close to the DP transition

$$H_R = (m + up - vq) p q. \quad (29)$$

The bare values of the constants are given by $m = \mu - \lambda$, $u = \mu$, and $v = \sigma/2$. The corresponding action, Eq. (3), acquires the form^{1,6,7}

$$S = \int dt d^d x (p(\partial_t - D\nabla^2) q - mpq - up^2 q + vpq^2), \quad (30)$$

which may be recognized as Reggeon field theory action^{67,68}.

There are many other reaction sets, in addition to Eq. (25), with the same “triangular” topology of the phase portrait. Some of the examples are: $A \rightarrow \emptyset$; $A \rightarrow (m+1)A$; $2A \rightarrow A$, $m > 1$, or $A \rightarrow 2A$; $2A \rightarrow \emptyset$, etc. In the vicinity of the phase transition they all exhibit the topology of the phase portrait depicted in Fig. 4(a-c). Therefore they all may be described by the model Hamiltonian (29). Accordingly they all belong to the same DP universality class.

Naive scaling dimensions of the action (30) are $z = 2$, while $[p] + [q] = d$. Since one expects⁶⁹ both vertices u and v to have the same scaling dimensions, one finds $[p] = [q] = d/2$. As a result, the bare scaling dimensions of the vertices are $[m] = 2$, while $[u] = [v] = 2 - d/2$. Therefore below the critical dimension $d_c = 4$ the non-linear vertices u and v are relevant and the mean-field treatment is expected to fail.

The one loop corrections to the naive scaling are given by the triangular diagrams, like the one depicted in Fig. 4(d). Such diagrams are logarithmically divergent in $d = 4$, as expected. The straightforward calculations^{1,7} (see Ref. [18] for the two-loop approximation) lead to the following set of the RG equations:

$$\partial_t m = (2 - Suv) m; \quad (31)$$

$$\partial_t u = (\epsilon/2 - 2Suv) u; \quad (32)$$

$$\partial_t v = (\epsilon/2 - 2Suv) v, \quad (33)$$

where $\epsilon = 4 - d$ and the differentiation is over the logarithm of the scaling factor. We have introduced factor

$$S = \partial_t \frac{1}{4} \int_{\Lambda e^{-t}}^{\Lambda} \frac{d^d k}{D^2 k^4} = \frac{\Lambda^{d-4}}{32\pi^2 D^2}, \quad (34)$$

which may be absorbed in the proper redefinition of the running constants.

According to Eqs. (32), (33), $\partial_t(u/v) = 0$, meaning that the slope of the “accidental” zero-energy line, $q = (m + up)/v$, remains intact upon renormalization procedure. In fact, this statement is exact because of the symmetry⁶⁹. As a result, the overall topology of the phase portrait is preserved by the RG. For $d < 4$ the RG equations Eqs. (31) – (33) predict the non-trivial fixed point given by: $m^* = 0$, $u^* = \sqrt{\epsilon\mu/(4S\sigma)}$ and $v^* = \sqrt{\epsilon\sigma/(4S\mu)}$. Substituting these values into Eq. (29), one finds the fix-point reaction Hamiltonian, $H_R^*(p, q)$, corresponding to the DP universality class phase transitions. Its phase portrait is depicted in Fig. 4(b). Linearizing the RG equations (31) – (33) near the fixed point, one finds $\partial_t m = \nu_{\perp}^{-1} m$ with the critical exponent $\nu_{\perp} = (2 - \epsilon/4)^{-1} \approx 1/2 + \epsilon/16$. The other critical exponents may be deduced in the standard way^{1,7}, e.g. $\beta \approx 1 - \epsilon/6$.

B. k-particle contact processes

As mentioned in the introduction, the DP universality class is extremely robust. This fact is due to the stability of the “triangular” topology of the phase portrait near the transition. One may try to change this topology by, say, requiring four or more zero-energy trajectories to intersect in the same point. It is clear, however, that in general one must fine-tune more than one parameter to reach such a scenario. Even if bare reaction rates are specially chosen to let it happen, the tuning is not expected to survive upon RG integration of fluctuations. Therefore crossing of more than three lines is possible only in a multi-critical transition point. The only way to go beyond the DP is if a different topology of the phase portrait is imposed by an additional *symmetry*.

In this section we discuss a class of models where the *minimum* number of particles needed to initiate *any* reaction is restricted to be $k > 1$. According to Eq. (10) *all* terms in the corresponding reaction Hamiltonian must





FIG. 5: Phase portrait of the 2CPD system in the active phase (cf. with DP, Fig. 3). The zero-energy line $q = 0$ is double degenerate and is depicted by the double line. At the transition points A, B, C coalesce.

contain \hat{q}^k or higher power. In terms of the phase portrait it means that the generic $q = 0$ zero-energy trajectory is k -times degenerate. To emphasize the difference with non-degenerate lines, we denote such k -degenerate trajectories by k closely spaced parallel lines. An important fact is that the degeneracy is preserved by the RG transformations. Indeed, the fluctuations cannot initiate a reaction with less than k incoming particles, if it is not in the original reaction set. We denote such models as k -particle contact process with diffusion (kCPD).

To be specific, let us consider the case of $k = 2$, which is represented by e.g. the following set of reactions:



The corresponding reaction Hamiltonian takes the form:

$$\begin{aligned} H_R &= \frac{\lambda}{2}(1-p^2)q^2 + \frac{\mu}{2}(p^3-p^2)q + \frac{\sigma}{3}(p^2-p^3)q^3 \\ &= \left(\frac{\lambda}{2}(1+p) - \frac{\mu}{2}p^2 + \frac{\sigma}{3}p^2q \right) (1-p)q^2. \end{aligned} \quad (36)$$

The phase portrait of the 2CPD system is depicted in Fig. 5. The lines of zero energy are generic $p = 1$ and *double degenerate* $q = 0$ trajectories. The “accidental” trajectory is given by $q = 3(\mu p^2 - \lambda(1+p))/(2\sigma p^2)$. There is an active phase with the average density $\langle n \rangle = (3\mu - 6\lambda)/2\sigma$ for $\mu > 2\lambda$. It corresponds to the point B in Fig. 5. The system may be driven to the extinction by tuning the control parameter $m = \mu/2 - \lambda$ to zero. Therefore the $\mu/2 = \lambda$ point corresponds to the continuous phase transition. At the transition point the *four* (we count $q = 0$ line twice) zero-energy lines are intersecting in the single point $(1, 0)$.

Focusing on the transition region, one may shift the momentum variable $p-1 \rightarrow p$ and model the zero-energy trajectories by the straight lines, Fig. 6(a). The resulting model Hamiltonian, applicable close to the transition, is

$$H_R = (m + up - vq) p q^2. \quad (37)$$



FIG. 6: Generic phase portraits of (a) 2CPD models and (b) 3CPD models in the active phase.

The bare values of the constants are given by $m = \mu/2 - \lambda$, $u = \mu - \lambda/2$, and $v = \sigma/3$. Since near the transition $m \approx 0$ and thus $\mu \approx 2\lambda > 0$, one finds for the bare value $u \approx 3\lambda/2 > 0$. Apart from Eq. (35), there are infinitely many other reaction sets, which have the same topology of the phase portraits. Therefore the phase transition of these other models is described by the same model Hamiltonian (37). Examples include: $2A \rightarrow A$; $2A \rightarrow 4A$; $4A \rightarrow \emptyset$, etc.

In an analogous way, one may show that the phase portrait of a generic kCPD process, such as e. g. $kA \rightarrow \emptyset$; $kA \rightarrow (k+1)A$; $(k+1)A \rightarrow kA$, contains triangle of k -times degenerate $q = 0$ line, $p = 0$ line (after the shift, Eq. (28)), and the “accidental” $q = (m + up)/v$ diagonal line, see Fig. 6(b). Thus it may be described by a model Hamiltonian of the form:

$$H_R = (m + up - vq) p q^k, \quad (38)$$

where m is the control parameter of the transition and u and v are positive constants.

To find scaling exponents near the transition one assigns bare dimensions $z = 2$, and $[p] + [q] = d$. There are no perturbative corrections to the propagator (for $k \geq 2$) and thus one does not expect these exponents to be changed in the ϵ -expansion. Since both u and v vertices has to be equally (ir)relevant on the mean-field level, one has to choose the bare dimensions as $[p] = [q] = d/2$. As a result, the bare scaling dimensions of the vertices are $[m] = 2 - (k-1)d/2$, and $[u] = [v] = 2 - kd/2$. Therefore the critical dimension is expected to be $d_c = 4/k$. The corresponding mean-field exponents of the kCPD transitions at $d > d_c$ are $\beta = 1$ and $\nu_{\perp} = k/2$. Only $k = 2$ and $k = 3$ process may exhibit non-trivial behavior in physically relevant dimensions⁷⁰. We shall analyze these two cases below.

The 2CPD transition is described by the action (3) with the reaction Hamiltonian (37) (see e.g. Ref. [8]). Its critical dimension is $d_c = 2$. The one loop renormalization is given by two-vertex loops, which are logarithmically divergent in $d = 2$. One arrives⁶⁴ at the following set of RG equations:

$$\partial_l m = (1 + \epsilon/2 + \tilde{S}u) m; \quad (39)$$

$$\partial_l u = (\epsilon + \tilde{S}u) u; \quad (40)$$

$$\partial_l v = (\epsilon + 3\tilde{S}u) v, \quad (41)$$

where $\epsilon = 2 - d$ and

$$\tilde{S} = \partial_t \frac{1}{2} \int_{\Lambda e^{-t}}^{\Lambda} \frac{d^d k}{Dk^2} = \frac{\Lambda^{d-2}}{4\pi D}. \quad (42)$$

Notice that the sign of the perturbative corrections in Eqs. (39)–(41) is *opposite* to that in the corresponding DP RG equations (31)–(33). As a result, the weak coupling fix-point appears to be absolutely unstable for $d \leq d_c$ (for positive initial u). Solving Eq. (40), one finds that the coupling constant u diverges once the RG reaches a certain spatial scale $\xi^{-1} = \Lambda e^{-t}$, where in $d < 2$:

$$\xi = \left(\frac{\epsilon D}{u_0} \right)^{1/(2-d)}, \quad (43)$$

(here u_0 is an initial value of u). In $d = 2$ one finds that $\xi = \Lambda^{-1} \exp(4\pi D/u_0)$. This indicates that some new physics shows up at the scale ξ . In Appendix A we suggest that the system may develop anomalous averages, similar to those in the BCS theory. The corresponding “coherence length” appears to be exactly ξ . (See also Ref. [8,64] for further discussion).

The similar situation is encountered for $k = 3$. The critical dimension is $d_c = 4/3$. The RG analysis of Eq. (38) with $k = 3$ shows that the only vertex which acquires perturbative corrections is v . The corresponding RG equations are:

$$\partial_t m = (2 - d) m; \quad (44)$$

$$\partial_t u = (2 - 3d/2) u; \quad (45)$$


$$\partial_t v = (2 - 3d/2) v - 3\tilde{S} m u. \quad (46)$$

Once again, the coupling constant u grows indefinitely and ϵ -expansion fails to predict critical exponents.

V. PARITY CONSERVING MODELS

A. PC model

Another symmetry, which is known to alter the universality class is parity. Consider the simplest possible parity conserving (PC) reaction set



The corresponding reaction Hamiltonian takes the form:

$$H_R = (up - vq)(p^2 - 1)q, \quad (48)$$

where the bare values of the constants are given by $u = \mu$ and $v = \sigma/2$. The corresponding action is invariant under the following transformations⁵:

$$p \rightarrow -p; \quad q \rightarrow -q, \quad (49)$$

which may be traced back to the conservation of parity. As a result, the phase portrait, Fig. 7, possesses the reflection point at the origin. This symmetry is preserved




FIG. 7: Phase portrait of the parity conserving model, Eq. (47). Notice the reflection symmetry around the origin.

upon RG transformations. Therefore the “accidental” zero-energy line ($q = up/v$, according to Eq. (48)) is bound to be an odd function and thus pass through the origin. Its shape, however, may change in the process of renormalization. Consequently the phase transition can not be described by the coalescence of three points A, B, C and its nature is different from the DP class.

According to the mean-field equation $\partial_t q = 2\mu q - \sigma q^2$, the model is always in the active phase with the number of particles $\langle n \rangle = 2\mu/\sigma = u/v$. The only way to drive the mean-field dynamics towards the extinction is to send $u = \mu \rightarrow 0$. In other words, the critical point is $u_c = 0$. One may discuss, though, the scaling of particle density with $u - u_c = u$ and this way define the “magnetization” exponent β :

$$\langle n \rangle \sim u^\beta, \quad (50)$$

where the mean-field value of the exponent is $\beta = 1$.

To consider fluctuations one notices that it is not possible to perform the shift of momentum, Eq. (28), and focus on the immediate vicinity of the $(1, 0)$ point on the phase plane. Because of the symmetry (49) one has to keep the entire interval $p \in [-1, 1]$ under consideration, see Fig. 7. Therefore one must choose the scaling dimension $[p] = 0$. Since the bare scaling requires $[p] + [q] = d$, the naive scaling dimension of q is $[q] = d$. As a result, one finds (since $z = 2$) $[u] = 2$ and $[v] = 2 - d$ and the critical dimension is $d_c = 2$. One may worry that since $[p] = 0$, it is not possible to restrict the consideration to the low order polynomial in p . Instead, one has to keep all the powers of p resorting to the functional RG. We perform such a procedure in Appendix B and show that it actually justifies the use of the truncated reaction Hamiltonian (48).

The one loop RG calculation, utilizing the two-vertex loop (which is logarithmically divergent in $d = 2$) yields the following RG equations^{5,7}:

$$\partial_t u = (2 - 3\tilde{S}v)u; \quad (51)$$

$$\partial_t v = (\epsilon - \tilde{S}v)v, \quad (52)$$

where $\epsilon = 2 - d$ and \tilde{S} is given by Eq. (42). For $\epsilon > 0$ there is a non-trivial stable fix-point $v^* = \epsilon/\tilde{S}$. In the vicinity of this fix-point the relevant parameter u scales as $\partial_l u = (2 - 3\epsilon)u$ and thus its new scaling dimension is

$$[u] = 2 - 3\epsilon = 3d - 4. \quad (53)$$

This leads to the non-mean-field exponent β given by


$$\beta = d/[u] \approx 1 + \epsilon. \quad (54)$$

The fact that $\beta > \beta^{MF} = 1$ means that the actual density in $d < 2$ is less than the mean-field prediction. The fluctuations drive the system closer to the extinction. Cardy and Täuber⁵ suggested that for $d < d'_c \approx 4/3$ there is the extinct phase at finite u and the transition to the active phase takes place at some $u_c > 0$.

B. Generalized PC models

One may invent other models conserving parity. For example Cardy and Täuber⁵ considered the class of parity conserving models $2A \rightarrow \emptyset$, and $A \rightarrow (m+1)A$ with even m . The corresponding reaction Hamiltonian is $H_R = (up h_m(p) - vq)(p^2 - 1)q$, where $h_m(p) = (p^m - 1)/(p^2 - 1)$ is an even polynomial. Its phase portrait is topologically identical to Fig. 7. Thus one expects this reaction set to belong to the same universality class as PC model. This was indeed the conclusion of Ref. [5].


To find a different topology of the phase portrait and therefore a new universality class one needs to impose an additional symmetry. Following section IV B, we shall consider parity conserving reactions with minimal number $k > 1$ needed to initiate all reactions. For example, consider a parity conserving set of reactions with even k :



The reaction Hamiltonian is

$$H_R = (up^{k-1} - v h_k(p)q) p(p^2 - 1)q^k, \quad (56)$$

where $h_k(p)$ is an even polynomial of the degree $k - 2$, and $u = \mu/k!$, $v = \sigma/(k+1)!$. The corresponding phase portrait is depicted in Fig. 8(a). For an odd k a representative set of reactions is



with the reaction Hamiltonian

$$H_R = (up^k - v h_{k+1}(p)q) (p^2 - 1)q^k, \quad (58)$$

where $h_{k+1}(p)$ is an even polynomial of degree $k - 1$. The corresponding phase portrait is depicted in Fig. 8(b). These phase portraits are topologically stable upon RG transformations and thus represent a set of distinct universality classes. We call them k -particle parity conserving (kPC) classes.




FIG. 8: Phase diagrams of kPC models. (a) 2PC model; the inset shows the diagram leading to the logarithmic corrections in $d = 1$. (b) 3PC model. (c) 2PC model with all reactions including only even number of particles. This topology is not stable and evolves towards (a). (d) 2PC with $2A \rightarrow \emptyset$, the corresponding topology is essentially equivalent to the 2CPD model, Fig. 5.


Assigning the scaling dimensions as in the PC model: $[p] = 0$, $[q] = d$ and $z = 2$, one finds $[u] = 2 - (k - 1)d$ and $[v] = 2 - kd$. At the critical dimension v turns to be relevant and thus $d_c = 2/k$. Therefore at any physical dimension the kPC behavior is described by the mean-field. The only exception is the 2PC model, Fig. 8(a), which acquires logarithmic corrections to the mean-field scaling at $d = 1$. The renormalization is due to the two-loop diagrams build with the help of p^3q^3 vertex, see the inset in Fig. 8(a).

Other attempts to generalize the PC universality class appears to be unstable against RG transformations. For example, consider a parity conserving set which contains only *even* number of reagents: $2A \rightarrow 4A$ and $4A \rightarrow \emptyset$. The corresponding reaction Hamiltonian is given by $H_R = (up^2 - v(1 + p^2)q^2)(p^2 - 1)q^2$. Its phase portrait is depicted in Fig. 8(c). In addition to the PC symmetry, Eq. (49), the Hamiltonian and the phase portrait possess the

$$q \rightarrow -q \quad (59)$$

symmetry. However, this is *not* the symmetry of the full action, Eq. (3). Therefore this symmetry is not stable against the RG transformations. Indeed, e.g., using three vertexes p^2q^2 , one may generate p^3q^3 vertex which violates the symmetry (59). As a result the system belongs to the 2PC class and its phase portrait drifts towards Fig. 8(a).

One may add a competing death reaction to the $kA \rightarrow (k+2)A$ process of kPC for $k \geq 2$, such as $kA \rightarrow (k-2)A$. For example,




Because of the competition, one expects the absorbing state transition to happen at $m = 2\mu - \lambda = 0$. The

Hamiltonian is $H_R = (m - u + up^2 - vpq)(p^2 - 1)q^2$, it obeys the PC symmetry (49). The phase portrait is plotted in Fig. 8(d). One notices that in the vicinity of the transition the *local* topology is indistinguishable from Figs. 5 and 6(a). Therefore the transition belongs to the same universality class as 2CPD (see section IV B). This fact was already noticed in numerical simulations⁴⁴, but, to the best of our knowledge, remained unexplained. The identity of the phase portrait's topologies in the vicinity of transition immediately explains the universality. One can show that the other kPC processes with the competition in the k -particle channel belong to the same universality classes as corresponding kCPD models.

VI. REVERSIBLE REACTIONS

All the phase transitions, discussed above, are associated with a deformation and rearrangement of a characteristic *triangular* structure (possibly with one degenerate side) on the phase plane. There is one more possibility for the stable transition's topology which is a *rectangular* structure. All models, exhibiting rectangular topology, consist of a *single* reaction which is allowed to go both directions with different rates.

Consider, for example, a reversible reaction (same as DP, Eq. (25), but without $A \rightarrow 0$)



The corresponding reaction Hamiltonian is:

$$\begin{aligned} H_R &= \mu(p^2 - p)q + \frac{\sigma}{2}(p - p^2)q^2 \\ &= (p^2 - p)(uq - vq^2), \end{aligned} \quad (62)$$

where $u = \mu$ and $v = \sigma/2$. The phase portrait is depicted in Fig. 9(a) and has characteristic rectangular shape comprised by the generic lines $p = 1$ and $q = 0$ along with the *two* "accidental" ones $p = 0$ and $q = u/v$. The mean-field predicts the average density to be $\langle n \rangle = u/v$. The control parameter is $u = \mu$ with the critical value $u_c = 0$. Consequently the mean-field "magnetization" exponent is $\beta = 1$. One may ask if it can be modified by the fluctuations.

To answer this question, one notices that the phase portrait topology is stable against renormalization. I.e. no terms violating the rectangular structure are generated. It may be checked either by considering possible diagrams, or realizing that the action possesses the symmetry

$$p \rightarrow \frac{v}{u}q, \quad q \rightarrow \frac{u}{v}p, \quad t \rightarrow -t. \quad (63)$$

To keep the entire interval $p \in [0, 1]$ unchanged upon rescaling, one has to choose the scaling dimensions as $[p] = 0$ and $[q] = d$, then $[u] = 2$ and $[v] = 2 - d$. From here one concludes that the critical dimension is $d_c = 2$.




FIG. 9: (a-c) Phase portraits of reversible reactions. (a) $A \leftrightarrow 2A$; (b) $2A \leftrightarrow 3A$. (c) parity conserving $A \leftrightarrow 3A$; (d) A topology with $q \rightarrow -q$ symmetry unstable against RG transformations, see section VII.

The RG equations are

$$\partial_t u = (2 - \tilde{S}v)u; \quad (64)$$

$$\partial_t v = (\epsilon - \tilde{S}v)v, \quad (65)$$

where $\epsilon = 2 - d$ and \tilde{S} is given by Eq. (42). For $\epsilon > 0$ there is a stable fix-point at $v^* = \epsilon/\tilde{S}$. In its vicinity u scales as $\partial_t u = (2 - \epsilon)u$, thus its new scaling dimension is $[u] = 2 - \epsilon = d$ (see also Appendix B). As a result, the exponent is given by $\beta = d/[u] = 1 + O(\epsilon^2)$. At least in this order, it is not affected by the fluctuations. It would be interesting to know if $\beta = 1$ is exact.

A generic reversible reaction $kA \leftrightarrow mA$, with $m > k$ is described by $H_R = (p^m - p^k)(uq^k - vq^m)$. Its zero-energy lines are $q = 0$ and $p = 1$, both k -times degenerate, along with non-degenerate $p = 1$ and $q = \langle n \rangle = (u/v)^{1/(m-k)}$. For the parity conserving case ($m - k$ even) also $p = -1$ and $q = -\langle n \rangle$, Fig. 9(c). By a proper rescaling of p and q the Hamiltonian may be brought to the symmetric separable form $H_R = -f(p)f(q)$. The corresponding action is symmetric against $p \leftrightarrow q$ and $t \rightarrow -t$. Therefore the rectangular structure is stable in the course of renormalization. The topological structure is fully determined by the index k and the parity.

Therefore, one may identify two more families of universality classes. One is parity non-conserving, represented by the reversible reaction



(higher number of offsprings, $k + 1 + 2n$, does not change the universality class) with the reaction Hamiltonian:

$$H_R = (p - 1)(u - vq)p^k q^k. \quad (67)$$

We denote it as k -particle reversible (kR). The action possesses the symmetry (63), rendering stability of the rectangular topology. The upper critical dimension is

$d_c = 2/k$. An example is $2A \leftrightarrow 3A$, see Fig. 9(b), with $d_c = 1$.

The parity conserving reversible reactions



(higher number of offsprings, $k+2n$, does not alter the universality class) with the reaction Hamiltonian:

$$H_R = (p^2 - 1)(u - vq^2)p^k q^k. \quad (69)$$

We denote it as kRPC. The corresponding action is symmetric against two symmetries Eqs. (49) and (63). They impose stability of the rectangular topology symmetric with respect to reflection around the origin. The critical dimension of kRPC family is $d_c = 2/(k+1)$. An example is $A \leftrightarrow 3A$, see Fig. 9(c) with $d_c = 1$. All other reactions of this type have $d_c < 1$ and thus are fully described by the mean-field treatment (see, however, Ref. [71]).

VII. CONCLUSIONS

We have argued that the universality classes of phase transitions in reaction-diffusion models may be classified according to the topological structures of the corresponding phase spaces. This structure is fully encoded in the web of the zero-energy trajectories. The simplest and most stable structure is given by three mutually intersecting lines. By changing a single control parameter the three intersection points may be made to coalesce, see Fig. 4. At such a value of the control parameter the system undergoes the phase transition into the absorbing phase. The corresponding universality class is known as directed percolation.

There is only a limited number of ways to organize a *continuous* phase transition, governed by a *single* control parameter, which utilizes topology different from the DP. We have identified *four* families of such transitions, which are stabilized by an additional symmetry or symmetries:

(i) A generic reaction set constrained by the requirement that *all* reactions need at least k incoming particles. The corresponding phase portrait is bound to have the $q = 0$ line to be k times degenerate, Fig. 6. This property is robust against RG transformations. Indeed, no vertexes with fewer than k external q “legs” can be generated. The triangular topology with one k -degenerate line, Fig. 6, defines a family of universality classes. We call them kCPDs ($k = 1$ is DP). Their upper critical dimension is $d_c = 4/k$.

(ii) A set of reactions which conserve parity. In this case the Hamiltonian and the action are invariant under the transformation (49). It dictates the reflection symmetry of the corresponding phase portraits, Figs. 7, 8. The symmetry is preserved upon renormalization. In addition to the parity conservation one may require the minimal k incoming particles for every reaction. This generate phase portraits depicted in Fig. 8. There is one universality class for every k , termed as kPC , Fig. 8(a,b)

($k = 1$ is PC). Their upper critical dimension is $d_c = 2/k$. To realize a kPC transition, the reaction starting from the minimal k number of particles must go only up, e.g. $kA \rightarrow (k+2)A$. By adding down-going reaction, e.g. $kA \rightarrow (k-2)A$, (for $k \geq 2$) the model is transformed into the kCPD class (despite of the parity conservation).

(iii) A single reaction which is allowed to go both directions, with different rates: $kA \leftrightarrow (k+1+2n)A$. The corresponding reaction Hamiltonian and the action are symmetric under the exchange transformation, Eq. (63). The phase portrait has the stable rectangular structure with k -degenerate $p = 0$ and $q = 0$ lines, Fig. 9(a,b). Upon decreasing the creation rate the rectangle collapses onto the interval $p = [0, 1]$. We call such transitions kR. Their critical dimension is $d_c = 2/k$ (same as kPC, but exponents are different in $d < d_c$).

(iv) A single reaction which is allowed to go both directions and conserves parity: $kA \leftrightarrow (k+2n)A$. The corresponding reaction Hamiltonian and the action are symmetric under the *two* symmetry transformations, Eqs. (49) and (63). The corresponding phase portrait has the stable “checkered” structure, Fig. 9(c), which collapses when sending creation rate to zero. We denote such transitions as kRPC. Their critical dimension is $d_c = 2/(k+1)$.

Altogether we identify five non-trivial universality classes with $d_c > 1$: PC, PCPD, 3CPD, PC, 1R. In addition there are four marginal classes with $d_c = 1$: 4CPD, 2PC, 2R, 1RPC.

We have not found any other stable, topologically distinct structures in the phase space. Consider, for example, topology depicted in Fig. 9(d). The phase portrait is symmetric under the transformation $q \rightarrow -q$. A corresponding reaction set consists of reactions *all* starting from *odd* number of particles: e.g. $A \rightarrow \emptyset$; $A \rightarrow 2A$; $3A \rightarrow \emptyset$. The corresponding Hamiltonian (after the shift (28) and neglecting irrelevant terms) is $H_R = (m + up - vq^2)pq$. By changing the parameter m , e.g. by increasing the rate of annihilation $A \rightarrow \emptyset$, one may bring the system to the phase transition into the extinction phase. Naively, such a transition is associated with the vertex of the parabola crossing the point $(1, 0)$, see Fig. 9(d). Such a topology is different from all considered above and could represent a new universality class.

However, the $q \rightarrow -q$ symmetric phase portrait, Fig. 9(d), is *unstable* upon renormalization. Indeed, combining two vertexes pq^3 and p^2q in the loop, one generates pq^2 term, which violates the symmetry. This term represents the induced reaction $2A \rightarrow A$, originating from two reactions $A \rightarrow 2A$ followed by $3A \rightarrow \emptyset$. In other words $q \rightarrow -q$ is *not* a symmetry of the action and therefore it is not preserved by the RG. As a result, the initially symmetric zero-energy line $p = (vq^2 - m)/u$ is shifted and deformed upon renormalization. The topology drifts towards that of the DP, Fig. 4. The latter is the actual universality class of the reaction set, considered here. To keep the vertex of the parabola right at the $q = 0$ line, one has to fine-tune at least one addi-

tional control parameter, besides m . This is a case of a tricritical transition point^{35,36,37,38,39}.

In the present work we have restricted ourselves to the one-component reactions. It is desirable to extent the strategy to reaction–diffusion models with several reagents. Each new reagent brings an additional reaction coordinate and corresponding momentum. E.g. a two-component model requires 4 dimensional phase space with 3d surfaces of constant energy. The corresponding classical dynamics may be not-integrable, making the phase space topology rather complicated. The situation may be simplified by the presence of conservation laws. For example, reaction $A + B \rightarrow 2B$ conserves number of particles. This leads to the classical dynamics with an additional integral of motion, besides energy. It is most clearly seen after a canonical transformation⁷² $p = e^P$, $q = Q e^{-P}$, which leads to the integral of motion $Q_A + Q_B = \text{const}$. Even with such simplifications understanding of the full phase space dynamics of multi-component systems remains a challenge.

This work was supported by the NSF grant DMR-0405212 and by A. P. Sloan foundation.

APPENDIX A: 2CPD MODEL

Here we consider the 2CPD model (same as PCPD) which was shown to exhibit the run-away RG flow for $d \leq 2$, see section IV B. We shall argue that this behavior of the RG indicates rearrangement of the groundstate, such that the vacuum supports the anomalous averages of the type $\langle q^2 \rangle$, similar to the BCS theory.

In a vicinity of the phase transition the action of the model is given by Eq. (3) with the reaction Hamiltonian (37). To ensure convergence of the functional integral, it is convenient to perform rotation of the integration contour in the complex p -plane: $p \rightarrow ip$. This way one arrives at the action of the form:

$$S = \int dt d^d x \left(ip (\partial_t q - D \nabla^2 q - m q^2 + v q^3) + up^2 q^2 \right). \quad (\text{A1})$$

We note, in parenthesis, that this is the Martin-Siggia-Rose action of the following Langevin process with the multiplicative noise η :

$$\partial_t q = D \nabla^2 q + m q^2 - v q^3 + q \eta(x, t), \quad (\text{A2})$$

where $\langle \eta(x, t) \eta(x', t') \rangle = 2u \delta(x - x') \delta(t - t')$.

We shall assume now that the vacuum of the theory supports the anomalous average value:

$$\langle q^2(x, t) \rangle = \Delta, \quad (\text{A3})$$

which is to be determined from the self-consistency condition. Neglecting the non-linear fluctuation terms, the action acquires the form:

$$S = \int dt d^d x \left(ip (\partial_t - D \nabla^2 + v \Delta) q + u \Delta p^2 - im \Delta p \right). \quad (\text{A4})$$

With this Gaussian action one can evaluate $\langle q^2(x, t) \rangle$ and impose the condition, Eq. (A3). This leads to the self-consistency equation:

$$\Delta = u \Delta \int \frac{d^d k}{D k^2 + v \Delta} + \frac{m^2}{v^2}. \quad (\text{A5})$$

Without the first term on the r.h.s. $\Delta = (m/v)^2$, which is simply the mean-field prediction for $\langle q^2(x, t) \rangle$. In $d \leq 2$ this equation has a non-trivial solution even at $m = 0$. In particular at $d = 2$ one finds

$$\Delta^{(d=2)} = \frac{\Lambda^2 D}{v} e^{-4\pi D/u}, \quad (\text{A6})$$

where $\Lambda \sim 1/a$ is the momentum cutoff. In $d < 2$ one obtains

$$\Delta^{(d<2)} \sim \frac{D}{v} \left(\frac{u}{(2-d)D} \right)^{2/(2-d)}, \quad (\text{A7})$$

As a result, one finds that the ‘‘order parameter’’ Δ is associated with the ‘‘coherence length’’ ξ , given by:

$$v \Delta = D \xi^{-2}, \quad (\text{A8})$$

Notice that ξ is exactly the characteristic spatial scale for the breakdown of the RG treatment of section IV B, see Eq. (43). This consideration suggests that the divergence of the RG flow is associated with the development of the anomalous average, Eq. (A3).

APPENDIX B: FUNCTIONAL RENORMALIZATION GROUP

In some problems one can not focus on the immediate vicinity of the $(1, 0)$ point in the phase plane (p, q) . Instead, one has to keep under consideration the entire interval $p \in [0, 1]$, or even $p \in [-1, 1]$. This happens e.g. in parity conserving models, because of the $p \rightarrow -p$, $q \rightarrow -q$ symmetry. To keep the p -interval intact upon the renormalization, one must choose the scaling dimension $[p] = 0$. Since the bare scaling requires $[p] + [q] = d$, one is left with the naive scaling dimension $[q] = d$. With such scaling dimensions one may restrict the reaction Hamiltonian to lowest powers in q (typically the first and second), but one must keep *all* powers of p . As a result, one has to employ the functional RG treatment (cf. Ref. [64]).

The generic reaction Hamiltonian for the absorbing-state models, mentioned above, is

$$H_R = f(p) q - g(p) q^2, \quad (\text{B1})$$

where $f(p) = \sum_n f_n p^n$ and $g(p) = \sum_n g_n p^n$ are polynomials of p . From normalization condition (12) follows that

$$f(1) = g(1) = 0. \quad (\text{B2})$$

Specific models may possess additional symmetries which dictate further restrictions on the polynomials $f(p)$ and $g(p)$, e.g. for the PC models $f(p)$ is odd, while $g(p)$ is even, due to the parity conservation: $p \rightarrow -p$, $q \rightarrow -q$. We shall keep presentation general, imposing this additional symmetries at a latter stage. Because of the assigned bare scaling dimensions $[p] = 0$, $[q] = d$, and $z = 2$ one finds $[f_n] = 2$ and $[g_n] = 2 - d$. Thus the formal critical dimension is $d_c = 2$.

The one loop renormalization is given by the two-vertex loops and leads to the following set of the RG equations:

$$\partial_l f_n = 2f_n - \frac{1}{2}\tilde{S} \sum_{m,k} m(m-1) \delta_{n,m+k-2} f_m g_k, \quad (\text{B3})$$

$$\partial_l g_n = \epsilon g_n - \frac{1}{2}\tilde{S} \sum_{m,k} m(m-1) \delta_{n,m+k-2} g_m g_k, \quad (\text{B4})$$

where $\epsilon = 2 - d$ and \tilde{S} is given by Eq. (42). Factors $\frac{1}{2}m(m-1)$ describe combinatorial number of pairs which may form the loop. The delta-symbols enforce the proper number of the external (slow) “legs”. Equations (B3), (B4) may be written as coupled partial differential equations for the functions $f(p, l) = \sum_n f_n(l) p^n$ and $g(p, l) = \sum_n g_n(l) p^n$:

$$\partial_l f = 2f - \frac{1}{2}\tilde{S} g \partial_p^2 f, \quad (\text{B5})$$

$$\partial_l g = \epsilon g - \frac{1}{2}\tilde{S} g \partial_p^2 g, \quad (\text{B6})$$

For $\epsilon > 0$ Eq. (B6) predicts the non-trivial stable fix-point polynomial $g^*(p)$, satisfying $\partial_p^2 g^* = 2\epsilon/\tilde{S}$. In view of Eq. (B2) the proper solution is

$$g^*(p) = \frac{\epsilon}{\tilde{S}} (p-1)(p+\kappa), \quad (\text{B7})$$

where parameter κ is not specified at this stage. Substituting this in Eq. (B5) one finds:

$$\partial_l f = 2f - \frac{\epsilon}{2} (p-1)(p+\kappa) \partial_p^2 f, \quad (\text{B8})$$

Since this is a Schrödinger type equation, one may look for its solution in the form

$$f(p, l) = \sum_n e^{(2-E_n)l} \phi_n(p), \quad (\text{B9})$$

where the eigen-functions $\phi_n(p)$ are solutions of the stationary equation

$$E_n \phi_n(p) = \frac{\epsilon}{2} (p-1)(p+\kappa) \partial_p^2 \phi_n(p). \quad (\text{B10})$$

Notice that if one chooses $\phi_n(j)$ to be a n -th degree polynomial, the r.h.s. of Eq. (B10) is also a polynomial of the same degree. It is clear then that one may always find a solution $\phi_n(p)$ as a polynomial of the degree n . This means that if one started from a polynomial of some degree N , the higher powers will not be generated by the RG. That is, the sum in Eq. (B9) is always confined to $1 \leq n \leq N$.

To find the eigen-energies E_n one needs to compare coefficients of the leading power of p on the both sides of Eq. (B10). This leads to $E_n = n(n-1)\epsilon/2$. From here and Eq. (B9) one finds that the scaling dimensions of the coefficients of the $f(p)$ polynomial at the non-trivial fix-point, Eq. (B7), are

$$[f_n] = 2 - E_n = 2 - n(n-1)\epsilon/2. \quad (\text{B11})$$

The conclusion is that it is sufficient to keep the polynomials $g(p)$ to be of the second degree, cf. Eq. (B7), while $f(p)$ to be of the lowest possible degree consistent with the symmetries of the model.

In the PC model of section V A $g(p)$ must be even and thus $\kappa = 1$, leading to $g(p) = g_2(p^2 - 1)$. On the other hand, $f(p)$ must be odd and at least of the degree $n = 3$ ($n = 1$ odd polynomial can not satisfy Eq. (B2)). Therefore it can be chosen to be $f(p) = f_3(p^3 - p)$. At the fix-point $g_2^* = \epsilon/\tilde{S}$ and $[f_3] = 2 - 3\epsilon$. This justifies treatment of section V A with the identification $v = g_2$, and $u = f_3$.

In the reversible model $A \leftrightarrow 2A$ of section VI $g(p) = v(p^2 - p)$ and thus $\kappa = 0$. On the other hand, $f(p) = u(p^2 - p)$ which is the proper stationary eigenfunction of Eq. (B10): $\phi_2(p)$. As a result, no other terms in $f(p)$ polynomial are generated upon renormalization. This is consistent with the robustness of the rectangular structure. Since $u = f_2$, its scaling dimension according to Eq. (B11) is $[u] = 2 - \epsilon$.

¹ For review see e. g. H. Hinrichsen, Adv. Phys. **49**, 815 (2000).

² J. Marro and R. Dickman, *Nonequilibrium phase transitions in lattice models* (Cambridge University Press) 1999.

³ G. Ódor, Rev. Mod. Phys. **76**, 663 (2004).

⁴ S. Lübeck, International Journal of Modern Physics B **18**, 3977 (2004).

⁵ J. Cardy and U. C. Täuber, Phys. Rev Lett. **77**, 4780

(1996). J. Cardy and U. C. Täuber, J. Stat. Phys. **90**, 1 (1998).

⁶ H.-K. Janssen, U. C. Täuber, Ann. Phys. (NY) **315** 147 (2005).

⁷ U. C. Täuber, M. Howard and B. P. Vollmayr-Lee, J. Phys. A: Math. Gen. **38**, 79 (2005).

⁸ M. Henkel and H. Hinrichsen, J. Phys. A **37**, R117 (2004).

⁹ M. Bramson and J. L. Lebowitz J. Stat. Phys. **65**, 941

- (1991).
- ¹⁰ J. L. Cardy, in *Proceedings of mathematical beauty of physics*, ed by J.-B. Zuber, Adv. Ser. in Math. Phys. **24**, 113 (1997)
 - ¹¹ D. C. Mattis and M. L. Glasser, Rev. Mod. Phys. **70**, 979 (1998).
 - ¹² D. Ben-Avraham and S. Havlin, *Diffusion and Reactions in Fractals and Disordered Systems*, Cambridge University Press, (2000).
 - ¹³ G. M. Schütz *Phase Transitions and Critical Phenomena* vol 19, ed C. Domb and J. L. Lebowitz (London: Academic Press) (2001).
 - ¹⁴ R. A. Blythe and A. J. Bray Phys. Rev. E **67**, 041101 (2003).
 - ¹⁵ L. Canet, B. Delamotte, O. Deloubrière, and N. Wschebor, Phys. Rev. Lett. **92**, 195703 (2004), L. Canet, H. Chaté, and B. Delamotte, Phys. Rev. Lett. **92**, 255703 (2004).
 - ¹⁶ P. Grassberger, Z. Phys. B **47**, 365 (1982).
 - ¹⁷ J. L. Cardy and P. Grassberger, J. Phys. **A 18**, L267–L271 (1985).
 - ¹⁸ H. K. Janssen, Z. Phys. B **42**, 151 (1981).
 - ¹⁹ T. E. Harris, Ann. Prob. **2** 969 (1974).
 - ²⁰ W. Kinzel, in G. Deutscher, R. Zallen, and J. Adler, editors, *Ann. Isr. Phys. Soc.*, v. 5, Bristol, Adam Hilger (1983).
 - ²¹ T. M. Liggett, *Interacting particle systems*. Springer, Heidelberg, (1985).
 - ²² I. Jensen and A. J. Guttmann, Nucl. Phys. B (Proc. Suppl.) **47**, 835 (1996).
 - ²³ G. Grinstein and M. A. Muñoz, *The statistical mechanics of absorbing states, Lecture Notes in Physics*, Springer **493**, 223 (1997).
 - ²⁴ S. Lübeck, R. D. Willmann, Nucl. Phys. B **718**, 341 (2005).
 - ²⁵ D. Zhong and D. Ben-Avraham, Phys. Lett. A **209** 333, (1995).
 - ²⁶ L. Canet, H. Chaté, B. Delamotte, I. Dornic, and M. A. Muñoz, Phys. Rev. Lett. **95**, 100601 (2005).
 - ²⁷ I. Dornic, H. Chaté, J. Chave and H. Hinrichsen, Phys. Rev. Lett. **87** 5701 (2001).
 - ²⁸ M. Droz, A. L. Ferreira and A. Lipowski, Phys. Rev. E **67** 056108 (2003).
 - ²⁹ M Rossi, R. Pastor-Satorras and A. Vespignani, Phys. Rev. Lett. **85** 1803 (2000).
 - ³⁰ K. Park, H. Hinrichsen, I-M. Kim, Phys. Rev. E **66**, 025101(R) (2002).
 - ³¹ S. Lübeck, Phys. Rev. E **65** 046150 (2002).
 - ³² Z. Racz, Lecture Notes, Les Houches, cond-mat/0210435 (2002).
 - ³³ S. Lübeck and PC. Heger, Phys. Rev. Lett. **90** 230601 (2003).
 - ³⁴ J.D. Noh and H. Park, Phys. Rev. E **69**, 016122 (2004).
 - ³⁵ H. K. Janssen, J. Phys. A: Math. Gen. **20** 5733 (1987)
 - ³⁶ T. Ohtsuki and T. Keyes, Phys. Rev. A **35** 2697 (1987).
T. Ohtsuki and T. Keyes, Phys. Rev. A **36** 4434 (1987).
 - ³⁷ H. K. Janssen, M. Mülller and O. Stenull, Phys. Rev. E **70** 026114 (2004).
 - ³⁸ S. Lübeck, J. Stat. Phys. **123**, 193 (2006)
 - ³⁹ P. Grassberger, J. Stat. Mech. **1**, P01004 (2006).
 - ⁴⁰ F. van Wijland, Phys. Rev. Lett. **89**, 190602 (2002).
 - ⁴¹ M.J. Howard and U.C. Täuber, J. Phys. A **30**, 7721 (1997).
 - ⁴² E. Carlon, M. Henkel, and U. Schollwöck, Phys. Rev. E **63** 036101 (2001).
 - ⁴³ H. Hinrichsen, Phys. Rev. E **63** 036102 (2001).
 - ⁴⁴ K. Park, H. Hinrichsen, and I-M. Kim, Phys. Rev. E **63** R065103 (2001).
 - ⁴⁵ M. Henkel M and H. Hinrichsen, J. Phys. A **34** 1561 (2001).
 - ⁴⁶ M. Henkel and U. Schollwöck, J. Phys. A **34** 3333 (2001).
 - ⁴⁷ G. Ódor, M.C. Marques and M.A. Santos, Phys. Rev. E **65** 056113 (2002).
 - ⁴⁸ R. Dickman and M.A.F. de Menezes, Phys. Rev. E **66** 045101 (2002).
 - ⁴⁹ J. Kockelkoren and H. Chaté, Phys. Rev. Lett. **90** 125701 (2003).
 - ⁵⁰ G. Ódor 2003, Phys. Rev. E **67** 016111 (2003).
 - ⁵¹ M. Paessens and G.M. Schütz, newblock J. Phys. A **37** 4709 (2004).
 - ⁵² F. Baumann, S. Stoimenov and M. Henkel, J. Phys. A: Math. Gen. **39**, 4095 (2006).
 - ⁵³ G.T. Barkema and E. Carlon 2003, Phys. Rev. E **68** 036113 (2003).
 - ⁵⁴ S.-C. Park and H. Park, Phys. Rev. E **71**, 016137 (2005).
 - ⁵⁵ H. Hinrichsen, e-print cond-mat/0501075.
 - ⁵⁶ I. Dornic, H. Chaté, M. A. Muñoz, Phys. Rev. Lett. **94**, 100601 (2005).
 - ⁵⁷ L. V. Keldysh, Zh. Eksp. Teor. Fiz. **47**, 1515 (1964).
 - ⁵⁸ P. C. Martin, E. D. Siggia, and H. A. Rose, Phys. Rev. A **8**, 423 (1973).
 - ⁵⁹ DeDominics, J. Physique (Paris) Colloq. **37**, C2247 (1976).
 - ⁶⁰ M. Doi, J. Phys. **A 9**, 1479–1495 (1976).
 - ⁶¹ L. Peliti, Journal de Physique **46**, 1469–1483 (1984).
 - ⁶² A. Kamenev, in *Strongly Correlated Fermions and Bosons in Low-Dimensional Disordered Systems* I. V. Lerner, et. al. editors, pp. 313-340, Kluwer Academic Publishers, Dordrecht, Boston, London, (2002).
 - ⁶³ A. Kamenev, in *Nanophysics: Coherence and Transport*, H. Bouchiat, et al. (editors); pp. 177-246, Elsevier, Amsterdam, (2005).
 - ⁶⁴ H.-K. Janssen, F. van Wijland, O. Deloubriere, and U. C. Täuber, Phys. Rev. E **70**, 056114 (2004).
 - ⁶⁵ B.P. Lee, J. Phys. A **27**, 2633 (1994); B.P. Lee and J.L. Cardy, J. Stat. Phys. **80**, 971 (1995).
 - ⁶⁶ V. Elgart, A. Kamenev, Phys. Rev. E **70**, 041106 (2004).
 - ⁶⁷ J. L. Cardy and R. L. Sugar, J. Phys. **A 13**, L423–L427 (1980).
 - ⁶⁸ P. Grassberger, in *Nonlinearities in complex systems, proceedings of the 1995 Shimla conference on complex systems*, edited by S. Puri et al., Narosa Publishing, New Dehli, (1997).
 - ⁶⁹ In fact the action Eq. (30) after a proper rescaling possesses the symmetry $p \rightarrow -q$, $q \rightarrow -p$ and $t \rightarrow -t$, which shows the equivalence of u and v vertices.
 - ⁷⁰ In principle, $k = 4$ may exhibit a marginally non-mean-field behavior in $d = 1$, but we shall not try to investigate it here.
 - ⁷¹ G. Ódor, Braz. J. of Phys. **33** 431 (2003).
 - ⁷² A. N. Jordan, E. V. Sukhorukov, and S. Pilgram, J. Math. Phys. **45**, 4386 (2004).