

# Ways to solve of the evolution equation in the homogeneous free molecular condensation under dynamic conditions

Victor Kurasov

## Abstract

The ways to construct solution of the evolution equation in the homogeneous free molecular condensation under dynamic conditions are presented in parametric form.

Taking the evolution equation in homogeneous case in the power approximation [1] one can come to

$$y^{(IV)} = Ay^m$$

where

$$y = \zeta/\zeta_0$$

is the ratio of supersaturations ( $\zeta_0$  is a base for approximation) and  $m$  is the big integer parameter. Here  $A$  is the intensity of droplets formation at peak or at  $\zeta = \zeta_0$ .

This equation is studied with the help of methods from [2].

Integration gives

$$2y'y''' - (y'')^2 = \frac{2A}{m+1}y^{m+1} + 4/3C$$

with the arbitrary constant  $C$ . The substitution

$$w = (y')^{3/2}$$

brings the last equation to

$$w''_{yy} = \left(\frac{3A}{2m+2}y^{m+1} + C\right)w^{-5/3}$$

Now we shall determine  $C$ . The value  $y = 0$  corresponds to the zero supersaturation. There (and before) there is no formation of droplets. It means that here the supersaturation is ideal one. The ideal supersaturation is considered as allowing the linearization. Then  $y' = 0$   $y'' = 0$ ,  $y''' = 0$  at  $y = 0$ . Then  $C = 0$ .

Then

$$w''_{yy} = \frac{3A}{2m+2} y^{m+1} w^{-5/3}$$

The last equation is the Emden-Fowler equation. The transformation

$$\xi = \frac{2(m+1) - 5/3 + 3}{-5/3 - 1} y^{\frac{m+1+2}{-5/3-1}} w$$

$$u = y^{\frac{(m+1)+2}{-5/3-1}} (y w'_y + \frac{(m+1)+2}{-5/3-1} w)$$

leads to the Abel equation

$$u u'_\xi - u = -\frac{(m+1+2)(m+1-5/3+1)}{(2(m+1)-5/3+3)^2} \xi + \frac{3A}{2m+2} \left( \frac{-5/3-1}{2(m+1)-5/3+3} \right)^2 \xi^{-5/3}$$

The value of parameter  $m$  is the big parameter in the theory of nucleation. Then one can see the behavior of coefficients

$$\frac{(m+1+2)(m+1-5/3+1)}{(2(m+1)-5/3+3)^2} \rightarrow m^0 \quad \text{at } m \rightarrow \infty$$

$$\frac{3A}{2m+2} \left( \frac{-5/3-1}{2(m+1)-5/3+3} \right)^2 \rightarrow m^{-3} \quad \text{at } m \rightarrow \infty$$

Then the last equation be reduced to

$$u u'_\xi - u = -\xi + \epsilon \xi^{-5/3}$$

with small  $\epsilon$ . The last equation evidently can be solved with ordinary perturbation technique.

In the zero approximation the solution can be written in a parametric form.

$$\xi = C \exp\left(-\int \frac{\tau d\tau}{\tau^2 - \tau + 1}\right)$$

$$u = \tau C \exp\left(-\int \frac{\tau d\tau}{\tau^2 - \tau + 1}\right)$$

or

$$u = \tau\xi$$

Beside the power approximation the exponential approximation for the rate of nucleation is widely used. Since the power approximation goes to the exponential one after the limit  $m \rightarrow \infty$ , the same constructions can be used for the evolution equation in the exponential approximation.

Although the ways to construct solution are given the current solution can not be effectively used. It is more convenient to use the solution of the integral equation directly in the exponential approximation where it has the universal form [3].

One can also study the case of nucleation on the surface. When the embryous growth occurs in the free molecular regime, one can quite analogously come to the following equation

$$y''' = Ay^m$$

With the help of substitution

$$v(y) = (y')^2$$

it can be reduced to

$$v''_{yy} = 2Ay^m v^{-1/2}$$

This is the equation of the Emden-Fauler type

$$y'' = Ax^n y^m$$

Then the already mentioned transition

$$\xi = \frac{2n + m + 3}{m - 1} x^{\frac{n+2}{m-1}} y$$

$$u = x^{\frac{n+2}{m-1}} (xy'_z + \frac{n+2}{m-1} y)$$

leads to the Abel equation

$$uu'_\xi - u = -\frac{(n+2)(n+m+1)}{(2n+m+3)^2} \xi + A \left( \frac{m-1}{2n+m+3} \right)^2 \xi^m$$

Here  $n$  is the initial big parameter  $m$  and  $m$  here is  $-1/2$ . Then

$$\frac{(n+2)(n+m+1)}{(2n+m+3)^2} \rightarrow 1/4$$

$$\left(\frac{m-1}{2n+m+3}\right)^2 \sim n^{-2} \rightarrow 0$$

So, approximately

$$uu'_\xi - u = -\frac{(n+2)(n+m+1)}{(2n+m+3)^2}\xi$$

which has been already discussed above.

The same constructions can be used for the exponential approximation.

Here the generalization for the case of decay of metastable phase can be directly made.

One can not regard these constructions as some concrete precise formulas but only as a way to seek a solution. Even if the solution is presented it is more convenient to use the initial integral form and to construct the solution numerically or approximately by methods discussed in [3] and the papers cited there.

## References

- [1] Kurasov V.B., Distribution of embryos in the process of condensation. Deponed in VINITI N 2809-B95 from 24.10.1995, 15 p.
- [2] A.Polyanin, V. Zaitsev, Handbook of exact solutions for ordinary differential equations, Second edition, Chapman and Hall/CRC, London-NewYork, 2003, 802p.
- [3] Kurasov V.B. The form of the embryos size spectrum in the first order phase transition, Theoretical and Mathematical Physics, v. 131, N 3, p.503-528