

# Stable and unstable vortices in multicomponent Bose-Einstein condensates

Juan J. García-Ripoll and Víctor M. Pérez-García,  
*Departamento de Matemáticas, Escuela Técnica Superior de Ingenieros Industriales  
 Universidad de Castilla-La Mancha, 13071 Ciudad Real, Spain*  
 (February 10, 2019)

We study the stability and dynamics of vortices in a two-species condensate as prepared in the recent JILA experiment ( M. R. Matthews *et al.*, Phys. Rev. Lett. 83 (1999) 2498). We find that of the two available configurations, in which one specie has vorticity  $m = 1$  and the other one has  $m = 0$ , only one is linearly stable, which agrees with experimental results. However, it is found that in the unstable case, the vortex is not destroyed by the instability but suffers a revival process in which it is transferred from one of the condensate species to the other. This prediction can be checked in current experiments.

PACS number(s): 03.75. Fi, 67.57.Fg, 67.90.+z, 02.70 Hm

Vortices appear in many different physical contexts ranging from classical phenomena such as fluid mechanics [1] and nonlinear Optics [2] to purely quantum phenomena such as superconductivity and superfluidity [3]. Only in the last two years more than 100 papers have been published in Physical Review Letters concerning vortices, which is only a naive way to measure one very important object in physics.

In fact a vortex is the simplest topological defect one can construct: in a closed path around a vortex the phase of the involved field undergoes a  $2\pi$  winding and stabilizes a zero value of the field placed in what is called the vortex core. The vortex is usually stabilized by topological constraints since removing the phase singularity implies an effect on the boundaries of the system which is difficult to achieve using only local perturbations.

In particular, the concept of a vortex is central to our understanding of superfluidity and quantized flow. This is why after the experimental realization of Bose-Einstein condensates (BEC) with ultracold atomic gases [4] the question of whether atomic BEC's are superfluids has triggered the analysis of vortices. The main goals have been to propose a robust mechanism to generate vortices [5] and detect them [6]. Finally another research area has been the analysis of vortex stability [7–9].

Although most of the theoretical effort has concentrated on single condensate systems, the final experimental realization of BEC vortices [11] following a proposal of [12] was attained with a two-species Rb condensate. In Ref. [11] it was found that the vortex is stable in only one of the two possible configurations (the so-called  $|1\rangle$  state), while the other possibility leads to some kind of instability. It is our intention in this paper to study in detail this instability mechanism, which is purely dynamic and prove that it is not due to dissipation.

Our results, to be discussed in detail later, can be well illustrated in Fig. 1, which presents real time evolution of different initial configurations. When the vortex is initially put in the  $|1\rangle$  state nothing happens [Fig 1(a)], because of the linear stability of this type of vortices, to be proven later. Instead, when the vortex is put in a  $|2\rangle$  specie instability develops [Fig 1(b)].




FIG. 1. (a) Time evolution of a stable vortex in a  $|1\rangle$  state. Different plots are equispaced by 10 time units. (b) Instability of a vortex initially put in a  $|2\rangle$  state (c) Evolution of the  $|1\rangle$  component in the same simulation as (b), following a nontrivial evolution which includes capture of the vortex for some time before transferring it again to the  $|2\rangle$  component

The study of the simultaneous evolution of both components reveals that the vortex is transferred to  $|1\rangle$  and eventually goes back to  $|2\rangle$ , but it is never ejected out of the condensate. This prediction, together with an stability analysis which agrees with experimental results, are the main conclusions of our paper.

*The model.*- In this work we will use the zero temperature approximation, in which collisions between the condensed and non condensed atomic clouds are neglected. In the two species case this leads to a pair of coupled Gross-Pitaevskii equations (GPE)

$$i\hbar \frac{d}{dt} \psi_1 = \left[ -\frac{\hbar^2 \nabla^2}{2m} + V_1 + \tilde{u}_{11} |\psi_1|^2 + \tilde{u}_{12} |\psi_2|^2 \right] \psi_1, \quad (1a)$$

$$i\hbar \frac{d}{dt} \psi_2 = \left[ -\frac{\hbar^2 \nabla^2}{2m} + V_2 + \tilde{u}_{21} |\psi_1|^2 + \tilde{u}_{22} |\psi_2|^2 \right] \psi_2. \quad (1b)$$

To simplify the formalism and in analogy with experiments we assume that both potentials are of the form  $V_1(\vec{r}) = V_2(\vec{r}) = \frac{1}{2} m \omega^2 r^2$ .

Next we change to a new set of units based on the trap characteristic length,  $a_0 = \sqrt{\hbar/m\omega}$ , and period,  $\tau = 1/\omega$  defined as  $x \rightarrow x/a_0, t \rightarrow t/\tau, u_{ij} = 4\pi N a_{ij}/a_0$ . Since experimental results are obtained with systems in which the number of particles is the same for each component,  $N_1 = N_2 = N$ , we can impose the normalization

$$\int |\psi_1(\vec{r})|^2 = \int |\psi_2(\vec{r})|^2 \equiv 1. \quad (2)$$

Although the experiments at JILA [11] have been performed in spherically symmetric traps, it has been shown in a previous work [9] that in the transition from a spherical trap to a pancake one no new unstable modes are introduced. This motivates us to use a simpler model in which the dependency on the axial direction has been integrated out following the ideas of [14]. To scan all the range of realistic values for the interaction strengths,  $u_{ii}$  we have studied the following line of the parameter space,

$$U = g \begin{pmatrix} 1.00 & 0.97 \\ 0.97 & 0.94 \end{pmatrix}. \quad (3)$$

The proportion between the off- and on-diagonal elements have been chosen as to match the experimental measures for Rb [10]. If we assume that the number of particles is distributed equally along the three spatial directions, we can estimate the global factor as  $g \sim N^{2/3} a_{11}/a_0$ ,  $a_{11}$  is the scattering length of the first component. However, as we will show later our results are essentially independent on the number of particles.

In a previous work with pinned vortices [13] we showed that the onset of dynamical stability begins around  $u_{11} u_{22} = u_{12}^2$ . In our present case, which corresponds to the experimental values of JILA experiment [11], we take values close to that regime, since  $u_{11} u_{22}/u_{12}^2 \simeq 0.9999$ . It is also remarkable that, since  $u_{11} > u_{12} > u_{22}$ , we are in a regime in which the first component chases the second one, which rejects mixing with the chaser. Thus a desired configuration has little overlap between both species. This will have remarkable consequences for the unstable configurations.

*Search of solutions.*- We are interested in stationary configurations in which each component has a well defined value of the angular momentum. They are of the form

$$\psi_i(r, \theta) = e^{-i\mu_i t} e^{im_i \theta} \phi_i(r). \quad (4)$$

where  $\phi_i(r)$  is the ground state for vorticity  $m_i$ . These functions satisfy a nonlinear set of coupled PDE

$$\mu_i \psi_i = -\frac{1}{2} \nabla^2 \psi_i + \frac{1}{2} r^2 \psi_i + \sum_j u_{ij} |\psi_j|^2 \psi_i, \quad (5)$$

with  $i = 1, 2$ . Our focus will be on three particular cases,  $(m_1, m_2) = (0, 0), (1, 0), (0, 1)$ , which are the three states with the lowest energy and correspond to the ground state and single vortex states for  $|1\rangle$  and  $|2\rangle$  respectively.

To find the radial dependence of the wave functions,  $\phi_i(r)$ , we expand them approximately on a finite subset of the harmonic oscillator basis,

$$\phi_i(r) \simeq \sum_{n=0}^{n=N} c_n P_n^{(m_i)}(r) e^{-r^2/2}. \quad (6)$$

Next we use a steepest descent technique combined with a Newton method to search the ground states for each of the  $(m_1, m_2)$  families of configurations. The details of this method can be seen in Ref. [9]. As a result one gets the eigenfunctions and energies [Fig. 2] of each family




FIG. 2. Energies of the three relevant states in dimensionless units. (a) Energy of the ground state  $(m_1, m_2) = (0, 0)$  (b) Energies of the  $(1, 0)$  (solid) and  $(0, 1)$  (dashed) vortices as a function of the interaction strength.

*Linear stability theory.*- The main question we will study in what follows is the stability of the previously found states under small perturbations (e.g. in the initial data, small amounts of noise, etc.). Let us define the excitations through

$$\psi_i(r, \theta) = e^{-i\mu_i t + im_i \theta} (\phi_i(r) + e^{-i\lambda t + in\theta} \alpha_i(r)), \quad (7a)$$

$$\bar{\psi}_i(r, \theta) = e^{i\mu_i t - im_i \theta} (\bar{\phi}_i(r) + e^{-i\lambda t - in\theta} \beta_i(r)). \quad (7b)$$

It is easy to show that Eqs. (7) correspond to the right combination of modes for the kind of coupling of (1). Then we introduce Eqs. (7) into Eq. (1) and keep, in the linear approximation, the  $\mathcal{O}(\alpha)$  and  $\mathcal{O}(\beta)$  terms. Finally we obtain the following type of equation

$$J\partial_t\vec{W} = H\vec{W}. \quad (8)$$

Here  $\vec{W}$  is a 4-vector,  $H$  is an hermitian operator, and  $J$  is an anti-hermitian operator. We will search a Jordan basis such that

$$\lambda\vec{W} = iJH\vec{W}. \quad (9)$$

The lack of such a Jordan form leads to polynomial instabilities, while the presence of non-real eigenvalues is a signal of the exponentially growing instabilities. All other modes give us the frequencies of the linear response of the system to small perturbations.

This analysis is formally equivalent to the Bogoliubov stability analysis of the states under consideration. However, as it was shown in [9], for our perturbation theory, which is of order  $O(\alpha^2, \beta^2)$  for the energy, the exponentially unstable modes lay along directions of constant energy. In other words, a Bogoliubov expansion cannot account for the instability, since we cannot recover the equations for those modes up from the Hamiltonian –they can only be reached by perturbing directly on the GPE. It is then important to confirm our results for the stability with numerical simulations of the whole system.

*Linear stability results.*– In Fig. 3 we show the results of the preceding diagonalization. As it was expected, all of the eigenvalues for the ground state,  $(m_1, m_2) = (0, 0)$ , are positive numbers, which implies that it is at least a local minimum of our energy functional –indeed, it can easily be shown that it is an absolute minimum and thus globally stable.

Next we have the  $(1, 0)$  family, which has a single negative eigenvalue plus an infinite set of positive ones. This shows that there exists a path in the configuration space along which the energy decreases. The energetically favorable direction belongs to a  $m = 0$  perturbation on the first component which corresponds to the vortex walking out of the condensate [8]. This means that under energy dissipation mechanisms the vortex could be unstable. But as there exist no complex eigenvalues in this case we conclude that the lifetime of the configuration is only limited by dissipation, i.e. that under non-dissipative perturbations the vortex should remain stable. This is clear when we perform real time evolution of an initially perturbed condensate with this initial configuration as shown in Fig. 1(a) and also consistent with the experiments of Ref. [11]. It is also remarkable that the fact that there is a negative eigenvalue in the linearized spectrum of the  $(1, 0)$  vortex contradicts the belief that the second component could act as a pinning potential, since the configuration under study is neither an absolute nor a local minimum of the energy functional.

Finally we focus on the  $(0, 1)$  family. Here we find an infinite number of modes with positive energy, plus a pair of them with complex eigenvalues. Qualitatively, these unstable modes are similar to those of the  $(1, 0)$ . The difference is that, due to the imaginary part of the eigenvalue, which is of order  $\text{Im}(\lambda) = 0.04$ , they exhibit a self sustained growth that breaks the system. In a few words: single charged vortices in  $|2\rangle$  are unstable under a generic perturbation of initial data. This coincides with the JILA experiments in which the vortex in  $|2\rangle$  was found to be unstable.




FIG. 3. Linearized spectra around: (a) The ground state  $(0,0)$ . (b) A vortex in component  $|1\rangle$ ,  $(m_1, m_2) = (1, 0)$  and (c) A vortex in component  $|2\rangle$ , with  $(m_1, m_2) = (0, 1)$

*Does the vortex break?.*– The linear stability analysis cannot be used to make any definitive conclusions on the behavior of the vortex out of the perturbative limit. To get further information we have performed a set of numerical simulations in which the  $(0, 1)$  state was slightly perturbed by displacing very little the first component with respect to the second one.

The outcome of these simulations is shown in Fig. 1(b),(c). There we see a first stage in which the first component, which resembles a ball, and the vortex oscillate synchronously (the hole in  $|2\rangle$  pins the peak of  $|1\rangle$ ). These oscillations grow in amplitude until the linked system spirals out up to a point at which they form a sort of ying-yang which rotates clockwise, one chasing the other. Finally the first component develops a tail and later a hole which traps the second component. That

hole is a vortex, which somehow has been transferred from  $|2\rangle$  to  $|1\rangle$ . Though it is not completely periodic, this mechanism does exhibit some recurrence and the vortex returns to  $|2\rangle$ .

This phenomenon does not contradict any topological rule or conservation law, in fact the angular momentum of each component is no longer a conserved quantity, and the topological charge of each specie needs not survive through the evolution. Instead, what it is conserved is the total angular momentum

$$L_z = i\hbar \int \bar{\psi}_1 \partial_\theta \psi_1 + i\hbar \int \bar{\psi}_2 \partial_\theta \psi_2 = L_z^{(1)} + L_z^{(2)}. \quad (10)$$

In Fig. 4 the evolution of the total angular momentum and that of each component  $L_z^{(j)}$  is plotted with time. It is seen how there is a complex interchange of angular momentum between both components. The intermediate states are topologically nontrivial states since the phase singularity is being transferred from one component to the other. A more detailed analysis of this process will be presented elsewhere.




FIG. 4. Angular momentum evolution: Solid line corresponds to the evolution of the total angular momentum  $L_z$ , which is well conserved by our numerical scheme, the dotted line corresponds to the evolution of  $L_z^{(1)}$  and the dashed line corresponds to the evolution of  $L_z^{(2)}$

*Conclusions and discussion.*- We have analyzed the stability of vortices in multi-component atomic Bose-Einstein condensates. Our results agree well with the experimental results of [11] proving that a vortex in a  $|1\rangle$  is a dynamically stable object although it is not a global minimum of the energy. In contrast, a vortex in  $|2\rangle$  state is dynamically and energetically unstable and tends to spiral out of the condensate. Since this process is purely dynamic and does not involve transfer of particles to the thermal component one cannot get rid of the vortex angular momentum and this leads to a complex dynamics in which the vortex is transferred to the  $|1\rangle$  state and goes back recurrently to  $|2\rangle$ . In fact if the instability found in the experiments where due to dissipation through a mechanism similar to that proposed in Ref. [15] it would affect both the  $(1,0)$  and  $(0,1)$  type of vortices in a similar way, which is not the case. In support to our theory we can see from Fig. 3 of Ref. [11] that the vortex does not completely escape but

some kind of defect is formed in the periphery of the  $|2\rangle$  component, a behavior similar to that found in our real time simulations [Fig 2(c)].

This work has been partially supported by the DGICYT under grant PB96-0534.

- 
- [1] P. G. Saffman, "Vortex dynamics", Cambridge University Press (1997).
  - [2] Y. Kivshar, B. Luther-Davis, Phys. Rep. **298**, 81 (1998)
  - [3] L. P. Pitaevskii, Sov. Phys. JETP **13**, 451 (1961).
  - [4] M. H. Anderson *et al.*, Science **269**, 198 (1995); C. C. Bradley *et al.*, Phys. Rev. Lett. **75**, 1687 (1995); K. B. Davis *et al.*, Phys. Rev. Lett. **75**, 3969 (1995).
  - [5] R. Dum, J. I. Cirac, M. Lewenstein, P. Zoller, Phys. Rev. Lett. **80** (1998) 2973; K.-P. Marzlin, W. Zhang, Phys. Rev. Lett. **79**, 4728 (1997); K.-P. Marzlin, W. Zhang, Phys. Rev. A **57** (1998) 3801; Caradoc-Davies, R. Ballagh, K. Burnett, Phys. Rev. Lett. **83** (1999) 895; Nakanahra *et al.*, cond-mat/9905374; Jackson, McCann, Adams, Phys. Rev. Lett. **80** (1998) 3903; Petrosyan, You, Phys. Rev. A **59** (1999) 3903.
  - [6] F. Zambelli, S. Stringari, Phys. Rev. Lett. **81** (1998) 1754; Lundhl, C. Pethick, Smith, Phys. Rev. A **58** (1998) 4816; Goldstein, E. Wright, Meystre, Phys. Rev. A **58** (1998) 576; E. L. Bolda, D. Walls, Phys. Rev. Lett. **81** (1998) 5477; F. Dalfovo, M. Modugno, cond-mat/9907102.
  - [7] R. J. Dodd, K. Burnett, M. Edwards, and C. W. Clark, Phys. Rev. A **56** (1997) 587; T. Isochima, K. Machida, Jour. Phys. Soc. Jpn. **66** (1997) 3502; A. A. Svidzinsky, A. L. Fetter, Phys. Rev. A **58** (1998) 3168; A. L. Fetter, J. Low Temp. Phys. **113**, 189 (1998); H. Pu *et al.*, **59** (1999) 1533; T. Isochima, K. Machida, Phys. Rev. A **59** (1999) 2203.
  - [8] D. S. Rokhsar, Phys. Rev. Lett. **79** (1997) 2164.
  - [9] J. J. García-Ripoll, V. M. Pérez-García, Phys. Rev. A (to appear) (cond-mat/9903353).
  - [10] D. S. Hall, M. R. Matthews, J. R. Ensher, C. E. Wieman, and E. A. Cornell, Phys. Rev. Lett. **81** (1998) 1539.
  - [11] M. R. Matthews, B. P. Anderson, P. C. Haljan, C. E. Wiemann, E. A. Cornell, Phys. Rev. Lett. **83** (1999) 2498.
  - [12] J. Williams and M. Holland, Nature (London) (to be published), cond-mat/9909163.
  - [13] J. J. García-Ripoll, J. I. Cirac, J. Anglin, V. M. Pérez-García, P. Zoller, Phys. Rev. A (to appear), cond-mat/9811340
  - [14] V. M. Pérez-García, H. Michinel, and H. Herrero, Phys. Rev. A **47**, 4114 (1998).
  - [15] P. O. Fedichev, G. V. Schlyapnikov, Phys. Rev. Lett. **60** R1779 (1999).