# Bianchi VIII Empty Futures

John D. Barrow and Yves Gaspar DAMTP Centre for Mathematical Sciences Cambridge University Wilberforce Road, Cambridge CB3 0WA UK

November 1, 2018

#### Abstract

Using a qualitative analysis based on the Hamiltonian formalism and the orthonormal frame representation we investigate whether the chaotic behaviour which occurs close to the initial singularity is still present in the far future of Bianchi VIII models. We describe some features of the vacuum Bianchi VIII models at late times which might be relevant for studying the nature of the future asymptote of the general vacuum inhomogeneous solution to the Einstein field equations.

## 1 Introduction

Many studies have been made of the asymptotic states of spatially homogeneous cosmological models. Studies of the early-time behaviour have focussed upon the possibility of chaotic behaviour in the most general cases [2], [4], [5], while investigations of the late-time asymptotics have focussed upon the relative generality of cosmological models which evolve to become at least as isotropic as the observed universe [1]. In this paper we are going to consider some aspects of the late-time evolution of one of the four most general classes of spatially homogeneous universe, see [3], [6], [7], [8], and [18] for related studies. The Bianchi type VIII universe has not attracted the same interest from cosmologists as some other homogeneous models because it does not contain an isotropic Friedmann universe as a special case. However, it can become arbitrarily close to isotropy. It is primarily of interest because of its status as one of the most general homogeneous universes and hence in some sense, yet to be made precise, may offer a leading order approximation to some part of the general cosmological solution of Einstein's equations at late times, far from the initial singularity. In fact, if space is topologically compact it becomes the most general of the ever-expanding Bianchi type universes [9].

In what follows we will study certain aspects of the late-time behaviour of vacuum Bianchi type VIII models. In particular, we will try to determine whether the chaotic Mixmaster behaviour which takes place close to the initial singularity still takes place in the far future as  $t \to +\infty$ . We will show how a qualitative analysis based on the Hamiltonian formalism can reveal some basic features of the future asymptote of vacuum Bianchi VIII models. These provide a basis for studying the nature of the late-time behaviour of the general solution to the vacuum Einstein field equations.

It is also of interest to note that late-time cosmological behaviour in general relativity corresponds to early times in the so-called Pre Big Bang scenario suggested by symmetries of superstring theory. It has been argued [10] that plane-wave space-times are the generic initial states for these models. So in this context the study of the asymptotic future behaviour of general vacuum space-times is also of interest. Also, recent work [11] has pointed out that inhomogeneous models of low-energy bosonic field equations for all superstring models and M-theory (D = 11 supergravity) exhibit an oscillatory approach to the initial singularity, so it would be interesting as well to study how these oscillatory singularities transform in type VIII universes at late times using a similar qualitative analysis, see also [12].

We will first introduce the field equations for the vacuum Bianchi type VIII model and then discuss the orthonormal frame approach and the Hamiltonian formalism to analyse the possible future asymptotic states.

### 2 The field equations for Bianchi type VIII

In the metric approach, where the metric components are the basic variables of the gravitational field, we can introduce group invariant and timeindependent frame vectors  $e_a$  such that the line element is given by

$$ds^2 = -dt^2 + g_{ab}W^aW^b$$

where  $W^a$  are time-independent 1-forms dual to the frame vectors  $e_a$ . In the case of vacuum Bianchi type VIII we can introduce three time-dependent expansion scale factors as

$$g_{ab} = diag(a^2, b^2, c^2)$$

and Einstein's field equations become [6]

$$\frac{(\dot{a}bc)}{abc} = \frac{1}{2a^2b^2c^2}[(b^2+c^2)^2-a^4]$$
(1)

$$\frac{(a\dot{b}c)}{abc} = \frac{1}{2a^2b^2c^2}[(a^2+c^2)^2-b^4]$$
(2)

$$\frac{(ab\dot{c})}{abc} = \frac{1}{2a^2b^2c^2}[(a^2 - b^2)^2 - c^4]$$
(3)

$$\frac{\ddot{a}}{a} + \frac{\ddot{b}}{b} + \frac{\ddot{c}}{c} = 0 \tag{4}$$

Here, the dot denotes differentiation with respect to the synchronous time t. If we replace the synchronous time t by a variable  $\eta$  defined by

$$dt = (abc) d\eta$$

then the field equations (1)-(4) become

$$2(\ln a)'' = (b^2 + c^2)^2 - a^4$$
(5)

$$2(\ln b)'' = (a^2 + c^2)^2 - b^4 \tag{6}$$

$$2(\ln c)'' = (a^2 - b^2)^2 - c^4 \tag{7}$$

$$(\ln a^2)'' + (\ln b^2)'' + (\ln c^2)'' = (\ln a^2)'(\ln b^2)' + (\ln b^2)'(\ln c^2)' + (\ln c^2)'(\ln a^2)'$$
(8)

where ' denotes differentiation with respect to  $\eta$ .

These equations are invariant under the transformation  $\eta \to -\eta$ . Now, as is widely exploited in the literature, [4], [2], to a good approximation we have  $abc \sim t$  as  $t \to 0$ , which implies

$$\eta \propto \ln(t)$$

and the invariance of the field equations (5)-(8) under  $\eta \to -\eta$  corresponds to invariance under

$$t \to \frac{1}{t}$$

in (1)-(4): an apparent duality transformation relating early and late-time behaviour. If this symmetry were exact it would imply that the presence of chaotic behaviour at early times  $(t \to 0)$  means that there must also be chaotic behaviour as  $t \to +\infty$ , as suggested in ref. [3]. A crucial point in the above argument is the accuracy of the assumption that

$$abc \sim t$$

One can use the field equations (1)-(3) to test this assumption, by looking at the second derivative of the metric volume element *abc*. Taking the sum of the equations (1)-(3) we get

$$2(abc)(abc)^{..} = (a^2 - b^2)^2 + 2a^2c^2 + 2c^2b^2 + c^4.$$
(9)

Since the right-hand side is the sum of four positive terms, if  $abc \neq 0$  we have

$$(abc)^{\cdot\cdot} > 0 \tag{10}$$

This shows that the relation  $abc \propto t$  cannot be exact and so the duality symmetry argument cannot be applied to relate late and early time behaviour for the vacuum Bianchi type VIII model. However, the inequality (10) will be useful in deducing the late-time behaviour as  $t \to +\infty$ . There is another consequence of (10) worth pointing out : if abc is a power law

$$abc \sim t^n$$

then (10) implies n > 1. Since  $dt = (abc)d\eta$ , we would have

$$\eta \propto \frac{1}{1-n} t^{1-n}, \qquad n \neq 1$$

Note that near n = 1 there is a structural instability with respect to the variation of t with respect to  $\eta$ . When n > 1 we have  $t \to 0$  as  $\eta \to -\infty$  and  $t \to +\infty$  when  $\eta \to 0$ . However, this differs form the case where n = 1, where  $\eta \propto \ln(t)$  (which is a good approximation close to the initial singularity) since here  $\eta \to +\infty$  as  $t \to +\infty$ . If n < 1, then we would have  $\eta \to 0$  as  $t \to 0$  and  $\eta \to +\infty$  as  $t \to +\infty$ . Tiny variations on either side of n = 1 would make  $\eta$  a bad choice of variable for an early-time analysis of Mixmaster behaviour. This may be of importance for numerical studies of Mixmaster behaviour [13], which are notoriously difficult to carry out because of the slow time variation of the oscillations compared with the growth in volume.

Let  $u^a$  be the 4-velocity of the Hubble flow, the Hubble parameter is defined by

$$H = \frac{1}{3}u^a{}_{;a}$$

This parameter can be expressed as follows

$$H = \frac{\dot{l}}{l}$$

where l(t) corresponds to a geometric-mean expansion scale factor. The deceleration parameter q is defined by

$$q = -\frac{l\ddot{l}}{\dot{l}^2}$$

The Hubble parameter then satisfies the following equation

$$\frac{dH}{dt} = -(1+q)H^2 \tag{11}$$

Defining the a dimensionless shear parameter by

$$\Sigma^2 = \frac{\sigma^2}{3H^2}$$

with  $\sigma^2 = \frac{1}{2}\sigma_{ab}\sigma^{ab}$ , where  $\sigma_{ab}$  is the shear tensor, we have the following simple relation between the shear parameter and the deceleration parameter in vacuum:

$$q = 2\Sigma^2. \tag{12}$$

Equations (11) and (12) can now be used to express the Hubble parameter as

$$H(t) = \frac{1}{t + 2\int \Sigma^2(t)dt}$$
(13)

For the isotropic ( $\Sigma = 0$ ) Milne universe we have  $H(t) = t^{-1}$  so equation (13) expresses the departure from the Milne expansion rate for vacuum homogeneous cosmologies.

We specialise to the Bianchi VIII universe and calculate the Hubble scalar defined above for  $u^a = (1, 0, 0, 0)$  using the Christoffel symbols for Bianchi type VIII models [14], so:

$$H(t) = \frac{1}{3} \frac{d(abc)}{dt} \frac{1}{abc}$$
(14)

Equation (13) then gives the following expression for abc:

$$abc = \exp\left(\int \frac{3}{t + 2s(t)} dt\right) \tag{15}$$

with  $s(t) = \int \Sigma^2 dt$ 

From the above equation we can get (abc), as

$$(abc)^{..} = \exp\left(\int \frac{3}{t+2s(t)} dt\right) \frac{1}{\left(t+2s(t)\right)^2} [9-3(1+2\Sigma^2)]$$
(16)

For this to agree with the inequality (10) we require

$$9 - 3(1 + 2\Sigma^2) > 0$$

which implies

$$\Sigma^2 < 1 \tag{17}$$

As we will see this is consistent with bounds on  $\Sigma$  given by equation (28) of the next section.

### 3 The orthonormal frame formalism

We use the orthonormal frame formalism of Ellis and MacCallum [15]. The orthonormal frame  $e_a$  is chosen to be invariant under the action of the isometry group with  $e_0 = u$ , so that the commutation functions  $\gamma^c{}_{ab}$  defined by

$$[e_a, e_b] = \gamma^c{}_{ab}e_c$$

depend on the synchronous time t and are the basic variables of the gravitational field. With this choice of frame, the line-element takes the form

$$ds^2 = -dt^2 + \delta_{ab}\omega^a(t)\omega^b(t),$$

where  $\omega^a$  are time dependent 1-forms dual to the vector fields  $e_a$ .

The physical state of the models can be described using the components of the shear tensor  $\sigma_{ab}$  and the curvature anisotropies  $n_{ab}$  which occur in the expression of the commutation functions, see Wainwright and Ellis [16] and [17], [18]. In the above frame we define

$$\sigma_{ab} = diag(\sigma_{11}, \sigma_{22}, \sigma_{33})$$

and

$$n_{ab} = diag(n_1, n_2, n_3)$$

and the linear combinations

$$\sigma_{+} = \frac{1}{2}(\sigma_{22} + \sigma_{33})$$
$$\sigma_{-} = \frac{1}{2\sqrt{3}}(\sigma_{22} - \sigma_{33})$$
$$n_{+} = \frac{1}{2}(n_{2} + n_{3})$$
$$n_{-} = \frac{1}{2\sqrt{3}}(n_{2} - n_{3})$$

Now scale out the overall expansion of the universe by introducing the expansion-normalised variables

$$\Sigma_{\pm} = \frac{\sigma_{\pm}}{H} \tag{18}$$

$$N_a = \frac{n_a}{H} \tag{19}$$

The cosmological model is then fully described by the dimensionless variables  $(\Sigma_+, \Sigma_-, N_1, N_2, N_3)$ . The variables  $\Sigma_{\pm}$  describe the anisotropy of the Hubble flow and the  $N_a$  describe the spatial curvature of the group orbits. If we define a dimensionless curvature parameter by

$$K = -\frac{R^{(3)}}{6H^2}$$

then we can deduce from the generalised Friedmann equation the relation

$$1 = \Sigma^2 + K \tag{20}$$

Note that the curvature parameter K is related to the variables  $N_a$  by

$$K = \frac{1}{12} [N_1^2 + N_2^2 + N_3^2 - 2(N_1N_2 + N_2N_3 + N_3N_1)]$$

We can introduce a dimensionless time variable  $\tau$  defined by

$$\frac{dt}{d\tau} = \frac{1}{H} \tag{21}$$

and related to the mean expansion scale factor, l(t), by

$$l = l_0 e^{\tau} \tag{22}$$

These variables yield the following simplified evolution equations for the vacuum type VIII state variables :

$$\Sigma_{\pm}' = -(2-q)\Sigma_{\pm} - S_{\pm}$$
(23)

$$N_1' = (q - 4\Sigma_+)N_1 \tag{24}$$

$$N_2' = (q + 2\Sigma_+ + 2\sqrt{3}\Sigma_-)N_2 \tag{25}$$

$$N_{3}' = (q + 2\Sigma_{+} - 2\sqrt{3}\Sigma_{-})N_{3}$$
(26)

where ' denotes derivatives with respect to  $\tau$  and the functions  $S_{\pm}$  are given by :

$$S_{+} = \frac{1}{6} [(N_{2} - N_{3})^{2} - N_{1}(2N_{1} - N_{2} - N_{3})]$$
$$S_{-} = \frac{1}{2\sqrt{3}} (N_{2} - N_{3})(N_{1} - N_{2} - N_{3})$$

This system, together with equation (20), is an autonomous system of first-order ordinary differential equations for which a monotone function can be found with respect to the time variable  $\tau$ . Different Bianchi types lead to different restrictions on the ranges of the variables  $N_a$ . For Bianchi type VIII we require  $N_1 < 0$ ,  $N_2 > 0$ ,  $N_3 > 0$  and the set defined by these constraints is an invariant set for the evolution equations. On this invariant Bianchi type VIII set there exists the following monotone increasing function :

$$Z = (N_1 N_2 N_3)^2 \tag{27}$$

As a consequence of eq. (20) the following bounds can placed on the state variables [16]:

$$0 \le \Sigma^2 \le 1 \tag{28}$$

$$-1 \le \Sigma_{\pm} \le +1 \tag{29}$$

$$0 \le N_1^2 \le 12$$
 (30)

$$0 \le N_1 (N_2 - N_3) \le 12 \tag{31}$$

$$0 \le -2N_1(N_2 + N_3) \le 12 \tag{32}$$

$$0 \le (N_2 - N_3)^2 \le 12 \tag{33}$$

Therefore, the invariant set S for Bianchi type VIII is unbounded in such way that  $N_2 + N_3$  can assume arbitrarily large values. Together with the existence of the monotone increasing function Z, given by equation (27), this means we do not expect Bianchi type VIII orbits to approach an equilibrium point as  $\tau \to +\infty$ .

### 4 Hamiltonian Cosmology

As discussed in section 2, in the metric approach, we can introduce a set of group-invariant and time-independent one-forms  $W^a$ , so that with an arbitrary time variable t' the metric has the form

$$ds^{2} = -N(t')dt'^{2} + g_{ab}(t')W^{a}W^{b}$$
(34)

In vacuum, the Bianchi type VIII metric is diagonal and we write the metric components in the form [4]

$$g_{ab} = diag(e^{2\beta_1}, e^{2\beta_2}, e^{2\beta_3}) \tag{35}$$

with

$$\beta_1 = \beta^0 - 2\beta^+$$
$$\beta_2 = \beta^0 + \beta^+ + \sqrt{3}\beta^-$$
$$\beta_3 = \beta^0 + \beta^+ - \sqrt{3}\beta^-$$

In the Hamiltonian approach the spatial Ricci curvature defines a potential function  $V(\beta_+, \beta_-)$ ,

$$V(\beta_{+},\beta_{-}) = -e^{-2\beta^{0}}R^{(3)}$$

The variable  $\beta^0$  is related to the overall length scale for the metric (34) and to the Hubble parameter by the relations

$$l = l_0 e^{\beta^0} \tag{36}$$

$$H = N^{-1} \dot{\beta^0} \tag{37}$$

The evolution of the Bianchi type VIII universe is then described by the motion of a point in the mini-superspace  $(\beta^0, \beta^+, \beta^-)$  evolving in the potential  $V(\beta^+, \beta^-)$ . The variable  $\beta^0$  can be chosen to be the time variable so that  $\dot{\beta}^0 = 1$  and  $N = H^{-1}$ . Equations (22) and (36) imply that we have  $\beta^0 = \tau$ .

This Hamiltonian approach allows us to study the early and late-time asymptotic behaviour of Class A Bianchi-type universes qualitatively. For vacuum Bianchi type I, the potential function vanishes,  $V(\beta^{\pm}) = 0$ , so that the evolution is a straight line in  $\beta^{\pm}$ -space. For Bianchi type VIII the equipotential curves of  $V(\beta^{\pm})$  are approximately triangular in shape, [14], with an infinite open channel which starts at the vertex on the  $\beta^+$  axis and becomes increasingly narrow as  $\beta_+ \to +\infty$ . As  $\tau \to +\infty$  (evolution away from the initial singularity) the walls of the triangle contract. If the universe point moves initially in a straight line (corresponding to Kasner-like behaviour) it will bounce off a potential wall and then move along another straight line (a transition to another Kasner-like behaviour). The triangular shape of the potential allows the evolution to be chaotic, which corresponds to the Mixmaster behaviour and is familiar from studies of the evolution near the singularity ( $\tau \to -\infty$ ). But, what happens as  $\tau \to +\infty$ ? The Hamiltonian picture implies that the triangular potential contracts, so this would enable the universe point to bounce off the walls in a chaotic sequence, ad infinitum. We shall argue however from the inequality (10), or the bound (28), and the fact that  $N_2 + N_3$  can take arbitrarily large values, that this chaotic future evolution cannot occur as  $\tau \to +\infty$ , contrary to the claim of Halpern, [3].

In order to show this, we introduce two further equations, relating the  $\Sigma_{\pm}$  to the  $\beta^{\pm}$  variables. Since [16]

$$\Sigma_{\pm} = \frac{d\beta^{\pm}}{d\tau},\tag{38}$$

where  $\Sigma_{\pm}$  coincides with expression calculated in the orthonormal frame formalism, using equation (12), we have the relation

$$\frac{q}{2} = \Sigma^2 = {\Sigma_+}^2 + {\Sigma_-}^2 = \left(\frac{d\beta^+}{d\tau}\right)^2 + \left(\frac{d\beta^-}{d\tau}\right)^2 \tag{39}$$

The Bianchi type VIII model exhibits Mixmaster behaviour towards the past and this chaotic evolution is reflected in the infinite sequence of Kasner era's and bounces off the triangular potential walls in the Hamiltonian picture [2], [4], [14]. When bounces occur, the function

$$\left(\frac{d\beta^+}{d\tau}\right)^2 + \left(\frac{d\beta^-}{d\tau}\right)^2 \tag{40}$$

exhibits a series of peaks (a bounce produces an abrupt change in  $\beta^+$  or  $\beta^-$ ). By equation (39) this implies that  $\Sigma^2$  should exhibit a series of peaks as well. A graph of the function  $\Sigma$  is given in Wainwright and Ellis [16]. Now, if the triangular potential contracts as  $\tau \to +\infty$ , the interval between bounces decreases as well. This means that time derivatives of  $\beta_{\pm}$  could diverge (the peaks of  $\beta_{\pm}(\tau)$  would become increasingly narrow) and so the function (40) would be unbounded. This would imply, by equation (39), that  $\Sigma^2$  would be unbounded, which contradicts the inequality (17) or (28). In this picture of evolution in an eternally contracting potential, one can imagine two solutions which avoid this problem.

First, it could be that  $\beta_{\pm}(\tau)$  is of the form

$$\beta_{\pm}(\tau) = e^{-a\tau} osc(be^{a\tau}) + c,$$

with a, b, c constants. In the above the osc function could have a periodic or chaotic oscillatory behaviour. Clearly this could give increasingly rapid chaotic oscillations of decreasing amplitude and using equation (38) one finds a bounded, eternally chaotic, oscillating  $\Sigma_{\pm}$ . However, defining the function  $p_0(\tau)$  by

$$\frac{dp_0}{d\tau} = (2-q)p_0$$

one can show the following relations in the Hamiltonian approach [16]

$$\tilde{N}_a = \frac{12e^{2\beta_a}}{(-p_0)} \tag{41}$$

where  $\tilde{N}_a = n_a N_a$ , (no sum) with  $n_1 = C^{1}_{23}$ ,  $n_2 = C^{2}_{31}$  and  $n_3 = C^{3}_{12}$ . Using equations (41), (35) and (12), one finds,

$$|N_1 N_2 N_3| \sim \exp(6\tau - 6 \int (1 - \Sigma^2) d\tau)$$
 (42)

We see that if  $\Sigma_{\pm}$  oscillates eternally and satisfies (17) or (28), then (42) implies that  $|N_1N_2N_3|$  increases as  $\tau \to +\infty$  (see end of section 3). Now, equation (30) implies that  $N_1$  is bounded. Since  $\beta_{\pm}$  tend to constant values, the expression for  $\beta_1$  in (35) and equation (41) with a = 1 implies that

$$\frac{e^{2\beta_0}}{(-p_0)}$$

is bounded, and this means that equations (35) and (41) with a = 2 and a = 3 will imply that  $N_2$  and  $N_3$  are each bounded. Hence the Bianchi type *VIII* invariant set would be bounded and this clearly contradicts the fact that  $N_2 + N_3$  diverges as  $\tau \to +\infty$  because of equation (42).

Second, the universe point could slow down, but then we could end up reaching a point where  $\beta_{\pm} \rightarrow \text{constant}$  and  $\Sigma^2 \rightarrow 0$ . One can easily see that in this case equation (23) will imply that equation (20) will not be satisfied. Indeed,  $\Sigma_{-} = 0$  will imply that  $N_2 = N_3$  for Bianchi VIII , so equation (23) requires  $\Sigma_{-}' = 0$ . Equations (24),..., (26), (41) and (35) imply that the  $N_a$  become constants, but then (23) requires  $S_+ = 0$ , otherwise  $\Sigma_+$  does not satisfy the bound (17) or (28). In order to have  $S_+ = 0$  we must satisfy either  $N_1 = 0$  or  $N_1 = N_2 = N_3$ . The first case implies that the curvature parameter K = 0, and this is inconsistent with the Friedmann equation (20), and the second possibility implies that  $N_1 = N_2 = N_3 = 0$  because of the sign restrictions mentioned in the previous section for Bianchi type VIII. This also leads to K = 0 and contradicts (20) as well. Note that this argument is not necessarily true in the non-vacuum situation (see appendix).

So what are the alternative possibilities consistent with the Hamiltonian picture? The infinitely long channel along the  $\beta^+$  axis offers a solution. The universe point could evolve eternally between the two walls of the infinite channel, for instance there could be an infinity of oscillations between the two walls of the channel. This evolution would correspond to a non-chaotic oscillation of the universe point about the axisymmetric solution (see below) characterised by

$$\beta^- = 0$$

In this case, because the channel becomes increasingly narrow as  $\beta_+ \rightarrow +\infty$ , we expect  $\beta_-$  to undergo increasingly rapid oscillations of decreasing amplitude as  $\tau \rightarrow +\infty$ . Similar behaviour of the universe point when interacting with a channel of the Bianchi IX potential can be found in Ryan and Shepley [14] for a so-called 'mixing bounce'. Equation (42) also implies that  $N_2 + N_3 \rightarrow +\infty$  as  $\tau \rightarrow +\infty$  and so the orbits in the state space escape to infinity.

Non-chaotic oscillatory behaviour for Bianchi type VIII seems also to be observed in the behaviour of the Weyl curvature of other Bianchi type universes according to Coley [19] and in Wainwright et al. [20]. In this interesting work the late-time behaviour of Bianchi type  $VII_0$  models with perfect fluid source was investigated, (see also refs. [1], [6], [22]. If the universe contains matter with pressure p and density  $\rho$ , obeying an equation of state

$$p = (\gamma - 1)\rho$$

with  $\gamma$  constant, then if  $\gamma < \frac{4}{3}$  the universe isotropizes with respect to shear but not with respect to the Weyl curvature. So even if the shear decreases at late times, the anisotropic Weyl curvature dominates the dynamics, producing a form of self-similarity breaking at late times. If  $\frac{4}{3} < \gamma \leq 2$ , then  $\Sigma^2$  is oscillatory as  $\tau \to +\infty$ , with  $\beta_-$  a decreasing oscillatory function, while the Weyl scalar diverges. It was further argued that since an open subset of a more general class of models (i.e. Bianchi type *VIII*) will display features of Bianchi *VII*<sub>0</sub> models, the non-self-similar behaviour of the Weyl curvature could occur at some stage during the evolution of Bianchi *VIII* models.

Let us introduce the dimensionless electric and magnetic parts of the Weyl curvature

$$\tilde{E}_{ab} = E_{ab} \frac{1}{H^2}$$
$$\tilde{H}_{ab} = H_{ab} \frac{1}{H^2}$$

For all Bianchi class A models, in the standard frame  $\tilde{E}_{ab}$  and  $\tilde{H}_{ab}$  are diagonal and we define:

$$\tilde{E}_{+} = \frac{1}{2}(\tilde{E}_{22} + \tilde{E}_{33}) \tag{43}$$

$$\tilde{E}_{-} = \frac{1}{2\sqrt{3}} (\tilde{E}_{22} - \tilde{E}_{33}) \tag{44}$$

and likewise for  $\tilde{H}_+$  and  $\tilde{H}_-$ . By defining

$$N_{+} = \frac{1}{2}(N_{2} + N_{3}) \tag{45}$$

$$N_{-} = \frac{1}{2}(N_2 - N_3). \tag{46}$$

We have the following useful relations between the state variables introduced in the orthonormal frame approach and the electric and magnetic parts of the Weyl curvature :

$$\tilde{E}_{+} = \Sigma_{+} + (\Sigma_{+}^{2} - \Sigma_{-}^{2}) + S_{+}$$
(47)

$$\tilde{E}_{-} = \Sigma_{-} - 2\Sigma_{+}\Sigma_{-} + S_{-} \tag{48}$$

$$\tilde{H}_{+} = -\frac{3}{2}N_{1}\Sigma_{+} - 3N_{-}\Sigma_{-}$$
(49)

$$\tilde{H}_{-} = -3N_{-}\Sigma_{+} + \frac{1}{2}(N_{1} - 4N_{+})\Sigma_{-}.$$
(50)

If  $N_2 + N_3$  becomes arbitrarily large, then by equation (50), we can have

$$\lim_{\tau \to +\infty} \tilde{H}_{-} = +\infty \tag{51}$$

and the Weyl scalar, given by

$$W = \tilde{E}_{+}^{2} + \tilde{E}_{-}^{2} + \tilde{H}_{+}^{2} + \tilde{H}_{-}^{2},$$

can become arbitrarily large as  $\tau \to +\infty$ , see Wainwright [7] for a discussion of the self-similarity breaking. Similar behaviour for the Weyl curvature was found in perfect fluid Bianchi type  $VII_0$  universes with equation of state parameter satisfying  $1 < \gamma \leq 2$  [20].

# 5 The axisymmetric (LRS) vacuum Bianchi type VIII model

Our qualitative analysis shows that the late-time behaviour of vacuum Bianchi type VIII models will be non-chaotic. Evolution could occur which exhibits oscillations about the axisymmetric (LRS) vacuum Bianchi type VIII model with  $\Sigma_{-} = 0$ . The asymptotic late-time behaviour of this solution is known [21]. The Bianchi type VIII LRS model is described by the following restrictions on the state variables, and these define an invariant set for the evolution equations (23)-(26) :

$$N_1 < 0$$

$$N_2 = N_3 > 0$$
$$\Sigma_- = 0$$

This state space is unbounded, since  $N_2$  can take arbitrarily large values. However, we can use the following bounds on  $N_1N_2$  (see inequalities (28)-(33)):

$$0 \le -N_1 N_2 \le 3$$

We introduce a new variable Z defined by Wainwright, [21],

$$Z^2 = -N_1 N_2$$

It can then be shown that the system of evolution equations (23)-(26) admits the following equilibrium point :

$$N_1 = 0$$
$$\Sigma_+ = \frac{1}{2}$$
$$Z = \frac{3}{2}$$

This corresponds to a flat plane-wave asymptote, which has W = 0. More precisely, it is the Bianchi type *III* form of flat space-time, with metric

$$ds^{2} = -dt^{2} + t^{2}(dx^{2} + e^{2x}dy^{2}) + dz^{2}.$$
(52)

The Bianchi type  $VI_h$  diagonal plane-wave metric with h = -1 reduces to this line element for a particular choice of parameters.

Since the above equilibrium point solution, (52), is of LRS Bianchi type III with  $n^{\alpha}{}_{\alpha} = 0$ , it is a particular case of the following general metric which includes Bianchi types  $I, VI_0, VIII, V$  and  $VI_h$  with  $n^{\alpha}{}_{\alpha} = 0$  [15]

$$ds^{2} = -dt^{2} + (X^{2} + Y^{2}w^{2})dx^{2} - 2Y^{2}we^{-(a_{0}+q_{0})x}dxdy + Y^{2}e^{-2(a_{0}+q_{0})x}dy^{2} + Z^{2}e^{2(a_{0}-q_{0})x}dz^{2}$$
(53)

where  $a_0$  and  $q_0$  constants and X(t), Y(t), Z(t), w(t) are functions of the synchronous time t, and

$$w(t) = -2b \int XY^{-3}Z^{-1}dt$$

where b is constant. The line element (52) has,

$$b = 0, X = Y = t, Z = 1, a_0 = q_0 = -\frac{1}{2}.$$

Now since this line element is a future attractor for the LRS vacuum Bianchi type VIII, and since (53) contains non-LRS solutions, it is interesting to perform a stability analysis of this line element for non-LRS perturbation modes. The perturbation technique for special cases of (53) has already been developed by Barrow and Sonoda [22]. In this approach one defines a new time variable as

$$\frac{d}{du} = X(t)\frac{d}{dt} = '$$

as well as the new variables  $F = \frac{X'}{X}$ ,  $G = \frac{Y'}{Y}$  and  $H = \frac{Z'}{Z}$  so that the field equations for (53) in vacuum become

$$F' = 2(a_0^2 + q_0^2) - \frac{2b^2 X^2}{Y^4 Z^2} - F(G + H),$$
(54)

$$G' = 2(a_0^2 + a_0 q_0) + \frac{2b^2 X^2}{Y^4 Z^2} - G(G + H),$$
(55)

$$H' = 2(a_0^2 - a_0 q_0) - H(G + H),$$
(56)

with the first integral constraints

$$\frac{2b^2 X^2}{Y^4 Z^2} = 2F(G+H) + 2GH - (6a_0^2 + 2q_0^2)$$
(57)

and

$$2Fa_0 = q_0(G - H) + G + H.$$
 (58)

The unperturbed solution is the line element (52) with  $F_0 = 1$ ,  $G_0 = 1$ and  $H_0 = 0$ . We require equations for the following small perturbations, f, g, h:

$$F = F_0 + f,$$
  

$$G = G_0 + g,$$
  

$$H = H_0 + h.$$

Using the equations (54)-(58), one gets the following (the equation for F is eliminated using the constraint (58)):

$$g' = -4g - 4h \tag{59}$$

$$h' = -h \tag{60}$$

The eigenvalues associated with this problem are  $\lambda_1 = -4$  and  $\lambda_2 = -1$  and so the solution (52) is asymptotically stable under small non-LRS perturbations consistent with (53), thus supporting the prediction that the universe point will evolve about the axisymmetric solution with  $\beta_- = 0$  in the narrow infinite channel along the  $\beta_+$ -axis. Taking into account the results of the previous section, this means we expect the future asymptotic state of the full non-LRS vacuum Bianchi type *VIII* to have

$$\Sigma = \frac{1}{2}.$$

The fact that  $\Sigma_{-}$  vanishes is a consequence of the following : since  $\beta_{-} \to 0$ , the line element (34) becomes axisymmetric and so the field equations (5), (6) and (7) lead to the LRS solution for the scale factors, thus  $\beta_{+} \to \frac{\tau}{2}$ , however from equation (41) one finds that if  $\Sigma_{-}$  oscillates and does not decrease, then the curvature parameter K can be unbounded, which contradicts the generalised Friedmann equation (20). Also, although  $\Sigma_{-}$  decreases, it can be shown to undergo increasingly rapid oscillations : if  $\beta_{-}$  is bounded, oscillatory and decreasing, then equations (23), (38) and (41) imply that  $\beta_{-}$  ( and thus  $\Sigma_{-}$  ) undergoes increasingly rapid oscillations. If the universe point oscillates about the axisymmetric solution, then the metric shares a feature of some gravitational-wave space-times, namely that the solution oscillates about a given form of flat space-time as  $\tau \to +\infty$ .

### 6 Conclusion

A qualitative analysis based on the Hamiltonian formalism and the orthonormal frame approach gives the following information about the future asymptotic state of vacuum Bianchi type VIII models :

a. The chaotic Mixmaster solution close to the initial singularity evolves at late times into a simpler non-chaotic solution, contrary to the claims of ref. [3], and corresponding to the fact that the universe point evolves along the infinite open channel of the Bianchi VIII potential and oscillates about the LRS Bianchi type VIII solution in such a way that the shear parameter exhibits increasingly rapid oscillations as  $\tau \to +\infty$ .

b. The form of the line element (34) tends to the Bianchi III form of flat space-time because  $\beta_{-} \rightarrow 0$ . It is interesting to compare this with asymptotic states of other vacuum Bianchi types : Bianchi types IV,  $VI_h$ , and  $VII_h$  tend to plane-wave space-times; Bianchi type  $VII_0$  tends to the special equilibrium point  $N_1 = 0$ ,  $N_2 = N_3 = \text{constant}$ , with  $\Sigma_+ = -1$  and  $\Sigma_- = 0$ , which in the state space is a line emanating from a point corresponding to the Taub form of flat space; finally, Bianchi type V tends to the Milne form of flat space. So it appears that different forms of flat and plane-wave space-times play an important role in the future state of these eternally expanding vacuum Bianchi models and one might expect that they occur in a leading-order approximation to part of the general inhomogeneous vacuum solution to the Einstein field equations at late times.

### 7 Acknowledgements

We would like to thank John Wainwright for sending us helpful information about the late-time behaviour of LRS Bianchi type *VIII* models.

### 8 Appendix

For a non-vacuum homogenous cosmology the Friedmann equation (20) becomes

$$1 = \Sigma^2 + K + \Omega \tag{61}$$

where  $\Omega$  is the density parameter defined by  $(8\pi G = 1)$ 

$$\Omega = \frac{\rho}{3H^2}$$

with  $\rho$  the energy density of a perfect fluid with equation of state  $p = (\gamma - 1)\rho$ . It appears that if  $0 < \gamma \leq \frac{2}{3}$ , the LRS Bianchi *VIII* models admits the flat Friedmann model as future attractor i.e.  $\Sigma = N_1 = N_2 = N_3 = 0$  and  $\Omega = 1$ , so that the argument developed in section 4 does not hold. This is simply the well known cosmic no hair attractor of power-law inflationary universes whose matter content violates the strong energy condition. In fact, for non-vacuum Bianchi type *VIII*, the analysis of section 4 can be adapted as follows when the strong energy condition holds. Note first that the deceleration parameter is now given by

$$q = 2\Sigma^2 + \frac{1}{2}(3\gamma - 2)\Omega$$

and so we have  $q \ge 0$  if  $3\gamma - 2 \ge 0$ . For all  $\gamma$  satisfying this inequality, one can apply the analysis of section 4 and the universe point will again evolve in the infinite channel as  $\tau \to +\infty$ .

We end this appendix by a remark about the stiff fluid case with  $\gamma = 2$ . This case is of interest namely because the stiff fluid is equivalent to the presence of a massless scalar field. The evolution equation for  $\Omega$  is

$$\frac{d\Omega}{d\tau} = [2q - (3\gamma - 2)]\Omega \tag{62}$$

Now when  $\gamma = 2$ , the field equations (1),..., (3) are identical for vacuum and non-vacuum cases. So, the inequality (17) is valid and this corresponds to

q < 2

and so equation (62) implies that  $\Omega \to 0$  as  $\tau \to +\infty$ , so that the evolution tends to a vacuum solution.

## References

- [1] Collins C.B. and Hawking S.W., 1973 Astrophys. J. 180, 317
- [2] Belinskii V.A., Khalatnikov, I.M. and Lifshitz E.M., 1970 Adv. Phys. 19, 525

- [3] Halpern P. 1987 Gen. Rel. Grav. 19, 73
- [4] Misner C.W., 1969 Phys. Rev. Lett. 22, 1071
- [5] Barrow, J.D. 1982 Phys. Reports 85,1
- [6] Doroshkevich A.G., Lukash V.N., Novikov I.D. 1973 Zh. Eksp. Teor. Fiz. 64, 1457
- [7] Wainwright J. 2000 Gen. Rel. Grav. 32, 1041
- [8] Uggla, C., Jantzen, R.T., Rosquist, K. and von Zur-Mühlen, H., 1991 Gen. Rel. Grav. 23, 947
- [9] J.D. Barrow and H. Kodama, 2001 paper in preparation
- [10] Kunze K.E. 1999 Class. Quantum Grav. 16, 3795-3806
- [11] Damour T., Henneaux M., hep-th/0006171
- [12] Barrow, J.D. and M. Dabrowski, 1998 Phys. Rev. D57, 1
- [13] Hobill D., Burd A., and Coley, A.A., (eds.) 1994 Deterministic Chaos in General Relativity, Plenum, NY
- [14] Ryan M.P. and Shepley L.C. 1975 Homogeneous relativistic cosmologies, Princeton U P, Princeton NJ
- [15] Ellis G.F.R. and MacCallum M.A.H. 1969 Comm. Math. Phys. 12, 108
- [16] Wainwright J. and Ellis G.F.R. (eds.) 1997 Dynamical Systems in Cosmology, Cambridge U P, Cambridge
- [17] Bogoyavlensky O.I. 1985 Methods in the qualitative theory of dynamical systems in astrophysics and gas dynamics, Berlin, Springer-Verlag
- [18] Bogoyavlensky O.I. and Novikov S.P. 1982 Sel. Math. Sov. 2, 159
- [19] Coley A., gr-qc/9812010
- [20] Wainwright J., Hancock M.J., Uggla C. 1999, Class. Quantum Grav. 16, 2577

- [21] J. Wainwright, private communication.
- $\left[22\right]$ Barrow J.D., Sonoda D.H. 1986 Phys. Rep. 139, 1