New approach to calculating the News

Nigel T. Bishop & Shrirang S. Deshingkar Department of Mathematics, Applied Mathematics & Astronomy, University of South Africa, P.O. Box 392, Unisa 0003, South Africa

We consider the problem of finding the gravitational radiation output, or news, within the context of a numerical simulation of a spacetime by means of the null-cone, or characteristic, approach to numerical relativity. We develop a method for computing the news that uses an explicit coordinate transformation to the Bondi gauge in a neighborhood of \mathcal{I}^+ . The method has been implemented computationally. We present results of applying the method to certain test problems, demonstrating second order convergence of the news to the analytic value.

PACS numbers: 04.25.Dm, 04.30.-w, 95.85.Sz

I. INTRODUCTION

In many simulations in numerical relativity, an important objective is to compute the emitted gravitational radiation. Within ADM-type approaches, one uses computed data near the outer boundary of the simulation to estimate the radiation at future null infinity (\mathcal{I}^+) in the form of an expansion of spherical harmonics. The underlying theory was developed some time ago [1, 2, 3], and the computational implementation is discussed, for example, in [4, 5, 6]. However, this paper uses the characteristic, or null-cone, approach with Bondi-Sachs coordinates, in which \mathcal{I}^+ is included in the compactified grid, and therefore the gravitational radiation, or news, can in principle be found exactly [7].

Even so, the computation of the news is not a straightforward matter. The news takes a very simple form, $N = \frac{1}{2}J_{,\ell u}$ (the meaning of these symbols will be given later), when the null-cone coordinate system satisfies the Bondi gauge conditions. However, these are gauge conditions at \mathcal{I}^+ , and in any physical simulation the gauge freedoms in the null-cone coordinates are set in the interior of the spacetime. In general, a computed metric will not satisfy the Bondi gauge conditions. Thus, a procedure for computing the news in a general Bondi-Sachs coordinate system has been developed [8].

A motivation for this work is that some difficulties with the news computation of [8] have been reported [9]. The cause of the difficulties is not known, and it may be that they can be resolved by technical improvements to the method and code [10], but as yet no such results have been reported. Our method has also not been tested for the problem of [9], and so it is not known whether it will be of assistance.

However, our primary motivation is that the method presented here is both technically and conceptually much simpler than the previous method; and also it contains a means to monitor the accuracy of results obtained.

We construct a coordinate transformation between the general gauge and a gauge that satisfies the Bondi conditions. The coordinate transformation on \mathcal{I}^+ is known [8], but constructing the transformation off \mathcal{I}^+ is more difficult. We do so by means of a Taylor series approach, the

point being that in order to find the news $N = \frac{1}{2}J_{,\ell u}$, we need to know the transformed metric only to first order in ℓ . Even so, the algebraic calculations are quite lengthy and require the use of computer algebra.

Our news calculation has been implemented computationally, and used in problems in which an analytic result is known. Results are presented below, and it is seen that the errors are second-order convergent to zero.

The paper starts with a summary of relevant results and notation for the characteristic formulation of numerical relativity (Sec. II). The coordinate transformation and computer algebra results are given in Sec. III, and then the procedure for computing the news – at both analytic and computational levels – is described in Sec. IV. The computational tests and results are presented in Sec. V. The Conclusion, Sec. VI, gives a comparison between the method in this paper and that of [8]; and also discusses possibilities for further work.

II. NOTATION

The formalism for the numerical evolution of Einstein's equations, in null cone coordinates, is well known [7, 8] (see also [11, 12, 13, 14]). For the sake of completeness, we give a summary of those aspects of the formalism that will be used here. We start with coordinates based upon a family of outgoing null hypersurfaces. We let u label these hypersurfaces, x^A (A=2,3), label the null rays and r be a surface area coordinate. In the resulting $x^{\alpha}=(u,r,x^A)$ coordinates, the metric takes the Bondi-Sachs form [14, 15]

$$\begin{split} ds^2 &= -\left(e^{2\beta}(1+\frac{W}{r}) - r^2h_{AB}U^AU^B\right)du^2\\ -2e^{2\beta}dudr - 2r^2h_{AB}U^Bdudx^A + r^2h_{AB}dx^Adx^B, \quad (1) \end{split}$$

where $h^{AB}h_{BC} = \delta_C^A$ and $det(h_{AB}) = det(q_{AB})$, with q_{AB} a unit sphere metric. We work in stereographic coordinates $x^A = (q, p)$ for which the unit sphere metric is

$$q_{AB}dx^{A}dx^{B} = \frac{4}{P^{2}}(dq^{2} + dp^{2}),$$
 (2)

where

$$P = 1 + q^2 + p^2. (3)$$

We also introduce a complex dyad q_A defined by

$$q^A = \frac{P}{2}(1,i), \quad q_A = \frac{2}{P}(1,i)$$
 (4)

with $i = \sqrt{-1}$. For an arbitrary Bondi-Sachs metric, h_{AB} can then be represented by its dyad component

$$J = h_{AB}q^A q^B / 2, (5)$$

with the spherically symmetric case characterized by J = 0. The full nonlinear h_{AB} is uniquely determined by J, since the determinant condition implies that the remaining dyad component

$$K = h_{AB} q^A \bar{q}^B / 2 \tag{6}$$

satisfies $1 = K^2 - J\bar{J}$. We introduce the spin-weighted field

$$U = U^A q_A, \tag{7}$$

as well as the (complex differential) eth operators \eth and $\bar{\eth}$ (see [16] for full details).

The news calculation is performed in conformally compactified coordinates. Specifically, $(u, r, x^A) \rightarrow (u, \ell, x^A)$ where $\ell = 1/r$. In (u, ℓ, x^A) coordinates, the compactified metric is $d\hat{s}^2 = \ell^2 ds^2$ where ℓ is a conformal factor with future null infinity \mathcal{I}^+ given by $\ell = 0$. The compactified metric will be denoted by $\hat{g}^{\alpha\beta}$, and the general compactified Bondi-Sachs metric is

$$\hat{g}^{11} = e^{-2\beta} V_a, \quad \hat{g}^{1A} = e^{-2\beta} U^A,$$

$$\hat{g}^{10} = e^{-2\beta}, \quad \hat{g}^{AB} = h^{AB}, \quad \hat{g}^{0A} = \hat{g}^{00} = 0, \tag{8}$$

where $V_a = \ell^2(1 + \ell W)$. In addition to the compactified Bondi-Sachs metric in a general gauge Eq. (8), we will refer to the compactified Bondi-Sachs metric in the Bondi gauge, and such quantities will be denoted with a suffix $_{[B]}$. The Bondi gauge compactified metric and coordinates are $\hat{g}_{[B]}^{\alpha\beta}$ and $(u_{[B]}, \ell_{[B]}, x_{[B]}^A)$, respectively. On \mathcal{I}^+ , i.e. $\ell_{[B]} = 0$, $\hat{g}_{[B]}^{\alpha\beta}$ satisfies the Bondi gauge conditions

$$\hat{g}_{[B]}^{11} = 0, \ \hat{g}_{[B]}^{1A} = 0, \ \hat{g}_{[B]}^{01} = 1, \ \hat{g}_{[B]}^{AB} = q_{[B]}^{AB}, \ \ (9)$$

where $q_{[B]}^{AB}$ is a unit sphere metric with respect to the Bondi angular coordinates $x_{[B]}^{A}$.

III. COORDINATE TRANSFORMATION

We define a coordinate transformation, near \mathcal{I}^+ , between (u, ℓ, x^A) and $(u_{[B]}, \ell_{[B]}, x_{[B]}^A)$:

$$u_{[B]} = u_{[B]0} + A^{u}\ell + C^{u}\ell^{2}, \quad \ell_{[B]} = \omega\ell + C^{\ell}\ell^{2},$$

$$x_{[B]}^{A} = x_{[B]0}^{A} + A^{A}\ell + C^{A}\ell^{2}, \quad (10)$$

where ω , $x_{[B]0}^A$, A^A , $u_{[B]0}$, A^u are all functions of x^A and u only. It will turn out that the C^{α} are irrelevant to the news calculation, but they are included so that the Jacobian (constructed by differentiating the above) is manifestly correct to first order in ℓ . We also introduce complex quantities

$$A = q_A A^A \text{ and } X = q_A x_{[B]0}^A,$$
 (11)

both of which are defined to have spin weight 1. The general metric quantities are expanded as a Taylor series to first order in ℓ , with the $_0$ quantities being the values at $\ell=0$, i.e. on \mathcal{I}^+ , and the $_\ell$ quantities being the first ℓ derivative evaluated at $\ell=0$, i.e. $(\partial_\ell)_{\ell=0}$.

$$\beta = \beta_0 + \ell \beta_\ell, \ U = U_0 + \ell U_\ell, \ J = J_0 + \ell J_\ell,$$

$$K = K_0 + \ell K_\ell, \ V_a = \ell V_{a\ell}.$$
 (12)

The same notation will be used for the metric quantities in the Bondi gauge; these are related to quantities in the general gauge by

$$\hat{g}_{[B]0}^{\alpha\beta} + \ell \hat{g}_{[B]\ell}^{\alpha\beta} = \hat{g}_{[B]}^{\alpha\beta} = \omega^{-2} \frac{\partial x_{[B]}^{\alpha}}{\partial x^{\mu}} \frac{\partial x_{[B]}^{\beta}}{\partial x^{\nu}} \hat{g}^{\mu\nu}, \tag{13}$$

with the factor ω^{-2} appearing because there is also an implicit change of compactification factor from ℓ to $\ell_{[B]}$.

We have used Maple to find $\hat{g}_{[B]}^{\alpha\beta}$ to first order in ℓ , and in doing so we have imposed the conditions (c.f. Eqs. (39) and (40) in [8]).

$$(\partial_u + U_0^B \partial_B) x_{[B]0}^A = 0 (14)$$

and

$$(\partial_u + U_0^B \partial_B) u_{[B]0} = \omega e^{2\beta_0}. \tag{15}$$

We found that the Bondi gauge conditions

$$\hat{g}_{[B]0}^{11} = 0, \ \hat{g}_{[B]0}^{1A} = 0, \ \hat{g}_{[B]0}^{01} = 1$$
 (16)

are satisfied identically. The remaining Bondi gauge condition is

$$\hat{g}_{[B]0}^{AB} = q_{[B]}^{AB}. \tag{17}$$

Taking the determinant of Eq. (17) leads to an explicit expression for ω

$$\omega = \frac{P}{P_{[B]}} \sqrt{|q_{[B],q} p_{[B],p} - q_{[B],p} p_{[B],q}|}, \tag{18}$$

where $P_{[B]} = 1 + q_{[B]}^2 + p_{[B]}^2$. The remaining content of Eq. (17) can be written as

$$J_{[B]0} = -\hat{g}_{[B]0}^{AB} q_{A[B]} q_{B[B]} / 2 = 0, \tag{19}$$

where $q_{A[B]}$ is the complex dyad appropriate to the metric $q_{[B]}^{AB}$. The Maple calculation finds

$$J_{[B]0} = 4P^{2} \Big[4J_{0}X^{2}\bar{\zeta}^{2} - 2K_{0}\eth X\bar{\eth}X + 4J_{0}X\bar{\eth}X\bar{\zeta}$$
$$-4K_{0}X\bar{\eth}X\bar{\zeta} + \bar{J}_{0}(\bar{\eth}X)^{2} + J_{0}(\bar{\eth}X)^{2} \Big]$$
$$/[(4 + P^{2}X\bar{X})^{2}\omega^{2}] = 0, \tag{20}$$

where $\bar{\zeta} = q - ip$. Eq. (20) is not used to determine any of the transformation parameters, but rather as a check on the accuracy of the computation, as described in Sec. IV.

In addition to satisfying the Bondi gauge conditions, $\hat{g}_{[B]}^{\alpha\beta}$ is also a Bondi-Sachs metric and must satisfy the conditions

$$\hat{g}_{[B]}^{00} = 0, \ \hat{g}_{[B]}^{0A} = 0.$$
 (21)

Now, $\hat{g}^{00}_{[B]0}=0$ and $\hat{g}^{00}_{[B]\ell}=0$ can be expressed as linear equations in A^u and C^u respectively, but these values will not be needed. Also, $\hat{g}^{0A}_{[B]0}=0$ and $\hat{g}^{0A}_{[B]\ell}=0$ can be expressed as linear equations in A^A and C^A respectively: the expression for C^A will not be needed, but the expression for A^A will be used in the news calculation and, in complex form,

$$A = [J_0 \bar{\eth} u_{[B]} (2X\bar{\zeta} + \bar{\eth}X) + \bar{J}_0 \bar{\eth} u_{[B]} \bar{\eth}X -K_0 (\bar{\eth} u_{[B]} (2X\bar{\zeta} + \bar{\eth}X) + \bar{\eth} u_{[B]} \bar{\eth}X)]/(2\omega).$$
(22)

The computer algebra calculation evaluates the inverse Jacobian at $\ell=0$, from which we find a result that will be used in the news calculation

$$\frac{\partial}{\partial u_{[B]}}(u,\ell,x^A) = \frac{e^{-2\beta_0}}{\omega}(1,0,U_0^A).$$
 (23)

The computer algebra yields an expression for $J_{[B]\ell}$: Defining

$$T_1 = J_0 \bar{\eth} X - K_0 \bar{\eth} X,$$

$$T_2 = J_0 (\bar{\eth} X)^2 + \bar{\eth} X (\bar{J}_0 \bar{\eth} X - 2K_0 \bar{\eth} X),$$
(24)

then

$$J_{[B]\ell} = P^{2}(4(1 - P_{[B]})AT_{2} + 2XP_{[B]}(J_{\ell}(\bar{\eth}X)^{2} + \bar{J}_{\ell}(\bar{\eth}X)^{2} + 2(\bar{\eth}AT_{1} + 2\bar{\zeta}XJ_{0}\bar{\eth}A + \bar{\eth}A(\bar{J}_{0}\bar{\eth}X - K_{0}\bar{\eth}X) - 2\bar{\zeta}XK_{0}\bar{\eth}A - K_{\ell}\bar{\eth}X(2\bar{\zeta}X + \bar{\eth}X)) + 4\bar{\zeta}J_{\ell}X(\bar{\zeta}X + \bar{\eth}X)) - 4AX(-4\bar{\zeta}^{2}J_{0}X + P_{[B]}(e^{-2\beta_{0}}(2\bar{\zeta}U_{\ell}X + (U_{\ell}\bar{\eth}X + \bar{U}_{\ell}\bar{\eth}X) + 2A_{,u} + AV_{a\ell} + (U_{0}\bar{\eth}A + \bar{U}_{0}\bar{\eth}A) + 2\bar{\zeta}AU_{0}) + 2\bar{\zeta}T_{1})) - P^{2}X^{2}\bar{A}(4\bar{\zeta}X(\bar{\zeta}J_{0}X + T_{1}) + T_{2}) + 16\bar{\zeta}AXT_{1})/(8XP_{[B]}^{3}\omega^{2}).$$
 (25)

IV. PROCEDURE FOR COMPUTING THE NEWS

We can now describe the procedure used for calculating the news. First, we look at the process from an analytic viewpoint, and then we give some details of the computational implementation. As discussed in Sec. I, we assume that the Bondi-Sachs metric, in the general gauge, is known throughout the spacetime.

A. Analytic procedure

- 1. Solve Eq. (14) to find $x_{[B]0}^A=x_{[B]0}^A(u,x^A)$.
- 2. Eq. (18) gives $\omega = \omega(u, x^A)$.
- 3. Use Eq. (20) to find by how much $J_{[B]0}$ differs from zero, and thus obtain an estimate of the accuracy of the calculation.
- 4. Solve Eq. (15) to find $u_{[B]0} = u_{[B]0}(u, x^A)$.
- 5. Eq. (21) gives $A = A(u, x^A)$.

6. We note that

$$\left(\frac{\partial J_{[B]}}{\partial \ell_{[B]}}\right)_{\ell_{[B]}=0} = \frac{\partial x^{\mu}}{\partial \ell_{[B]}} \frac{\partial J_{[B]}}{\partial x^{\mu}}
= \frac{\partial \ell}{\partial \ell_{[B]}} \frac{\partial J_{[B]}}{\partial \ell} = \frac{J_{[B]\ell}}{\omega},$$
(26)

because $J_{[B]}$ is zero on \mathcal{I}^+ , and so derivatives in directions other than ℓ vanish. Then the news $N = N(u, x^A)$ is

$$N = \frac{1}{2} \frac{\partial^{2} J_{[B]}}{\partial \ell_{[B]} \partial u_{[B]}} = \left(\frac{\partial}{\partial u_{[B]}}\right) \frac{J_{[B]\ell}}{2\omega}$$
$$= \left(\frac{\partial u}{\partial u_{[B]}} \frac{\partial}{\partial u} + \frac{\partial x^{A}}{\partial u_{[B]}} \frac{\partial}{\partial x^{A}}\right) \frac{J_{[B]\ell}}{2\omega}. \tag{27}$$

The terms $\partial u/\partial u_{[B]}$ etc. are evaluated at $\ell=0$ and were given in Eq. (23) above.

7. Finally, we need to express the news as a function of Bondi coordinates, i.e. to transform $N(u, x^A)$ to $N(u_{[B]}, x_{[B]}^A)$.

B. Computational implementation

The news module has been written to interface directly with the null gravity code in its current form [17]. Thus

we use a compactified radial coordinate x = r/(R+r), with R = 1. There are n_x points in the x direction in the range [0.5, 1] (corresponding to $1 < r < \infty$). We use stereographic angular coordinates, with the number of grid points in each angular direction n_n .

Instead of solving equations (14) and (15), we convert them to ordinary differential equations as in reference [8], and solve them by means of the second order method

$$(x^{A(n+1)} - x^{A(n)})_i = \Delta u (U^A_0)_i^{(n+\frac{1}{2})}, \tag{28}$$

$$(u_{[B]}^{(n+1)} - u_{[B]}^{(n)})_i = \Delta u \left(\omega e^{2\beta 0}\right)_i^{(n+\frac{1}{2})}.$$
 (29)

At the initial time-slice we match the Bondi coordinates to the Bondi-Sachs coordinates: i.e. at $u=0,\ u_{[B]}=0$ and $x_{[B]}^A=x^A$. This also gives A=0 and $J_{[B]\ell}=J_\ell$ on the initial surface. The solution of Eq. (28) leads to $x^A=x^A(u,x_{[B]}^A)$, and this needs to be inverted to give $x_{[B]}^A=x_{[B]}^A(u,x^A)$. At this stage the inversion has not been implemented in general, because it is not needed in the test examples considered below.

Second derivatives do not appear directly in the formulas leading to the news. First angular (eth) derivatives are implemented using the central difference second order method – with second order accurate upwind/downwind at the boundaries. The first time derivative of A is needed to calculate $J_{[B]\ell}$, and the first time derivative of $J_{[B]\ell}/\omega$ appears in the Bondi news function. For both the quantities, apart from the first two time-steps, we use one sided-downwind second order accurate differentiation scheme. For the initial steps we use the standard first order scheme.

V. COMPUTATIONAL TESTS AND RESULTS

We tested our method for calculating the Bondi news function against two exact solutions: linearized Robinson-Trautman, and Schwarzschild in rotating coordinates. We discretize this analytic solutions on the required computational grid and determine the various metric components and use them as an input for the news module. The news module as such sees it as input from an equivalent numerical evolution code.

A. Linearized Robinson-Trautman solution

The Robinson-Trautman solution represents a distorted black hole emitting purely outgoing radiation. The radiation decays exponentially, and asymptotically the solution becomes Schwarzschild.

The Robinson-Trautman metric can be put in the form

$$ds^{2} = -(\mathcal{K} - \frac{2}{rW})du^{2} - 2Wdudr - 2rW_{,A}dudx^{A} + r^{2}q_{AB}dx^{A}dx^{B}, (30)$$

where $\mathcal{K} = \mathcal{W}^2[1 - L^2(\log \mathcal{W})]$, L^2 is the angular momentum operator (i.e., the angular part of the Laplacian operator on the unit sphere), and $\mathcal{W}(u, x^A)$ has to satisfies the nonlinear differential equation

$$12\partial_u(\log \mathcal{W}) = \mathcal{W}^2 L^2 \mathcal{K}. \tag{31}$$

We recover the Schwarzschild solution by putting W =constant in the above metric.

Linearized solutions to the Robinson-Trautman equation (31) are obtained by setting $W = 1 + \phi$ and dropping nonlinear terms in ϕ ,

$$12\partial_u \phi = L^2 (2 - L^2) \phi. \tag{32}$$

For a spherical harmonic perturbation $\phi = \lambda(u)Y_{\ell m}$ this leads to the exponential decay $\lambda = \lambda(0)e^{-u\,\ell(\ell+1)(\ell^2+\ell-2)/12}$. The (linear) perturbation can be a linear sum of various spherical harmonics with small amplitudes.

We consider a linearized solution of the form:

$$\phi = \Re[(\lambda_{22}e^{-2u}Y_{22} + \lambda_{33}e^{-10u}Y_{33})] \tag{33}$$

with $\lambda_{22}=3\times 10^{-7}$ and $\lambda_{33}=7\times 10^{-7}$. The metric components are, $J=0,\,\beta=0.5\phi,\,U=\eth\phi/r$ (so U=0 at $r=\infty$) and W=1. Giving these quantities as inputs at the grid points at each time level, we call the news module to calculate the Bondi news function. We calculate the L_2 norm $J_{[B]0}$ (which should be zero), as well as the L_2 norm of the error in the Bondi news function, whose analytic value is

$$N(u_B, x^A) = \frac{1}{2} \mathcal{W}^{-1} \eth^2 \mathcal{W}. \tag{34}$$

The computation was performed for the following grid sizes

- (a) $\Delta q = \Delta p = 0.100, \Delta x = 1/60, \Delta u = 0.04$
- (b) $\Delta q = \Delta p = 0.050, \Delta x = 1/120, \Delta u = 0.02$
- (c) $\Delta q = \Delta p = 0.025, \Delta x = 1/240, \Delta u = 0.01$
- (d) $\Delta q = \Delta p = 0.020, \Delta x = 1/300, \Delta u = 0.008(35)$

In this case as $\zeta_{[B]} = \zeta$ we find that $||J_{[B]0}||_2$ constant with u and we plot $||J_{[B]0}||_2$ at u=1.6 for the discretizations (35): the observed convergence rate is 1.99. The L_2 norm of the error in the news is plotted against u for the discretizations (35) in Fig. 2 We see convergence that, at earlier times, is approximately second order, but at later times saturation is observed. The convergence rate between different grids as u increases is given in Table (I). Further, Fig. 2 shows that the error for the finest grid 35 (d) saturates at about e^{-24} , the next finest grid 35(c) saturates at $e^{-24.5}$, grid 35(b) saturates at $e^{-26.5}$, while grid 35(a) saturates at $e^{-28.5}$. We observe that the saturation level scales approximately with discretization length as Δ^{-3} . These matters are discussed further in Sec. VI.




FIG. 1: Convergence of $||J_{[B]0}||_2$ at u=1.6 for the linearized Robinson-Trautman case.




FIG. 2: L_2 norm of the error in the news as a function of time u for various discretizations for the linearized Robinson-Trautman case.

The Y_{33} component initially dominates the total news function, which should therefore decay approximately as e^{-10u} i.e. with slope 10 on a log scale; while at later times the Y_{22} component dominates and the news should decay as e^{-2u} . If the percentage of error remains constant in the

convergence rate at u	grids (a) & (b)	(b) & (c)	(c) & (d)
0.4	1.955	1.992	1.996
0.8	1.929	1.982	1.966
1.2	1.928	1.966	1.578
1.6	1.928	1.907	0.4746

TABLE I: Change of convergence rate with u in the L_2 norm of news function for linearized Robinson-Trautman for different grid resolutions in equation 35.

news function, the L_2 norm of the error should also show a similar trend with increasing u. We see from Fig. 1 that this is actually the case. Until saturation effects become evident, a little before u=2, the norm of the error in the news decreases as expected at all grid resolutions.

The error in the news is significantly higher at the first two time-steps, presumably because we are using a first order scheme since there is not enough past data to evaluate second order accurate time derivatives (actually, at the second time-step one can implement a second order scheme, but its error is larger).

B. Schwarzschild in rotating coordinates

By the transformation $\varphi \to \tilde{\varphi} + \kappa u$ of the azimuthal coordinate, the Schwarzschild line element in null coordinates can be written as

$$ds^{2} = -\left(1 - \frac{2m}{r} - \kappa^{2} r^{2} \sin^{2}\theta\right) du^{2} - 2dudr$$
$$+2\kappa r^{2} \sin^{2}\theta du d\varphi + r^{2} q_{AB} dx^{A} dx^{B}. \tag{36}$$

In this case we set $\kappa=4$ This coordinate change gives a nontrivial value for U at \mathcal{I}^+ and thus it becomes a useful test to check that the numerically calculated Bondi news function remains zero. In this case the the values of the various metric components are $\beta = 0$, J = 0, $U = 2I\zeta/(1+\zeta\bar{\zeta}), W = 1.$ Giving these as inputs we calculate $||J_{[B]0}||_2$ and the news function $||N||_2$, both of which have zero as the analytic value. We again found that $||J_{[B]0}||_2$ hardly varies with u, and in Fig. 3 we plot $||J_{B|0}||_2$ at u=1.6 for the discretizations (35): the observed convergence rate is again 1.99. The L_2 norm of the news is plotted against u for the discretizations (35) in Fig. 4. We find convergence to zero at u = 1.6 at a rate 4.96 averaged over the grids (35) (with only minor variations according to the specific grid used). The higher than expected rate is presumably due to some cancellation effect in the truncation errors. It is interesting that the magnitude of the error in the news builds up quite rapidly with time. Since the error remains convergent to zero, this must mean that the truncation error constants are growing with time.




FIG. 3: Convergence of $||J_{[B]0}||_2$ at u=1.6 for the rotating Schwarzschild case.




FIG. 4: L_2 norm of the error in the news as a function of time u for various discretizations for the rotating Schwarzschild case.

VI. CONCLUSIONS

We have presented a method for calculating the news, based upon an explicit coordinate transformation to the Bondi gauge at \mathcal{I}^+ . The method has been implemented

computationally and validated against known analytic solutions. There are similarities and differences between the method and that of [8]

- Both methods require J=0 on \mathcal{I}^+ at u=0.
- Both methods solve evolution equations for $x_{[B]0}^A$ (Eq. (14)) and $u_{[B]0}$ (Eq. (15)).
- The method of [8] solves evolution equations for δ (Eq. ([8]-36)) and ω (Given after Eq. ([8]-31)). In our method, δ is not required, and ω is found explicitly (Eq. (18)).
- The formulas used here are much simpler than those of [8].
- There is a means to monitor the accuracy of results obtained: we simply find $J_{[B]0}$ and compare the value to zero.

The saturation and reduction in convergence rate reported in Sec. VA, is typically an indication that roundoff error becomes significant as the magnitude of the signal decays. The saturation level scales approximately as Δ^{-3} , which corresponds to the finite difference representation of a third derivative. This is consistent with the fact that here the news calculation implicitly involves third derivatives: $x_{[B]0}^A$ and $u_{[B]0}$ are differentiated to obtain A (Eq. 22)); then A is differentiated to obtain $J_{[B]\ell}$ (Eq. (25)); and finally $J_{[B]\ell}$ is differentiated to obtain the news N (Eq. (27)). The observed scaling of saturation as Δ^{-3} is for the specific case of the Robinson-Trautman solution, which has the special property $x_{[B]0}^A = x_0^A$; thus, in a general case the effect may be more severe. For future applications, it is important to be aware that round-off error may be a problem.

It will be interesting to apply our news calculation to the problem considered in [9] (for which the method of [8] was unsuccessful), but that is deferred to further work. Also deferred to further work is the possibility of extending the coordinate transformation to higher order in ℓ , and so being able to extract the Bondi mass and angular momentum aspects – the difficulty will be that the computer algebra is likely to lead to very complicated formulas.

Acknowledgements

We benefited from the hospitality of the Max-Planck-Institut für GravitationsPhysik, Albert-Einstein-Institut and the Center for Gravitational Wave Physics at Pennsylvania State University (funded by the NSF cooperative agreement PHY-0114375). The work was supported by the National Research Foundation, South Africa under Grant number 2053724. We thank Sascha Husa and Jeffrey Winicour for discussions.

- [1] T.Regge and J. Wheeler, Phys. Rev. 108, 1063 (1957).
- [2] F. Zerilli, Phys. Rev. Lett. 24, 737 (1970).
- [3] V. Moncrief, Ann. Phys. (N.Y.) 88, 323 (1974).
- [4] M. Rupright, A. Abrahams, and L. Rezzolla, Phys. Rev. D 48, 044005 (1998).
- [5] K. Camarda and E. Seidel, Phys.Rev. D 59, 064019 (1999).
- [6] E. Seidel and W.-M. Suen, Jnl. Comp. and Appl. Math. 109, 493 (1999).
- [7] N. T. Bishop, R. Gómez, L. Lehner, and J. Winicour, Phys. Rev. D 54, 6153 (1996).
- [8] N. T. Bishop, R. Gómez, L. Lehner, M. Maharaj, and J. Winicour, Phys. Rev. D 56, 6298 (1997).
- [9] N. T. Bishop, R. Gómez, S. Husa, L. Lehner, and J. Winicour, gr-qc/0301060 (2003).

- [10] Y. E. Zlochower, Ph.D. thesis, University of Pittsburgh (2002).
- [11] N. T. Bishop, Class. Quantum Grav. 10, 333 (1993).
- [12] N. T. Bishop, C. J. S. Clarke, and R. A. d'Inverno, Class. Quantum Grav. 7, L23 (1993).
- [13] R. A. Isaacson, J. S. Welling, and J. Winicour, J. Math. Phys. 24, 1824 (1983).
- [14] H. Bondi, M. J. G. van der Burg, and A. W. K. Metzner, Proc. R. Soc. London A269, 21 (1962).
- [15] R. K. Sachs, Proc. R. Soc. London A270, 103 (1962).
- [16] R. Gómez, L. Lehner, P. Papadopoulos, and J. Winicour, Class. Quantum Grav. 14, 977 (1997).
- [17] R. Gómez, Phys. Rev. D 64, 024007 (2001).