

GRAVITATIONAL COLLAPSE OF BAROTROPIC SPHERICAL FLUIDS

ROBERTO GIAMBÒ, FABIO GIANNONI, GIULIO MAGLI, AND PAOLO PICCIONE

ABSTRACT. The gravitational collapse of spherical, barotropic perfect fluids is analyzed here. For the first time, the final state of these systems is characterized without resorting to simplifying assumptions - such as self-similarity - using a new approach based on non-linear o.d.e. techniques. Formation of naked singularities is shown to occur for solutions such that the mass function is sufficiently regular in a neighborhood of the spacetime singularity.

1. INTRODUCTION

The final state of gravitational collapse is an open problem of classical gravity. It is, in fact, commonly believed that a collapsing star that it is unable to radiate away - via e.g. supernova explosion - a sufficient amount of mass to fall below the neutron star limit, will certainly and inevitably form a black hole, so that the singularity corresponding to diverging values of energy and stresses will be safely hidden - at least to faraway observers - by an event horizon. However, this is nothing more than a conjecture - what Roger Penrose first called a "Cosmic Censorship" conjecture [34] - and has never been proved. Actually, it is easy to see that one just cannot prove the conjecture as a statement on the mathematical evolution of *any* collapsing system via Einstein field equations, because in this case what is conjectured is baldly false: it is indeed an easy exercise producing counterexamples using e.g. negative energy densities or "ad hoc" field configurations. Thus, to go beyond the conjecture what is needed is a set of hypotheses, possibly based on sound physical requirements, which would allow the proof of a mathematically rigorous theorem. However, what turned out to be the truth in the last twenty years of research is that such a theorem (and, in fact, even the hypotheses of the theorem) is/are extremely difficult to be stated (see e.g. [23]).

In the meanwhile, many examples of spherically symmetric solutions exhibiting naked singularities and satisfying the principles of physical reasonableness have been discovered.

Spherically symmetric naked singularities can be divided into two groups: those occurring in scalar fields models [6, 8] and those occurring in astrophysical sources modeled with continuous media, which are of exclusive interest here (see [22] for a recent review). The first (shell focusing) examples of naked singularities were discovered in dust models, numerically by Eardley and Smarr [10] and analytically by Christodoulou [5]. Today, the gravitational collapse of dust is known in full details [25].

The dust models can, of course, be strongly criticized from the physical point of view. In fact, they have the obvious drawback that stresses are expected to develop during the collapse, possibly influencing its dynamics. In particular, such models are an unsound description of astrophysical sources in the late stage of the

collapse even if the latter does *not* form a singularity: one can, for instance, regard a white dwarf or a neutron star as being an extremely compact *planet*, composed by a solid crust and a liquid (super)fluid core: such objects are sustained by enormous amounts of (generally anisotropic) stresses. It is, therefore, urgent to understand models of gravitational collapse with stresses.

Recently, several new results have been obtained in this direction by considering systems sustained by anisotropic stresses (see e.g. [16, 18, 20, 21]). Besides the details of the physics of the collapse of such systems, the general pattern arising from all such examples is that existence of naked singularities persists in presence of stresses: actually, we have recently shown that the mechanism responsible for the formation or whatsoever of a naked singularity is *the same* in all such cases [13].

In spite of the aforementioned physical relevance of anisotropic systems, it is beyond any doubt of exceeding interest the case of *isotropic* stresses, i.e. the gravitational collapse of perfect fluids. In fact, for instance, the perfect fluid model is (in part for historical reasons) the preferred model used in most approximations of stellar matter of astrophysical interest. Unfortunately, although *local* existence and uniqueness for the solution of the Einstein field equations has been proved [28, 35], very few sound analytical models of gravitational collapse of perfect fluids are known and, as a consequence, the problem of the final state of gravitational collapse of perfect fluids in General Relativity is still essentially open. Exceptions are the solutions describing shear-free fluids (see e.g. [26, 27]) and those obtained by matching of shock waves [36]; in both cases, however, the collapse is synchronous (i.e. the singularity is of the Friedmann-Robertson-Walker type) and therefore such solutions say little about Cosmic Censorship [3, 24].

There is a unique perfect fluid class of solutions which has been investigated in full details. This is the case of self-similar fluids, which has been treated by many authors since the pioneering work by Ori and Piran [33] (for a recent review see [4]). Self-similarity is compatible with the field equations if the equation of state is of the form $p = \alpha\epsilon$ (where p is the pressure, ϵ the energy density, and α a constant). In this case the field equations reduce to ordinary differential equations and therefore can be analyzed with the powerful techniques of dynamical systems. Ori and Piran found that self similar perfect fluids generically form naked singularities; more precisely, they showed numerically that for any α in a certain range there are solutions with naked singularities. Recently, Harada added some numerical examples which remove the similarity hypotheses [17].

These results clearly go in the direction of disproving *any* kind of censorship at least in spherical symmetry, since they show that naked singularities have to be expected in perfect fluids with physically sound equations of state. However, although being extremely relevant as a "laboratory", the self-similar ansatz is a oversimplifying assumption, and the general case of perfect fluid collapse remained untractable up today, essentially due to the lack of exact solutions.

We are going to circumvent this problem here using a combination of two new ingredients. The first is the fact that, in a suitable system of coordinates (the so-called area-radius coordinates) we are able to reduce the field equations to a single, quasi linear, second order partial differential equation. As a consequence, the metric for a barotropic spherical fluid can be written, in full generality, in terms of only

one unknown function. In this way the behavior of the null radial geodesics near the singular point can be analyzed in terms of the Taylor expansion of such a function. The second ingredient is a new framework for doing this analysis based on techniques for singular non linear ordinary differential equations [13, 14].

Our results show that the endstate of all barotropic perfect fluids solutions describing complete gravitational collapse for which the mass function is sufficiently regular in a neighborhood of the center is a naked singularity.

2. REDUCTION OF THE FIELD EQUATIONS TO A QUASI-LINEAR P.D.E.

Consider a spherically symmetric perfect fluid. The general line element in co-moving coordinates can be written as

$$(2.1) \quad ds^2 = -e^{2\nu} dt^2 + e^{2\lambda} dr^2 + R^2(d\theta^2 + \sin^2 \theta d\phi^2)$$

where ν , λ and R are function of r and t (we shall use a dot and a prime to denote derivatives with respect to t and r respectively). Denoting by ϵ and p the energy density and the isotropic pressure of the fluid, Einstein field equations can be written as

$$(2.2a) \quad \Psi' = 4\pi\epsilon R^2 R',$$

$$(2.2b) \quad \dot{\Psi} = -4\pi p R^2 \dot{R},$$

$$(2.2c) \quad \dot{R}' = \dot{R}\nu' + R'\dot{\lambda},$$

$$(2.2d) \quad p' = -(\epsilon + p)\nu',$$

where $\Psi(r, t)$ is the *Misner-Sharp mass*, defined in such a way that the equation $R = 2\Psi$ spans the boundary of the *trapped region*, i.e. the region in which outgoing null rays re-converge:

$$(2.3) \quad \Psi(r, t) = \frac{R}{2} [1 - g^{\mu\nu}(\partial_\mu R)(\partial_\nu R)] = \frac{R}{2} [1 - (R')^2 e^{-2\lambda} + (\dot{R})^2 e^{-2\nu}],$$

The curve $t_h(r)$ describing this boundary, i.e. the function defined implicitly by

$$(2.4) \quad R(r, t_h(r)) = 2\Psi(r, R(r, t_h(r))),$$

is called *apparent horizon* and will play a fundamental role in what follows.

Initial data for the field equations can be assigned on any Cauchy surface ($t = 0$, say). Physically, the arbitrariness on the data refers to the initial distribution of energy density and the initial velocity profile, and is therefore described by two functions of r only. Data for R do not carry physical information and we parameterize the initial surface in such a way that $R(r, 0) = r$.

The data must be complemented with the information about the physical nature of the collapsing material. In the present paper we shall consider only barotropic perfect fluids, i.e. fluids for which the equation of state can be given in the standard thermodynamical form: the pressure p equals minus the derivative w.r. to the specific volume v of the specific energy density $e(v)$. We are going to work however with the matter density $\rho = 1/v$ and with the energy density $\epsilon(\rho) = \rho e(1/\rho)$. Therefore we are going to use in the sequel the equation of state of the fluid in the

form (slightly less familiar than $p = -de/dv$):

$$(2.5) \quad p = \rho \frac{d\epsilon}{d\rho} - \epsilon$$

Using the comoving description of the fields the matter density is proportional to the determinant of the 3-metric, i.e.

$$(2.6) \quad \rho = \frac{e^{-\lambda}}{4\pi ER^2}$$

where $E = E(r)$ is an arbitrary positive function.

In order to simplify reading, we are going to develop in full details in the next sections the special - although physically very relevant - case of the linear equation of state

$$(2.7) \quad p = \alpha \epsilon,$$

where α is a constant parameter. However, in the final section, we will show how the results can be easily extended to (virtually) *all* the - physically valid - barotropic equations of state.

In terms of the matter density eq. (2.5) implies $\epsilon = \rho^{\alpha+1}$ up to a multiplicative constant which however can be absorbed in the definition of $E(r)$. For such fluids the field equation (2.2d) integrates to

$$(2.8) \quad e^\nu = \rho^{-\alpha}$$

up to a multiplicative function of time only which can be taken equal to one by a reparameterization of t .

We are now going to show that the remaining field equations simplify considerably (and actually the problem of the final state becomes tractable) if another system of coordinates, the area-radius ones, are used. The advantages of this system were first recognized by Ori [32], who used it to obtain the general exact solution for charged dust. Subsequently, the area-radius framework has been successfully applied to models of gravitational collapse and cosmic censorship (see e.g. [13, 18, 30]).

Area-radius coordinates are obtained using R in place of the comoving time. Denoting by subscripts derivatives w.r. to the new coordinates, we have $\Psi' = \Psi_{,r} + R' \Psi_{,R}$, $\dot{\Psi} = \dot{R} \Psi_{,R}$. Substituting in eqs (2.2a), (2.2b) we obtain R' and ρ in terms of the mass:

$$(2.9) \quad R' = -\frac{\alpha}{\alpha + 1} \frac{\Psi_{,r}}{\Psi_{,R}}.$$

$$(2.10) \quad \rho = \left(-\frac{\Psi_{,R}}{4\pi\alpha R^2} \right)^{\frac{1}{\alpha+1}},$$

In writing the above formulae we have excluded the case $\alpha = 0$. This case corresponds to the dust (Tolman-Bondi) solutions which is already very well known and will not be considered further in the present paper (see [25] and references therein).

Equation (2.3) can be used to express the velocity $u = |\dot{R}e^{-\nu}|$ as

$$(2.11) \quad u^2 = \frac{2\Psi}{R} + Y^2 - 1.$$

where we have introduced the function

$$(2.12) \quad Y = R' e^{-\lambda},$$

using (2.8), (2.9) and (2.10) we have

$$(2.13) \quad Y = \frac{E \psi_{,r}}{(\alpha + 1) \rho^\alpha}.$$

This function plays the role of an “effective potential” for the motion of the shells. Notice that u is known when Y and Ψ are; Y is known when $E(r)$ is given and Ψ is known. Thus, in particular, the initial velocity profile $u(r, r)$ is known when the functions $\Psi_0(r) = \Psi(r, r)$ and $Y_0(r) = Y(r, r)$ are known. It is therefore convenient to use Y_0 as the second arbitrary function, eliminating E :

$$(2.14) \quad Y(r, R) = \frac{\Psi_{,r}(r, R)}{\Psi_{,r}(r, r)} \left[\frac{\Psi_{,R}(r, r) R^2}{\Psi_{,R}(r, R) r^2} \right]^{\frac{\alpha}{\alpha+1}} Y_0(r),$$

where (2.10) and (2.13) have been used.

We conclude that the metric for a barotropic perfect fluid in area-radius coordinates can be written in terms of the data and of the function Ψ and its first derivatives as follows:

$$(2.15) \quad ds^2 = -\frac{1}{u^2} \left[dR^2 - 2R' dR dr + \left(\frac{R'}{Y} \right)^2 \left(1 - \frac{2\Psi}{R} \right) dr^2 \right] + R^2 (d\theta^2 + \sin^2 \theta d\phi^2)$$

where u , R' and Y are given by formulae (2.10), (2.9) and (2.14) above. By a tedious but straightforward calculation the remaining field equation can be rearranged as a second order equation for Ψ . Remarkably enough, this equation is quasi-linear. In fact, the following holds true:

Theorem 2.1. *The Einstein field equations for a spherical barotropic fluid in the coordinate system (2.15) are equivalent to the following, second order PDE:*

$$(2.16) \quad a \Psi_{,RR} + 2b \Psi_{,rR} + c \Psi_{,rr} = d,$$

where a, b, c, d are functions of $r, R, \Psi, \Psi_r, \Psi_R$ given by:

$$(2.17a) \quad a = \frac{1}{(\alpha + 1) \Psi_{,R}} \left[1 - \alpha \left(\frac{Y}{u} \right)^2 \right],$$

$$(2.17b) \quad b = \left(\frac{Y}{u} \right)^2 \frac{1}{\Psi_{,r}},$$

$$(2.17c) \quad c = -\frac{(\alpha + 1) \Psi_{,R}}{\alpha \Psi_{,r}^2} \left(\frac{Y}{u} \right)^2,$$

$$(2.17d) \quad d = \frac{1}{R} \left[-\frac{2\alpha}{\alpha+1} \left(1 + \left(\frac{Y}{u} \right)^2 \right) + \frac{\alpha\Psi + R\Psi_{,R}}{\alpha u^2 R} + \frac{(\alpha+1)\Psi_{,R}}{\alpha\Psi_{,r}} \left(\frac{Y'_0}{Y_0} - \frac{1}{\alpha+1} \frac{\Psi''_0}{\Psi'_0} - \frac{2\alpha}{(\alpha+1)r} \right) \left(\frac{Y}{u} \right)^2 R \right].$$

Remark 2.2. Equation (2.16) must be supplemented with a set of data on the surface $R = r$. Since

$$(2.18) \quad ac - b^2 = -\frac{1}{\alpha} \left(\frac{Y}{u\Psi_{,r}} \right)^2,$$

the character of the equation is determined by the sign of α . In particular, the equation is hyperbolic for positive pressures and elliptic for the negative ones (recall that $\alpha = 0$ is excluded). For physical reasons, however, we consider here only the hyperbolic case (see next section). The initial data for equation (2.16) are thus given, in principle, by two functions. The value of Ψ on the data corresponds to the physical freedom of assigning the initial mass distribution, while the first derivative can be calculated using eq. (2.9) evaluated on the data. On $R = r$ one has $R' = 1$ and therefore:

$$(2.19) \quad \Psi_{,R}(r, r) = -\frac{\alpha}{\alpha+1} \Psi_{,r}(r, r).$$

Remark 2.3. Equation (2.16) becomes degenerate at the sonic point, when the relative velocity of the fluid equals the speed of sound. The behavior of the solutions at the sonic point is quite complicated, and not all the solutions can be extended. The problem of characterizing the structure of the space of the solutions is extremely interesting. As far as the present authors are aware, such an analysis has been carried out in full details only in the self-similar case [1, 33, 11]. In the present paper, however, we are interested only in singularities which arise from the gravitational interaction, and therefore we are going to consider only solutions for which the mass function is sufficiently regular in a neighborhood of the spacetime singularity.

3. FORMATION OF SINGULARITIES

3.1. Physical requirements. We are going to impose here strict requirements of physical reasonableness. First of all, we impose the dominant energy condition, namely, energy density must be positive and the modulus of the pressure cannot exceed the energy density (so that $-1 \leq \alpha \leq 1$). We consider, however, only the case of positive pressure. It must, in fact, be taken into account that, while tensions are common in anisotropic materials, a perfect fluid can hardly be considered as physical in presence of a negative isotropic pressure.

Therefore, $\alpha > 0$ and (2.10) imply that

$$(3.1) \quad \Psi_{,R}(r, R) < 0, \quad \forall r > 0, \quad \forall R \in [0, r],$$

and since we want $R' > 0$ to avoid shell-crossing singularities, it must also be, from (2.9),

$$(3.2) \quad \Psi_{,r}(r, R) > 0, \quad \forall r > 0, \quad \forall R \in [0, r].$$

We also require the existence of a regular Cauchy surface ($t = 0$, say) carrying the initial data for the fields. This requirement is fundamental, since it assures that the singularities eventually forming will be a genuine outcome of the dynamics. It is easy to show that, with the equation of state used here, it is equivalent to require the matter density to be finite and non vanishing on the data. Due to eqs. (2.10) and (2.19) we get

$$(3.3) \quad \lim_{r \rightarrow 0^+} \frac{\Psi_{,r}(r, r)}{r^2} \in (0, +\infty).$$

For physical reasonableness, we also require that the initial energy $\epsilon(r, r)$ is a not increasing function.

Moreover, we require regularity of the metric at the center that is, in comoving coordinates:

$$(3.4) \quad R(0, t) = 0, \quad e^{\lambda(0, t)} = R'(0, t),$$

for each $t \geq 0$ up to the time of singularity formation t_0 (see (3.18) below). Using (2.12), we can translate relations (3.4) in area–radius coordinates requiring that $\lim_{r \rightarrow 0^+} Y(r, R(r)) = 1$ for each infinitesimal curve $R(r)$ strictly bounded between 0 and r . In particular

$$(3.5) \quad \lim_{r \rightarrow 0^+} Y(r, r\tau) = 1, \quad \forall \tau \in (0, 1].$$

3.2. Taylor expandability of the mass function. In the present paper we assume Taylor expandability of the mass function at $(0, 0)$. The following holds true:

Proposition 3.1. *The Taylor expansion of the mass function $\Psi(r, R)$ has the following structure*

$$(3.6) \quad \Psi(r, R) = \frac{h}{2} \left(r^3 - \frac{\alpha}{\alpha + 1} R^3 \right) + \sum_{i+j=3+k} \Psi_{ij} r^i R^j + \dots$$

where $k \geq 2$ and h is a positive constant.

Proof. Regularity condition (3.3) and (2.19) imply the Taylor expansion to start from third order terms, except perhaps for a quadratic term which however has to vanish on the data. Therefore, one certainly has

$$(3.7) \quad \Psi(r, R) = \frac{1}{2} m_0 (R - r)^2 + \sum_{i+j=3} \Psi_{ij} r^i R^j + \dots$$

We divide the proof into 3 parts.

1). The quantity m_0 cannot be non-zero.

Let us observe that the function $Y_0(r)$ defined in (2.14) is such that

$$\lim_{r \rightarrow 0^+} Y_0(r) = 1$$

(using (3.5) with $\tau = 1$). If m_0 is non vanishing it is easily seen, from (2.10) and (2.14) respectively, that

$$(3.8) \quad Y(r, r\tau) = c(\tau) \frac{1}{r} + \dots,$$

plus some bounded terms, for some $c(\tau)$ finite and non zero. The above expression is, however, in contradiction with (3.5), and therefore m_0 vanishes.

2. It must be

$$(3.9) \quad \Psi(r, R) = \frac{h}{2} \left(r^3 - \frac{\alpha}{\alpha+1} R^3 \right) + \dots$$

For the sake of convenience we set, for each $n \geq 0$,

$$(3.10) \quad A_n(\tau) = \sum_{i+j=3+n} i \Psi_{ij} \tau^j, \quad B_n(\tau) = \sum_{i+j=3+n} j \Psi_{ij} \tau^{j-1},$$

so that are Taylor expandable and the r^{n+2} 's coefficients of Taylor expansions of $\Psi_{,r}(r, r\tau)$ and $\Psi_{,R}(r, r\tau)$ are $A_n(\tau)$ and $B_n(\tau)$ respectively. We recall that (3.3) implies $A_0(1) > 0$, and, from (2.19), $B_0(1) < 0$ follows. Using (2.14) we get

$$Y(r, r\tau) = \frac{A_0(\tau)}{A_0(1)} \left[\frac{B_0(1)\tau^2}{B_0(\tau)} \right]^{\frac{\alpha}{\alpha+1}} + o(1),$$

at least for each $\tau \in (0, 1]$ such that $B_0(\tau) \neq 0$ (but this polynomial can possibly vanish only for two values of τ), and then (3.5) holds if

$$(3.11) \quad \frac{A_0(\tau)}{A_0(1)} \left[\frac{B_0(1)\tau^2}{B_0(\tau)} \right]^{\frac{\alpha}{\alpha+1}} = 1, \quad \forall \tau \in (0, 1] \text{ with } B_0(\tau) \neq 0.$$

But

$$\frac{B_0(1)\tau^2}{B_0(\tau)} = \tau^2 \frac{\Psi_{12} + 2\Psi_{21} + 3\Psi_{03}}{\Psi_{21} + 2\Psi_{12}\tau + 3\Psi_{03}\tau^2},$$

and therefore if Ψ_{21} was not vanishing, the above quantity would tend to zero as $\tau \rightarrow 0$, which is in contradiction with (3.11). Then $\Psi_{21} = 0$. A similar argument applies to Ψ_{12} to show that this quantity is zero as well. Finally, relation (2.19) imposes a constraint on $A_n(1)$ and $B_n(1)$:

$$(3.12) \quad -\alpha A_n(1) = (\alpha + 1) B_n(1), \quad \forall n \geq 0.$$

Using this equation for $n = 0$ and setting $h := 2A_0(1)$ we finally get formula (3.9).

3. The Taylor expansion of Ψ does not contain fourth order terms, i.e. the integer k in (3.6) is strictly greater than 1.

We need the Taylor expansions of the initial data $\Psi_0(r) = \Psi(r, r)$ and $Y_0(r) = Y(r, r)$. The first is easily obtained from eq. (3.6):

$$(3.13) \quad \Psi_0(r) = \frac{h}{2(\alpha+1)} r^3 + \dots$$

While the second has the form

$$(3.14) \quad Y_0(r) = 1 + c_l r^l + \dots \quad l \geq 2.$$

In fact the center must be initially at rest (due to the regularity condition $R(0, t) = 0$) and this implies $\lim_{r \rightarrow 0} u(r, r) = 0$. Choosing this initial velocity $u(r, r)$ as a differentiable function actually implies $u(r, r) = O(r)$. Using formulae (2.11) and (3.13) we easily get (3.14).

Using relations $A_0(\tau) = A_0(1)$, $B_0(\tau) = B_0(1)\tau^2$, coming from (3.9), together with (3.12), (3.6), we can calculate the Taylor expansion of $Y(r, r\tau)$ near $r = 0$ as

follows:

$$\begin{aligned}
Y(r, r\tau) &= \\
&= \left[\frac{A_0(1) + A_k(\tau)r^k + \dots}{A_0(1) + A_k(1)r^k + \dots} \right] \left[\frac{B_0(1)\tau^2 + B_k(1)\tau^2 r^k + \dots}{B_0(1)\tau^2 + B_k(\tau)r^k + \dots} \right]^{\frac{\alpha}{\alpha+1}} (1 + c_l r^l + \dots) \cong \\
&\cong \left(1 + \frac{A_k(\tau) - A_k(1)}{A_0(1)} r^k \right) \left(1 + \frac{\alpha}{\alpha+1} \left(\frac{B_k(1)}{B_0(1)} - \frac{B_k(\tau)}{B_0(1)\tau^2} \right) r^k \right) (1 + c_l r^l) \cong \\
&\cong \left(1 + \frac{2}{3h} (A_k(\tau) - A_k(1)) r^k \right) \left(1 - \frac{2}{3h} \left(B_k(1) - \frac{B_k(\tau)}{\tau^2} \right) r^k \right) (1 + c_l r^l) \cong \\
&\cong \left(1 + \frac{2}{3h} \left(A_k(\tau) - A_k(1) - B_k(1) + \frac{B_k(\tau)}{\tau^2} \right) r^k \right) (1 + c_l r^l),
\end{aligned}$$

and therefore

$$(3.15) \quad Y(r, r\tau) = 1 + c_l r^l + y_k(\tau) r^k + \dots$$

where

$$(3.16) \quad y_k(\tau) = \frac{2}{3h} \left(A_k(\tau) - A_k(1) + \frac{B_k(\tau)}{\tau^2} - B_k(1) \right).$$

Let us now suppose by contradiction that Taylor expansion of Ψ contains fourth order terms, that is $k = 1$. Let us take into account PDE (2.16), where we have explicitly written u^2 using (2.11):

$$\begin{aligned}
&\left(\frac{2\Psi}{R} + Y^2 - 1 \right) \left[\frac{\Psi_{,RR}}{(\alpha+1)\Psi_{,R}} + \frac{2\alpha}{(\alpha+1)R} \right] = \\
&= Y^2 \left[\frac{\alpha\Psi_{,RR}}{(\alpha+1)\Psi_{,R}} - \frac{2\Psi_{,rR}}{\Psi_{,r}} + \frac{(\alpha+1)\Psi_{,R}\Psi_{,rr}}{\alpha\Psi_{,r}^2} - \right. \\
&\quad \left. - \frac{2\alpha}{(\alpha+1)R} - \frac{(\alpha+1)\Psi_{,R}}{\alpha\Psi_{,r}} \left(\frac{1}{\alpha+1} \frac{\Psi_0''}{\Psi_0'} + \frac{2\alpha}{(\alpha+1)r} - \frac{Y_0'}{Y_0} \right) \right] + \\
&\quad \quad \quad + \frac{\alpha\Psi + R\Psi_{,R}}{\alpha R^2}.
\end{aligned}$$

Evaluating this equation along the lines $R = r\tau$, using (3.9) and (3.13)–(3.16), is a simple but tedious calculation, that results in equating to zero a polynomial $P(\tau)$. Then its coefficients must vanish, which implies $\Psi_{04} = \Psi_{13} = \Psi_{22} = \Psi_{31} = 0$. The constraint (3.12) for $n = 1$ finally shows that also $\Psi_{40} = 0$, and therefore there are no fourth order terms in the Taylor expansion of Ψ and the theorem is proved. Incidentally, let us observe that we have also shown that (see (3.16)) $y_1(\tau) = 0$, and therefore, for each τ in $(0, 1]$, one has

$$(3.17) \quad Y(r, r\tau) = 1 + y_2(\tau)r^2 + o(r^2).$$

□

Remark 3.2. A tedious but straightforward calculation shows that the Taylor expansion (3.6) is compatible with (2.16) "in the Cauchy-Kowaleski sense" at *any* order, that is, the equation allows the iterative calculation of all the higher order terms once the data are chosen. Of course, we stress that this is *not* a proof of global existence up to singularity formation (the coefficients of the equation are not analytic)

but only a - fundamental - consistency check for solutions here assumed a priori as regular.

3.3. Central singularity formation. The singularity forms whenever the denominator in (2.10) vanishes, that is $R = 0$. This kind of singularity is called a *shell-focusing* singularity (we are going to exclude here the so called *shell-crossing* singularities at which the particle flow-lines intersect each other). In comoving coordinates (r, t) , the locus of the zeroes of $R(r, t)$ defines implicitly a singularity curve $t_s(r)$ via $R(r, t_s(r)) = 0$. The quantity $t_s(r)$ represents the comoving time at which the shell labeled r becomes singular. The singularity forms if $t_s(r)$ is finite for each shell. In physically viable cases the curve $t_s(r)$ is strictly increasing and the center is the first point which can become singular. Obviously

$$(3.18) \quad \lim_{r \rightarrow 0^+} t_s(r) = t_0 \in (0 + \infty),$$

because we exclude the presence of a singularity on the data.

A perfect fluid solution need not form a singularity: one can have oscillating, regular spheres as well. However, it is very well known from various models of gravitational collapse (like e.g. shearfree perfect fluids [2]) that if the data are taken in a region sufficiently close to the center and the “effective potential” is not repulsive, the center itself collapses in a finite amount of comoving time. This is the case also for barotropic perfect fluids, as we are going to show.

Since $\dot{R} = -e^\nu u$, it is

$$(3.19) \quad t_s(r) = \int_0^r \rho^\alpha(r, \sigma) \sqrt{\frac{\sigma}{2\Psi(r, \sigma) + \sigma(Y^2(r, \sigma) - 1)}} d\sigma = \\ = \int_0^1 \left[-\frac{\Psi_{,R}(r, r\tau)}{4\pi\alpha r^2 \tau^2} \right]^{\frac{\alpha}{\alpha+1}} \frac{\sqrt{\tau} r^{3/2}}{(2\Psi(r, r\tau) + r\tau(Y^2(r, r\tau) - 1))^{1/2}} d\tau,$$

where the equation (2.8) has been used. The singularity forms if the above integral gives a real, finite value of $t_s(r)$ and of its limit t_0 (3.18). Reality of the integrand corresponds to the requirement that the considered regions of the R, r plane are allowed by the dynamics, i.e. $u^2 \geq 0$. Near the center, it suffices to consider this inequality on the $R = r\tau$ lines. Thus we study

$$(3.20) \quad 2\Psi(r, r\tau) + r\tau(Y^2(r, r\tau) - 1) \geq 0, \quad \forall \tau \in [0, 1].$$

Using (3.6) and (3.17) we get

$$(3.21) \quad 2\Psi(r, r\tau) + r\tau(Y^2(r, r\tau) - 1) \cong \\ \cong h \left(1 - \frac{\alpha}{\alpha+1} \tau^3 \right) r^3 + 2c_l \tau r^{l+1} + 2\tau y_k(\tau) r^{k+1} + \dots$$

We recall that $k, l \geq 2$ by Proposition 3.1. If both l and k are *strictly* greater than 2, (3.21) implies that (3.20) is automatically satisfied. In the case $l = 2$ and/or $k = 2$ the condition holds if

$$(3.22) \quad h \left(1 - \frac{\alpha}{\alpha+1} \tau^3 \right) + 2\delta_{l2} c_2 \tau + 2\delta_{k2} \tau y_2(\tau) > 0, \quad \forall \tau \in (0, 1),$$

where δ_{ij} is Kronecker delta. Once (3.22) is satisfied, $h > 0$ implies that integral (3.19) is finite for each $r > 0$, and it converges to a finite non-zero value as r goes to zero, obtaining the following

Proposition 3.3. *If regular initial data for a $p = \alpha\epsilon$ perfect fluid have initial density and velocity profiles such that*

- (1) l and k are strictly greater than 2, or
- (2) $l = 2$ and/or $k = 2$ but (3.22) is satisfied,

then the matter flow lines form a central singularity in a finite amount of comoving time.

4. THE SPECTRUM OF ENDSTATES

4.1. The apparent horizon. A key role in the study of the nature of a singularity is played by the apparent horizon $t_h(r)$ defined in (2.4) (see for instance [23]). The apparent horizon is the boundary of the trapped surfaces, and therefore represents the comoving time at which the shell labeled r becomes trapped. In area-radius coordinates this boundary is defined by $R_{hor} = 2\Psi(r, R_{hor})$. Since $\Psi_{,R}(0, 0) = 0$, implicit function theorem ensures that the curve R_{hor} is defined in a right neighborhood of $r = 0$. In what follows, we shall need the behavior of this curve near $r = 0$. It is easy to check that R_{hor} is strictly increasing and such that $R_{hor}(r) < r$. Moreover it is $R_{hor}(r) \cong 2\Psi(r, 0)$, since from (2.4) it is

$$R_{hor} = 2\Psi(r, 0) + 2R_{hor}\Psi_{,R}(r, 0) + R_{hor}^2 g(r, R_{hor}),$$

where g is bounded and $\Psi_{,R}(r, 0)$ is infinitesimal. Therefore, due to eq. (3.6), we conclude that

$$(4.1) \quad R_h(r) = hr^3 + \dots$$

Next section is devoted to the study of the nature of the central ($R = r = 0$) singularity. We restrict ourselves to this singularity since, in barotropic perfect fluid models with positive pressures, it is the only one that can be naked. This is easily seen using comoving coordinates. Indeed, a singularity cannot be naked if it occurs after the formation of the apparent horizon (i.e. it must be $t_h(r) \geq t_s(r)$). A necessary condition for this is that the singularity must be massless ($\Psi(r, t_s(r)) = 0$). But, due to equation (2.2b), in presence of a positive pressure the mass is strictly increasing in a collapsing ($\dot{R} < 0$) situation, while it is zero at the regular centre. The situation can be completely different if negative pressures are allowed: in this case non central singularities can be naked as well [9].

4.2. Nakedness of the central singularity. At the center ($R = r = 0$) the apparent horizon and the singularity form simultaneously and the necessary condition for nakedness is satisfied. The singularity will be (locally) naked if there exists a radial lightlike future pointing local solution $R_g(r)$ of the geodesic equation with initial condition $R(0) = 0$ "travelling before the apparent horizon", that is - in area radius coordinates - $R_g(r) > R_{hor}(r)$ for $r > 0$. We will study in full details only the existence of *radial* null geodesics emanating from the singularity. It can in fact be proved that, if a singularity is radially censored (that is, no radial null geodesics escape), then it is censored [31, 13].

The equation of radial null geodesics in the coordinate system (r, R) is easily found from (2.15) setting $ds^2 = 0$ together with $d\theta = d\phi = 0$:

$$(4.2) \quad \frac{dR}{dr} = -\frac{\alpha}{\alpha + 1} \frac{\Psi_{,r}}{\Psi_{,R}} \left(1 - \frac{u}{Y}\right).$$

Our main result can be stated as follows:

Theorem 4.1. *For any choice of initial data $Y_0(r)$, $\Psi_0(r)$ for the Einstein field equations such that*

- (1) *the conditions stated in proposition 3.3 hold (i.e. the central singularity forms in a finite amount of comoving time), and*
- (2) *the mass function is regular in a neighbor of the spacetime singularity,*

there exists solutions of (4.2) that extend back to the central singularity, which is therefore locally naked.

Proof. To show the existence of singular geodesics we use a simple technique developed earlier [13]. First of all, we recall that a function $y_0(r)$ is called a subsolution (respectively supersolution) of an ordinary differential equation of the kind $y' = f(r, y)$ if it satisfies $y'_0 \leq f(r, y_0)$ (respectively \geq). Now, it can be shown [12] that the apparent horizon $R_h(r)$ is a supersolution of the geodesic equation (4.2). The singularity is certainly naked if it is possible to find a subsolution $R_+(r)$ of the same equation which stays over the horizon. In fact, choose a point (r_0, R_0) in the region $S = \{(r, R) : r > 0, R_{hor}(r) < R < R_+(r)\}$. At this point the (regular) Cauchy problem with datum $R(r_0) = R_0$ admits a unique local solution $R_g(r)$. Now the extension of this solution in the past cannot escape from S since either it would cross the supersolution from above or it would cross the subsolution from below. Thus it must extend back to the singularity with $\lim_{r \rightarrow 0^+} R_g(r) = 0$.

We now proceed to show that a subsolution always exist. For this aim, it suffices to consider lines $R_x(r) = xr$ where x is a positive parameter. Due to eq. (4.1), $R_x < R_{hor}$ for sufficiently small r . From (3.6) it is

$$(4.3) \quad -\frac{\alpha}{\alpha + 1} \frac{\Psi_{,r}(r, xr)}{\Psi_{,R}(r, xr)} \cong \frac{1}{x^2},$$

while (2.11), (3.5), (3.6) and (3.17) easily imply

$$(4.4) \quad \frac{u(r, xr)}{Y(r, xr)} = O(r).$$

Then $R_x(r) = xr$ is the sought subsolution of (4.2) if

$$x < \frac{1}{x^2},$$

that is for any $x < 1$. □

5. EXTENSION TO THE GENERAL BAROTROPIC CASE

We are going to show in the present section that our main results - in particular, the formation of naked singularities as endstates of all the solutions for which the mass function is regular in a neighborhood of the spacetime singularity - actually

hold for the general, barotropic equation of state $\epsilon = \epsilon(\rho)$. Of course, not all functions are allowed, and we shall assume the following - quite natural - hypotheses on $\epsilon(\rho)$.

Assumption 5.1. We assume $\epsilon = \epsilon(\rho)$ to be a \mathcal{C}^1 function in $[\bar{\rho}, +\infty)$ (where $\bar{\rho} \geq 0$), such that $\epsilon(\rho) \geq 0$ ($= 0$ iff $\rho = \bar{\rho}$). Recalling (2.5), that is $p(\rho) = \rho \frac{d\epsilon}{d\rho} - \epsilon$, we also assume $p(\rho)$ is a strictly positive \mathcal{C}^1 function with $\frac{dp}{d\rho} > 0$, except at most for a bounded interval $[\bar{\rho}, \rho_1]$, possibly coinciding with a single point, where $p(\rho)$ can vanish.

Remark 5.2. Observe that:

(1) The assumptions made imply that

$$(5.1) \quad \frac{d\epsilon}{d\rho}(\rho) > 0 \text{ if } \rho > \bar{\rho}.$$

and therefore $\epsilon(\rho)$ is a strictly increasing positive function.

(2) Differentiating (2.5) we have, where it makes sense,

$$(5.2) \quad \frac{dp}{d\rho} = \rho \frac{d^2\epsilon}{d\rho^2},$$

then $\epsilon(\rho)$ is strictly convex for ρ sufficiently large, and so

$$(5.3) \quad \lim_{\rho \rightarrow +\infty} \frac{d\epsilon}{d\rho}(\rho) = +\infty.$$

(3) The assumptions made also shows the existence of $\lim_{\rho \rightarrow \infty} p(\rho)$. If the limit was finite, say l , it would be $\frac{d\epsilon}{d\rho}(\rho) < \frac{1}{\rho}(l + \epsilon(\rho))$, and then $\epsilon(\rho) < \rho + l$ by a simple comparison argument in o.d.e., which is in contradiction with (5.3). Then

$$(5.4) \quad \lim_{\rho \rightarrow \infty} p(\rho) = +\infty.$$

Remark 5.3. We stress that the above mentioned hypotheses are quite natural from the physical point of view. Besides obviously including the $p = \alpha\epsilon$ equation of state considered so far, they include, for instance, the equation of state of the perfect gas $p(\rho) = K_2\rho$ for which $\epsilon(\rho) = K_1\rho + K_2\rho \log \rho$ where K_1 and K_2 are positive constants (in this case one obviously has $\rho_1 = \bar{\rho} = e^{-K_1/K_2}$).

Einstein's equation (2.2b) reads

$$(5.5) \quad p = -\frac{\Psi_{,R}}{4\pi R^2}.$$

Using it, together with (2.2a), (2.5), (2.6) and the coordinate change formulae $\Psi' = \Psi_{,r} + R' \Psi_{,R}$ and $\dot{\Psi} = \dot{R} \Psi_{,R}$, we obtain the general counterparts for equations (2.9) and (2.13), namely

$$(5.6) \quad R' = -\frac{\Psi_{,r}}{\Psi_{,R}} \frac{p}{\epsilon + p} = \frac{\Psi_{,r}}{4\pi R^2(\epsilon + p)},$$

and

$$(5.7) \quad Y(r, R) = \frac{E(r)\Psi_{,r}(r, R)}{(d\epsilon/d\rho)(\rho(r, R))}.$$

Using these formulae, we can again express the metric in the form (2.15). The crucial point is now that we can express $\frac{d\epsilon}{d\rho}(\rho)$ as a function of $p(\rho)$ and, as a consequence, the dynamics of the system is, also in the general case, expressed in terms of the mass function and its derivatives only. To this end, consider the parameterized curve in \mathfrak{R}^2

$$\mathfrak{R}^+ \ni \rho \mapsto \left(\xi(\rho) = p(\rho), \zeta(\rho) = \frac{d\epsilon}{d\rho}(\rho) \right).$$

This curve is globally the graphic of a function $\zeta = \zeta(\xi)$, recalling that ξ is a non-decreasing function of ρ (by the assumptions on $p(\rho)$), and that, by (5.2), $\frac{(d\zeta/d\rho)}{(d\xi/d\rho)} = \rho \neq 0$. Since also ζ is increasing by (1) of Remark 5.2, there exists $\lim_{\rho \rightarrow \bar{\rho}^+} \zeta(\rho) = \zeta_0$ finite. Denoting by ξ_0 the finite number $\lim_{\rho \rightarrow \bar{\rho}^+} \xi(\rho)$, the function may be prolonged up to the point (ξ_0, ζ_0) . Let also observe that this function is C^1 , for each $\xi > \xi_0$, where indeed $\frac{d\xi(\rho)}{d\rho}$ is strictly positive by the assumptions made on $p(\rho)$.

Using this result, and recalling (2.5), one finds that Y in (5.7) (and then u in (2.11)) can be expressed as functions of the data and of the mass function $\Psi(r, R)$ and its derivatives. Then, again, with some calculations one obtains a second order PDE that must be satisfied by Ψ . As in the case treated so far, we consider only Taylor expandable solutions of this equation, and proceed to analyze the structure of the lower order terms of the mass profile.

First of all, since $R' \equiv 1$ on the data surface $R = r$, the expression for the initial energy is:

$$\epsilon(r, r) = \frac{\Psi_{,r}(r, r) + \Psi_{,R}(r, r)}{4\pi r^2}.$$

Imposing the regularity condition $\lim_{r \rightarrow 0^+} \epsilon(r, r) \in (0, +\infty)$ and making reference to the notation used in Section 3.2 we get

$$A_0(1) + B_0(1) > 0.$$

Actually, $A_0(1) \neq 0$. Otherwise, $\epsilon(r, r) + p(r, r) \rightarrow 0$ as $r \rightarrow 0^+$, since $\epsilon(r, r) + p(r, r) = \frac{\Psi_{,r}(r, r)}{4\pi r^2} = \frac{A_0(1)}{4\pi} + o(1)$. But $(\epsilon + p)(\rho)$ is a strictly increasing and non negative function of ρ , then it would be $\rho(r, r) \rightarrow \bar{\rho}$, which would imply $\epsilon(r, r) \rightarrow 0$, that is a contradiction.

As in section 3.1, for physical reasonableness we suppose the initial energy $\epsilon(r, r)$ (and therefore $\rho(r, r)$) to be a non increasing function of r . This implies that we can consider, without loss of generality, the case in which also $B_0(1) \neq 0$. In fact, if $B_0(1)$ vanishes by (2.5) it has to be $p(r, r) \rightarrow 0$ as $r \rightarrow 0^+$. This fact, recalling the assumptions made on the pressure, shows that $p(r, r)$ (that is a non increasing function of r) must be identically zero. But ρ (and therefore p) must diverge at the spacetime singularity, and therefore there exists an hypersurface, such that p is non zero but the energy ϵ is still regular, where we can re-assign the initial data on. On this hypersurface, the pressure must converge to a finite non-zero value as $r \rightarrow 0^+$. Then we will suppose $B_0(1) \neq 0$. Finally, we note that positivity of pressure on the data further implies that $A_0(1) > 0$ and $B_0(1) < 0$.

We are now ready to investigate lower order terms in the mass function. Recall that regularity of pressure along the initial data implies that Ψ cannot contain first

order terms (see (5.5)). Then, as in (3.7) we set

$$(5.8) \quad \Psi(r, R) = \frac{1}{2}m_0(R-r)^2 + \sum_{i+j=3} \Psi_{ij}r^iR^j + \dots$$

We denote by $\epsilon_0 > 0$ the limit $\lim_{r \rightarrow 0^+} \epsilon(r, r)$. By (1) of Remark 5.2, there exists a unique $\rho_0 > 0$ such that $\epsilon(\rho_0) = \epsilon_0$, and clearly $\rho_0 = \lim_{r \rightarrow 0} \rho(r, r)$. We also denote by β_0 the positive number $\frac{d\epsilon}{d\rho}(\rho_0)$. Using (3.5) at $\tau = 1$ we have

$$(5.9) \quad E(r) = \frac{\beta_0}{A_0(1)} \frac{1}{r^2} + \dots,$$

plus higher order terms. Let us show, as in Proposition 3.1 (claim 1) that $m_0 = 0$. Otherwise, using (3.5) at $\tau < 1$, together with (5.7) and (5.9), one finds

$$\frac{d\epsilon}{d\rho}(\rho(r, r\tau)) = \frac{E(r)\Psi_{,r}(r, r\tau)}{Y(r, r\tau)} = \frac{\epsilon_1}{r} + \dots,$$

plus higher order terms, whereas, using (2.5), it is also

$$p(\rho(r, r\tau)) = \frac{p_1}{r} + \dots$$

plus higher order terms again. Then, since ϵ_1 and p_1 are strictly positive constants, it is $\lim_{\rho \rightarrow \infty} \frac{(d\epsilon/d\rho)(\rho)}{p(\rho)} > 0$. But recalling (5.2) $\lim_{\rho \rightarrow \infty} \frac{(d^2\epsilon/d\rho^2)(\rho)}{(dp/d\rho)(\rho)} = \lim_{\rho \rightarrow \infty} \frac{1}{\rho} = 0$, coming to a contradiction by de L'Hospital theorem. Thus m_0 vanishes.

Observe now that, for a fixed τ ,

$$p(r, r\tau) = \frac{B_0(\tau)}{4\pi\tau^2} + \dots = -\frac{1}{4\pi} \left(\frac{\Psi_{21}}{\tau^2} + 2\frac{\Psi_{12}}{\tau} + 3\Psi_{03} \right) \equiv p_0(\tau).$$

If $(\Psi_{21}, \Psi_{12}) \neq (0, 0)$, then $p_0(\tau) \rightarrow \infty$ as $\tau \rightarrow 0$, and so $\rho(r, r\tau)$ (and therefore $\frac{d\epsilon}{d\rho}(\rho(r, r\tau))$) is sufficiently large, for τ near to 0. This leads to a contradiction, since (5.8) with $m_0 = 0$ and (5.9) show that $Y(r, r\tau) \cong \frac{\mu}{(d\epsilon/d\rho)(r, r\tau)}$, for some non-zero constant μ independent of τ , but (3.5) must hold. Then, again, $B_0(\tau) = B_0(1)\tau^2$, and $p_0(\tau) = p_0(1) \equiv p_0 > 0$.

Taken together, the above arguments show that the lower order terms of the mass have the structure, analogue to (3.9),

$$(5.10) \quad \Psi(r, R) = \frac{h}{2} \left(r^3 - \frac{p_0}{\epsilon_0 + p_0} R^3 \right) + \dots$$

We now proceed analyzing the nature of the singularity forming at the center. For this purpose, we recall that the denominator in the righthandside of (5.7) can be written as a function $\zeta \left(-\frac{\Psi_{,R}}{4\pi R^2} \right)$ (where we have also used (5.5)), which is certainly \mathcal{C}^1 at the point $p_0 > 0$ defined above. Then

$$\begin{aligned} \zeta(p(r, r\tau)) &= \zeta \left(-\frac{B_0(1)}{4\pi} \right) + \frac{d\zeta}{d\xi} \left(-\frac{B_0(1)}{4\pi} \right) \left(-\frac{B_k(\tau)}{4\pi\tau^2} r^k + \dots \right) = \\ &= \zeta(p_0) - \frac{B_k(\tau)}{4\pi\tau^2} \frac{d\zeta}{d\xi}(p_0) r^k + \dots, \end{aligned}$$

therefore the Taylor development for the function $Y(r, r\tau)$ can be written in a similar manner as in (3.15). It is now to check that, due to the regular behavior of the

equation of state near the center, arguments similar to those used in section 3.3 can be used to show that data leading to complete gravitational collapse always exist.

For such data we consider the behavior of the null radial geodesic equation

$$(5.11) \quad \frac{dR}{dr} = R' \left(1 - \frac{u}{Y} \right).$$

It is easy to check that some of the curves $R = r\tau$ are super-horizon subsolution of this equation, and therefore the singularity is naked. First of all, we notice that (4.4) still holds. Now using (2.6), (3.5) and (5.9), one has

$$R'(r, r\tau) \cong e^\lambda(r, r\tau) = \frac{1}{4\pi\rho(r, r\tau)E(r)r^2\tau^2} \cong \frac{A_0(1)}{4\pi\rho(r, r\tau)\beta_0\tau^2} \cong \frac{\rho_0}{\rho(r, r\tau)} \frac{1}{\tau^2},$$

where last equality follows from $R'(r, r) = 1$. Since $p(r, r\tau) \rightarrow p_0 > 0$, this implies that $\rho(r, r\tau)$ is bounded from above, uniformly with respect to τ , as r goes to 0. Then there exists at least one τ satisfying $\tau^3 < \frac{\rho_0}{\rho(r, r\tau)}$, and again $R(r) = r\tau$ is a super-horizon subsolution of null radial geodesic equation. We therefore conclude this section with the analogue of Theorem 4.1, that is:

Theorem 5.4. *Under the hypotheses made on the equation of state in the assumption 5.1, for any choice of initial data for the Einstein field equations such that*

- (1) *the central singularity forms in a finite amount of comoving time, and*
- (2) *the mass function is regular in a neighbor of the spacetime singularity,*

there exists solutions of (5.11) that extend back to the central singularity, which is therefore locally naked.

6. DISCUSSION AND CONCLUSIONS

Our main theorem shows, that all the solutions of the Einstein field equations describing the complete collapse of barotropic perfect fluids for which the mass function is regular in a neighborhood of the regular center up to singularity formation form naked singularities. Besides of spherical symmetry, this result is independent on any simplifying assumption (such as self-similarity).

As a consequence, the problem of the cosmic censorship conjecture in spherical symmetry is solved at least for the cases of collapsing matter continua which are of main astrophysical interest: a cosmic censor simply does not exist for such materials. Dust and anisotropic media form naked singularities for "generic" choices of the data (see e.g. [13] and references therein) while perfect fluids "always" form naked singularities.

The next problem facing future research is to go beyond spherical symmetry. Trends in this direction became quite clear in recent years, together with the overwhelming evidence for naked singularities. First, physically observable effects of naked singularity formation, such as gravitational radiation, have to be analyzed [19]. On the other end, the case for a (by necessity, non spherically symmetric) cosmic censorship waits to be investigated.

REFERENCES

- [1] O. I. Bogoyavlenski, *Methods in the Qualitative Theory of Dynamical Systems in Astrophysics and Gas Dynamics*, (Springer-Verlag, 1985).
- [2] H. Bondi, *Mon. Not. Roy. Astr. Soc.* **142** 333 (1969).

- [3] E. Brinis, S. Jhingan and G. Magli, *Class. Quantum Gravity* **17** (2000), 4481-4490
- [4] B. J. Carr, A. A. Coley, M. Goliath, U. S. Nilsson, C. Uggla, *Class.Quant.Grav.* **18** (2001) 303.
- [5] D. Christodoulou, *Comm. Math. Phys.* **93** 171 (1984)
- [6] D. Christodoulou, *Ann. Math.* **140** 607 (1994)
- [7] D. Christodoulou, *Arch. Rat. Mech. Anal.* **130**, 343 (1995)
- [8] D. Christodoulou, *Ann. Math.* **149** 183 (1999)
- [9] F. I. Cooperstock, S. Jhingan, P. S. Joshi and T. P. Singh, *Class. Quantum Grav.* **14** 2195 (1997).
- [10] D. M. Eardley and L. Smarr, *Phys. Rev. D* **19** 2239 (1979)
- [11] T. Foglizzo and R. N. Henriksen *Phys. Rev D* **48** 4645 (1993)
- [12] R. Giambò, F. Giannoni, G. Magli, *Class. Quantum Grav.* **19** L5 (2002).
- [13] R. Giambò, F. Giannoni, G. Magli, P. Piccione, gr-qc/0110027, *Comm. Math. Phys.*, at press.
- [14] R. Giambò, F. Giannoni, G. Magli, P. Piccione, gr-qc/0212082, *Class. Quantum Grav.*, at press
- [15] S. Gonçalves, *Phys. Rev. D* **63** 64017 (2001)
- [16] S. Gonçalves, S. Jhingan, G. Magli, *Phys. Rev. D* **65**, 64011 (2002)
- [17] T. Harada, *Phys. Rev. D* **58** 104015 (1998)
- [18] T. Harada, H. Iguchi and K. Nakao, *Phys. Rev. D* **58** R041502 (1998)
- [19] T. Harada, H. Iguchi and K. Nakao, *Prog.Theor.Phys.* 107 (2002) 449
- [20] T. Harada, H. Iguchi and K. Nakao, *Class.Quant.Grav.* **16** (1999) 2785
- [21] S. Jhingan and G. Magli *Phys. Rev. D* **61** 124006 (2000)
- [22] S. Jhingan and G. Magli, *Gravitational collapse of fluid bodies and cosmic censorship: analytic insights*, In *Recent developments in General Relativity* B. Casciaro, D. Fortunato, A. Masiello, M. Francaviglia ed., Springer Verlag (Berlin) (2000)
- [23] P. S. Joshi, *Global aspects in gravitation and cosmology*, (Clarendon press, Oxford, 1993)
- [24] P.S. Joshi, N. Dadhich, and R. Maartens, gr-qc/0109051
- [25] P. S. Joshi, T. P. Singh, *Class. Quantum Grav.* **13** 559 (1996)
- [26] D. Kramer, H. Stephani, E. Herlt, M. MacCallum, *Exact Solutions of the Einstein's Field Equations*, Cambridge University Press (1980)
- [27] A. Krasinski, *Inhomogeneous cosmological models*, (Cambridge Univ. Press, 1997)
- [28] S. Kind and J. Ehlers, *Class. Quantum Grav.* **10** 2123 (1993)
- [29] G. Magli, *Class. Quantum Grav.* **14** 1937 (1997)
- [30] G. Magli, *Class. Quantum Grav.* **15** 3215 (1998)
- [31] F. C. Mena, B. C. Nolan, *Class. Quantum Grav.* **18** 4531 (2001)
- [32] A. Ori, *Class. Quantum Grav.* **7** 985 (1990)
- [33] A. Ori and T. Piran, *Phys. Rev. D* (1990) **42** 1068.
- [34] R. Penrose, *Nuovo Cimento* **1** 252 (1969).
- [35] A. D. Rendall, *J. Math. Phys.*, **33**, 1047 (1992).
- [36] J. Smoller and B. Temple, *Arch. Rat. Mech. Anal.* **138**, 239 (1997).

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI CAMERINO, ITALY
E-mail address: roberto.giambo@unicam.it, fabio.giannoni@unicam.it

DIPARTIMENTO DI MATEMATICA, POLITECNICO DI MILANO, ITALY
E-mail address: magli@mate.polimi.it

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI CAMERINO, ITALY
 (ON LEAVE FROM DEPARTAMENTO DE MATEMÁTICA,
 UNIVERSIDADE DE SÃO PAULO, BRAZIL)
E-mail address: paolo.piccione@unicam.it, (piccione@ime.usp.br)