

Quasinormal behavior of the d -dimensional Schwarzschild black hole and higher order WKB approach

R.A.Konoplya

Department of Physics, Dniepropetrovsk National University

St. Naukova 13, Dniepropetrovsk 49050, Ukraine

konoplya@ff.dsu.dp.ua

Abstract

We find the low lying scalar quasinormal modes of d -dimensional Schwarzschild black hole. It proves that the real part of the complex mode is proportional to the number of dimensions d . In addition the WKB formula for computing QN modes, developed to the 3th order beyond the eikonal approximation, is extended to the 6th order here. This gives us accurate and economic way to compute the quasinormal frequencies. The asymptotic formula for large multipole number l and arbitrary d is derived.

When perturbing a black hole, it undergoes damping oscillations which are characterized by some complex eigenvalues of the wave equations. These values (*quasinormal frequencies*) are complex, and their real parts give the oscillation frequencies, while the imaginary ones determine the damping rates of the modes. The quasinormal modes (QN) of black holes (BH's) depend only on a black hole parameters and not on a way in which they were excited. QN's are called, therefore, "footprints" of a black hole. Quasinormal modes are of considerable interest now: they are studied within the context of Anti-de-Sitter/Conformal Field Theory (AdS/CFT) correspondence (see for example [1]- [10] and references therein), because of the possibility of observing quasinormal ringing of astrophysical BH's (see [12] for a review), when considering thermodynamic properties of black holes in one loop quantum gravity [13]-[16], in the context of possible connection with critical collapse [1], [30], [5], [17]. Quasinormal modes of multidimensional black hole were considered in [18] and also in [19] within context of one loop quantum gravity.

The metric of the Schwarzschild black hole in d -dimensions has the form:

$$ds^2 = f(r)dt^2 - f^{-1}(r)dr^2 + r^2 d\Omega_{d-2}^2, \quad (1)$$

where

$$f(r) = 1 - \left(\frac{r_0}{r}\right)^{d-3} \quad (2)$$

The scalar perturbation equation of this black hole can be reduced to the Schrodinger wave-like equation:

$$\frac{d^2\psi}{dx^2} + Q(x)\psi(x) = 0, \quad dx = \frac{dr}{f(r)} \quad (3)$$

where "the potential" $-Q(x)$ is constant at the event horizon ($x = -\infty$), and at infinity ($x = +\infty$) and it rises to maximum at some intermediate $x = x_0$:

$$Q(x) = \omega^2 - f(r) \left(\frac{l(l+d-3)}{r^2} + \frac{(d-2)(d-4)}{4r^2} f(r) + \frac{d-2}{2r} f'(r) \right), \quad (4)$$

The quasinormal modes satisfy the boundary conditions:

$$\phi(x) \sim c_{\pm} e^{\mp\omega x} \quad as \quad x \rightarrow \pm\infty \quad (5)$$

Consider radiation of a given frequency ω incident on the black hole from infinity and let $R(\omega)$ and $T(\omega)$ be the reflection and transmission amplitudes respectively. Extend $R(\omega)$ to the complex frequency plane such that $Re(z) \neq 0$, and $T(z)/R(z)$ is regular. Then the quasinormal modes correspond to the singularities of $R(z)$. We have a direct analogy with the problem of scattering near the peak of the potential barrier in quantum mechanics, where ω^2 plays a role of energy, and the two turning points divide the space into three regions, at which boundaries the corresponding solutions should be matched.

For calculation of the QNMs of SBH in various dimensions we used the WKB formula extended to the 6th order beyond the eikonal approximation. First analytical method for calculations of BH QNMs was apparently proposed by Bahram Mashhoon who used the Poschl-Teller potential to estimate the QN frequencies [20]. In [21] there was proposed a semi-classical method for computing QNM's based on the WKB treatment [22]. Then it was extended to the third order beyond the eikonal approximation in [22]. To extend the 3th order WKB formula of [22] we used the techniques of Iyer and Will. Note that

since the coefficients M_{ij} that connect amplitudes near the horizon with those at infinity depend only on some parameter related to the overtone number n , the WKB corrections may be found to higher orders, simply by solving the interior (between the turning points) problem to higher orders, and without performing an explicit match of the solutions to WKB solutions in the exterior (outside turning points) regions to the same order. The result has a simple form:

$$\frac{iQ_0}{\sqrt{2Q_0''}} - \Lambda_2 - \Lambda_3 - \Lambda_4 - \Lambda_5 - \Lambda_6 = n + \frac{1}{2}, \quad (6)$$

where the correction terms $\Lambda_4, \Lambda_5, \Lambda_6$, can be found in the Appendix 1. Note that Λ_4 coincides with preliminary formula (A3) of [22] in proper designations. The QNMs of $d = 4$ Schwarzschild black hole for fields of different spin are considered in Appendix 2 and may serve as a test of the obtained corrections, since one can be sure that each next WKB order gives the QN value of ω^2 which is closer to the numerical values than the corresponding QNMs given by a previous order.

An alternative, pure algebraic approach to finding higher order WKB corrections was proposed by O.Zaslavskii [24], using a quantum anharmonic oscillator problem where WKB correction terms come from perturbation theory corrections to the potential anharmonicity. When considering BH QNMs the WKB approach was effectively used also in many problems (see for example [25]-[33] and references therein).

The 6th WKB order formula used here gives quite accurate results for low overtones, since the previous orders serve us to see the convergence of the WKB values of ω^2 to an accurate numerical result. Namely we can observe that for $l = 2, 3, 4, ..$ for the fundamental overtone the 6th order values differs from its 5th order value by fractions of a percent or less at not very large d (we are restricted here by $d = 4, 5, ...15$). It proves out that the real part of ω is proportional to the dimension d and for the fundamental overtone we obtain the following approximate relation:

$$\omega_{Re} \approx 0.244d, \quad l = 2 \quad (7)$$

$$\omega_{Re} \approx 0.275d, \quad l = 3 \quad (8)$$

$$\omega_{Re} \approx 0.290d, \quad l = 4. \quad (9)$$

Here we take $\omega = \omega_{Re} - i\omega_{Im}$.

The more d and n are, the less accuracy of the WKB approach is. Therefore we did not present the higher overtone and the case of very large d . Nevertheless, when l is considerably greater than n we can check with the WKB formula that the above qualitative properties remain for higher n ; in particular, ω_{Re} is again proportional to d . It is possible that the asymptotic value for ω_{Re} as $n \rightarrow \infty$ will be proportional to d as well.

For large l the well-known approximate formula reads

$$\omega_{Re} = \frac{1}{3\sqrt{3}} \left(l + \frac{1}{2} \right), \quad \omega_{Im} = \frac{1}{3\sqrt{3}} \left(n + \frac{1}{2} \right) \quad (10)$$




Figure 1: Rew as a function of the dimension d ; $l = 2, 3, 4$, from the bottom to the top; $n = 0$.




Figure 2: Imw as a function of the dimension d ; $l = 2, 3, 4$, from the top to the bottom; $n = 0$.

To obtain its d -dimensional generalization we find a value r_{max} at which the effective potential V attains its maximum, provided l is large

$$r_{max} \approx 2^{\frac{d-4}{d-3}} (d-1)^{\frac{1}{d-3}}, \quad d = 4, 5, 6, \dots \quad (11)$$

Then let us make use of this value r_{max} when dealing with the first order WKB formula. After expansion in terms of small values of $1/l$ we obtain

$$\omega_{Re} \approx \frac{d+2l-3}{4} \left(\frac{2}{d-1} \right)^{\frac{1}{d-3}} \sqrt{\frac{d-3}{d-1}} \quad (12)$$

$$\omega_{Im} \approx \frac{d-3}{4} \left(\frac{2}{d-1} \right)^{\frac{2n-1}{d-3}} \frac{1}{\sqrt{d-1}} \quad (13)$$

When $d = 4$ these formulas go over into (11).

References

- [1] G.T.Horowitz and V.Hubeny *Phys.Rev.* **D62** 024027 (2000)
- [2] V.Cardoso and J.P.S.Lemos *Phys.Rev.* **D63** 124015 (2001)
- [3] D. Birmingham, I. Sachs, and S.N. Solodukhin, *Phys. Rev. Lett.* **88** 151301 (2002)
- [4] D. Birmingham, I. Sachs, and S.N. Solodukhin, hep-th/0212308 (2002)
- [5] R.A.Konoplya, *Phys.Rev.* **D66** 084007 (2002)
- [6] R.A.Konoplya, *Phys.Rev.* **D66** 044009 (2002)
- [7] A. O. Starinets, *Phys.Rev.D66*, 124013 (2002)
- [8] A. Nunez, A. O. Starinets, hep-th/0302026
- [9] R.Aros, C.Martinez, R.Troncoso, and J.Zanelli, hep-th/0211024;
- [10] I.G.Moss and J.P.Norman, *Class.Quant.Grav.* **19**, 2323 (2002) (2002)
- [11] S. Musiri, G. Siopsis, hep-th/0301081
- [12] K.Kokkotas and B.Schmidt, "Quasi-normal modes of stars and black holes", *Living.Reviews.Relativ.*, **2**, 2 (1999)
- [13] O. Dreyer, gr-qc/0211076.
- [14] G. Kunstatter, gr-qc/0211076.
- [15] V. Cardoso, J.P.S. Lemos, gr-qc/0301078
- [16] L. Motl, gr-qc/0212096.
- [17] W.T.Kim and J.J.Oh, *Phys.Lett.***B514** 155 (2001)
- [18] V. Cardoso, O.J.C. Dias, J.P.S. Lemos hep-th/0212168
- [19] L. Motl, A. Neitzke, hep-th/0301173
- [20] V.Ferrari and B.Mashhoon, *Phys. Rev. Lett.* **52** 1361 (1984); H-J.Blome and B.Mashhoon, *Phys.Lett.* **100A** 231 (1984)
- [21] B.F.Schutz and C.M.Will *Astrophys.J.Lett* **291** L33 (1985)
- [22] S.Iyer and C.M.Will, *Phys.Rev.* **D35** 3621 (1987)
- [23] E.Leaver, *Proc.R.Soc.London* **A402**, 285 (1985)
- [24] O.B.Zaslavskii, *Phys.Rev.* **D43** 605 (1991)
- [25] S.Iyer, *Phys.Rev.* **D35** 3632 (1987)
- [26] K.Kokkotas, and B.F.Schutz, *Phys.Rev.* **D37** 3378 (1988)

- [27] K.Kokkotas, *Nuovo.Cimento.* **B108** 991 (1993)
- [28] L.E.Simone and C.M.Will, *Class.Quant.Grav.* **9**, 963 (1992)
- [29] N.Andersson and H.Onozawa, *Phys.Rev.* **D54** 7470 (1996)
- [30] R.A.Konoplya, *Phys.Lett.* **B 550**, 117 (2002)
- [31] R.A.Konoplya, *Gen.Relativ.Grav.* **34**, 329 (2002)
- [32] V.Ferrari, M.Pauri and F.Piazza, *Phys.Rev.* **D63** 064009 (2001)
- [33] H.Onozawa, T.Okamura, T.Mishima, and H.Ishihara, *Phys.Rev.* **D53** 7033 (1996)

1 Appendix 1: Correction terms for WKB formula

Here we shall follow the designations: Q_0 means the value of the potential Q at its pick, while Q_i is the i th derivative of Q with respect to the tortoise coordinate x . Then Q_i^j is the j th power of the i th derivative of Q .

$$\begin{aligned}
\Lambda_4 = & \frac{1}{597196800\sqrt{2}Q_2^7\sqrt{Q_2}}(2536975Q_3^6 - 9886275Q_2Q_3^4Q_4 + 5319720Q_2^2Q_3^3Q_5 - \\
& 225Q_2^2Q_3^2(-40261Q_4^2 + 9688Q_2Q_6) + 3240Q_2^3Q_3(-1889Q_4Q_5 + 220Q_2Q_7) - \\
& 729Q_2^3(1425Q_4^3 - 1400Q_2Q_4Q_6 + 8Q_2(-123Q_5^2 + 25Q_2Q_8))) + \\
& \frac{(n+1/2)^2}{4976640\sqrt{2}Q_2^7\sqrt{Q_2}}(348425Q_3^6 - 1199925Q_2Q_3^4Q_4 + 57276Q_2^2Q_3^3Q_5 - \\
& 45Q_2^2Q_3^2(-20671Q_4^2 + 4552Q_2Q_6) + 1980Q_2^3Q_3(-489Q_4Q_5 + 52Q_2Q_7) - \\
& 27Q_2^3(2845Q_4^3 - 2360Q_2Q_4Q_6 + 56Q_2(-31Q_5^2 + 5Q_2Q_8))) + \\
& \frac{(n+1/2)^4}{2488320\sqrt{2}Q_2^7\sqrt{Q_2}}(192925Q_3^6 - 581625Q_2Q_3^4Q_4 + 234360Q_2^2Q_3^3Q_5 - \\
& 45Q_2^2Q_3^2(-8315Q_4^2 + 1448Q_2Q_6) + 1080Q_2^3Q_3(-161Q_4Q_5 + 12Q_2Q_7) - \\
& 27Q_2^3(625Q_4^3 - 440Q_2Q_4Q_6 + 8Q_2(-63Q_5^2 + 5Q_2Q_8))) \tag{14}
\end{aligned}$$

$$\begin{aligned}
\Lambda_5 = & \frac{(n+1/2)}{57330892800Q_2^{10}}(2768256Q_{10}Q_2^7 - 1078694575Q_3^8 + 5357454900Q_2Q_3^6Q_4 - \\
& 2768587920Q_2^2Q_3^5Q_5 + 90Q_2^2Q_3^4(-88333625Q_4^2 + 12760664Q_2Q_6) - \\
& 4320Q_2^3Q_3^3(-1451425Q_4Q_5 + 91928Q_2Q_7) - 27Q_2^4(7628525Q_4^4 - 9382480Q_2Q_4^2Q_6 + \\
& 64Q_2^2(19277Q_2^6 + 37764Q_2Q_7) + 576Q_2Q_4(-21577Q_5^2 + 2505Q_2Q_8)) + \\
& 540Q_2^3Q_3^2(6515475Q_4^3 - 3324792Q_2Q_4Q_6 + 16Q_2(-126468Q_5^2 + 12679Q_2Q_8)) - \\
& 432Q_2^4Q_3(5597075Q_4^2Q_5 - 854160Q_2Q_4Q_7 + 8Q_2(-145417Q_5Q_6 + 6685Q_2Q_9))) +
\end{aligned}$$

$$\begin{aligned}
& \frac{(n+1/2)^3}{477757440Q_2^{10}}(31104Q_{10}Q_2^7 - 42944825Q_3^8 + 193106700Q_2Q_3^6Q_4 - \\
& -90039120Q_2^2Q_3^5Q_5 + 30Q_2^2Q_3^4(-8476205Q_4^2 + 1102568Q_2Q_6) - \\
& 4320Q_2^3Q_3^3(-41165Q_4Q_5 + 2312Q_2Q_7) - 9Q_2^4(445825Q_4^4 - 472880Q_2Q_4^2Q_6 + \\
& 64Q_2^2(829Q_6^2 + 1836Q_2Q_7) + 4032Q_2Q_4(-179Q_5^2 + 15Q_2Q_8)) + \\
& 180Q_2^3Q_3^2(532615Q_4^3 - 241224Q_2Q_4Q_6 + 16Q_2(-9352Q_5^2 + 799Q_2Q_8)) - \\
& 144Q_2^4Q_3(392325Q_4^2Q_5 - 51600Q_2Q_4Q_7 + 8Q_2(-8853Q_5Q_6 + 335Q_2Q_9))) + \\
& \frac{(n+1/2)^5}{1194393600Q_2^{10}}(10368Q_{10}Q_2^7 - 66578225Q_3^8 + 272124300Q_2Q_3^6Q_4 - \\
& 112336560Q_2^2Q_3^5Q_5 + 9450Q_2^2Q_3^4(-33775Q_4^2 + 3656Q_2Q_6) - \\
& 151200Q_2^3Q_3^3(-1297Q_4Q_5 + 56Q_2Q_7) - 27Q_2^4(89075Q_4^4 - 83440Q_2Q_4^2Q_6 + \\
& 64Q_2^2(131Q_6^2 + 396Q_2Q_7) + 576Q_2Q_4(-343Q_5^2 + 15Q_2Q_8)) + \\
& 540Q_2^3Q_3^2(188125Q_4^3 - 71400Q_2Q_4Q_6 + 16Q_2(-3052Q_5^2 + 177Q_2Q_8)) - \\
& 432Q_2^4Q_3(118825Q_4^2Q_5 - 11760Q_2Q_4Q_7 + 8Q_2(-2303Q_5Q_6 + 55Q_2Q_9))) \quad (15)
\end{aligned}$$

$$\begin{aligned}
\Lambda_6 = & \frac{-i}{202263389798400Q_2^{12}\sqrt{2Q_2}}(-171460800Q_{12}Q_2^9 + 1714608000Q_{11}Q_2^8Q_3 - \\
& 10268596800Q_{10}Q_2^7Q_3^2 + 970010662775Q_3^{10} + 3772137600Q_{10}Q_2^8Q_4 - \\
& 6262634175525Q_2Q_3^8Q_4 + 13782983196150Q_2^2Q_3^6Q_4^2 - 11954148125850Q_2^3Q_3^4Q_4^3 + \\
& 3449170577475Q_2^4Q_3^2Q_4^4 - 144528059025Q_2^5Q_4^5 + 3352602187200Q_2^2Q_3^7Q_5 - \\
& 12300730092000Q_2^3Q_3^5Q_4Q_5 + 11994129604800Q_2^4Q_3^3Q_4^2Q_5 - 2624788605600Q_2^5Q_3Q_4^3Q_5 + \\
& 2580769643760Q_2^4Q_3^4Q_5^2 - 3453909784416Q_2^5Q_3^2Q_4Q_5^2 + 438440697072Q_2^6Q_4^2Q_5^2 + \\
& +260524397952Q_2^6Q_3Q_5^3 - 1475306441280Q_2^3Q_3^6Q_6 + 4329682610400Q_2^4Q_3^4Q_4Q_6 - \\
& 2865128172480Q_2^5Q_3^2Q_4Q_6 + 233443879200Q_2^6Q_3^4Q_6 - 1660199804928Q_2^5Q_3^3Q_5Q_6 + \\
& 1281705296256Q_2^6Q_3Q_4Q_5Q_6 - 87403857408Q_2^7Q_5^2Q_6 + 231105873600Q_2^6Q_3^2Q_6^2 - \\
& 68412859200Q_2^7Q_4Q_6^2 + 552968700480Q_2^4Q_3^5Q_7 - 1231789749120Q_2^5Q_3^3Q_4Q_7 + \\
& 470726303040Q_2^6Q_3Q_4^2Q_7 + 413953400448Q_2^6Q_3^2Q_5Q_7 - 126242178048Q_2^7Q_4Q_5Q_7 - \\
& 91489305600Q_2^7Q_3Q_6Q_7 + 5619715200Q_2^8Q_7^2 - 175752294480Q_2^5Q_3^4Q_8 + \\
& 271759652640Q_2^6Q_3^2Q_4Q_8 - 39736040400Q_2^7Q_4^2Q_8 - 73378363968Q_2^7Q_3Q_5Q_8 + \\
& 9773265600Q_2^8Q_6Q_8 + 47107126080Q_2^6Q_3^3Q_9 - 43345290240Q_2^7Q_3Q_4Q_9 + 7400248128Q_2^8Q_5Q_9) - \\
& \frac{(n+1/2)^2i}{687970713600Q_2^{12}\sqrt{2Q_2}}(-4551552Q_{12}Q_2^9 + 60279552Q_{11}Q_2^8Q_3 - \\
& 425036160Q_{10}Q_2^7Q_3^2 + 73727194625Q_3^{10} + 116743680Q_{10}Q_2^8Q_4 - \\
& 443649208275Q_2Q_3^8Q_4 + 901144103850Q_2^2Q_3^6Q_4^2 - 711096726150Q_2^3Q_3^4Q_4^3 +
\end{aligned}$$

$$\begin{aligned}
& 182164306725Q_2^4Q_3^2Q_4^4 - 6289615575Q_2^5Q_4^5 + 222467624400Q_2^2Q_3^7Q_5 - \\
& 746418445200Q_2^3Q_3^5Q_4Q_5 + 653423900400Q_2^4Q_3^3Q_4^2Q_5 - 124319674800Q_2^5Q_3Q_4^3Q_5 + \\
& 143980943040Q_2^4Q_3^4Q_5^2 - 169712521920Q_2^5Q_3^2Q_4Q_5^2 + 18188188416Q_2^6Q_4^2Q_5^2 + \\
& 11240861184Q_2^6Q_3Q_5^3 - 91198200240Q_2^3Q_3^6Q_6 + 241513732080Q_2^4Q_3^4Q_4Q_6 - \\
& 140030897040Q_2^5Q_3^2Q_4^2Q_6 + 9200103120Q_2^6Q_4^3Q_6 - 84218693760Q_2^5Q_3^3Q_5Q_6 + \\
& 55248386688Q_2^6Q_3Q_4Q_5Q_6 - 3173043456Q_2^7Q_5^2Q_6 + 10464952896Q_2^6Q_3^2Q_6^2 - \\
& 2403421632Q_2^7Q_4Q_6^2 + 31637744640Q_2^4Q_3^5Q_7 - 62649953280Q_2^5Q_3^3Q_4Q_7 + \\
& 20409822720Q_2^6Q_3Q_4^2Q_7 + 18860532480Q_2^6Q_3^2Q_5Q_7 - 4693344768Q_2^7Q_4Q_5Q_7 - \\
& 3625731072Q_2^7Q_3Q_6Q_7 + 188054784Q_2^8Q_7^2 - 9155635200Q_2^5Q_3^4Q_8 + \\
& 12238024320Q_2^6Q_3^2Q_4Q_8 - 1405278720Q_2^7Q_4^2Q_8 - 2866700160Q_2^7Q_3Q_5Q_8 + \\
& 303295104Q_2^8Q_6Q_8 + 2210705280Q_2^6Q_3^3Q_9 - 1685525760Q_2^7Q_3Q_4Q_9 + 235488384Q_2^8Q_5Q_9) -
\end{aligned}$$

$$\begin{aligned}
& \frac{(n+1/2)^4 i}{20065812480Q_2^{12}\sqrt{2Q_2}} (-66528Q_{12}Q_2^9 + 1245888Q_{11}Q_2^8Q_3 - \\
& 11158560Q_{10}Q_2^7Q_3^2 + 4668804525Q_3^{10} + 2116800Q_{10}Q_2^8Q_4 - \\
& 25898331375Q_2Q_3^8Q_4 + 47959232650Q_2^2Q_3^6Q_4^2 - 33861927750Q_2^3Q_3^4Q_4^3 + \\
& 7454763225Q_2^4Q_3^2Q_4^4 - 184988475Q_2^5Q_4^5 + 11891917800Q_2^2Q_3^7Q_5 - \\
& 36105463800Q_2^3Q_3^5Q_4Q_5 + 27953667000Q_2^4Q_3^3Q_4^2Q_5 - 4457716200Q_2^5Q_3Q_4^3Q_5 + \\
& 6285855240Q_2^4Q_3^4Q_5^2 - 6471756144Q_2^5Q_3^2Q_4Q_5^2 + 565259688Q_2^6Q_4^2Q_5^2 + \\
& 380939328Q_2^6Q_3Q_5^3 - 4375251160Q_2^3Q_3^6Q_6 + 10317018600Q_2^4Q_3^4Q_4Q_6 - \\
& 5113813320Q_2^5Q_3^2Q_4^2Q_6 + 238888440Q_2^6Q_4^3Q_6 - 3203871552Q_2^5Q_3^3Q_5Q_6 + \\
& 1758685824Q_2^6Q_3Q_4Q_5Q_6 - 88566912Q_2^7Q_5^2Q_6 + 335466432Q_2^6Q_3^2Q_6^2 - \\
& 55073088Q_2^7Q_4Q_6^2 + 1351294560Q_2^4Q_3^5Q_7 - 2341442880Q_2^5Q_3^3Q_4Q_7 + \\
& 626542560Q_2^6Q_3Q_4^2Q_7 + 619520832Q_2^6Q_3^2Q_5Q_7 - 123524352Q_2^7Q_4Q_5Q_7 - \\
& 96574464Q_2^7Q_3Q_6Q_7 + 4048704Q_2^8Q_7^2 - 341160120Q_2^5Q_3^4Q_8 + \\
& 386210160Q_2^6Q_3^2Q_4Q_8 - 30837240Q_2^7Q_4^2Q_8 - 78073632Q_2^7Q_3Q_5Q_8 + \\
& 5848416Q_2^8Q_6Q_8 + 70415520Q_2^6Q_3^3Q_9 - 43424640Q_2^7Q_3Q_4Q_9 + 5255712Q_2^8Q_5Q_9) -
\end{aligned}$$

$$\begin{aligned}
& \frac{(n+1/2)^6 i}{300987187200Q_2^{12}\sqrt{2Q_2}} (-72576Q_{12}Q_2^9 + 1886976Q_{11}Q_2^8Q_3 - \\
& 22135680Q_{10}Q_2^7Q_3^2 + 27463538375Q_3^{10} + 2903040Q_{10}Q_2^8Q_4 - \\
& 141448688325Q_2Q_3^8Q_4 + 240655765350Q_2^2Q_3^6Q_4^2 - 152907158250Q_2^3Q_3^4Q_4^3 + \\
& 28724479875Q_2^4Q_3^2Q_4^4 - 413669025Q_2^5Q_4^5 + 59058073200Q_2^2Q_3^7Q_5 - \\
& 164264209200Q_2^3Q_3^5Q_4Q_5 + 113654696400Q_2^4Q_3^3Q_4^2Q_5 - 15166342800Q_2^5Q_3Q_4^3Q_5 +
\end{aligned}$$

$$\begin{aligned}
& 26061194880Q_2^4Q_3^4Q_5^2 - 23876233920Q_2^5Q_3^2Q_4Q_5^2 + 1767189312Q_2^6Q_4^2Q_5^2 + \\
& 1292433408Q_2^6Q_3Q_5^3 - 18902165520Q_2^3Q_3^6Q_6 + 40256773200Q_2^4Q_3^4Q_4Q_6 - \\
& 17116974000Q_2^5Q_3^2Q_4^2Q_6 + 483582960Q_2^6Q_3^4Q_6 - 11384150400Q_2^5Q_3^3Q_5Q_6 + \\
& 5285056896Q_2^6Q_3Q_4Q_5Q_6 - 246903552Q_2^7Q_5^2Q_6 + 992779200Q_2^6Q_3^2Q_6 - \\
& 101860416Q_2^7Q_4Q_6^2 + 4966859520Q_2^4Q_3^5Q_7 - 7661606400Q_2^5Q_3^3Q_4Q_7 + \\
& 1683037440Q_2^6Q_3Q_4^2Q_7 + 1861574400Q_2^6Q_3^2Q_5Q_7 - 316141056Q_2^7Q_4Q_5Q_7 - \\
& 235146240Q_2^7Q_3Q_6Q_7 + 8895744Q_2^8Q_7^2 - 1042372800Q_2^5Q_3^4Q_8 + \\
& 1016789760Q_2^6Q_3^2Q_4Q_8 - 52436160Q_2^7Q_4^2Q_8 - 189060480Q_2^7Q_3Q_5Q_8 + \\
& 9217152Q_2^8Q_6Q_8 + 175190400Q_2^6Q_3^3Q_9 - 87816960Q_2^7Q_3Q_4Q_9 + 10378368Q_2^8Q_5Q_9) \quad (16)
\end{aligned}$$

All six WKB corrections printed in MATHEMATICA are available from the author in electronic form upon request.

2 Appendix 2: QNMs of a 4-dimensional Schwarzschild black hole

"The potential" $Q(x)$ in case of a Schwarzschild black hole has the form

$$Q(x) = \omega^2 - \left(1 - \frac{1}{r}\right) \left(\frac{l(l+1)}{r^2} + \frac{1-s^2}{r^3}\right), \quad (17)$$

where $s = 0$ corresponds to scalar perturbations, $s = 1/2$ - neutrino perturbations, $s = 1$ - electromagnetic perturbations, $s = 3/2$ - gravitino perturbations, $s = 2$ - gravitational perturbations. The quasinormal frequencies at 3th and 6th WKB orders and in comparison with numerical results [23] are presented in the tables I-V.

multipole, overtone	numerical	3th order WKB	6th order WKB
$l = 0, n = 0$	0.1105 - 0.1049 <i>i</i>	0.1046 - 0.1152 <i>i</i>	0.1105 - 0.1008 <i>i</i>
$l = 1, n = 0$	0.2929 - 0.0977 <i>i</i>	0.2911 - 0.0980 <i>i</i>	0.2929 - 0.0977 <i>i</i>
$l = 1, n = 1$	0.2645 - 0.3063 <i>i</i>	0.2622 - 0.3074 <i>i</i>	0.2645 - 0.3065 <i>i</i>
$l = 2, n = 0$	0.4836 - 0.0968 <i>i</i>	0.4832 - 0.0968 <i>i</i>	0.4836 - 0.0968 <i>i</i>
$l = 2, n = 1$	0.4639 - 0.2956 <i>i</i>	0.4632 - 0.2958 <i>i</i>	0.4638 - 0.2956 <i>i</i>
$l = 2, n = 2$	0.4305 - 0.5086 <i>i</i>	0.4317 - 0.5034 <i>i</i>	0.4304 - 0.5087 <i>i</i>

Table I. Schwarzschild QN frequencies for scalar perturbations ($s = 0$).

multipole, overtone	numerical	3th order WKB	6th order WKB
$l = 1, n = 0$	—	$0.2803 - 0.0969i$	$0.2822 - 0.0967i$
$l = 1, n = 1$	—	$0.2500 - 0.3049i$	$0.2525 - 0.3040i$
$l = 2, n = 0$	—	$0.4768 - 0.9639i$	$0.4772 - 0.0963i$
$l = 2, n = 1$	—	$0.4565 - 0.2947i$	$0.4571 - 0.2945i$
$l = 2, n = 2$	—	$0.4244 - 0.5016i$	$0.4231 - 0.5070i$
$l = 3, n = 0$	—	$0.6706 - 0.0963i$	$0.6708 - 0.0963i$
$l = 3, n = 1$	—	$0.6557 - 0.2917i$	$0.6560 - 0.2917i$
$l = 3, n = 2$	—	$0.6299 - 0.4931i$	$0.6286 - 0.4950i$
$l = 3, n = 3$	—	$0.5970 - 0.6997i$	$0.5932 - 0.7102i$

Table II. Schwarzschild QN frequencies for Dirac perturbations ($s = 1/2$).

multipole, overtone	numerical	3th order WKB	6th order WKB
$l = 1, n = 0$	$0.2483 - 0.0925i$	$0.2459 - 0.0931i$	$0.2482 - 0.0926i$
$l = 1, n = 1$	$0.2145 - 0.2937i$	$0.2113 - 0.2958i$	$0.2143 - 0.2941i$
$l = 2, n = 0$	$0.4576 - 0.0950i$	$0.4571 - 0.0951i$	$0.4576 - 0.0950i$
$l = 2, n = 1$	$0.4365 - 0.2907i$	$0.4358 - 0.2910i$	$0.4365 - 0.2907i$
$l = 2, n = 2$	$0.4012 - 0.5016i$	$0.4023 - 0.4959i$	$0.4009 - 0.5017i$
$l = 3, n = 0$	$0.6569 - 0.0956i$	$0.6567 - 0.0956i$	$0.6569 - 0.0956i$
$l = 3, n = 1$	$0.6417 - 0.2897i$	$0.6415 - 0.2898i$	$0.6417 - 0.2897i$
$l = 3, n = 2$	$0.6138 - 0.4921i$	$0.6151 - 0.4901i$	$0.6138 - 0.4921i$
$l = 3, n = 3$	$0.5779 - 0.7063i$	$0.5814 - 0.6955i$	$0.5775 - 0.7065i$

Table III. Schwarzschild QN frequencies for electromagnetic perturbations ($s = 1$).

multipole, overtone	numerical	3th order WKB	6th order WKB
$l = 1, n = 0$	—	$0.1817 - 0.0866i$	$0.1739 - 0.08357i$
$l = 1, n = 1$	—	$0.1354 - 0.2812i$	$0.1198 - 0.2813i$
$l = 2, n = 0$	—	$0.4231 - 0.926i$	$0.4236 - 0.0925i$
$l = 2, n = 1$	—	$0.4000 - 0.2842i$	$0.4007 - 0.2838i$
$l = 2, n = 2$	—	$0.3636 - 0.4853i$	$0.3618 - 0.4919i$
$l = 3, n = 0$	—	$0.6332 - 0.0945i$	$0.6333 - 0.0944i$
$l = 3, n = 1$	—	$0.6173 - 0.2864i$	$0.6175 - 0.2863i$
$l = 3, n = 2$	—	$0.5898 - 0.4846i$	$0.5884 - 0.4868i$
$l = 3, n = 3$	—	$0.5547 - 0.6882i$	$0.5505 - 0.7000i$

Table IV. Schwarzschild QN frequencies for gravitino perturbations ($s = 3/2$).

multipole, overtone	numerical	3th order WKB	6th order WKB
$l = 2, n = 0$	$0.3737 - 0.0890i$	$0.3732 - 0.0892i$	$0.3736 - 0.0890i$
$l = 2, n = 1$	$0.3467 - 0.2739i$	$0.3460 - 0.2749i$	$0.3463 - 0.2735i$
$l = 2, n = 2$	$0.3011 - 0.4783i$	$0.3029 - 0.4711i$	$0.2985 - 0.4776i$
$l = 3, n = 0$	$0.5994 - 0.0927i$	$0.5993 - 0.0927i$	$0.5994 - 0.0927i$
$l = 3, n = 1$	$0.5826 - 0.2813i$	$0.5824 - 0.2814i$	$0.5826 - 0.2813i$
$l = 3, n = 2$	$0.5517 - 0.4791i$	$0.5532 - 0.4767i$	$0.5516 - 0.4790i$
$l = 3, n = 3$	$0.5120 - 0.6903i$	$0.5157 - 0.6774i$	$0.5111 - 0.6905i$
$l = 4, n = 0$	$0.8092 - 0.0942i$	$0.8091 - 0.0942i$	$0.8092 - 0.0942i$
$l = 4, n = 1$	$0.7966 - 0.2843i$	$0.7965 - 0.2844i$	$0.7966 - 0.2843i$
$l = 4, n = 2$	$0.7727 - 0.4799i$	$0.7736 - 0.4790i$	$0.7727 - 0.4799i$
$l = 4, n = 3$	$0.7398 - 0.6839i$	$0.7433 - 0.6783i$	$0.7397 - 0.6839i$
$l = 4, n = 4$	$0.7015 - 0.8982i$	$0.7072 - 0.8813i$	$0.7006 - 0.8985i$

Table V. Schwarzschild QN frequencies for graviton perturbations ($s = 2$).