

Surface terms and the Gauss-Bonnet Hamiltonian

Antonio Padilla*

*Theoretical Physics, Department of Physics,
University of Oxford, 1 Keble Road, Oxford OX1 3NP, U.K.*

Abstract

We derive the gravitational Hamiltonian starting from the Gauss-Bonnet action, keeping track of all surface terms. This is done using the language of orthonormal frames and forms to keep things as tidy as possible. The surface terms in the Hamiltonian give a remarkably simple expression for the total energy of a spacetime. This expression is consistent with energy expressions found in hep-th/0212292. However, we can apply our results whatever the choice of background and whatever the symmetries of the spacetime.

arXiv:gr-qc/0303082v1 20 Mar 2003

*a.padilla1@physics.ox.ac.uk

1 Introduction

The energy of gravitational systems has attracted much interest down the years. In particular, for Einstein gravity,

$$S = \kappa \int R - 2\Lambda, \quad (1)$$

expressions were found for the energy of asymptotically flat [1] and asymptotically AdS [2] spacetimes. Some time later, Hawking and Horowitz [3] gave a general derivation of the gravitational Hamiltonian, keeping careful track of surface terms. When evaluated on a solution, this Hamiltonian gave an expression for the total energy. This agreed with the earlier expressions found in [1,2]. However, this new expression could be applied regardless of the asymptotic behaviour of the solution.

Recently, Deser and Tekin have found energy expressions for higher derivative gravities [4,5]. This initial work has focused on test spacetimes that are *asymptotically* maximally symmetric, with background spacetimes (vacua), defined to be maximally symmetric *everywhere*. In this paper, we will perform the analogue of Hawking and Horowitz's calculation by deriving the Hamiltonian for Gauss-Bonnet (GB) gravity: a special, but important class of higher derivative gravities. We will obtain an expression for the energy by once again evaluating this Hamiltonian on a solution. As with [3], our expression will be consistent with earlier results [4,5], but can be easily applied even if the background is not maximally symmetric. This greater flexibility allows for a more natural choice of vacua in some cases, as we will illustrate with an example.

Before going any further, let us briefly mention what GB gravity is, and why it is important. GB gravity is given by the addition of the Gauss-Bonnet invariant to the Einstein-Hilbert action (1),

$$S = \kappa \int R - 2\Lambda + \alpha (R^2 - 4R_{AB}R^{AB} + R_{ABCD}R^{ABCD}). \quad (2)$$

In $D = 4$ dimensions, the GB term is a topological invariant and does not enter the dynamics. This ceases to be the case in $D > 4$ dimensions. Furthermore, in Einstein gravity, the vacuum field equations are given by a linear combination of the Einstein tensor and the metric. In four dimensions, this is the most general combination of tensors that satisfies the following conditions [6]:

- it is symmetric.
- it depends only on the metric and its first two derivatives.
- it has vanishing divergence.
- it is linear in the second derivatives of the metric¹.

¹In $D = 4$, this condition is actually implied by the other three.

However, it has become unfashionable to restrict ourselves to just four dimensions. If we go to $D = 5$ or 6 , it turns out that these conditions are satisfied by a linear combination of the metric, the Einstein tensor, and the *Lovelock tensor* [6, 7]. The Lovelock tensor arises from the variation of the Gauss-Bonnet term in the above action (2). In this sense, GB gravity is the natural generalisation of Einstein gravity to higher dimensions.

Perhaps the most compelling reason to study GB gravity is its appearance in String theory. Consider the slope (ie. α') expansion for the heterotic string. At lowest order, it is given by the Einstein-Hilbert action (1). The first stringy corrections give rise to quadratic curvature terms [8, 9]. For this effective theory to be ghost-free, the quadratic curvatures must be in the combination given in the GB action (2) [10, 11]. This link to String theory has generated a lot of research into braneworlds in GB gravity [12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28]. Braneworld models are themselves inspired by String theory [29, 30, 31] so it is natural to ask what effect any stringy corrections might have on their cosmology. From a holographic point of view, we might expect such higher curvature terms in the bulk to correspond to next to leading order corrections in the $1/N$ expansion of the CFT on the boundary/brane [21]. Calculating the GB Hamiltonian will allow us to investigate the GB version of “exact” braneworld holography [32, 33]. This will be discussed in a future article [34].

The rest of this paper is organised as follows: in section 2 we will give a detailed derivation of the GB Hamiltonian. We will use orthonormal frames and differential forms to keep things as tidy as possible. Some readers may wish to ignore the details of this derivation and go directly to the energy expression at the end of the section. In section 3 we will show that this expression is consistent with existing literature [35, 4, 5]. We will illustrate the flexibility of our approach with a special example in section 3.2. Section 4 contains some concluding remarks. There are two appendices: appendix A contains specific proofs, whereas appendix B gives a summary of the notation used in this paper. Appendix B should be of use to anyone skipping the details in various places.

2 Derivation of the Gauss-Bonnet Hamiltonian

2.1 The action

2.1.1 The bulk

The Gauss-Bonnet action (2) is most elegantly written in terms of differential forms. Suppose our D -dimensional spacetime, \mathcal{M} , has metric,

$$g = \eta_{AB} E^A \times E^B \tag{3}$$

where $\{E^A\}$ is an orthonormal basis of 1-forms, and indices are raised/lowered using $\eta_{AB} = \text{diag}(-1, +1, \dots, +1)$. We write $\{X_A\}$ for the dual basis of vectors.

We will find it useful to define the following forms,

$$e_{A_1 \dots A_m} = \frac{1}{(D-m)!} \epsilon_{A_1 \dots A_m A_{m+1} \dots A_D} E^{A_{m+1}} \wedge \dots \wedge E^{A_D}, \quad (4)$$

where $\epsilon_{A_1 \dots A_D}$ is the totally antisymmetric tensor with $\epsilon_{0 \dots (D-1)} = 1$. Notice that the scalar D -form, e , is the volume measure on \mathcal{M} . We assume zero torsion so that the connection 1-form, $\omega^A{}_B$, is given by

$$0 = dE^A + \omega^A{}_B \wedge E^B. \quad (5)$$

Meanwhile, the curvature 2-form, $\Omega^A{}_B$, can be expressed in terms of the connection 1-form as follows,

$$\Omega^A{}_B = d\omega^A{}_B + \omega^A{}_C \wedge \omega^C{}_B = \frac{1}{2} R^A{}_{BCD} E^C \wedge E^D. \quad (6)$$

The right hand equation above gives the Riemann tensor, $R^A{}_{BCD}$. The Ricci tensor is then defined by $R_{BD} = R^C{}_{BCD}$ and the Ricci scalar by $R = \eta^{AB} R_{AB}$. We now rewrite the GB action (2), in terms of these newly defined forms,

$$S = \kappa \int_{\mathcal{M}} -2\Lambda e + \Omega^{AB} \wedge e_{AB} + \alpha \Omega^{AB} \wedge \Omega^{CD} \wedge e_{ABCD}. \quad (7)$$

Given the right hand equation in (6), and the useful identity [36],

$$E^B \wedge e_{A_1 \dots A_m} = \delta_{A_m}^B e_{A_1 \dots A_{m-1}} - \delta_{A_{m-1}}^B e_{A_1 \dots A_{m-2} A_m} + \dots + (-1)^{m-1} \delta_{A_1}^B e_{A_2 \dots A_m}, \quad (8)$$

it can be checked that the actions (2) and (7) are indeed the same.

2.1.2 The boundary

If \mathcal{M} has a boundary, $\partial\mathcal{M}$, we need to define boundary conditions on $\partial\mathcal{M}$. We usually demand that the geometry of the boundary is fixed. If we are near $\partial\mathcal{M}$, we can go to Gaussian normal coordinates, (y, x^α) , and write the metric as

$$g = \pm dy^2 + \eta_{\alpha\beta} E^\alpha \times E^\beta. \quad (9)$$

where $E^\alpha = E^\alpha{}_\beta(y, x^\gamma) dx^\beta$. Here the boundary is given by $y = 0$, so the induced metric on $\partial\mathcal{M}$ is given by

$$h = \eta_{\alpha\beta} E_0^\alpha \times E_0^\beta, \quad E_0^\alpha = E^\alpha{}_\beta(0, x^\gamma) dx^\beta. \quad (10)$$

The $\pm dy^2$ in (9) corresponds to a spacelike (timelike) boundary. From now on, we write E^y for the extension of dy into the bulk. The dual vector X_y is the extension of the inward pointing normal vector $\partial/\partial y$.

The boundary condition now states that

$$\delta E^\alpha = \delta \omega^{\alpha\beta} = 0 \quad \text{on } \partial\mathcal{M} \quad (11)$$

Notice that $\delta E^y = 0$ is automatic on $\partial\mathcal{M}$, but crucially we are allowed $\delta\omega^{y\alpha} \neq 0$. If we vary the action (7) with respect to the connection, we get [36],

$$\delta_\omega S = \kappa \int_{\partial\mathcal{M}} \delta\omega^{AB} \wedge e_{AB} + 2\alpha\delta\omega^{AB} \wedge \Omega^{CD} \wedge e_{ABCD}. \quad (12)$$

Since $\delta\omega^{y\alpha}$ is allowed to be non-zero, we see that $\delta_\omega S$ does not necessarily vanish on shell, which violates the action principle. The problem is that our action and boundary conditions are inconsistent. To cure this, we need to add a boundary integral. In Einstein gravity, this is the well known Gibbons-Hawking term [37]. The generalisation to higher derivative gravities was worked out by Myers [36]. For GB gravity it is given by

$$S_{\text{boundary}} = -\kappa \int_{\partial\mathcal{M}} \theta^{AB} \wedge e_{AB} + 2\alpha\theta^{AB} \wedge \left(\Omega^{CD} - \frac{2}{3}\theta^C{}_E \wedge \theta^{ED} \right) \wedge e_{ABCD}, \quad (13)$$

where θ^{AB} is the second fundamental form [38, 39], defined as follows. Consider the product metric which agrees with g on the boundary, $\partial\mathcal{M}$,

$$g_0 = \pm dy^2 + \eta_{\alpha\beta} E_0^\alpha \times E_0^\beta. \quad (14)$$

This metric gives a new connection, ω_0^{AB} , and we define the second fundamental form as

$$\theta^{AB} = \omega^{AB} - \omega_0^{AB}. \quad (15)$$

It is clear from this definition that $\theta^{y\alpha} = \omega^{y\alpha}$ and $\theta^{\alpha\beta} = 0$ on $\partial\mathcal{M}$. Furthermore, we can relate $\theta^{y\alpha}$ to the more familiar extrinsic curvature, $K^\alpha{}_\beta$ of $\partial\mathcal{M}$ in \mathcal{M} ,

$$\theta^{y\alpha} = -(\pm)K^\alpha = -(\pm)K^\alpha{}_\beta E^\beta \quad \text{on } \partial\mathcal{M}. \quad (16)$$

Again the (\pm) corresponds to a spacelike (timelike) boundary.

We conclude that a fully consistent action for Gauss-Bonnet gravity is given by,

$$S = S_{\text{bulk}} + S_{\text{boundary}} \quad (17)$$

where S_{bulk} and S_{boundary} are given by equations (7) and (13) respectively.

2.1.3 The background

If \mathcal{M} were spatially compact, equation (17) would give a well defined action. However, for spatially noncompact spacetimes, this action diverges [3]. To get round this we need to choose a reference background, $\bar{\mathcal{M}}$ with metric \bar{g} . This background should be a static solution to the field equations [3], but does not have to be maximally symmetric. The boundary conditions are unchanged which means that $\partial\bar{\mathcal{M}}$ should have the same geometry as $\partial\mathcal{M}$. We can then define the background action, \bar{S} , in the same way as before. The physical action is the difference,

$$I = \Delta S = S - \bar{S}. \quad (18)$$

In the Hamiltonian picture, this background can be thought of as defining a background energy or zero energy solution. For example, for an asymptotically AdS spacetime, we would probably choose the background to be pure AdS, but we do not have to. Any asymptotically AdS black hole spacetime would be equally valid.

2.2 Splitting space and time

Although the physical Hamiltonian will be constructed from the action (18), it is clear that it will just be the difference of the Hamiltonian constructed from S and that constructed from \bar{S} [3]. For the time being we will concentrate on the former.

2.2.1 Foliations of \mathcal{M}

To proceed, we need to deconstruct the spacetime \mathcal{M} by separating space from time in the following way. First, we choose a timelike vector field, $\partial/\partial t$. Now introduce a family of spacelike hypersurfaces $\{\Sigma_t\}$ labelled by the parameter t . This family is a foliation of the full spacetime. We assume that the hypersurfaces have no inner boundaries and do not intersect each other. They meet the timelike part of the boundary (call this B) orthogonally, and in the far past/future, they coincide with the spacelike part of the boundary (call this Σ_∞). Therefore the total boundary, $\partial\mathcal{M} = B \cup \Sigma_\infty$.

We can write the metric for \mathcal{M} in ADM form [1],

$$g = -N^2 dt^2 + \gamma_{ab}(t, x^a)(dx^a + N^a dt)(dx^b + N^b dt), \quad (19)$$

where N is the lapse function, N^a the shift vector, and $\gamma_{ab}(t, x^a)$ the induced metric on Σ_t . It is natural to choose the following orthonormal basis of 1-forms,

$$E^\perp = N dt, \quad E^a = E^a{}_b(dx^b + N^b dt) \quad (20)$$

where $\delta_{ab}E^a{}_c E^b{}_d = \gamma_{cd}$. We would like to emphasize some notation at this point. Lowercase latin indices label components in Σ_t , whereas uppercase latin indices label components in \mathcal{M} . For example, $\{E^a\}$ is an orthonormal basis for Σ_t whereas $\{E^A\} = E^\perp \cup \{E^a\}$ is an orthonormal basis for \mathcal{M} .

The dual basis of vectors is given by [40]

$$X_\perp = \frac{1}{N} \left(\frac{\partial}{\partial t} - N^a \frac{\partial}{\partial x^a} \right), \quad X_a = E_a{}^b \frac{\partial}{\partial x^b}. \quad (21)$$

X_\perp is the vector normal to Σ_t , and is not necessarily equal, or even parallel to $\partial/\partial t$. We should also note that X_\perp is tangent to B , as Σ_t and B are orthogonal.

Normally, the next step is to use the Gauss-Codazzi equations [41] to rewrite the bulk part of the action (see for example [3]). This has infact been done for GB gravity [42] although the contribution from surface terms was ignored. In this paper we are using the language of orthonormal frames and differential forms. We therefore need to know how to translate the Gauss-Codazzi equations into this language. This is explained in [43], although we will review the main points presently.

The Gauss-Codazzi equations describe the decomposition of the bulk Riemann tensor into spatial tensors defined on Σ_t . In the language of forms, it is the curvature 2-form that we wish to decompose. We start by decomposing the connection, ω^{AB} .

$$\omega^{\perp a} = -H^a + a^a E^\perp, \quad (22)$$

$$\omega^{ab} = \tilde{\omega}^{ab} + \tilde{l}^{ab} E^\perp. \quad (23)$$

Here, we have two 0-forms: a vector, a^a , and an antisymmetric tensor, \tilde{l}^{ab} . The 1-form $H^a = H^a_b E^b$, where H^a_b is the extrinsic curvature of Σ_t in \mathcal{M} . The other 1-form, $\tilde{\omega}^{ab}$, is the connection for the induced metric $\gamma = \delta_{ab} E^a \wedge E^b$. All four forms live entirely in Σ_t and know nothing about E^\perp .

We will also need to know the decomposition of the exterior derivative. For a general p -form, A , in \mathcal{M} , we write

$$A = \tilde{A} + E^\perp \wedge A_\perp, \quad (24)$$

where \tilde{A} and A_\perp are p and $(p - 1)$ -forms respectively, living entirely on Σ_t . The exterior derivative is given by

$$dA = \tilde{d}\tilde{A} + E^\perp \wedge \left[\mathbb{S}_\perp \tilde{A} - \frac{1}{N} \tilde{d}(N A_\perp) \right]. \quad (25)$$

A couple of things need to be explained here. Firstly, anything with a tilde is intrinsic to Σ_t , as opposed to \mathcal{M} . So for example, in (25), d is the exterior derivative on \mathcal{M} , whereas \tilde{d} is the exterior derivative on Σ_t .

We also need to define \mathbb{S}_\perp . For a *scalar* p -form, A , the Lie derivative with respect to the spacetime vector Y satisfies

$$\mathcal{L}_Y A = d(i_Y A) + i_Y(dA), \quad (26)$$

where $i_Y A$ is the *interior product* of Y and A [44]. For \tilde{A} living entirely on Σ_t , we can define the *projection* of $\mathcal{L}_Y \tilde{A}$:

$$\mathbb{S}_Y A = P \left[\mathcal{L}_Y \tilde{A} \right] = P \left[d(i_Y \tilde{A}) + i_Y(d\tilde{A}) \right]. \quad (27)$$

Here, the projection operator P simply takes the part that does not contain E^\perp . If Y is a spatial vector, $\mathbb{S}_Y \tilde{A}$ is just the intrinsic Lie derivative on Σ_t . However, we are interested in \mathbb{S}_\perp for which $Y = X_\perp$, so we need this more obscure/confusing definition.

The operator \mathbb{S}_\perp can also act on tensor valued forms. Normally, Lie derivatives recognise the tensor structure of forms. However, by definition, \mathbb{S}_\perp ignores it and acts on each component as if it were a scalar. So for example,

$$\mathbb{S}_\perp H^a = P \left[d(i_\perp H^a) + i_\perp(dH^a) \right] \quad (28)$$

$$\mathbb{S}_\perp \tilde{\omega}^{ab} = P \left[d(i_\perp \tilde{\omega}^{ab}) + i_\perp(d\tilde{\omega}^{ab}) \right] \quad (29)$$

We shall now give the frame-form version of the Gauss-Codazzi equations. As we said earlier, this is the decomposition of the curvature 2-form.

$$\Omega^{\perp a} = -\tilde{\nabla} H^a + E^\perp \wedge \left[-\mathbb{S}_\perp H^a - \frac{1}{N} \tilde{\nabla}(N a^a) - \tilde{l}^{ab} H_b \right] \quad (30)$$

$$\Omega^{ab} = \tilde{\Omega}^{ab} + H^a \wedge H^b + E^\perp \wedge \left[\mathbb{S}_\perp \tilde{\omega}^{ab} - \frac{1}{N} \tilde{\nabla}(N \tilde{l}^{ab}) + H^a a^b - H^b a^a \right] \quad (31)$$

where $\tilde{\Omega}^{ab} = \tilde{d}\tilde{\omega}^{ab} + \tilde{\omega}^a_c \wedge \tilde{\omega}^{cb}$ is the curvature 2-form for Σ_t . The operator $\tilde{\nabla}$ is the *covariant exterior derivative* (on Σ_t). It acts on an arbitrary tensor valued p -form, $A^{a_1 \dots a_n}_{b_1 \dots b_m}$, in the following way.

$$\begin{aligned} \tilde{\nabla} A^{a_1 \dots a_n}_{b_1 \dots b_m} &= \tilde{d} A^{a_1 \dots a_n}_{b_1 \dots b_m} + \sum_{i=1}^n \tilde{\omega}^{a_i}_c \wedge A^{\dots a_{i-1} c a_{i+1} \dots}_{b_1 \dots b_m} \\ &\quad - \sum_{j=1}^m \tilde{\omega}^c_{b_j} \wedge A^{a_1 \dots a_n}_{\dots b_{j-1} c b_{j+1} \dots} \end{aligned} \quad (32)$$

Note that $\tilde{\nabla}$ reduces to \tilde{d} when acting on scalars.

In principle we could also decompose the torsion 2-form (see [43]). However, we have set torsion to zero, which means that every component of the torsion decomposition must vanish. This gives the following conditions:

$$H^{ab} = H^{ba}, \quad (33)$$

$$a_a E^a = \frac{\tilde{d}N}{N}, \quad (34)$$

$$\tilde{\nabla} E^a = 0 \quad (35)$$

$$\$_\perp E^a = -H^a - \tilde{l}^a_b E^b. \quad (36)$$

For a far more detailed explanation of everything in this section, refer to [43].

2.2.2 Foliations of B

Since we intend to keep careful track of surface terms, we will need a foliation of B , as well as \mathcal{M} . On B , the foliation is given by the family of surfaces $\{S_t\}$. For a given value of t , S_t is the intersection of B and Σ_t .

We need to understand how bulk quantities project on to B and then S_t . Near B the metric can be written in Gaussian normal coordinates,

$$g = dz^2 + \eta_{\mu\nu} E^\mu \times E^\nu. \quad (37)$$

In analogy with section 2.1.2, we write E^z for the extension of dz into the bulk, and X_z for the extension of the inward pointing normal $\partial/\partial z$. Notice that we are using greek indices, μ, ν etc to label components in B .

Near B , we can decompose the bulk connection and bulk curvature form into well defined forms on B .

$$\omega^{z\mu} = K^\mu, \quad (38)$$

$$\omega^{\mu\nu} = \omega_0^{\mu\nu} + l_0^{\mu\nu} E^z, \quad (39)$$

$$\Omega^{z\mu} = \nabla_0 K^\mu + E^z \wedge \{\dots\}, \quad (40)$$

$$\Omega^{\mu\nu} = \Omega_0^{\mu\nu} - K^\mu \wedge K^\nu + E^z \wedge \{\dots\}. \quad (41)$$

Here, the suffix “0” labels anything intrinsic to B (in analogy with tilde in the last section). The 1-form, $K^\mu = K^\mu_\nu E^\nu$, where K^μ_ν is the extrinsic curvature of B in \mathcal{M} .

The slight differences in signs compared with equations (22) to (31), are due to the fact that B is timelike whereas Σ_t is spacelike. We will not care what is contained in $\{\dots\}$ as the integration of E^z over B is zero.

We are now ready to project on to S_t . We will use latin indices like i, j, k etc to label components in S_t . Furthermore, anything wearing a hat is intrinsic to S_t . Using equations (22), (23), (38) and (39), we can deduce the decomposition of the connection.

$$\omega^{\perp z} = b_i E^i + c E^\perp \quad (42)$$

$$\omega^{\perp i} = -\widehat{H}^i + a^i E^\perp + b^i E^z \quad (43)$$

$$\omega^{zi} = \widehat{K}^i + b^i E^\perp \quad (44)$$

$$\omega^{ij} = \omega^{ij} + \tilde{l}^{ij} E^\perp + l_0^{ij} E^z \quad (45)$$

where $\widehat{\omega}^{ij}$ is the connection on S_t . The 0-forms c and b_i are given by,

$$c = a^z = -K^\perp, \quad (46)$$

$$b^i = -H^{zi} = -K^{\perp i} = l_0^{\perp i} = \tilde{l}^{zi}, \quad (47)$$

where we have used the symmetry of H^{ab} and $K^{\mu\nu}$. We have also defined,

$$\widehat{H}^i = H^i_j E^j, \quad (48)$$

$$\widehat{K}^i = K^i_j E^j. \quad (49)$$

Here, we should interpret H^i_j and K^i_j as the extrinsic curvatures of S_t in B and Σ_t respectively.

To decompose the curvature form, we need to introduce the operator $\widehat{\$}_\perp$. This is the analogue of $\$_\perp$ on B . More precisely, if \widehat{A} is a p -form living on S_t ,

$$\widehat{\$}_\perp \widehat{A} = P \left[d_0 \left(i_\perp \widehat{A} \right) + i_\perp \left(d_0 \widehat{A} \right) \right] \quad (50)$$

where d_0 is the exterior derivative on B . $\$_\perp$ and $\widehat{\$}_\perp$ can be related to one another as we will now explain. Suppose \tilde{A} is an arbitrary p -form in Σ_t . Near S_t , we can write $\tilde{A} = \widehat{A} + A_z \wedge E^z$, where \widehat{A} and A_z are p and $(p-1)$ -forms respectively, living on S_t . By the proof given in appendix A, we can say

$$\$_\perp \tilde{A} = \widehat{\$}_\perp \widehat{A} + E^z \wedge \{\dots\}. \quad (51)$$

We are nearly ready to decompose the curvature form. We will use equations (30), (31), (40) and (41), along with the following results,

$$\$_\perp H^z = -\widehat{\$}_\perp (b_i E^i) + E^z \{\dots\}. \quad (52)$$

$$\$_\perp \tilde{\omega}^{zi} = \widehat{\$}_\perp \widehat{K}^i + E^z \{\dots\}. \quad (53)$$

$$\tilde{\nabla} H^z = -\widehat{d} (b_i E^i) - \widehat{H}^i \wedge \widehat{K}_i + E^z \{\dots\} \quad (54)$$

$$\frac{1}{N} \tilde{\nabla} (N a^z) = \frac{1}{N} \widehat{d} (N c) + a^i \widehat{K}_i + E^z \{\dots\} \quad (55)$$

$$\frac{1}{N} \tilde{\nabla} (N \tilde{\omega}^{zi}) = \frac{1}{N} \widehat{\nabla} (N b^i) - \widehat{K}_j \tilde{l}^{ij} + E^z \{\dots\} \quad (56)$$

where $\widehat{\nabla}$ is the covariant exterior derivative on S_t . Recall that this reduces to the exterior derivative, \widehat{d} , when acting on scalars. Equations (52) and (53) are examples of equation (51). The proofs of equations (54), (55) and (56) are very similar to one another. As an example, equation (56) is proven in appendix A.

The curvature decomposition is given by,

$$\begin{aligned}\Omega^{\perp z} &= \widehat{d}(b_i E^i) + \widehat{H}^i \wedge \widehat{K}_i + E^z \wedge \{\dots\} \\ &\quad + E^\perp \wedge \left[\widehat{\$}_\perp(b_i E^i) - \frac{1}{N} \widehat{d}(Nc) - a^i \widehat{K}_i - b^i \widehat{H}_i \right]\end{aligned}\quad (57)$$

$$\begin{aligned}\Omega^{\perp i} &= -\widehat{\nabla} \widehat{H}^i + b_j E^j \wedge \widehat{K}^i + E^z \wedge \{\dots\} \\ &\quad + E^\perp \wedge \left[-\widehat{\$}_\perp \widehat{H}^i - \frac{1}{N} \widehat{\nabla}(Na^i) - \tilde{l}^{ij} \widehat{H}_j + c \widehat{K}^i - b^i b_j E^j \right]\end{aligned}\quad (58)$$

$$\begin{aligned}\Omega^{zi} &= \widehat{\nabla} \widehat{K}^i - b_j E^j \wedge \widehat{H}^i + E^z \wedge \{\dots\} \\ &\quad + E^\perp \wedge \left[\widehat{\$}_\perp \widehat{K}^i - \frac{1}{N} \widehat{\nabla}(Nb^i) + \widehat{K}_j \tilde{l}^{ij} - b_j E^j a^i - c \widehat{H}^i \right]\end{aligned}\quad (59)$$

$$\begin{aligned}\Omega^{ij} &= \widehat{\Omega}^{ij} + \widehat{H}^i \wedge \widehat{H}^j - \widehat{K}^i \wedge \widehat{K}^j + E^z \wedge \{\dots\} \\ &\quad + E^\perp \wedge \left[\widehat{\$}_\perp \widehat{\omega}^{ij} - \frac{1}{N} \widehat{\nabla}(N \tilde{l}^{ij}) + 2 \widehat{H}^{[i} a^{j]} + 2 \widehat{K}^{[i} b^{j]} \right]\end{aligned}\quad (60)$$

As long as we are near B , these are the frame-form version of the Gauss-Codazzi equations for a hypersurface of codimension two [45].

We have introduced a huge amount of notation in this section so we remind the reader that a summary of this can be found in appendix B.

2.3 The Hamiltonian

We are now ready to start calculating the Hamiltonian. However, we will find it convenient to continue working with the action until virtually the bitter end. When our action finally has the desired form we will switch to the Hamiltonian picture, and give an expression for the gravitational energy of a solution.

We will start with the bulk part of the action (7). Our aim is to write it so that it contains no derivatives of the lapse function or the shift vector. This is because these are ignorable coordinates, and should behave like Lagrange multipliers. They will be paired with the Hamiltonian and momentum constraints respectively, as is the case in Einstein gravity [3]. We also want to eliminate terms like $\$_\perp H^a$, which contain second time derivatives of the canonical variable E^a . We will need to use integration by parts to achieve these aims. This means that the bulk action (7) will contribute surface terms. In summary, we expect to write (7) as

$$S_{\text{bulk}} = S_{\text{bulk}}^* + S_{\text{leftover}}, \quad (61)$$

where S_{leftover} are the leftover surface terms, and

$$S_{\text{bulk}}^* = \kappa \int dt \int_{\Sigma_t} \pi_a \wedge \$_{\frac{\partial}{\partial t}} E^a - N \mathcal{H} - N^a \mathcal{H}_a. \quad (62)$$

Here π_a is the momentum conjugate to E^a , and \mathcal{H} and \mathcal{H}_a are the Hamiltonian and momentum constraints respectively.

At this stage, the surface part of the action is given by equation (13). We can split this into two parts,

$$S_{\text{boundary}} = S_B + S_\infty, \quad (63)$$

where S_B contains the integration over B and S_∞ the integration over Σ_∞ . In section 2.5, we will group these terms with S_{leftover} . This will give us a modified boundary term

$$S_{\text{boundary}}^* = S_{\text{boundary}} + S_{\text{leftover}}. \quad (64)$$

which will be closely related to the gravitational energy of a solution.

2.4 The bulk

As promised, we start with the bulk part of the action (7), in the hope of deriving S_{bulk}^* and S_{leftover} .

Consider the Gauss-Codazzi equations (30) and (31). They take the form,

$$\Omega^{\perp a} = F^a + E^\perp \wedge G^a, \quad \Omega^{ab} = F^{ab} + E^\perp \wedge G^{ab}, \quad (65)$$

where F^a and F^{ab} are 2-forms living on Σ_t , whereas G^a and G^{ab} are 1-forms living on Σ_t . We insert this into the bulk part of the action. Making use of the formula (8), we find,

$$\begin{aligned} S_{\text{bulk}} = & \kappa \int_{\mathcal{M}} E^\perp \wedge \left\{ -2\Lambda\zeta + 2G^b \wedge \zeta_b + F^{ab} \wedge \zeta_{ab} \right. \\ & \left. + \alpha [4(F^b \wedge G^{cd} + G^b \wedge F^{cd}) \wedge \zeta_{bcd} + F^{ab} \wedge F^{cd} \wedge \zeta_{abcd}] \right\} \end{aligned} \quad (66)$$

where we have introduced²

$$\zeta_{a_1 \dots a_m} = e_{\perp a_1 \dots a_m} = \frac{1}{(D-1-m)!} \epsilon_{a_1 \dots a_m a_{m+1} \dots a_{D-1}} E^{a_1} \wedge \dots \wedge E^{a_{D-1}}. \quad (67)$$

Note that (66) does not contain terms like $F^b \wedge \zeta_b$ or $F^b \wedge F^{cd} \wedge \zeta_{bcd}$, contrary to what we might have expected. This is because they are D -forms that depend only on $\{E^a\}$ and not E^\perp . $\{E^a\}$ is a basis on Σ_t , which is $(D-1)$ -dimensional, so we conclude that terms like $F^b \wedge \zeta_b$ and $F^b \wedge F^{cd} \wedge \zeta_{bcd}$ should vanish.

We now expand the bulk action (66) even further to give,

$$S_{\text{bulk}} = S_{\text{kinetic}} + S_1 + S_2 + S_3, \quad (68)$$

²Compare with equation (4). $\zeta_{a_1 \dots a_m}$ is just the analogue of $e_{A_1 \dots A_m}$ on Σ_t .

where

$$S_{\text{kinetic}} = \kappa \int_{\mathcal{M}} E^\perp \wedge \left\{ -2\mathbb{S}_\perp H^b \wedge \zeta_b - 4\alpha \left[\tilde{\nabla} H^b \wedge \mathbb{S}_\perp \tilde{\omega}^{cd} + \mathbb{S}_\perp H^b \wedge \left(\tilde{\Omega}^{cd} + H^c \wedge H^d \right) \right] \wedge \zeta_{bcd} \right\} \quad (69)$$

$$S_1 = \kappa \int_{\mathcal{M}} E^\perp \wedge \left\{ -2\Lambda \zeta + F^{ab} \wedge \zeta_{ab} + \alpha F^{ab} \wedge F^{cd} \wedge \zeta_{abcd} \right\} \quad (70)$$

$$S_2 = \kappa \int_{\mathcal{M}} E^\perp \wedge \left\{ -\frac{2}{N} \tilde{\nabla} (N a^b) \wedge \zeta_b - 4\alpha \left[\frac{1}{N} \tilde{\nabla} (N a^b) \wedge \left(\tilde{\Omega}^{cd} + H^c \wedge H^d \right) + 2\tilde{\nabla} H^b \wedge H^{[c} a^{d]} \right] \wedge \zeta_{bcd} \right\} \quad (71)$$

$$S_3 = \kappa \int_{\mathcal{M}} E^\perp \wedge \left\{ 4\alpha \left[\tilde{\nabla} H^b \wedge \frac{1}{N} \tilde{\nabla} (N \tilde{l}^{cd}) - \tilde{l}^{ba} H_a \wedge \left(\tilde{\Omega}^{cd} + H^c \wedge H^d \right) \right] \wedge \zeta_{bcd} \right\} \quad (72)$$

We will now attempt to rewrite each of the above expressions, bearing in mind the goals we mentioned at the beginning of this section.

2.4.1 S_{kinetic}

Let us start with S_{kinetic} . This contains terms like $\mathbb{S}_\perp H^a$, which need to be eliminated by integration by parts. Using the fact that

$$\mathbb{S}_\perp \zeta_{a_1 \dots a_m} = \mathbb{S}_\perp E^{a_{m+1}} \wedge \zeta_{a_1 \dots a_{m+1}}, \quad (73)$$

we find

$$S_{\text{kinetic}} = \kappa \int_{\mathcal{M}} E^\perp \wedge \left\{ \pi_a \wedge \mathbb{S}_\perp E^a + 4\alpha \left[H^b \wedge \mathbb{S}_\perp \tilde{\Omega}^{cd} - \tilde{\nabla} H^b \wedge \mathbb{S}_\perp \tilde{\omega}^{cd} \right] \wedge \zeta_{bcd} \right\} - \kappa \int_{\Sigma_\infty} 2H^b \wedge \zeta_b + 4\alpha H^b \wedge \Lambda^{cd} \wedge \zeta_{bcd} \quad (74)$$

Here we have,

$$\pi_a = -2\zeta_{ab} \wedge H^b - 4\alpha \zeta_{abcd} \wedge H^b \wedge \Lambda^{cd} \quad (75)$$

$$\Lambda^{cd} = \tilde{\Omega}^{cd} + \frac{1}{3} H^c \wedge H^d. \quad (76)$$

We will eventually show that π_a is indeed the momentum conjugate to E^a . If we note that $\theta^{\perp b} = -H^b$ is the only non-zero component of θ^{AB} on Σ_∞ , we can use equations (13) and (31) to show that,

$$S_\infty = \kappa \int_{\Sigma_\infty} 2H^b \wedge \zeta_b + 4\alpha H^b \wedge \Lambda^{cd} \wedge \zeta_{bcd}. \quad (77)$$

This will cancel off the second line in equation (74). In appendix A, we prove the relation,

$$E^\perp \wedge \mathbb{S}_\perp \tilde{\Omega}^{cd} = dt \wedge \tilde{\nabla} (N \mathbb{S}_\perp \tilde{\omega}^{cd}). \quad (78)$$

We also have the following identities,

$$\tilde{\nabla}\zeta_{a_1\dots a_m} \equiv 0 \equiv \tilde{\nabla}\tilde{\Omega}^{cd}. \quad (79)$$

The left hand side follows automatically from the zero-torsion condition (35), whereas the right hand side is just the frame-form version of the Bianchi identity. We can use (78) and (79) to show that,

$$\begin{aligned} \int_{\mathcal{M}} E^\perp \wedge \left[H^b \wedge \mathbb{S}_\perp \tilde{\Omega}^{cd} - \tilde{\nabla} H^b \wedge \mathbb{S}_\perp \tilde{\omega}^{cd} \right] \wedge \zeta_{bcd} \\ = \int dt \int_{\Sigma_t} \left[H^b \wedge \tilde{\nabla} (N \mathbb{S}_\perp \tilde{\omega}^{cd}) - \tilde{\nabla} H^b \wedge N \mathbb{S}_\perp \tilde{\omega}^{cd} \right] \wedge \zeta_{bcd} \\ = - \int dt \int_{\Sigma_t} \tilde{d} [N H^b \wedge \mathbb{S}_\perp \tilde{\omega}^{cd} \wedge \zeta_{bcd}] \\ = \int dt \int_{S_t} N H^b \wedge \mathbb{S}_\perp \tilde{\omega}^{cd} \wedge \zeta_{bcd} \end{aligned} \quad (80)$$

Note that we have applied Stokes' Theorem in the following way. If \tilde{A} is a $(D-2)$ -form on Σ_t , then

$$\begin{aligned} \int dt \int_{\Sigma_t} \tilde{d}\tilde{A} &= \int_{\mathcal{M}} dt \wedge \tilde{d}\tilde{A} = \int_{\mathcal{M}} dt \wedge d\tilde{A} \\ &= - \int_{\mathcal{M}} d(dt \wedge \tilde{A}) = - \int_B dt \wedge \tilde{A} = - \int dt \int_{S_t} \tilde{A} \end{aligned} \quad (81)$$

where we have used the fact that $d\tilde{A} = \tilde{d}\tilde{A} + E^\perp \{ \dots \}$ [43]. We now insert (77) and (80) into equation (74) to give,

$$S_{\text{kinetic}} = \kappa \int dt \int_{\Sigma_t} \pi_a \wedge N \mathbb{S}_\perp E^a + \kappa \int dt \int_{S_t} 4\alpha N H^b \wedge \mathbb{S}_\perp \tilde{\omega}^{cd} \wedge \zeta_{bcd} - S_\infty. \quad (82)$$

Notice that $\mathbb{S}_\perp \tilde{\omega}^{cd}$ has been removed from the bulk part of S_{kinetic} . This is because we have zero torsion. $\tilde{\omega}^{cd}$ is not an independent dynamical variable, so any time derivatives of it should indeed disappear from the bulk.

We are not yet finished with S_{kinetic} . Equation (82) still contains derivatives of the shift vector. From equation (21), we deduce that

$$\mathbb{S}_\perp E^a = \frac{1}{N} \left(\mathbb{S}_{\frac{\partial}{\partial t}} E^a - \mathbb{S}_{\vec{N}} E^a \right), \quad \vec{N} = N^a \frac{\partial}{\partial x^a} = E^a{}_b N^b X_a. \quad (83)$$

Since \vec{N} lives entirely on Σ_t , $\mathbb{S}_{\vec{N}}$ is just the intrinsic Lie derivative on Σ_t . Therefore,

$$\mathbb{S}_{\vec{N}} E^a = i_{\vec{N}}(\tilde{d}E^a) + \tilde{d}(i_{\vec{N}}E^a) = -N^b i_{[\frac{\partial}{\partial x^b}]}(\tilde{\omega}^a{}_c) E^c + \tilde{\nabla}(E^a{}_b N^b). \quad (84)$$

After some integration by parts we see that,

$$\int dt \int_{\Sigma_t} \pi_a \wedge \mathbb{S}_{\vec{N}} E^a = \int dt \int_{\Sigma_t} N^a \mathcal{H}_a - \int dt \int_{S_t} (-1)^D N^a \pi_b E^b{}_a, \quad (85)$$

where the momentum constraint,

$$\mathcal{H}_a = (-1)^{(D-1)} E^b{}_a \tilde{\nabla} \pi_b. \quad (86)$$

In deriving the momentum constraint, we have used the fact that,

$$\pi^b \wedge E^c = \pi^c \wedge E^b. \quad (87)$$

This is not obvious but can be shown using the symmetries of H_{ab} and the Riemann tensor³ (on Σ_t).

We are now ready to write down our final expression for S_{kinetic} ,

$$\begin{aligned} S_{\text{kinetic}} &= \kappa \int dt \int_{\Sigma_t} \pi_a \wedge \mathbb{S}_{\frac{\partial}{\partial t}} E^a - N^a \mathcal{H}_a \\ &+ \kappa \int dt \int_{S_t} 4\alpha N H^b \wedge \mathbb{S}_{\perp} \tilde{\omega}^{cd} \wedge \zeta_{bcd} + (-1)^D N^a \pi_b E^b{}_a - S_{\infty}. \end{aligned} \quad (88)$$

We now consider the remaining terms in the bulk action, starting with S_1 .

2.4.2 S_1

S_1 can be written

$$S_1 = -\kappa \int dt \int_{\Sigma_t} N \mathcal{H}, \quad (89)$$

where the Hamiltonian constraint, \mathcal{H} , is given by,

$$\mathcal{H} = 2\Lambda\zeta - F^{ab} \wedge \zeta_{ab} - \alpha F^{ab} \wedge F^{cd} \wedge \zeta_{abcd}. \quad (90)$$

Notice that the bulk part of $(S_{\text{kinetic}} + S_1)$ already has the form given in equation (62). We therefore expect S_2 and S_3 to give surface terms only.

2.4.3 S_2

Making use of the torsion-free and Bianchi identities (79), it is easy to show that,

$$S_2 = \kappa \int dt \int_{\Sigma_t} \tilde{d} \left[-2N a^b \zeta_b - 4\alpha N a^b \left(\tilde{\Omega}^{cd} + H^c \wedge H^d \right) \wedge \zeta_{bcd} \right] \quad (91)$$

By Stokes' Theorem we conclude that S_2 is indeed eliminated from the bulk,

$$S_2 = \kappa \int dt \int_{S_t} N \left[2a^b \zeta_b + 4\alpha a^b F^{cd} \wedge \zeta_{bcd} \right]. \quad (92)$$

³The Riemann tensor on Σ_t is given by $\tilde{R}^a{}_{bcd}$ where $\tilde{\Omega}^a{}_b = \frac{1}{2} \tilde{R}^a{}_{bcd} E^c \wedge E^d$.

2.4.4 S_3

Now consider S_3 . Integrating by parts we find that,

$$\begin{aligned} S_3 &= -\kappa \int dt \int_{\Sigma_t} 4\alpha N H_a \wedge \left[\tilde{l}^{ba} F^{cd} + \tilde{l}^{cd} \tilde{\Omega}^{ba} \right] \wedge \zeta_{bcd} \\ &\quad - \kappa \int dt \int_{S_t} 4\alpha N \tilde{\nabla} H^b \tilde{l}^{cd} \wedge \zeta_{bcd}, \end{aligned} \quad (93)$$

where we have used the fact that,

$$\tilde{\nabla}^2 H^b = \tilde{\Omega}^b_c \wedge H^c. \quad (94)$$

At first sight, it appears that S_3 contributes something to the bulk part of the action. However, in appendix A, we show that,

$$H_a \wedge \left[\tilde{l}^{ba} F^{cd} + \tilde{l}^{cd} \tilde{\Omega}^{ba} \right] \wedge \zeta_{bcd} = 0 \quad (95)$$

This ensures that we are only left with a surface term,

$$S_3 = \kappa \int dt \int_{S_t} 4\alpha N F^b \tilde{l}^{cd} \wedge \zeta_{bcd}. \quad (96)$$

Collecting together equations (88), (89), (92), and (96), we conclude that S_{bulk} takes the desired form

$$S_{\text{bulk}} = S_{\text{bulk}}^* + S_{\text{leftover}} \quad (97)$$

with S_{bulk}^* given by equation (62) and S_{leftover} given by

$$\begin{aligned} S_{\text{leftover}} &= -S_\infty + \kappa \int dt \int_{S_t} (-1)^D N^a \pi_b E^b_a \\ &\quad + N \left\{ 2a^b \zeta_b + 4\alpha \left[H^b \wedge \$_\perp \tilde{\omega}^{cd} + a^b F^{cd} + F^b \tilde{l}^{cd} \right] \wedge \zeta_{bcd} \right\}. \end{aligned} \quad (98)$$

2.5 The boundary

As expected, rewriting the bulk part of the action in the desired form (62) has altered the boundary part of the action. In particular, we have a leftover surface integral (98) that must be added to the original boundary part of the action (63). This gives the modified boundary action,

$$S_{\text{boundary}}^* = S_{\text{leftover}} + S_\infty + S_B. \quad (99)$$

In order to combine each term in the above equation, we need to write them in a common form. This will involve integrations over S_t , of well defined quantities on S_t .

Let us begin with $S_{\text{leftover}} + S_\infty$. From (37) we know that near B , $E^z_a = \delta^z_a$ and $N^z = 0$. This means that the momentum term in (98) gives

$$\kappa \int dt \int_{S_t} (-1)^D N^a \pi_b E^b_a = \kappa \int dt \int_{S_t} (-1)^D N^i \pi_j E^j_i. \quad (100)$$

We now use equations (46), (47) and (53) to rewrite the remaining terms. We find that,

$$\begin{aligned} S_{\text{leftover}} + S_\infty &= \kappa \int dt \int_{S_t} (-1)^D N^i \pi_j E^j_i + 2Nc\phi \\ &+ 4\alpha N \left[-b_k E^k \wedge \widehat{\mathbb{S}}_\perp \widehat{\omega}^{ij} + cF^{ij} + F^z \tilde{l}^{ij} - 2\widehat{H}^i \wedge \widehat{\mathbb{S}}_\perp \widehat{K}^j - 2a^i F^{zj} - 2F^i b^j \right] \wedge \phi_{ij} \end{aligned} \quad (101)$$

where $\phi_{i_1 \dots i_n}$ is the S_t analogue of $\zeta_{a_1 \dots a_n}$ and $e_{A_1 \dots A_n}$. More precisely,

$$\phi_{i_1 \dots i_n} = \zeta_{z i_1 \dots i_n} = e_{\perp z i_1 \dots i_n}. \quad (102)$$

Meanwhile, terms like F^z and F^{ij} can be deduced from equations (57) to (60),

$$F^z = \widehat{d}(b_k E^k) + \widehat{H}^k \wedge \widehat{K}_k \quad (103)$$

$$F^i = -\widehat{\nabla} \widehat{H}^i + b_k E^k \wedge \widehat{K}^i \quad (104)$$

$$F^{zj} = \widehat{\nabla} \widehat{K}^j - b_k E^k \wedge \widehat{H}^j \quad (105)$$

$$F^{ij} = \widehat{\Omega}^{ij} + \widehat{H}^i \wedge \widehat{H}^j - \widehat{K}^i \wedge \widehat{K}^j \quad (106)$$

Note that we have dropped all terms like $E^z \{ \dots \}$ as we are now integrating over B .

Now consider S_B . Since $\theta^{z\mu} = K^\mu$ is the only non-zero component of θ^{AB} on B , we can write

$$S_B = -\kappa \int_B 2K^\mu \wedge e_{z\mu} + 4\alpha K^\mu \wedge \left[\Omega^{\nu\sigma} + \frac{2}{3} K^\nu \wedge K^\sigma \right] \wedge e_{z\nu\sigma} \quad (107)$$

Again using (46) and (47), we can rewrite S_B in the following way,

$$\begin{aligned} S_B &= \kappa \int dt \int_{S_t} 2N \widehat{K}^i \wedge \phi_i - 2Nc\phi + 4\alpha N \widehat{K}^i \wedge \left[F^{jk} + \frac{2}{3} \widehat{K}^j \wedge \widehat{K}^k \right] \wedge \phi_{ijk} \\ &+ 4\alpha N \left\{ -2b^i \left[F^j - \frac{2}{3} b_k E^k \wedge \widehat{K}^j \right] + 2\widehat{K}^i \wedge \left[G^j + \frac{2}{3} \left(-c\widehat{K}^j + b^j b_k E^k \right) \right] \right. \\ &\quad \left. - c \left[F^{ij} + \frac{2}{3} \widehat{K}^i \wedge \widehat{K}^j \right] + b_k E^k \wedge \left[G^{ij} + \frac{4}{3} b^{[i} \widehat{K}^{j]} \right] \right\} \wedge \phi_{ij} \end{aligned} \quad (108)$$

where from (58) and (60),

$$G^j = -\widehat{\mathbb{S}}_\perp \widehat{H}^j - \frac{1}{N} \widehat{\nabla} (N a^j) - \tilde{l}^{jk} \widehat{H}_k + c \widehat{K}^j - b^j b_k E^k \quad (109)$$

$$G^{ij} = \widehat{\mathbb{S}}_\perp \widehat{\omega}^{ij} - \frac{1}{N} \widehat{\nabla} (N \tilde{l}^{ij}) + 2\widehat{H}^{[i} a^{j]} + 2\widehat{K}^{[i} b^{j]} \quad (110)$$

The expressions (101) and (108) now have a common form, so we can combine them to get

$$S_{\text{boundary}}^* = S_{\text{boundary},1} + S_{\text{boundary},2} + S_{\text{boundary},3} \quad (111)$$

where

$$S_{\text{boundary},1} = \kappa \int dt \int_{S_t} (-1)^D N^i \pi_j E^j{}_i + N \left\{ 2\widehat{K}^i \wedge \phi_i + 4\alpha K^i \wedge \left[F^{jk} + \frac{2}{3} \widehat{K}^j \wedge \widehat{K}^k \right] \wedge \phi_{ijk} \right\} \quad (112)$$

$$S_{\text{boundary},2} = \kappa \int dt \int_{S_t} 4\alpha N \left\{ \widehat{d}(b_k E^k) \tilde{l}^{ij} - \frac{1}{N} b_k E^k \wedge \widehat{\nabla}(N \tilde{l}^{ij}) - 2a^i \widehat{\nabla} \widehat{K}^j - \frac{2}{N} \widehat{K}^i \wedge \widehat{\nabla}(N a^j) \right\} \wedge \phi_{ij} \quad (113)$$

$$S_{\text{boundary},3} = \kappa \int dt \int_{S_t} 4\alpha N \left\{ -2\widehat{H}^i \wedge \widehat{\mathcal{S}}_{\perp} \widehat{K}^j - 2\widehat{K}^i \wedge \widehat{\mathcal{S}}_{\perp} \widehat{H}^j + \widehat{H}^k \wedge \widehat{K}_k \tilde{l}^{ij} - 2\widehat{K}^i \wedge \tilde{l}^{jk} \widehat{H}_k \right\} \wedge \phi_{ij} \quad (114)$$

As with our analysis of the bulk, we will focus on each term, one at a time, beginning with $S_{\text{boundary},1}$.

2.5.1 $S_{\text{boundary},1}$

There is not much to do here. We simply use the expression for F^{ij} given in equation (106), to write (112) as

$$S_{\text{boundary},1} = \kappa \int dt \int_{S_t} (-1)^D N^i \pi_j E^j{}_i + N \left\{ 2\widehat{K}^i \wedge \phi_i + 4\alpha \widehat{K}^i \wedge \left[\widehat{\Omega}^{jk} + \widehat{H}^j \wedge \widehat{H}^k - \frac{1}{3} \widehat{K}^j \wedge \widehat{K}^k \right] \wedge \phi_{ijk} \right\} \quad (115)$$

2.5.2 $S_{\text{boundary},2}$

Now consider equation (113). In analogy with the left hand side of (79), we have the zero torsion condition,

$$\widehat{\nabla} \phi_{i_1 \dots i_n} \equiv 0 \quad (116)$$

This enables us to write the integrand in (113) as a total derivative,

$$S_{\text{boundary},2} = \kappa \int dt \int_{S_t} 4\alpha \widehat{d} \left[N \left(b_k E^k \tilde{l}^{ij} - 2a^i \widehat{K}^j \right) \wedge \phi_{ij} \right]. \quad (117)$$

Since S_t is a boundary of Σ_t , it has no boundary of its own. This means that,

$$S_{\text{boundary},2} = 0 \quad (118)$$

2.5.3 $S_{\text{boundary},3}$

From the decomposition of zero torsion (36), we can infer that

$$\widehat{\mathbb{S}}_{\perp} E^i = -\widehat{H}^i - \tilde{l}^i_j E^j. \quad (119)$$

We now do integration by parts on the time derivatives in (114). This gives,

$$\begin{aligned} S_{\text{boundary},3} = \kappa \int dt \int_{S_t} 4\alpha N \left\{ -2\widehat{K}^i \wedge \widehat{H}^j \wedge \widehat{H}^k \wedge \phi_{ijk} \right. \\ \left. + \widehat{H}^k \wedge \widehat{K}_k \tilde{l}^{ij} \wedge \phi_{ij} - 2\widehat{K}^i \wedge \widehat{H}_k \tilde{l}^{jk} \wedge \phi_{ij} - 2\widehat{K}^i \wedge \widehat{H}^j \tilde{l}^k_l \wedge E^l \wedge \phi_{ijk} \right\}. \end{aligned} \quad (120)$$

where we have used (119) along with the fact that,

$$\widehat{\mathbb{S}}_{\perp} \phi_{i \dots i_m} = \widehat{\mathbb{S}}_{\perp} E^{i_{m+1}} \wedge \phi_{i \dots i_{m+1}} \quad (121)$$

We have also thrown away some far past/future integrals over S_{∞} for the same reason that we threw away the surface integrals in the previous section.

Using the useful formula (8), we can show that,

$$\begin{aligned} \widehat{H}^k \wedge \widehat{K}_k \tilde{l}^{ij} \wedge \phi_{ij} - 2\widehat{K}^i \wedge \widehat{H}_k \tilde{l}^{jk} \wedge \phi_{ij} - 2\widehat{K}^i \wedge \widehat{H}^j \tilde{l}^k_l \wedge E^l \wedge \phi_{ijk} \\ = \widehat{K}^i \wedge E_i \wedge \widehat{H}^j \tilde{l}^{kl} \wedge \phi_{jkl} = 0. \end{aligned} \quad (122)$$

where the last equality follows from the symmetry of \widehat{K}^{ij} . We conclude that we are only left with the following,

$$S_{\text{boundary},3} = \kappa \int dt \int_{S_t} 4\alpha N \left\{ -2\widehat{K}^i \wedge \widehat{H}^j \wedge \widehat{H}^k \wedge \phi_{ijk} \right\}. \quad (123)$$

We now collect together equations (115), (118) and (123) to deduce that,

$$\begin{aligned} S_{\text{boundary}}^* = \kappa \int dt \int_{S_t} (-1)^D N^i \pi_j E^j_i + N \left\{ 2\widehat{K}^i \wedge \phi_i \right. \\ \left. + 4\alpha \widehat{K}^i \wedge \left[\widehat{\Omega}^{jk} - \widehat{H}^j \wedge \widehat{H}^k - \frac{1}{3} \widehat{K}^j \wedge \widehat{K}^k \right] \wedge \phi_{ijk} \right\} \end{aligned} \quad (124)$$

2.6 The conjugate momentum

Now that we have the action in its correct form, it remains to calculate the momentum conjugate to E^a . This is given by,

$$p_a = \frac{\partial \mathcal{L}_{\text{bulk}}}{\partial \left(\widehat{\mathbb{S}}_{\frac{\partial}{\partial t}} E^a \right)} \quad (125)$$

where $\mathcal{L}_{\text{bulk}}$ is the bulk integrand. We could take $\mathcal{L}_{\text{bulk}}$ from S_{bulk}^* . However, it is convenient to temporarily undo the integration by parts that gave equation (85). In other words, we leave the derivatives of N^a in the bulk action. This is perfectly OK, because it does not affect the bulk dynamics, and therefore the value of the conjugate momentum. The bulk integrand is temporarily given by,

$$\mathcal{L}_{\text{bulk}} = \pi_a \wedge N \$_{\perp} E^a - N\mathcal{H} \quad (126)$$

where π_a and \mathcal{H} are given by equations (75) and (90) respectively. Using the zero torsion decomposition (36), we can say

$$\mathcal{L}_{\text{bulk}} = -N\pi_a \wedge \left(H^a + \tilde{l}^a{}_b E^b \right) - N\mathcal{H} = -N\pi_a \wedge H^a - N\mathcal{H} \quad (127)$$

where the right hand equation follows from (87), and the antisymmetry of \tilde{l}^{ab} .

Referring to equations (36) and (83), we use the chain rule to show that,

$$p_a = -\frac{1}{N} \frac{\partial \mathcal{L}_{\text{bulk}}}{\partial \widehat{H}^a} = \pi_a \quad (128)$$

This non-trivial result is due to the following cancelation,

$$\frac{\partial \pi_b}{\partial \widehat{H}^a} \wedge \widehat{H}^b + \frac{\partial \mathcal{H}}{\partial \widehat{H}^a} = 0. \quad (129)$$

We conclude that π_a is indeed the conjugate momentum. It should be thought of as a function of $\$_{\frac{\partial}{\partial t}} E^a$. To derive the Hamiltonian, we should invert this function. However, π_a is cubic in $\$_{\frac{\partial}{\partial t}} E^a$, so the inverse is multivalued. This is a well known property of higher derivative gravities. In the Hamiltonian picture, this could mean that we could jump from one solution to another. These ‘‘zigzagging’’ histories still provide an extremum of the action. In this paper, we will assume that at any given time, we have a unique solution. This is just the same as saying that we are not in the process of jumping from one solution to another. For more discussion on multivalued Hamiltonians in this context, refer to [42, 46].

2.7 The physical Hamiltonian

We have shown that we can write our action as,

$$S = S_{\text{bulk}}^* + S_{\text{boundary}}^* = \kappa \int dt \left[\int_{\Sigma_t} \mathcal{L}_{\text{bulk}}^* + \int_{S_t} \mathcal{L}_{\text{boundary}}^* \right] \quad (130)$$

where

$$\mathcal{L}_{\text{bulk}}^* = \pi_a \wedge \$_{\frac{\partial}{\partial t}} E^a - N\mathcal{H} - N^a \mathcal{H}_a \quad (131)$$

$$\begin{aligned} \mathcal{L}_{\text{boundary}}^* = N \left\{ 2\widehat{K}^i \wedge \phi_i + 4\alpha \widehat{K}^i \wedge \left[\widehat{\Omega}^{jk} - \widehat{H}^j \wedge \widehat{H}^k - \frac{1}{3} \widehat{K}^j \wedge \widehat{K}^k \right] \wedge \phi_{ijk} \right\} \\ + (-1)^D N^i \pi_j E^j{}_i. \end{aligned} \quad (132)$$

The corresponding Hamiltonian is defined as,

$$\begin{aligned}
H &= \kappa \int_{\Sigma_t} \pi_a \wedge \mathcal{E}^a - \mathcal{L}_{\text{bulk}}^* - \kappa \int_{S_t} \mathcal{L}_{\text{boundary}}^* \\
&= \kappa \int_{\Sigma_t} N\mathcal{H} + N^a \mathcal{H}_a - \kappa \int_{S_t} \mathcal{L}_{\text{boundary}}^*
\end{aligned} \tag{133}$$

To arrive at the physical Hamiltonian, we need to subtract off the background Hamiltonian,

$$\bar{H} = -\kappa \int_{S_t} \bar{\mathcal{L}}_{\text{boundary}}^* \tag{134}$$

Here we have used the fact that the background is a stationary solution to the field equations [3],

$$\bar{\mathcal{H}} = \bar{\mathcal{H}}_a = \bar{\pi}_a = 0. \tag{135}$$

The physical Hamiltonian is therefore given by,

$$H_{\text{phys}} = \kappa \int_{\Sigma_t} N\mathcal{H} + N^a \mathcal{H}_a - \kappa \int_{S_t} \Delta \mathcal{L}_{\text{boundary}}^* \tag{136}$$

where

$$\begin{aligned}
\Delta \mathcal{L}_{\text{boundary}}^* &= (-1)^D N^i \pi_j E^j{}_i + N \left\{ 2\Delta \widehat{K}^i \wedge \phi_i \right. \\
&\quad \left. + 4\alpha \left[\Delta \widehat{K}^i \wedge \left(\widehat{\Omega}^{jk} - \widehat{H}^j \wedge \widehat{H}^k \right) - \frac{1}{3} \Delta \left(\widehat{K}^i \wedge \widehat{K}^j \wedge \widehat{K}^k \right) \right] \wedge \phi_{ijk} \right\}
\end{aligned} \tag{137}$$

For any quantity Q in the test spacetime with corresponding quantity \bar{Q} in the background, $\Delta Q = Q - \bar{Q}$. Notice that we have $\Delta \widehat{\Omega}^{jk} = \Delta \widehat{H}^j = 0$. This is because the geometry of the boundary is the same in the test spacetime, as in the background.

If our test spacetime is a solution to the field equations, it satisfies the constraints

$$\mathcal{H} = \mathcal{H}_a = 0 \tag{138}$$

Its energy is then given by the value of the physical Hamiltonian,

$$\begin{aligned}
E &= -\kappa \int_{S_t} (-1)^D N^i \pi_j E^j{}_i + N \left\{ 2\Delta \widehat{K}^i \wedge \phi_i \right. \\
&\quad \left. + 4\alpha \left[\Delta \widehat{K}^i \wedge \left(\widehat{\Omega}^{jk} - \widehat{H}^j \wedge \widehat{H}^k \right) - \frac{1}{3} \Delta \left(\widehat{K}^i \wedge \widehat{K}^j \wedge \widehat{K}^k \right) \right] \wedge \phi_{ijk} \right\}
\end{aligned} \tag{139}$$

Given the technical complexity of Gauss-Bonnet gravity, we believe that this expression is remarkably simple. Note that for $\alpha = 0$, we recover the correct result for Einstein gravity, as of course we should.

For the benefit of anyone who has skipped the details of this section, we remind them that a summary of notation can be found in appendix B.

2.8 Using a coordinate basis

Although the final result (139) is neat and tidy, we might prefer to work in a coordinate basis, and express the Hamiltonian in terms of the familiar tensors of General Relativity. In this case, our canonical variable is the induced metric γ_{ab} . The conjugate momentum, π^{ab} , is given by [43],

$$\pi^{ab} d^{D-1}x = \frac{1}{2} \pi^a \wedge E^b. \quad (140)$$

With this in mind, we can verify that the Hamiltonian and momentum constraints given by equations (90) and (86) respectively, agree with the corresponding expressions in [42]. If z^A is the normal to the timelike boundary, B , and n^A the normal to Σ_t , the Hamiltonian (136) can be written

$$H_{\text{phys}} = \kappa \int_{\Sigma_t} d^{D-1}x [N\mathcal{H} + N^a \mathcal{H}_a] - \kappa \int_{S_t} d^{D-2}x \Delta \mathcal{L}_{\text{boundary}}^* \quad (141)$$

where now

$$\begin{aligned} \Delta \mathcal{L}_{\text{boundary}}^* = & 2N^i \pi_{ij} z^j + N \sqrt{\lambda} \left\{ 2\Delta K \right. \\ & \left. + 2\alpha(3!) \delta_i^{[l} \delta_j^m \delta_k^{n]} \left[\Delta K_l^i \left(\widehat{R}^{jk}_{mn} - 2H_m^j H_n^k \right) - \frac{2}{3} \Delta (K_l^i K_m^j K_n^k) \right] \right\} \end{aligned} \quad (142)$$

Here, \widehat{R}^{jk}_{mn} is the Riemann tensor on S_t , constructed out of the induced metric λ_{ij} . K_j^i and H_j^i are the extrinsic curvatures of S_t in Σ_t and B respectively, and K is the trace of K_j^i . More precisely,

$$K_{ij} = -\lambda_{(i}^l \lambda_{j)}^m \nabla_l z_m, \quad H_{ij} = -\lambda_{(i}^l \lambda_{j)}^m \nabla_l n_m, \quad K = \lambda_j^i K_j^i. \quad (143)$$

3 Comparison with previous energy expressions

3.1 Deser and Tekin

Now that we have derived an expression for the energy in GB gravity, we should compare it with previous results in the literature. In particular, Deser and Tekin [4,5] used a ‘‘conserved charge’’ technique to derive the energy of asymptotically maximally symmetric spacetimes above backgrounds of constant curvature. This method can be applied to generic higher derivative gravities, but we can check it is consistent with our result in the Gauss-Bonnet case.

Suppose our test spacetime \mathcal{M} , is asymptotically maximally symmetric. We choose our background, $\bar{\mathcal{M}}$ to be the maximally symmetric solution with curvature form,

$$\bar{\Omega}^{AB} = \frac{2\Lambda_{\text{eff}}}{(D-1)(D-2)} E^A \wedge E^B \quad (144)$$

The effective cosmological constant is given by,

$$\Lambda_{\text{eff}} = -\frac{1}{4\alpha^*} \left[1 \pm \sqrt{1 + 8\Lambda\alpha^*} \right], \quad \alpha^* = \alpha \frac{(D-3)(D-4)}{(D-1)(D-2)}. \quad (145)$$

which is real and multivalued for $1 + 8\Lambda\alpha^* > 0$.

Our aim is to calculate the energy associated with the timelike Killing vector $\partial/\partial t$. We can choose our foliation $\{\Sigma_t\}$ so that the shift vector vanishes. Since $\partial/\partial t$ is Killing, it is clear that $\widehat{H}^i = 0$. The expression for the energy reduces to,

$$E = -\kappa \int_{S_t} N \left\{ 2\Delta \widehat{K}^i \wedge \phi_i + 4\alpha \left[\Delta \widehat{K}^i \wedge \widehat{\Omega}^{jk} - \frac{1}{3} \Delta \left(\widehat{K}^i \wedge \widehat{K}^j \wedge \widehat{K}^k \right) \right] \wedge \phi_{ijk} \right\} \quad (146)$$

Since S_t lies in the asymptotic region, we can say,

$$\Delta \left(\widehat{K}^i \wedge \widehat{K}^j \wedge \widehat{K}^k \right) \wedge \phi_{ijk} \approx 3\Delta \widehat{K}^i \wedge (\bar{K}^j \wedge \bar{K}^k) \wedge \phi_{ijk}, \quad (147)$$

where \bar{K}^j is \widehat{K}^j evaluated on the background. The energy is now given by,

$$E = -\kappa \int_{S_t} N \left\{ 2\Delta \widehat{K}^i \wedge \phi_i + 4\alpha \Delta \widehat{K}^i \wedge \left(\widehat{\Omega}^{jk} - \bar{K}^j \wedge \bar{K}^k \right) \wedge \phi_{ijk} \right\} \quad (148)$$

Using the fact that $\bar{H}^j = 0$, it is clear from equations (60) and (144) that,

$$\widehat{\Omega}^{jk} - \bar{K}^j \wedge \bar{K}^k = \frac{2\Lambda_{\text{eff}}}{(D-1)(D-2)} E^j \wedge E^k. \quad (149)$$

This implies that the energy,

$$E = -\kappa (1 + 4\alpha^* \Lambda_{\text{eff}}) \int_{S_t} 2N \Delta \widehat{K}^i \wedge \phi_i = \pm \kappa \sqrt{1 + 8\Lambda\alpha^*} \int_{S_t} 2N \Delta \widehat{K}^i \wedge \phi_i \quad (150)$$

where we have used the S_t analogue of the useful formula (8), and the cosmological constant relation (145).

In order to make contact with [4, 5], we switch to a coordinate basis,

$$E = \pm \kappa \sqrt{1 + 8\Lambda\alpha^*} \int_{S_t} d^{D-2} x \sqrt{\lambda} 2N \Delta K \quad (151)$$

We now follow the procedure described in [3] for Einstein gravity. Let us start with the test spacetime. Near S_t , we can express the metric on Σ_t in Gaussian normal coordinates,

$$ds_{\Sigma_t}^2 = \gamma_{ab} dx^a dx^b = dz^2 + q_{ij}(z, x^k) dx^i dx^j. \quad (152)$$

where $q_{ij}(0, x^k) = \lambda_{ij}$ is the metric on S_t . Similarly, for the background, we can write the metric on $\bar{\Sigma}_t$ (near S_t), as,

$$ds_{\bar{\Sigma}_t}^2 = \bar{\gamma}_{ab} d\bar{x}^a d\bar{x}^b = d\bar{z}^2 + \bar{q}_{ij}(\bar{z}, \bar{x}^k) d\bar{x}^i d\bar{x}^j. \quad (153)$$

To ensure that the normals to S_t agree on the test spacetime and the background, we choose the diffeomorphism $z = \bar{z}$ and $x^i = \bar{x}^i$. In these coordinates,

$$K = -\frac{1}{2}q^{ij}q_{ij,z}, \quad \bar{K} = -\frac{1}{2}\bar{q}^{ij}\bar{q}_{ij,z}. \quad (154)$$

Since both metrics agree on the boundary, we note that $\Delta q_{ij} = 0$ there. Therefore on S_t ,

$$\Delta K = -\frac{1}{2}\lambda^{ij}(\Delta q_{ij})_{,z} = -\frac{1}{2}(\Delta q)_{,z}. \quad (155)$$

where $\Delta q = \lambda^{ij}\Delta q_{ij}$. This gives a final energy expression,

$$E = \mp\kappa\sqrt{1+8\Lambda\alpha^*} \int_{S_t} d^{D-2}x\sqrt{\lambda}N(\Delta q)_{,z}. \quad (156)$$

We now use Deser and Tekin's method [4, 5] to calculate⁴ the energy associated with the timelike Killing vector, t^A .

$$E_{\text{DT}} = \mp\kappa\sqrt{1+8\Lambda\alpha^*} \int_{S_t} dS_A n_B \{ t_C \bar{\nabla}^B h^{AC} - t_C \bar{\nabla}^A h^{BC} + t^B \bar{\nabla}^A h - t^A \bar{\nabla}^B h \\ + h^{BC} \bar{\nabla}^A t_C - h^{AC} \bar{\nabla}^B t_C + t^A \bar{\nabla}_C h^{BC} - t^B \bar{\nabla}_C h^{AC} + h \bar{\nabla}^B t^A \} \quad (157)$$

where $h^{AB} = g^{AB} - \bar{g}^{AB}$ and $h = \bar{g}_{AB}h^{AB}$. Here g_{AB} and \bar{g}_{AB} are the metrics on \mathcal{M} and $\bar{\mathcal{M}}$ respectively. We will choose to work in a synchronous gauge for which $n^A h_{AB} = 0$ [3]. As the metrics agree on the boundary we can also set $h^{AB} = 0$ on S_t . If S_t has *inward* pointing normal $\partial/\partial z$, the measure is given by $dS_A = -d^{D-2}x\sqrt{\lambda}\delta_A^z$. For vanishing shift vector, we have $t^A = Nn^A$, and the energy (157) simplifies to,

$$E_{\text{DT}} = \pm\kappa\sqrt{1+8\Lambda\alpha^*} \int_{S_t} d^{D-2}x\sqrt{\lambda}N(\partial_b h^{zb} - \partial^z h). \quad (158)$$

In the Gaussian Normal coordinates we have recently described, $h^{zb} = 0$ and $h = \Delta q$. Deser and Tekin's energy now reads,

$$E_{\text{DT}} = \mp\kappa\sqrt{1+8\Lambda\alpha^*} \int_{S_t} d^{D-2}x\sqrt{\lambda}N\partial^z(\Delta q). \quad (159)$$

This expression clearly agrees with the equation (156). We can conclude that although our derivation was very different to that in [4, 5], our results are consistent.

3.1.1 Application to GB black holes

One of the nice features of GB gravity (2) is that it contains static, spherically symmetric solutions [47, 35, 48, 49, 50, 51, 27, 46] of the form,

$$ds^2 = -V(r)dt^2 + \frac{dr^2}{V(r)} + r^2 d\Omega_{D-2}^2 \quad (160)$$

⁴In [4, 5], the authors do not explicitly write down an energy expression for the GB action with a bare cosmological constant Λ . However, they give enough information to easily derive equation (157).

where $d\Omega_{D-2}^2$ is the metric on a unit $(D-2)$ -sphere. We will assume that $1+8\Lambda\alpha^* > 0$, so that there are two possible branches for the potential,

$$V(r) = 1 + \frac{r^2}{2(D-1)(D-2)\alpha^*} \left(1 \pm \sqrt{1 + 8\Lambda\alpha^* + \frac{4(D-1)(D-2)\alpha^*\mu}{r^{D-1}}} \right) \quad (161)$$

Here $\mu \geq 0$ is a constant of integration that gives mass to the spacetime. The upper branch has a naked singularity at $r = 0$, whereas the lower branch is a real black hole with a unique event horizon surrounding the singularity [35]. We wish to calculate the mass, M , of these spacetimes above the appropriate maximally symmetric background,

$$ds^2 = -\bar{V}(r)dt^2 + \frac{dr^2}{\bar{V}(r)} + r^2 d\Omega_{D-2}^2 \quad (162)$$

where

$$\bar{V}(r) = 1 + \frac{r^2}{2(D-1)(D-2)\alpha^*} \left(1 \pm \sqrt{1 + 8\Lambda\alpha^*} \right). \quad (163)$$

The foliation in each case is given by surfaces of constant t , so we soon see that,

$$q_{ij} = r^2(z)\chi_{ij}, \quad \bar{q}_{ij} = \bar{r}^2(z)\chi_{ij} \quad (164)$$

where χ_{ij} is the metric on the unit $(D-2)$ -sphere, and

$$\frac{dr}{dz} = -\sqrt{\bar{V}}, \quad \frac{d\bar{r}}{dz} = -\sqrt{\bar{V}}. \quad (165)$$

If $r(0) = \bar{r}(0) = R$, on S_t , it follows that,

$$(\Delta q)_{,z} = -\frac{2(D-2)}{R}\Delta\sqrt{V(R)} \approx -\frac{(D-2)}{R\sqrt{V}}\Delta V(R) \quad \text{on } S_t. \quad (166)$$

For large R ,

$$\Delta V(R) \approx \pm \frac{1}{\sqrt{1+8\Lambda\alpha^*}} \frac{\mu}{R^{D-3}} \quad (167)$$

Finally, we note that $N = \sqrt{\bar{V}}$ and apply equation (156) to give,

$$M = \kappa\Omega_{D-2}(D-2)\mu \quad (168)$$

where Ω_{D-2} is the volume of the unit $(D-2)$ -sphere. This is the standard result. It is always valid for $\Lambda_{\text{eff}} \leq 0$. For $\Lambda_{\text{eff}} > 0$ our analysis is valid only if the de Sitter horizon is much larger than the black hole horizon [5].

3.2 A special case

In the last section, we assumed that $1+8\Lambda\alpha^* > 0$. Now consider what happens when $1+8\Lambda\alpha^* = 0$. We cannot make use of the expression (156) because it involves multiplying an infinite integral, by zero! We will not worry about how one would

modify the approach of [4, 5] to accomodate this. Instead, we will sell the approach developed in this paper. Let us focus on the 5-dimensional black hole with,

$$V(r) = 1 + \frac{r^2}{4\alpha} - \sqrt{\frac{\mu}{2\alpha}}. \quad (169)$$

To calculate the mass, we need to go back to equation (139). We also need to choose a background. In this example, the maximally symmetric solution with $\mu = 0$ is not necessarily the most natural choice. We might prefer μ to be chosen so that the horizon has zero area [35]. Whatever our choice, we illustrate the flexibility of this work by allowing for non-maximally symmetric backgrounds. We will keeps things general and say that the background potential is given by

$$\bar{V}(r) = 1 + \frac{r^2}{4\alpha} - \sqrt{\frac{\bar{\mu}}{2\alpha}}. \quad (170)$$

As before, our foliation is made up of surfaces of constant t , with boundary S_t given by $r = R$. To apply the energy expression (139), we need the following ingredients,

$$N = \sqrt{\bar{V}}, \quad N^i = 0, \quad \hat{K}^i = \frac{\sqrt{\bar{V}}}{R} E^i, \quad \hat{H}^i = 0, \quad \hat{\Omega}^{jk} = \frac{1}{R^2} E^j \wedge E^k. \quad (171)$$

Now use the useful formula (8) in (139), to derive the energy,

$$E = -\kappa\Omega_3 R^3 \sqrt{\bar{V}} \left\{ \frac{6}{R} \Delta(\sqrt{\bar{V}}) + \frac{24\alpha}{R^3} \left[\Delta(\sqrt{\bar{V}}) - \frac{1}{3} \Delta(V\sqrt{\bar{V}}) \right] \right\} \quad (172)$$

To keep things tidy, we write

$$V = y^2 - m, \quad \bar{V} = y^2 - \bar{m} \quad (173)$$

where

$$y^2 = 1 + \frac{R^2}{4\alpha}, \quad m = \sqrt{\frac{\mu}{2\alpha}}, \quad \bar{m} = \sqrt{\frac{\bar{\mu}}{2\alpha}}. \quad (174)$$

Now for large y ,

$$\sqrt{\bar{V}} = y \left[1 - \frac{m}{2y^2} + \mathcal{O}\left(\frac{1}{y^4}\right) \right] \quad (175)$$

$$\Delta(\sqrt{\bar{V}}) = -y \left[\frac{\Delta m}{2y^2} + \frac{\Delta(m^2)}{8y^4} + \mathcal{O}\left(\frac{1}{y^6}\right) \right] \quad (176)$$

$$\Delta(V\sqrt{\bar{V}}) = -3y^3 \left[\frac{\Delta m}{2y^2} - \frac{\Delta(m^2)}{8y^4} + \mathcal{O}\left(\frac{1}{y^6}\right) \right]. \quad (177)$$

If we plug this back into (172) we find,

$$E = 3\kappa\Omega_3 [2\alpha\Delta(m^2)] + \mathcal{O}\left(\frac{1}{y^2}\right) \quad (178)$$

Now we send R , or equivalently y , to infinity, to derive the black hole mass,

$$M = 3\kappa\Omega_3 \Delta\mu. \quad (179)$$

If we had chosen the background to be the black hole of zero size, we would have $\bar{\mu} = 2\alpha$. Our black hole mass would be given by $M = 3\kappa\Omega_3(\mu - 2\alpha)$, which agrees with the “minisuperspace” method employed in [35].

4 Discussion

In this paper, we have derived a neat and easy to use expression for the gravitational energy of a solution in Gauss-Bonnet gravity. This was done using a Hamiltonian approach, much like the one used by Hawking and Horowitz [3] for Einstein gravity. Given the technical complexity of the derivation, our final expression (139) is remarkably simple.

There have been other ways of calculating the energy of certain Gauss-Bonnet solutions [35, 4, 5]. We have shown that our Hamiltonian approach yields results that are consistent with these. Each approach has its advantages and disadvantages, as we will now discuss.

Consider the “conserved charge” method given in [4, 5]. The authors identify a conserved current associated with a timelike Killing vector. The gravitational energy corresponds to the “charge” of this current. This method can be applied to generic higher derivative gravities, of which Gauss-Bonnet gravity is just a special case. However, the background spacetimes, or vacua, are always assumed to be maximally symmetric everywhere. That is not to say that this method cannot be extended to a more general choice of background. This should clearly be a topic for future research. It would also be interesting to know how to apply this method to the special case discussed in section 3.2.

Similarly, we should also ask if we can extend our Hamiltonian approach to more general higher derivative gravities. It should be fairly easy to consider the Lovelock action [6],

$$S = \sum_{n=0}^{\lfloor \frac{D-1}{2} \rfloor} \alpha_n S_n, \quad S_n = \int_{\mathcal{M}} \Omega^{A_1 B_1} \wedge \dots \wedge \Omega^{A_n B_n} \wedge e_{A_1 B_1 \dots A_n B_n}. \quad (180)$$

The structure is very similar to GB gravity, with the surface terms for the action given in [36]. Life would be more difficult if we wanted to consider an arbitrary combination of Riemann tensors, as the surface terms are generally unknown.

The “minisuperspace” method used in [35] is closest in spirit to the Hamiltonian approach. The idea is to consider a static, spherically symmetric ansatz for the metric, and insert it back into the action. The action becomes one-dimensional, making it easier to fix the boundary term. When we turn to the Hamiltonian, and evaluate it on one of the black hole solutions given in sections 3.1.1 and 3.2, we derive the black hole mass. This method is very simple and easy to use, but somewhat limited. It can only be applied when the one-dimensional “minisuperspace” model is valid. This is OK for the black hole spacetimes discussed in [35], but a more general approach is clearly desirable.

The Hamiltonian approach developed in this paper is the appropriate generalisation. It can be applied whatever the symmetries of the solution, and without having to reduce the number of dimensions. In particular, we will use it to investigate the generalisation of braneworld holography [33] for Gauss-Bonnet gravity [34].

To sum up, we have derived an expression (139) for the energy of a solution to Gauss-Bonnet gravity. This can be applied whatever our choice of background, and

whatever the symmetries of our solution. This should, hopefully, give us a platform to investigate Gauss-Bonnet gravity more thoroughly.

Acknowledgements

I would like to thank James Gregory for many interesting discussions throughout this entire project. I would also like to thank him, and Simon Ross, for proof reading this article. Thanks also go to Christos Charmousis for introducing me to Gauss-Bonnet gravity during a very long viva! Final thanks go to Steven Gerrard and Michael Owen for scoring the goals that beat United. AP was funded by PPARC.

A Proofs

A.1 Proof of equation (51)

We start with

$$\tilde{A} = \hat{A} + A_z \wedge E^z, \quad (181)$$

as described in the paragraph leading up to equation (51). Now

$$\$_\perp \tilde{A} = \$_\perp \hat{A} + A_z \wedge \$_\perp E^z + E^z \wedge \{\dots\}. \quad (182)$$

From equation (36) we know that

$$\$_\perp E^z = -H^z{}_i E^i - l^z{}_i E^i + E^z \{\dots\} = E^z \{\dots\}. \quad (183)$$

The last result comes from equation (47). By definition

$$\$_\perp \hat{A} = P \left[d \left(i_\perp \hat{A} \right) + i_\perp \left(d \hat{A} \right) \right] = P \left[i_\perp \left(d \hat{A} \right) \right]. \quad (184)$$

In analogy with equation (25), we also have that

$$d \hat{A} = d_0 \hat{A} + E^z \wedge \{\dots\} \Rightarrow i_\perp \left(d \hat{A} \right) = i_\perp \left(d_0 \hat{A} \right) + E^z \wedge \{\dots\}. \quad (185)$$

So from equations (184) and (185) we deduce that

$$\$_\perp \hat{A} = P \left[i_\perp \left(d_0 \hat{A} \right) \right] = \hat{\$_\perp} \hat{A}. \quad (186)$$

Putting (183) and (186) into equation (182) gives the desired result,

$$\$_\perp \tilde{A} = \hat{\$_\perp} \hat{A} + E^z \wedge \{\dots\}. \quad (187)$$

A.2 Proof of equation (56)

By definition,

$$\begin{aligned} \frac{1}{N} \tilde{\nabla} (N \tilde{l}^{zi}) &= \frac{1}{N} \tilde{d} (N \tilde{l}^{zi}) + \tilde{\omega}^z{}_a \tilde{l}^{ai} + \tilde{\omega}^i{}_a \tilde{l}^{za} \\ &= \frac{1}{N} \tilde{d} (N \tilde{l}^{zi}) + \tilde{\omega}^z{}_j \tilde{l}^{ji} + \tilde{\omega}^i{}_j \tilde{l}^{zj}. \end{aligned} \quad (188)$$

In analogy with equation (25), we know that

$$\tilde{d}(N\tilde{l}^{zi}) = \widehat{d}(N\tilde{l}^{zi}) + E^z\{\dots\}. \quad (189)$$

From equations (22), (23), (44) and (45), the connection pieces can be written as

$$\tilde{\omega}^z_j = \widehat{K}_j + E^z\{\dots\}, \quad (190)$$

$$\tilde{\omega}^i_j = \widehat{\omega}^i_j + E^z\{\dots\}. \quad (191)$$

From equation (47), we know that $b^i = l^{zi}$. Therefore, combining this with equations (188) to (191) yields the following,

$$\begin{aligned} \frac{1}{N}\tilde{\nabla}(N\tilde{l}^{zi}) &= \frac{1}{N}\widehat{d}(Nb^i) + \widehat{\omega}^i_j b^j + K_j \tilde{l}^{ji} + E^z\{\dots\} \\ &= \frac{1}{N}\widehat{\nabla}(Nb^i) - K_j \tilde{l}^{ij} + E^z\{\dots\} \end{aligned} \quad (192)$$

where we have used the antisymmetry of \tilde{l}^{ij} in the last line.

A.3 Proof of equation (78)

We start by showing that

$$E^\perp \wedge \$_\perp(\tilde{d}\tilde{\omega}^{cd}) = E^\perp \wedge \tilde{d}(\$_\perp\tilde{\omega}^{cd}) + dt \wedge \tilde{d}N \wedge \$_\perp\tilde{\omega}^{cd}. \quad (193)$$

From equation (25), we know that

$$d\tilde{\omega}^{cd} = \tilde{d}\tilde{\omega}^{cd} + E^\perp \wedge \$_\perp\tilde{\omega}^{cd}. \quad (194)$$

This means that

$$\begin{aligned} E^\perp \wedge \$_\perp(\tilde{d}\tilde{\omega}^{cd}) &= E^\perp \wedge \$_\perp [d\tilde{\omega}^{cd} - E^\perp \wedge \$_\perp\tilde{\omega}^{cd}] \\ &= E^\perp \wedge [\$_\perp(d\tilde{\omega}^{cd}) - \$_\perp E^\perp \wedge \$_\perp\tilde{\omega}^{cd}]. \end{aligned} \quad (195)$$

Now Lie derivatives commute with exterior derivatives, which implies that

$$E^\perp \wedge \$_\perp(d\tilde{\omega}^{cd}) = E^\perp \wedge d(\$_\perp\tilde{\omega}^{cd}). \quad (196)$$

Since $\$_\perp$ is actually the *projection* of the Lie derivative (see section 2.2.1), the wedging with E^\perp is crucial in the above equation. We also know that $\$_\perp E^\perp = -\tilde{d}N/N$. Equation (195) now gives

$$\begin{aligned} E^\perp \wedge \$_\perp(\tilde{d}\tilde{\omega}^{cd}) &= E^\perp \wedge d(\$_\perp\tilde{\omega}^{cd}) + dt \wedge \tilde{d}N \wedge \$_\perp\tilde{\omega}^{cd} \\ &= E^\perp \wedge \tilde{d}(\$_\perp\tilde{\omega}^{cd}) + dt \wedge \tilde{d}N \wedge \$_\perp\tilde{\omega}^{cd} \end{aligned} \quad (197)$$

where we have once again used equation (25). Now consider,

$$\begin{aligned} dt \wedge \tilde{\nabla}(N\$_\perp\tilde{\omega}^{cd}) &= dr \wedge \left[\tilde{d}N \wedge \$_\perp\tilde{\omega}^{cd} + N\tilde{\nabla}(\$_\perp\tilde{\omega}^{cd}) \right] \\ &= dt \wedge \tilde{d}N \wedge \$_\perp\tilde{\omega}^{cd} + E^\perp \wedge \left[\tilde{d}(\$_\perp\tilde{\omega}^{cd}) + \tilde{\omega}^c_a \wedge \$_\perp\tilde{\omega}^{ad} + \tilde{\omega}^d_a \wedge \$_\perp\tilde{\omega}^{ca} \right] \\ &= E^\perp \wedge \left[\$_\perp(\tilde{d}\tilde{\omega}^{cd}) + \tilde{\omega}^c_a \wedge \$_\perp\tilde{\omega}^{ad} + \tilde{\omega}^d_a \wedge \$_\perp\tilde{\omega}^{ca} \right] \\ &= E^\perp \wedge \$_\perp \left(\tilde{d}\tilde{\omega}^{cd} + \omega^c_a \wedge \omega^{ad} \right) \\ &= E^\perp \wedge \$_\perp \tilde{\Omega}^{cd}. \end{aligned} \quad (198)$$

A.4 Proof of equation (95)

Since $H_a \wedge H^a = 0$, we can write

$$H_a \wedge \left[\tilde{l}^{ba} F^{cd} + \tilde{l}^{cd} \tilde{\Omega}^{ba} \right] \wedge \zeta_{bcd} = H_a \wedge \left[\tilde{l}^{ba} F^{cd} + \tilde{l}^{cd} \tilde{F}^{ba} \right] \wedge \zeta_{bcd}. \quad (199)$$

Rearranging some indices on the right hand side,

$$\begin{aligned} H_a \wedge \left[\tilde{l}^{ba} F^{cd} + \tilde{l}^{cd} \tilde{\Omega}^{ba} \right] \wedge \zeta_{bcd} &= \tilde{l}^{ab} F^{cd} [H_a \wedge \zeta_{bcd} + H_c \wedge \zeta_{dab}] \\ &= \tilde{l}^{ab} F^{cd} \wedge H_e \wedge E^e \wedge \zeta_{abcd}. \end{aligned} \quad (200)$$

Here we have used the formula given in equation (8). Since H_{ef} is symmetric, we know that $H_e \wedge E^e = 0$. Equation (95) now follows automatically.

B Summary of notation

In this section, we give a summary of some of the notation used throughout this paper, to provide a quick and easily accessible reference.

B.1 Overview of manifolds

Manifold	Description	Dimension	Other features
\mathcal{M}	Test spacetime	D	Indices labelled with A, B, C .
Σ_t	Spacelike surfaces labelled by t	$D - 1$	Indices labelled with a, b, c . Objects labelled with a “tilde”.
Σ_∞	Spacelike part of $\partial\mathcal{M}$	$D - 1$	Limit as $t \rightarrow \pm\infty$ of Σ_t .
B	Timelike part of $\partial\mathcal{M}$	$D - 1$	Indices labelled with μ, ν, σ . Objects labelled with suffix “0”
S_t	Intersection of Σ_t and B	$D - 2$	Indices labelled with i, j, k . Objects labelled with a “hat”
$\bar{\mathcal{M}}$	Background spacetime	D	Objects labelled with a “bar”.

B.2 Objects on \mathcal{M}

Object	Description	Notes
g_{AB}	The metric	$g = g_{AB}dx^A dx^B$ $= -N^2 dt^2 + \gamma_{ab}(dx^a + N^a dt)(dx^b + N^b dt)$
$R^A{}_{BCD}$	The Riemann tensor	
$\{E^A\}$	The orthonormal basis of 1-forms	E^A can be E^\perp , E^z or E^i .
$\{X_A\}$	The dual basis of vectors	X_A can be X_\perp , X_z or X_i .
ω^{AB}	The connection 1-form	See equations (22) and (23) for its decomposition.
Ω^{AB}	The curvature 2-form	See equations (30) and (31) for its decomposition.
$e_{A_1 \dots A_m}$	The “measure” forms	See equation (4) for their formal definition.
d	The exterior derivative	
∇	The covariant exterior derivative	See equation (32) for the analogue on Σ_t .
$i_Y A$	The interior product of Y and A	Y is a vector, A is a p -form. See [44]

B.3 Objects on Σ_t

Object	Description	Notes
X_\perp or n^A	The unit normal to Σ_t	$X_\perp = \frac{1}{N} \left(\frac{\partial}{\partial t} - N^a \frac{\partial}{\partial x^a} \right)$
γ_{ab}	The induced metric	$\gamma = \gamma_{ab} dx^a dx^b$
$\tilde{R}^a{}_{bcd}$	The Riemann tensor	
$\{E^a\}$	The orthonormal basis of 1-forms	E^a can be E^z or E^i .
$\tilde{\omega}^{ab}$	The connection 1-form	See equations (44) and (45) for its decomposition.
$\tilde{\Omega}^{ab}$	The curvature 2-form	See equations (59) and (60) for its decomposition.
$\zeta_{a_1 \dots a_m}$	The “measure” forms	See equation (67) for their formal definition.
\tilde{d}	The exterior derivative	
$\tilde{\nabla}$	The covariant exterior derivative	See equation (32) for its definition.
$\$_\perp$ and $\$_{\frac{\partial}{\partial t}}$	The “time” derivatives	See equation (27).

B.4 Objects on B

Object	Description	Notes
X_z or z^A	The unit normal to B	$X_z = \frac{\partial}{\partial z}$ near B
$\{E^\mu\}$	The orthonormal basis of 1-forms	E^μ can be E^\perp or E^i .
$\omega_0^{\mu\nu}$	The connection 1-form	See equations (43) and (45) for its decomposition.
$\Omega_0^{\mu\nu}$	The curvature 2-form	See equations (58) and (60) for its decomposition.
d_0	The exterior derivative	
∇_0	The covariant exterior derivative	See equation (32) for the analogue on Σ_t .

B.5 Objects on S_t

Object	Description	Notes
λ_{ij}	The induced metric	
$\widehat{R}^i{}_{jkl}$	The Riemann tensor	
$\{E^i\}$	The orthonormal basis of 1-forms	
$\widehat{\omega}^{ij}$	The connection 1-form	
$\widehat{\Omega}^{ij}$	The curvature 2-form	
$\phi_{i_1\dots i_m}$	The “measure” forms	See equation (102) for their formal definition.
\widehat{d}	The exterior derivative	
$\widehat{\nabla}$	The covariant exterior derivative	See equation (32) for the analogue on Σ_t .
$\widehat{\$}_\perp$ and $\widehat{\$}_{\frac{\partial}{\partial t}}$	The “time” derivatives	The analogue of $\$_\perp$ on S_t . See equation (50).

B.6 Extrinsic Curvatures

Symbols	Description
$\widehat{K}^i = K^i{}_j E^j$	$K^i{}_j$ is the extrinsic curvature of S_t in Σ_t
$\widehat{H}^i = H^i{}_j E^j$	$H^i{}_j$ is the extrinsic curvature of S_t in B
$K^\mu = K^\mu{}_\nu E^\nu$	$K^\mu{}_\nu$ is the extrinsic curvature of B in \mathcal{M}
$H^a = H^a{}_b E^b$	$H^a{}_b$ is the extrinsic curvature of Σ_t in \mathcal{M}

References

- [1] R. Arnowitt, S. Deser, and C. Misner in *Gravitation: An Introduction to Current Research*, L. Witten, ed. New York, Wiley, 1962.

- [2] L. F. Abbott and S. Deser, “Stability of gravity with a cosmological constant,” Nucl. Phys. **B195** (1982) 76.
- [3] S. W. Hawking and G. T. Horowitz, “The Gravitational Hamiltonian, action, entropy and surface terms,” Class. Quant. Grav. **13** (1996) 1487–1498, [gr-qc/9501014](#).
- [4] S. Deser and B. Tekin, “Gravitational energy in quadratic curvature gravities,” Phys. Rev. Lett. **89** (2002) 101101, [hep-th/0205318](#).
- [5] S. Deser and B. Tekin, “Energy in generic higher curvature gravity theories,” [hep-th/0212292](#).
- [6] D. Lovelock, “The Einstein tensor and its generalizations,” J. Math. Phys. **12** (1971) 498–501.
- [7] C. Lanczos Ann. Math. **39** (1938) 842.
- [8] P. Candelas, G. T. Horowitz, A. Strominger, and E. Witten, “Vacuum configurations for superstrings,” Nucl. Phys. **B258** (1985) 46–74.
- [9] M. B. Green and P. Vanhove, “D-instantons, strings and M-theory,” Phys. Lett. **B408** (1997) 122–134, [hep-th/9704145](#).
- [10] B. Zwiebach, “Curvature squared terms and string theories,” Phys. Lett. **B156** (1985) 315.
- [11] B. Zumino, “Gravity theories in more than four-dimensions,” Phys. Rept. **137** (1986) 109.
- [12] C. Charmousis and J.-F. Dufaux, “General Gauss-Bonnet brane cosmology,” Class. Quant. Grav. **19** (2002) 4671–4682, [hep-th/0202107](#).
- [13] P. Binetruy, C. Charmousis, S. C. Davis, and J.-F. Dufaux, “Avoidance of naked singularities in dilatonic brane world scenarios with a Gauss-Bonnet term,” Phys. Lett. **B544** (2002) 183–191, [hep-th/0206089](#).
- [14] S. C. Davis, “Generalised Israel junction conditions for a Gauss-Bonnet brane world,” [hep-th/0208205](#).
- [15] E. Gravanis and S. Willison, “Israel conditions for the Gauss-Bonnet theory and the Friedmann equation on the brane universe,” [hep-th/0209076](#).
- [16] C. Germani and C. F. Sopuerta, “String inspired braneworld cosmology,” Phys. Rev. Lett. **88** (2002) 231101, [hep-th/0202060](#).
- [17] C. Germani and C. F. Sopuerta, “Varying fundamental constants from a string-inspired brane world model,” [hep-th/0210086](#).

- [18] S. Nojiri, S. D. Odintsov, and S. Ogushi, “Friedmann-Robertson-Walker brane cosmological equations from the five-dimensional bulk (A)dS black hole,” *Int. J. Mod. Phys.* **A17** (2002) 4809–4870, [hep-th/0205187](#).
- [19] J. E. Lidsey, S. Nojiri, and S. D. Odintsov, “Braneworld cosmology in (anti)-de Sitter Einstein-Gauss-Bonnet-Maxwell gravity,” *JHEP* **06** (2002) 026, [hep-th/0202198](#).
- [20] S. Nojiri, S. D. Odintsov, and S. Ogushi, “Cosmological and black hole brane world universes in higher derivative gravity,” *Phys. Rev.* **D65** (2002) 023521, [hep-th/0108172](#).
- [21] S. Nojiri and S. D. Odintsov, “Brane-world cosmology in higher derivative gravity or warped compactification in the next-to-leading order of AdS/CFT correspondence,” *JHEP* **07** (2000) 049, [hep-th/0006232](#).
- [22] K. A. Meissner and M. Olechowski, “Brane localization of gravity in higher derivative theory,” *Phys. Rev.* **D65** (2002) 064017, [hep-th/0106203](#).
- [23] I. P. Neupane, “Consistency of higher derivative gravity in the brane background,” *JHEP* **09** (2000) 040, [hep-th/0008190](#).
- [24] I. P. Neupane, “Gravitational potential correction with Gauss-Bonnet interaction,” *Phys. Lett.* **B512** (2001) 137–145, [hep-th/0104226](#).
- [25] I. P. Neupane, “Localized gravity with higher curvature terms,” *Class. Quant. Grav.* **19** (2002) 5507–5523, [hep-th/0106100](#).
- [26] Y. M. Cho and I. P. Neupane, “Warped brane-world compactification with Gauss-Bonnet term,” [hep-th/0112227](#).
- [27] Y. M. Cho and I. P. Neupane, “Anti-de sitter black holes, thermal phase transition and holography in higher curvature gravity,” *Phys. Rev.* **D66** (2002) 024044, [hep-th/0202140](#).
- [28] Y. M. Cho, I. P. Neupane, and P. S. Wesson, “No ghost state of Gauss-Bonnet interaction in warped background,” *Nucl. Phys.* **B621** (2002) 388–412, [hep-th/0104227](#).
- [29] P. Horava and E. Witten, “Heterotic and type I string dynamics from eleven dimensions,” *Nucl. Phys.* **B460** (1996) 506–524, [hep-th/9510209](#).
- [30] L. Randall and R. Sundrum, “A large mass hierarchy from a small extra dimension,” *Phys. Rev. Lett.* **83** (1999) 3370–3373, [hep-ph/9905221](#).
- [31] L. Randall and R. Sundrum, “An alternative to compactification,” *Phys. Rev. Lett.* **83** (1999) 4690–4693, [hep-th/9906064](#).
- [32] J. P. Gregory and A. Padilla, “Exact braneworld cosmology induced from bulk black holes,” *Class. Quant. Grav.* **19** (2002) 4071–4083, [hep-th/0204218](#).

- [33] A. Padilla, “Braneworld cosmology and holography,” [hep-th/0210217](#).
- [34] J. P. Gregory and A. Padilla, “Gauss-Bonnet braneworld holography.” To appear.
- [35] J. Crisostomo, R. Troncoso, and J. Zanelli, “Black hole scan,” *Phys. Rev.* **D62** (2000) 084013, [hep-th/0003271](#).
- [36] R. C. Myers, “Higher derivative gravity, surface terms and string theory,” *Phys. Rev.* **D36** (1987) 392.
- [37] G. W. Gibbons and S. W. Hawking, “Action integrals and partition functions in quantum gravity,” *Phys. Rev.* **D15** (1977) 2752–2756.
- [38] T. Eguchi, P. B. Gilkey, and A. J. Hanson, “Gravitation, gauge theories and differential geometry,” *Phys. Rept.* **66** (1980) 213.
- [39] Y. Choquet-Bruhat, C. DeWitt-Morette, and M. Dillard-Bleick, *Analysis, Manifolds and Physics*. North-Holland, New York, 1982.
- [40] S. R. Lau, “Differential forms and wave equations for general relativity,” *Int. J. Mod. Phys.* **D7** (1998) 857, [gr-qc/9612020](#).
- [41] R. M. Wald, *General Relativity*, ch. 10. The University of Chicago Press, 1984.
- [42] C. Teitelboim and J. Zanelli, “Dimensionally continued topological gravitation theory in Hamiltonian form,” *Class. Quant. Grav.* **4** (1987) L125–L129.
- [43] J. Isenberg and J. Nester, “Canonical gravity,” in *General Relativity and Gravitation*, A. Held, ed., vol. 1, pp. 23–97. 1981.
- [44] M. Nakahara, *Geometry, Topology and Physics*, ch. 5. Institute of Physics Publishing, 1990.
- [45] L. P. Eisenhart, *Riemannian Geometry*, ch. IV. Princeton University Press, 1964.
- [46] J. Louko, J. Z. Simon, and S. N. Winters-Hilt, “Hamiltonian thermodynamics of a Lovelock black hole,” *Phys. Rev.* **D55** (1997) 3525–3535, [gr-qc/9610071](#).
- [47] D. G. Boulware and S. Deser, “String generated gravity models,” *Phys. Rev. Lett.* **55** (1985) 2656.
- [48] M. Banados, C. Teitelboim, and J. Zanelli, “Dimensionally continued black holes,” *Phys. Rev.* **D49** (1994) 975–986, [gr-qc/9307033](#).
- [49] R. C. Myers, “Black holes in higher curvature gravity,” [gr-qc/9811042](#).
- [50] R.-G. Cai, “Gauss-Bonnet black holes in AdS spaces,” *Phys. Rev.* **D65** (2002) 084014, [hep-th/0109133](#).

- [51] R.-G. Cai and K.-S. Soh, “Topological black holes in the dimensionally continued gravity,” *Phys. Rev.* **D59** (1999) 044013, [gr-qc/9808067](#).