

Anisotropy and inflation in Bianchi I brane worlds

Juan M. Aguirregabiria,¹ Luis P. Chimento,² and Ruth Lazkoz¹

*¹Fisika Teorikoa eta Zientziaren Historia Saila,
Zientzi Fakultatea, Euskal Herriko Unibertsitatea,
644 Posta Kutxatila, 48080 Bilbao, Spain*

*²Departamento de Física, Facultad de Ciencias Exactas y Naturales,
Universidad de Buenos Aires, Ciudad Universitaria,
Pabellón I, 1428 Buenos Aires, Argentina.*

(Dated: May 22, 2019)

Abstract

After a more general assumption on the influence of the bulk on the brane is done, we extend some conclusions by Maartens *et al.* [1] and Santos *et al.* [2] on the asymptotic behavior of Bianchi I brane worlds. As a consequence of the nonlocal anisotropic stresses induced by the bulk, in our models, the brane does not generically isotropize and the nonlocal energy does not vanish in the limit in which the mean radius goes to infinity. We have also found the intriguing possibility that the inflation due to the cosmological constant might be prevented by the interaction with the bulk. We show that the problem for the mean radius can be completely solved in our models, which include as particular cases those in the references above.

The main stream approach to try and solve the problems arising from the break down of Einstein's theory at high energies is that of considering it a particular limit of a more general theory, exactly as happened with Newton's theory in relation to that of Einstein's. One of this schemes is the brane gravity picture [3, 4], according to which matter fields are confined to a hypersurface (the brane) with three spatial dimensions embedded in a higher dimensional space (the bulk) on which gravity can act also.

Lots of efforts in this area of research have been directed to testing that scheme by deducing cosmological implications from it. The usual framework for these studies is the geometric approach of [5], which provides the gravitational field equations induced on the brane along with conservation equation for bulk degrees of freedom.

Studies concerning FRW universes [6] hinted that the dynamics of the early Universe in this new scenario is rather peculiar because of the modifications in the Hubble equation. Then, when anisotropic models were considered [7], it was shown that the discrepancies with respect to the standard scenario extended to the shear dynamics as well.

In the setup of general relativity, Wald showed that a positive cosmological constant leads to the isotropization of expanding homogeneous cosmological models [8]. The anisotropy dissipation in brane world inflation in the absence of effective cosmological constant was analyzed by Maartens, Sahni and Saini [1]. Taking into account the cosmological constant, Santos, Vernizzi and Ferreira derived a set of sufficient conditions that allows extending Wald's result to the brane world scenario [2] (see also [9] for an alternative view on the same problem). The last three works were done directly on the brane, where an additional hypothesis has to be done for the nonlocal anisotropic stresses induced by the bulk, for they are not given by any evolution equation on the brane. This naturally raises the question to what extent the results depend on the aforementioned hypothesis.

The goal of this work is to extend Refs. [1, 2] by considering a more general additional hypothesis. We will analyze, as done in those works, a Bianchi I brane embedded in a 5-dimensional bulk. We assume that the nonlocal stresses satisfy a condition which reduces to the one considered in Refs. [1, 2] in particular cases. We will show that the evolution equations on the brane can be integrated and conclude that generically we have exponential inflation or an asymptotic power law for the mean radius, but the models do not isotropize unless they belong to the class considered in the aforementioned works.

Let us consider a Bianchi I brane with the induced metric

$$ds^2 = -dt^2 + a_1^2(t) dx^2 + a_2^2(t) dy^2 + a_3^2(t) dz^2 \quad (1)$$

when the matter on the brane is a perfect fluid of density ρ and pressure p . We define as usual the mean radius $a \equiv (a_1 a_2 a_3)^{1/3}$ and the mean Hubble parameter $H \equiv \dot{a}/a = \frac{1}{3} \sum_i H_i$, with $H_i = \dot{a}_i/a_i$.

We will be using the equations induced on the brane derived by Shiromizu, Maeda and Sasaki [5] but follow the notation of Refs. [1, 2]. For the metric (1) the dynamics of the mean radius a in this model can be described by the following equations on the brane:

$$3H^2 = \frac{6}{\kappa^2 \lambda} \mathcal{U} + \frac{1}{2} \sigma^2 + \Lambda + \kappa^2 \rho \left(1 + \frac{\rho}{2\lambda}\right), \quad (2)$$

$$3\dot{H} + 3H^2 + \sigma^2 + \frac{6}{\kappa^2 \lambda} \mathcal{U} = \Lambda - \frac{\kappa^2}{2} (\rho + 3p) - \frac{\kappa^2}{2} (2\rho + 3p) \frac{\rho}{\lambda}, \quad (3)$$

$$\dot{\sigma}_{\mu\nu} + 3H\sigma_{\mu\nu} = \frac{6}{\kappa^2 \lambda} \mathcal{P}_{\mu\nu}, \quad (4)$$

$$\dot{\mathcal{U}} + 4H\mathcal{U} + \sigma^{\mu\nu} \mathcal{P}_{\mu\nu} = 0, \quad (5)$$

where $\kappa^2/8\pi$ is the effective Newton constant in the brane, \mathcal{U} the effective nonlocal energy density, $\sigma_{\mu\nu}$ the shear, and

$$\sigma^2 \equiv \sigma^{\mu\nu} \sigma_{\mu\nu} = \sum_{i=1}^3 (H_i - H)^2. \quad (6)$$

Λ is the effective cosmological constant and $\lambda \equiv 6\kappa^2/\tilde{\kappa}^4$, where $\tilde{\kappa}^2/8\pi$ is the fundamental Newton constant in the bulk. A dot denotes $u^\mu \nabla_\mu$, with $u^\mu = \partial/\partial t$. There is also the constraint that the projected spatial covariant derivative of $\mathcal{P}_{\mu\nu}$ vanishes:

$$D^\nu \mathcal{P}_{\mu\nu} = 0. \quad (7)$$

On the other hand, as a consequence of system (2)–(5), we get the conservation law

$$\dot{\rho} + 3H(\rho + p) = 0. \quad (8)$$

Since there is no evolution equation on the brane for the nonlocal anisotropic stress $\mathcal{P}_{\mu\nu}$, but only the constraint (7), some additional hypothesis is necessary in order to integrate the system (2)–(5).

In order to get some insight on the problem, we will restrict ourselves for a moment to the vacuum case $\rho = p = 0$, where we get from (2)–(3)

$$\mathcal{U} = \frac{\kappa^2 \lambda}{2} (\dot{H} + 3H^2 - \Lambda). \quad (9)$$

Let us consider under which conditions the stable asymptotic behavior may approach a power-law $a \sim t^k$, ($k > 0$), so $\dot{H}, H \rightarrow 0$, $\mathcal{U} \rightarrow -\kappa^2\lambda\Lambda/2$ and, because of (5),

$$\sigma^{\mu\nu}\mathcal{P}_{\mu\nu} \rightarrow 2\kappa^2\lambda\Lambda H. \quad (10)$$

We will show in the following that if one chooses the unknown $\mathcal{P}_{\mu\nu}$ so that this condition is satisfied, the asymptotic behavior described by a power-law is stable. Furthermore, this will happen even in the presence of a fluid.

So as to integrate the system (2)–(5) one often [1] assumes $\mathcal{U} = 0$ or the more general $\sigma^{\mu\nu}\mathcal{P}_{\mu\nu} = 0$, which has been used in Ref. [2] to discuss the stability of the de Sitter spacetime. Although the asymptotic behavior (10) may happen with many choices of $\sigma^{\mu\nu}\mathcal{P}_{\mu\nu}$, we will consider only the simplest one. In the following, we will assume that $\sigma^{\mu\nu}\mathcal{P}_{\mu\nu}$ is proportional to the Hubble parameter, so that it can be written as

$$\sigma^{\mu\nu}\mathcal{P}_{\mu\nu} = 2\alpha\kappa^2\lambda H, \quad (11)$$

which reduces to the cases discussed in [1, 2] when the constant α vanishes. (10) is recovered, even before the asymptotic behavior is reached, for $\alpha = \Lambda$. By making this hypothesis we use a less restrictive assumption to check the generality of the conclusions reached in [1, 2], while still being able to integrate the evolution equations.

Whether this assumption is compatible with a full 5-dimensional physical solution remains an open problem, exactly as happens with the $\alpha = 0$ case often used in the bibliography.

The nonlocal energy density \mathcal{U} is given by the conservation equation (5), which is easily solved under the assumption (11):

$$\mathcal{U} = \frac{\kappa^2\lambda u_0}{a^4} - \frac{1}{2}\kappa^2\lambda\alpha, \quad (12)$$

where u_0 is an arbitrary integration constant. We see here that for expanding universes \mathcal{U} does not asymptotically vanish as $a \rightarrow \infty$, except in the special case $\alpha = 0$, where it reduces to the result by Toporensky [10]. This suggests that the asymptotic vanishing of \mathcal{U} arising in the cases studied in [2] might be non generic, as a consequence of the interaction of the brane with the bulk through $\mathcal{P}_{\mu\nu}$.

By contracting (4) with $\sigma^{\mu\nu}$ and using (11), one readily gets

$$\sigma^2 = \frac{\sigma_0^2}{a^6} + 4\alpha, \quad (13)$$

for any constant σ_0 . If $\alpha = 0$ the brane world isotropizes as $a \rightarrow \infty$, as described in [1, 2]; but generically there will be a remaining anisotropy induced by the nonlocal anisotropic stresses.

If we assume the equation of state $p = (\gamma - 1)\rho$, the conservation law (8) is equivalent to

$$\rho = \frac{\rho_0}{\kappa^2 a^{3\gamma}}, \quad (14)$$

with an arbitrary constant ρ_0 .

If we insert (12), (13) and (14) in (2), we get the generalized Friedmann equation

$$3H^2 = \frac{6u_0}{a^4} + \frac{\sigma_0^2}{2a^6} + \frac{\rho_0}{a^{3\gamma}} + \frac{\rho_0^2}{2\kappa^2 \lambda a^{6\gamma}} + \Lambda - \alpha. \quad (15)$$

The orbits $(H(\mathcal{U}), \sigma(\mathcal{U}))$ were found in [2] for the special case $\alpha = 0$, but we can see that, for any α , the problem for $a(t)$, $\sigma(t)$ and $\mathcal{U}(t)$ may be completely solved from equation (15) by means of a quadrature and a function inversion. The integral may be explicitly computed (in terms of elementary or elliptic functions) for different sets of values for the constants u_0 , σ_0 , ρ_0 , γ and $\Lambda - \alpha$. For instance, in vacuum, $\rho_0 = 0$, if $\Lambda = \alpha$, the solution may be written by using the parameter $0 < w < \infty$ as

$$t = \frac{\sigma_0^2}{48\sqrt{2u_0^3}} (\sinh w - w), \quad (16)$$

$$a = \frac{\sigma_0}{2\sqrt{3u_0}} \sinh \frac{w}{2}, \quad (17)$$

while for $u_0 = 0$ and $\Lambda > \alpha$ we get

$$a = \left[\frac{\sigma_0^2}{2(\Lambda - \alpha)} \sinh^2 \left(\sqrt{3(\Lambda - \alpha)} t \right) \right]^{1/6}. \quad (18)$$

We can also obtain directly from (15) some interesting asymptotic results. Generically, for $\Lambda - \alpha > 0$, the brane inflates exponentially when $a \rightarrow \infty$, as happened for $\alpha = 0$, $\Lambda > 0$ [2]. But now there is another interesting possibility: the repulsion due to the cosmological constant may be neutralized by the bulk influence if $\alpha = \Lambda$, in which case the asymptotic behavior of a is a power law, and the same would happen for any selection of $\sigma^{\mu\nu} \mathcal{P}_{\mu\nu}$ that, like our (11), would asymptotically approach $2\kappa^2 \lambda \Lambda H$. For instance, if we additionally assume that $\gamma > 4/3$ (or that there is no fluid) and that $u_0 \neq 0$, asymptotically the brane will expand as $a \sim t^{1/2}$ while remaining anisotropic ($\sigma^2 \sim 4\Lambda$). For $\gamma < 4/3$, or $u_0 = 0$, one would have $a \sim t^{2/3\gamma}$.

By making the more general assumption (11) on the (unknown) influence of the bulk on the brane, we have shown that some conclusions on the asymptotic behavior of Bianchi I brane worlds in Refs. [1, 2] can be generalized. Due to the nonlocal stresses, in our models, the brane does not generically isotropize and the nonlocal energy does not vanish in the limit $a \rightarrow \infty$. We have also found that, although our models generically inflate, there also exist the possibility that the inflation due to the cosmological constant may be prevented by the interaction with the bulk. Finally, we have shown that the problem for the mean radius a (as well as for σ^2 and \mathcal{U}) can be completely solved in our models, which include as particular cases the Bianchi I branes for which the orbits and stability were analyzed in [2].

Acknowledgments

This work was supported by the University of Buenos Aires under Project X223, the Spanish Ministry of Science and Technology jointly with FEDER funds through research grant BFM2001-0988, and the University of the Basque Country through research grant UPV00172.310-14456/2002. Ruth Lazkoz's work is also supported by the Basque Government through fellowship BFI01.412.

-
- [1] R. Maartens, V. Sahni, and T. D. Saini, *Phys. Rev. D* **63**, 063509 (2001).
 - [2] M. G. Santos, F. Vernizzi, and P. G. Ferreira, *Phys. Rev. D* **64**, 063506 (2001).
 - [3] P. Horava and E. Witten, *Nucl. Phys. B* **460**, 506 (1996).
 - [4] L. Randall, R. Sundrum, *Phys. Rev. Lett.* **83**, 4690 (1999).
 - [5] T. Shiromizu, K. Maeda, and M. Sasaki, *Phys. Rev. D* **62**, 024012 (2000).
 - [6] P. Binétruy, C. Deffayet, and D. Langlois, *Nucl. Phys. B* **565**, 269 (2000).
 - [7] A. Campos and C. F. Sopuerta, *Phys. Rev. D* **63**, 104012 (2001).
 - [8] R. Wald, *Phys. Rev. D* **28**, 2118 (1982).
 - [9] B. C. Paul, *Phys. Rev. D* **66**, 124019 (2002).
 - [10] A. V. Toporensky, *Class. Quan. Grav.* **18**, 2311 (2001).