## Hubble Red Shift and the Anomalous Acceleration of Pioneer 10 and 11

Kostadin Trenčevski<sup>\*</sup> Faculty of Natural Sciences and Mathematics, P.O.Box 162, 1000 Skopje, Macedonia

PACS number: 98.80.Jk

## Abstract

The basic relationship between the Hubble constant H and the apparent anomalous acceleration  $a_P$ , which appears in the motion of the spacecraft Pioneer 10 and 11, is given by  $a_P = cH$  [1]. Using this equality, both Hubble red shift and the anomalous acceleration  $a_P$  are explained, assuming that the gravitational potential in the universe changes linearly:  $\Delta V = -c^2 H \Delta t$ . As a consequence after each second the time is faster  $1 + \frac{1}{3.76 \times 10^{17}}$  times.

Hubble in 1928 has discovered that the velocities of the galaxies with respect to the Earth are proportional to the corresponding distances from the Earth. Namely, if v is the velocity of a chosen star, and R is its distance from the Earth, then v = RH, where H is an universal constant called Hubble constant. Some recent measurements show that H is about 72 km/s/Mpc [4]. We shall assume that it is about 82 km/s/Mpc  $\approx (11, 9 \times 10^9 \text{ years})^{-1} \approx (3, 76 \times 10^{17} \text{ s})^{-1} \approx$  $2, 66 \times 10^{-18} \text{ s}^{-1}$ . It is accepted that the age of the universe is about  $\frac{1}{H}$ , and the universe is expanding. The velocities among the galaxies are sometimes up to c/5. These velocities are determined using the Doppler effect.

In this paper it is shown that the previous Hubble law can be explained in another way, as a consequence of the change of the gravitational potential in the universe. Hence it will follow that such large velocities among the galaxies are only apparent, because the main effect is the red shift, and the red shift appears according to additional gravitational potential V in the universe, which changes linearly (or almost linearly) with the time, i.e.  $\frac{\partial V}{\partial t} \approx const$ . Indeed, it

This paper was done under the auspices of a CNCSIS Grant.

 $<sup>^*</sup>$ Electronic address: kostatre@iunona.pmf.ukim.edu.mk

is detected red shift such that  $\nu = \nu_0 \left(1 - \frac{v}{c}\right)$ , and replacing v = HR, indeed it is observed that

$$\nu = \nu_0 \left( 1 - \frac{RH}{c} \right). \tag{1}$$

On the other hand, if the potential changes linearly with the time, after time t = R/c we have additional potential  $\frac{R}{c} \frac{\partial V}{\partial t}$ , and hence we have a shift such that

$$\nu = \nu_0 \left( 1 + \frac{1}{c^2} \frac{R}{c} \frac{\partial V}{\partial t} \right). \tag{2}$$

*Remark.* Note that the gravitational potential V is considered to be larger near gravitational bodies compared with the potential where there is no gravitation. For example, near the spherical body we accept that  $V = \frac{GM}{r}$ . If we accept that  $V = -\frac{GM}{r}$ , then the sign "+" in the formula (2) should be replaced by "-".

Comparing the formulas (1) and (2) we obtain

$$H = -\frac{1}{c^2} \frac{\partial V}{\partial t}.$$
 (3)

Since  $\frac{\partial V}{\partial t} < 0$ , in the past the time in the universe was slower than now, and in future it will be faster than now. Indeed, each second the time is faster approximately  $\lambda = (1 + H \times 1s) = 1 + \frac{1}{3.76 \times 10^{17}}$  times.

According to (3) we obtain that

$$\frac{\partial V}{\partial t} \approx -2, 4 \times 10^3 \frac{\mathrm{cm}^2}{\mathrm{s}^3}.$$
(4)

This shows that the potential energy of mass of 1kg arises for  $2, 4 \times 10^3 \frac{\text{kg} \cdot \text{cm}^2}{\text{s}^2} = 0,24\text{J}$  per second. Probably this energy comes from the dark energy which is about 67% in the universe [4].

On the other hand, in the last decades are considered the motions of the spacecraft Pioneer 10 and 11, by comparing the initial frequency of a signal which is sent from the Earth to the spacecraft and the frequency of the signal which comes back. The frequency of the received signal on the Earth does not fit with the predicted frequency modeled by using the Doppler effect, the position and motion of the spacecraft using the general relativity and a lot of perturbations (see Ref. [1, 2]). In the case of the spacecraft Pioneer 10 and 11 we have a similar situation like with the red shift from the galaxies, but much more complicated. The observations show that it appears almost a constant apparent unmodeled acceleration  $a_P \approx (8,74\pm1,33) \times 10^{-8} \text{cm/s}^2$ , which seems to act to the spacecraft toward the Sun. The acceleration  $a_P$  is introduced by formula (15) in [1], i.e.

$$[\nu_{obs}(t) - \nu_{model}(t)]_{DSN} = -\nu_0 \frac{2a_P t}{c}, \qquad \nu_{model} = \nu_0 \Big[ 1 - 2\frac{v_{model}(t)}{c} \Big], \quad (5)$$

where  $\nu_0$  is the reference frequency, the factor 2 is because of two way data,  $v_{model}$  is the modeled velocity of the spacecraft due to the gravitational and other forces, and  $\nu_{obs}$  is frequency of the re-transmitted signal observed by DSN antennae. After time 2t has been detected a small blue shift on the top of the red shift caused by the motion of the spacecraft outwards the Sun. Form (5), an unexplained blue shift

$$\nu = \nu_0 \left( 1 + \frac{2a_P t}{c} \right),\tag{6}$$

is detected, where 2t is the time of the light signal in two directions. In [1] it is mentioned that without using the acceleration  $a_P$ , the anomalous frequency shift can be interpreted by "clock acceleration"  $-a_t = -2, 8 \times 10^{-18} \text{s/s}^2$ . This causes just the blue shift given by (6). This model assumes that the time is uniform, but there is only technical problem with the clocks. Although this model gives good explanations for both frequency shift and the trajectories, it is rejected [1]. Further we shall use this model for comparison with our model via the time dependent potential.

Many people have noticed that the acceleration  $a_P$  and the Hubble constant H are related by (see Ref. [1])

$$a_P = cH,\tag{7}$$

and some possible explanations about the anomalous acceleration were given (for example [3, 5, 6, 7]). Indeed, if we assume that H=82km/s/Mpc, then from (7) for  $a_P$  we obtain  $8 \times 10^{-8}$  cm/s<sup>2</sup>. For this reason we assumed that H=82km/s/Mpc.

Now we shall explain the apparent acceleration  $a_P$  connected with the frequency shift. Let us denote by X, Y, Z, T our natural coordinate system, in the deformed space-time, and let us denote by x, y, z, t the normed coordinates of an imagine coordinate system, where the space-time is "uniform", except close to the gravitational objects. Then, according to the general relativity we have the following equalities

$$dx = \left(1 + \frac{V}{c^2}\right)^{-1} dX = \left(1 + tH\right) dX, \qquad dy = \left(1 + \frac{V}{c^2}\right)^{-1} dY = \left(1 + tH\right) dY,$$
$$dz = \left(1 + \frac{V}{c^2}\right)^{-1} dZ = \left(1 + tH\right) dZ, \qquad dt = \left(1 - \frac{V}{c^2}\right)^{-1} dT = \left(1 - tH\right) dT.$$
(8)

*Remark.* Note that it is sufficient in this paper to assume that (8) are satisfied. Then it is not necessary to speak about the time dependent gravitational potential. The coefficients 1 + tH and 1 - tH in (8) should be exponential functions, but neglecting  $H^2$  and smaller quantities we accept these linear functions.

From (8) we obtain

$$\left(\frac{dX}{dT}, \frac{dY}{dT}, \frac{dZ}{dT}\right) = \left(\frac{dx}{dt}, \frac{dy}{dt}, \frac{dz}{dt}\right)(1 - 2tH)$$

and by differentiating this equality by T we get

$$\left(\frac{d^2 X}{dT^2}, \frac{d^2 Y}{dT^2}, \frac{d^2 Z}{dT^2}\right) = \left(\frac{d^2 x}{dt^2}, \frac{d^2 y}{dt^2}, \frac{d^2 z}{dt^2}\right) - 3tH\left(\frac{d^2 x}{dt^2}, \frac{d^2 y}{dt^2}, \frac{d^2 z}{dt^2}\right) - 2H\left(\frac{dX}{dT}, \frac{dY}{dT}, \frac{dZ}{dT}\right)$$

In normed coordinates x, y, z, t there is no acceleration caused by the time dependent gravitational potential. Thus, in real coordinates (X, Y, Z, T) appears an additional acceleration

$$-3tH\left(\frac{d^2x}{dt^2},\frac{d^2y}{dt^2},\frac{d^2z}{dt^2}\right) - 2\left(H\frac{dX}{dT},H\frac{dY}{dT},H\frac{dZ}{dT}\right).$$

The first component  $-3tH\left(\frac{d^2x}{dt^2}, \frac{d^2y}{dt^2}, \frac{d^2z}{dt^2}\right)$  is smaller than the second component

$$-2\left(H\frac{dX}{dT}, H\frac{dY}{dT}, H\frac{dZ}{dT}\right) = -2\left(\frac{a_P}{c}\frac{dX}{dT}, \frac{a_P}{c}\frac{dY}{dT}, \frac{a_P}{c}\frac{dZ}{dT}\right),\tag{9}$$

and so it is of minor role in the explanation of the frequency shift of the Pioneer spacecraft. Indeed, the first acceleration is also important, but not so much in the case of the spacecraft. Further, in our simplified model will be used only the acceleration (9).

We shall explain why it is measured the blue shift given by (6), instead of the red shift like in (1).

For the sake of simplicity we assume that the spacecraft is moving radially in the solar system outwards the Sun, and the DSN antenna has a fixed position in the solar system, collinear with the Sun and the spacecraft. Assume that far from the Sun, when the spacecraft is on distance R from the DSN antenna, its Newtonian acceleration is approximately a constant acceleration equal to -a, i.e. a towards the Sun. When it is on distance R, for t = 0, assume that its velocity is  $v_0$ . Without loss of generality we can deal according to the Newton theory, when it is possible. Let us calculate  $\frac{\Delta \nu}{\nu_0} = (\nu_{obs} - \nu_0)/\nu_0$ , neglecting the terms containing  $H^2$ .

If  $v_0 = 0$  and a = 0, then analogous to (1), after time 2R/c in two directions we have red shift

$$\frac{\Delta\nu}{\nu_0} = -2H\frac{R}{c}.$$

Since the real acceleration is equal to -a - 2Hv, it satisfies the differential equation  $\frac{dv}{dt} = -a - 2Hv$  and hence  $\frac{d^2v}{dt^2} = -2H\frac{dv}{dt} \approx 2Ha$ . Thus, using the Taylor's series for the velocity v we get

$$v = v_0 + (-a - 2Hv_0)t + aHt^2.$$

Then for the observed shift according to our model we obtain

$$\left(\frac{\Delta\nu}{\nu_0}\right)_I = -2H\frac{R+v_0t-\frac{1}{2}at^2}{c} - 2\frac{v_0+(-a-2Hv_0)t+aHt^2}{c},$$

$$\left(\frac{\Delta\nu}{\nu_{0}}\right)_{I} = 2H\frac{R + v_{0}t - \frac{1}{2}at^{2}}{c} - 2\frac{v_{0} - at}{c} - 4H\frac{R}{c}.$$

On the other side, if we neglect the acceleration (9), and consider the blue shift (6), i.e. if we use "clock acceleration" model, then

$$\left(\frac{\Delta\nu}{\nu_{0}}\right)_{II} = 2H\frac{R + v_{0}t - \frac{1}{2}at^{2}}{c} - 2\frac{v_{0} - at}{c}.$$

Comparing the frequencies in both cases  $\nu_I$  and  $\nu_{II}$ , we see that

$$\frac{d}{dt} \left(\frac{\Delta \nu}{\nu_0}\right)_I = \frac{d}{dt} \left(\frac{\Delta \nu}{\nu_0}\right)_{II}, \quad \text{i.e.} \quad \frac{d\nu_I}{dt} = \frac{d\nu_{II}}{dt}.$$

Although  $\nu_I \neq \nu_{II}$ , it is more important that their derivatives are equal, because the determination of  $a_P$  can not be done by a single measurement, but statistically followed on long time intervals. This explains the blue shift given by (6), according to our model.

Appart from the previous explanation, we shall give now an explicit formula for the anomalous acceleration  $a_P$  in general case.

Let R = R(T) be the distance from the spacecraft to the DSN antenna. Let us calculate  $\frac{\Delta \nu}{\nu_0} = (\nu_{obs} - \nu_0)/\nu_0$ . According to our model, neglecting the terms containing  $H^2$ , this expression is equal to

$$\frac{\Delta\nu}{\nu_0} = -2H\frac{R}{c} - 2\frac{\frac{dR}{dT}}{c},$$

where the expression  $-2H\frac{R}{c}$  corresponds to the Hubble red shift from (2). On the other side, according to (5), where  $v_{model}$  it determined without using the Hubble constant H, we have

$$\frac{\Delta\nu}{\nu_0} = 2a_P \frac{R}{c^2} - 2\frac{\left(\frac{dR}{dT}\right)_{H=0}}{c}$$

From the last two equalities we are able to find the expression for  $a_P$ 

$$a_P = -c \left(\frac{1}{R}\frac{dR}{dT} - \frac{1}{R}\left(\frac{dR}{dT}\right)_{H=0} + H\right).$$

Using that  $\frac{dR}{dT} = \frac{dr}{dt}(1-2HT) = \left(\frac{dR}{dT}\right)_{H=0}(1-2HT)$  according to (8), we obtain finally

$$a_P = cH \left( 2\frac{dR}{dT}\frac{T}{R} - 1 \right). \tag{10}$$

This equality gives the required expression of the acceleration  $a_P$ . Now we are not able to determine the initial value of T, i.e. when we should start to measure the time T in (10). It depends on the estimation of the initial values for the motion of the spacecraft, i.e. it depends on the departure of the real initial values. Thus, for different spacecraft are obtained close but different values of  $a_P$ . Note also that for each initial value of T, when T tends to infinity (or R tends to infinity), then  $a_P$  tends to cH. In an ideal case, if the experiment is done in an inertial system, then dR/dT = R/T is a constant velocity and then  $a_P \equiv cH$ .

The change of the gravitational potential causes slight changes in the results of the known experiments about general relativity, but they are negligible. The change of the results in the test of Shapiro time delay is negligible if we consider radio signals on short distances like in the solar system. Indeed, the change of the potential in the universe is about  $2, 4 \times 10^3 \text{ cm}^2/\text{s}^2$  each second, while the gravitational potential on 1AU from the Sun is equal to  $30^2 \text{km}^2/\text{s}^2 = 9 \times 10^{12} \text{cm}^2/\text{s}^2$ . This potential difference will be achieved from the universe after 118 years, which is too long compared with the time needed for an experiment about Shapiro time delay. The change of the deflection of the light rays near the Sun is completely negligible up to  $c^{-2}$ . The angle between two perihelions is changed additionally of order  $\frac{\Theta^2 H^2}{\epsilon}$ , where  $\Theta$  is the orbit period of the planet and  $\epsilon$  is the eccentricity. This change of the angle is negligible with respect to the measured angle  $\frac{6GM\pi}{ac^2(1-\epsilon^2)}$ .

## References

- J. D. Anderson, P. A. Laing, E. L. Lau, A. S. Liu, M. M. Nieto, and S. G. Turyshev, Study of anomalous acceleration of Pioneer 10 and 11, *Phys. Rev.* D 65, 082004.
- [2] J. D. Anderson, P. A. Laing, E. L. Lau, A. S. Liu, M. M. Nieto, and S. G. Turyshev, Indication from Pioneer 10/11, Galileo and Ulysses Data of an Apparent Anomalous, Weak, Long-Range Acceleration, *Phys. Rev. Lett.*, 81, 2858 (1998).
- [3] W. B. Belayev, Cosmological model with movement in fifth dimension, eprint gr-qc/0110099 (2001).
- [4] W. L. Freedman, M. S. Turner, Measuring and understanding of the universe, *Rev. of Modern Phys.* 75, 1433-1447 (2003).
- [5] M. A. Ivanov, Gen. Rel. and Grav. 33, 479 (2001), eprint: astro-ph/0005084.
- [6] M. Milgrom, Ann. Phys. (NY) 229, 384 (1994), eprint: astro-ph/0112069.
- [7] G. Modanese, Nucl. Phys. B 556, 397 (1999), eprint: gr-qc/9903085.